Published in final edited form as:

J Org Chem. 2008 February 15; 73(4): 1575. doi:10.1021/jo701778k.

Conjugate Addition/Ireland–Claisen Rearrangements of Allyl Fumarates: Simple Access to Terminally Differentiated Succinates

Cory C. Bausch and Jeffrey S. Johnson*

Department of Chemistry, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599-3290

Abstract

The conjugate addition of dialkylzinc reagents to allyl fumarates with subsequent Ireland–Claisen rearrangement has been accomplished yielding substituted unsymmetrical succinic acid derivatives. This one-pot reaction creates two new carbon–carbon bonds at contiguous stereogenic centers. The reaction proceeds for several alkylzinc reagents and substituted allyl fumarates. The products contain distinguishable functional handles for further manipulation.

The conjugate addition of nonstabilized organometallic nucleophiles to α,β -unsaturated carbonyls is a staple of organic synthesis, but the fumarate and maleate electrophile subclass in this reaction family has received scant attention relative to simple enones and enoates. The comparative lack of research activity is surprising since maleic and fumaric acid derivatives represent cheap progenitors to functionalized succinic acids. Notable advances in this area have been reported. Ibuka and co-workers disclosed organocopper(I)–Lewis acid additions to fumarate esters that afforded good yields of the addition products; however, olefin reduction was a competing reaction pathway.¹ Hayashi and co-workers have reported enantioselective rhodium-catalyzed 1,4-additions of arylboronic acids to fumarates and maleates.² While these examples provide useful conjugate adducts, a nonobvious feature of the reported additions is that the carbonyls in the products are still functionally equivalent and therefore challenging to distinguish. In this paper, we describe the conjugate addition of dialkylzinc reagents to diallyl fumarates with subsequent Ireland–Claisen rearrangement to give unsymmetrical succinate products. A defining characteristic of the title reaction is chemodifferentiation of the two carbonyls that can be exploited in further selective manipulations.

jsj@unc.edu.

Supporting Information Available: Spectroscopic and analytical data for all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Multicomponent couplings initiated by conjugate addition and terminated by electrophilic trapping of the resulting enolate are classic complexity-building reactions that exploit that latent nucleophilic character of the α -carbon in Michael acceptors (Scheme 1).³ Conjugate addition/[3,3]-rearrangement reactions represent an important subcategory of these tandem processes.⁴ We envisioned that by using an allyl fumarate as the conjugate acceptor with an organometallic nucleophile, the metal enolate or derived silyl ketene acetal generated in situ would undergo a subsequent ester enolate Claisen rearrangement (Scheme 2).⁵ In accord with precedent, the use of a chlorotrialkylsilane was projected to both accelerate the conjugate addition and generate the requisite silyl ketene acetal.^{6,7}

Organometallic nucleophiles for conjugate addition to diallylfumarates⁸ were evaluated. For the purpose of the initial screen, we examined only the 1,4-addition. The two catalysts utilized were the commercially available CuBr·SMe₂ and easily synthesized bis(*N*-*tert*-butylsalicylideneamino)copper(II),⁹ hereafter referred to as Cu(*N*'Bu-sal)₂.

The Cu(*N*'Bu-sal)₂-catalyzed addition of ethylmagnesium bromide led to the desired 1,4-addition product along with products derived from 1,2-addition and S_N2'displacement (Table 1, entry 1). The conjugate addition of ethylmagnesium bromide also proceeds in the absence of copper (entry 2). An alternative organometallic reagent, diethylzinc, is considerably less reactive in THF (entry 3), but simply switching solvents to diethyl ether provides clean conjugate addition in 90 min (entries 4 and 5).

The desired addition proceeds in the absence of TMSCl and copper additives (entries 6–8), although with decreased efficiency. The observation of uncatalyzed addition using Et₂Zn, even at low temperature, dramatically highlights the difference in reactivity between fumarates and simple enoates. The identity of the copper source was inconsequential to the progress of the reaction. Upon heating of the reaction mixture, the presumed silylketene acetal intermediate underwent Ireland–Claisen rearrangement. The carboxylic acid product was esterified with dimethyl sulfate and potassium carbonate in acetone for ease of characterization. With the optimized conditions identified, the scope of the reaction was explored (Table 2).

The addition of diethylzinc (1.0 M in diethyl ether) to diallyl fumarate (entry 1) or diallyl maleate (entry 2) and subsequent rearrangement provides the disubstituted unsymmetrical succinate product in good yield and diastereocontrol. Diprenyl fumarate is also an effective reaction partner (entry 3), with S_N2'displacement as a competitive process that slightly decreases the yield. The diastereoselectivity is eroded when a bulkier nucleophile, diisopropylzinc (1.0 M in toluene), is used (entry 4).

Other diorganozinc reagents were prepared *via* transmetalation. By using *n*-butyllithium and ZnCl₂ to generate Bu₂Zn·(LiCl)₂ in situ, the desired addition and subsequent rearrangement was accomplished. The addition of Bu₂Zn·(LiCl)₂ to both diallyl and diprenyl fumarates proceeded in good yield, although with decreased diastereoselectivity (entries 5 and 6). The reagent Bu₂Zn·(MgX₂)₂ gave rise to the desired 1,4-addition, but with little or no rearrangement observed. It is conceivable that the magnesium salts inhibit the enolate silylation that is apparently required for the [3,3]-rearrangement.

The addition of Et₂Zn to either a *cis*- (entry 8) or *trans*-substituted (entry 7) allyl fumarate proceeds in good yield but gives an intractable mixture of diastereomers.

The monocarboxylic acid derived from addition of diethylzinc to diallyl fumarate (**1a**) was transformed in two steps to the crystalline secondary amide **3**. Its relative stereostructure was ascertained via X-ray crystallography.¹⁰ The structure found in Figure 1 reveals the anti disposition of the ethyl and allyl moieties on the succinate backbone. A stereochemical model consistent with this observation supposes that addition to the favored *s*-*trans* conformation of

the enoate occurs to give an (*E*)-enolate intermediate.¹¹ In a relevant study, Blanco and co-workers reported that conjugate addition to the *s*-trans conformation of an α,β -unsaturated enoate led to the *E*-silylketene acetal geometry upon trapping with chlorotrimethylsilane.^{7d} A transition structure for the sigmatropic rearrangement that minimizes allylic strain and involves approach of the allyl moiety syn to the ester group has been previously invoked by Martin and co-workers in their study of Ireland–Claisen rearrangements of substituted succinates (Scheme 3).¹²

The desymmetrized succinate products are useful due to the presence of several distinguishable functional handles that allow for further synthetic manipulation. Subjecting the carboxylic acid intermediate to *N*-bromosuccinimide¹³ in the presence of NaHCO₃ gave efficient bromolactonization yielding the γ -lactone (Scheme 4) in 80% yield and 9:1 diastereoselection. The diastereomers result from the initial addition/sigmatropic rearrangement, indicating that the cyclization proceeds with complete diastereoselectivity.¹⁴

Alternatively, the carboxylic acid can be employed in a Curtius rearrangement to furnish a β -amino ester with diphenylphosphoryl azide, benzyl alcohol, and triethylamine (Scheme 5).¹⁵ No purification of the intermediate acid was performed for either the bromolactonization or Curtius rearrangement.

The amide formed with the racemic rearranged product and (*S*)-phenylethylamine using DCC¹⁶ gives a secondary amide as a 1:1 mixture of diastereomers (Scheme 6). The diastereotopic methyl groups are clearly distinguishable by ¹H NMR spectroscopy (δ 0.87 vs 0.78, CDCl₃), which allows this simple derivative to be used for assaying enantioselective variants. Efforts to this end employing methods developed by Feringa¹⁷ and others have provided no enantiomeric enrichment to date. The significant uncatalyzed background rate is almost certainly a complication in these efforts. To achieve an enantioselective variant, a substantially less reactive nucleophile will need to be incorporated into this tandem sequence to suppress the background addition of the achiral nucleophile.

In summary, we have developed a tandem conjugate addition/Ireland–Claisen rearrangement of dialkylzinc reagents and allyl fumarates in good to excellent yields giving substituted succinate products. The reaction renders the two carboxyl groups non-equivalent; consequently, the products can be selectively manipulated in subsequent synthetic operations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding for this work was provided by the National Institutes of Health (National Institute of General Medical Sciences, GM068443). Research support from Eli Lilly, Amgen, and GSK is gratefully acknowledged. J.S.J. is an Alfred P. Sloan Fellow and a Camille Dreyfus Teacher–Scholar.

References

1. Ibuka T, Aoyagi T, Kitada K, Yoneda F, Yamamoto Y. *J Organomet Chem* 1985;287:C18–C22.
2. Shintani R, Ueyama K, Yamada I, Hayashi T. *Org Lett* 2004;6:3425–3427. [PubMed: 15355068]
3. (a) Noyori, R. *Asymmetric Catalysis in Organic Synthesis*. Wiley; New York: 1994. p. 298–322. (b) Guo HC, Ma JA. *Angew Chem, Int Ed* 2006;45:354–366.
4. (a) Aoki Y, Kuwajima I. *Tetrahedron Lett* 1990;31:7457–7460. (b) Chai Y, Hong SP, Lindsay HA, McFarland C, McIntosh MC. *Tetrahedron* 2002;58:2905–2928. (c) Eriksson M, Hjelmencrantz MN,

- Nilsson M, Olsson T. *Tetrahedron* 1995;51:12631–12644. For an alternative approach, see: (d) Lambert TH, MacMillan DWC. *J Am Chem Soc* 2002;124:13646–13647. [PubMed: 12431073]
5. Ireland RE, Mueller RH, Willard AK. *J Am Chem Soc* 1976;98:2868–2877.
6. (a) Horiguchi Y, Matsuzawa S, Nakamura E, Kuwajima I. *Tetrahedron Lett* 1986;27:4025–4028. (b) Matsuzawa S, Horiguchi Y, Nakamura E, Kuwajima I. *Tetrahedron* 1989;45:349–362.
7. (a) Perlmutter, P. *Conjugate Addition Reactions in Organic Synthesis*. Pergamon Press; Oxford: 1992. (b) Sakata H, Kuwajima I. *Tetrahedron Lett* 1987;28:5719–5722. (c) Sakata H, Aoki Y, Kuwajima I. *Tetrahedron Lett* 1990;31:1161–1164. (d) Magot-Cuvru A, Blanco L, Drouin J. *Tetrahedron* 1991;47:8331–8338.
8. See the Supporting Information for the synthesis procedure.
9. Sacconi L, Ciampolini M. *J Chem Soc* 1964:276–80.
10. CCDC 635154 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via. www.ccdc.cam.ac.uk/data_request/cif
11. Shida N, Kabuto C, Niwa T, Ebata T, Yamamoto Y. *J Org Chem* 1994;59:4068–4075. and references therein.
12. Pratt LM, Bowles SA, Courtney SF, Hidden C, Lewis CN, Martin FM, Todd RS. *Synlett* 1998:531–533.
13. Fueleop F, Palko M, Forro E, Dervarics M, Martinek TA, Sillanpaae R. *Eur J Org Chem* 2005:3214–3220.
14. The relative stereochemistry was determined by 2D COSY and NOESY spectra. The correlations are shown in the Supporting Information.
15. Shioiri T, Ninomiya K, Yamada S. *J Am Chem Soc* 1972;94:6203–6205. [PubMed: 5054412]
16. Sheehan JC, Hess GP. *J Am Chem Soc* 1955;77:1067–1068.
17. (a) Feringa BL. *Acc Chem Res* 2000;33:346–353. [PubMed: 10891052] (b) Arnold LA, Imbos R, Mandoli A, De Vries AHM, Naasz R, Feringa BL. *Tetrahedron* 2000;56:2865–2878. For a list of conditions that were examined for the enantioselective addition, see the Supporting Information.

Figure 1.
X-ray structure of amide **3**.

Scheme 1.
General Three-Component Coupling with Enolates

Scheme 2.
Proposed Conjugate Addition/[3,3]-Rearrangement

Scheme 3.
Proposed Transition State for Ireland-Claisen Rearrangement

Scheme 4.
Site-Selective Bromolactonization

Scheme 5.
Curtius Rearrangement to a β -Amino Ester

Scheme 6.
DCC Coupling of Carboxylic Acid and Amine

Table 1

Initial Results of Conjugate Addition to Diallyl Fumarate

entry	EtM	solvent	catalyst	results ^a
1	EtMgBr	THF	Cu(N ^t Bu-sal) ₂	mix of products ^b
2	EtMgBr	THF	None	85% conv
3	Et ₂ Zn	THF	Cu(N ^t Bu-sal) ₂	12% conv
4	Et ₂ Zn	Et ₂ O	Cu(N ^t Bu-sal) ₂	100% conv
5	Et ₂ Zn	Et ₂ O	CuBr·DMS	100% conv
6 ^c	Et ₂ Zn	Et ₂ O	Cu(N ^t Bu-sal) ₂	74% conv
7	Et ₂ Zn	Et ₂ O	none	80% conv
8 ^c	Et ₂ Zn	Et ₂ O	none	60% conv

^a Conversion determined by ¹H NMR spectroscopy.^b Undetermined mixture of 1,4-addition, 1,2-addition, and S_N2'.^c Reaction conducted without TMSCl.

Table 2

Scope of Tandem Michael Addition/Ireland–Claisen Rearrangement

entry	R_2Zn	substrate	product	yield (%) (dr) ^a
1	Et_2Zn	(<i>E</i>)- 1a ($R^1, R^2 = H$)	2a	83 (9:1)
2	Et_2Zn	(<i>Z</i>)- 1a ($R^1, R^2 = H$)	2a	74 (6.5:1)
3	Et_2Zn	(<i>E</i>)- 1b ($R^1, R^2 = Me$)	2b	68 (7.5:1)
4	iPr_2Zn	(<i>E</i>)- 1a ($R^1, R^2 = H$)	2c	70 (4.9:1)
5	Bu_2Zn^b	(<i>E</i>)- 1a ($R^1, R^2 = H$)	2d	88 (4.5:1)
6	Bu_2Zn^b	(<i>E</i>)- 1a ($R^1, R^2 = Me$)	2e	63 (2:1)
7	Et_2Zn	(<i>E</i>)- 1c ($R^1 = Me, R^2 = H$)	2f	75 (mix)
8	Et_2Zn	(<i>E</i>)- 1d ($R^1 = H, R^2 = Pr$)	2g	78 (mix)

^aYield of isolated products purified by column chromatography.^b1.5 equiv of Bu_2Zn (generated from 3.0 equiv of $BuLi$ and 1.5 equiv of $ZnCl_2$ in Et_2O), 4.0 equiv of TMSCl.