Published in final edited form as:

Biochemistry. 2008 March 25; 47(12): 3875–3882. doi:10.1021/bi7023783.

Effects of protein oxidation on the structure and stability of model discoidal high-density lipoproteins

Shobini Jayaraman, Donald L. Gantz, and Olga Gursky

Department of Physiology and Biophysics, Boston University School of Medicine, Boston, MA 02118

Abstract

High-density lipoproteins (HDLs) prevent atherosclerosis by removing cholesterol from macrophages and by providing anti-oxidants for low-density lipoproteins. Oxidation of HDLs affects their functions via the complex mechanisms that involve multiple protein and lipid modifications. To differentiate between the roles of oxidative modifications in HDL proteins and lipids, we analyzed the effects of selective protein oxidation by hypochlorite (HOCl) on the structure, stability and remodeling of discoidal HDLs reconstituted from human apolipoproteins (A-I, A-II or C-I) and phosphatidylcholines. Gel electrophoresis and electron microscopy revealed that, at ambient temperatures, protein oxidation in discoidal complexes promotes their remodeling into larger and smaller particles. Thermal denaturation monitored by far-UV circular dichroism and light scattering in melting and kinetic experiments shows that protein oxidation destabilizes discoidal lipoproteins and accelerates protein unfolding, dissociation and lipoprotein fusion. This is likely due to reduced protein affinity for lipid resulting from oxidation of Met and aromatic residues in the lipid-binding faces of amphipathic α -helices and to apolipoprotein cross-linking into dimers and trimers on the particle surface. We conclude that protein oxidation destabilizes HDL disk assembly and accelerates its remodeling and fusion. This result, which is not limited to model discoidal but also extends to plasma spherical HDL, helps explain the complex effects of oxidation on plasma lipoproteins.

Keywords

Thermal stability; lipoprotein remodeling and fusion; apolipoprotein A-I; reverse cholesterol transport; atherosclerosis

High-density lipoproteins (HDLs) remove excess cholesterol from the body and thereby prevent atherosclerosis. Plasma HDLs form a heterogeneous population of particles differing in their protein and lipid composition, shape, size, and functional properties. Nascent discoidal HDLs are comprised of a cholesterol-containing phospholipid bilayer and apolipoproteins wrapped around the particle perimeter in a beltlike amphipathic α -helical conformation (1). Mature spherical HDLs contain proteins, phospholipids and free (unesterified) cholesterol (FC) in their surface and apolar lipids (mainly cholesterol esters) in the core (2).

Apolipoprotein A-I (apoA-I, 28 kDa) is a key structural and functional component of HDL that comprises ~70 % of the total HDL protein content. Plasma levels of HDL and apoA-I are inversely related to the risk of atherosclerosis (3). HDLs protect against atherosclerosis by removing excess cholesterol from arterial macrophages and other peripheral cells via the reverse cholesterol transport (RCT) pathway (2). Other cardioprotective mechanisms have been proposed to involve the antioxidant and antiinflammatory properties of HDL and its

constituent proteins, including apoA-I (4-10). ApoA-I also plays key roles at various stages of RCT. It promotes efflux of cell cholesterol and phospholipids to HDL via the interactions with ATP-binding cassette transporter A1 (11); it provides a co-factor for lecithin:cholesterol acyltransferase (LCAT), which esterifies cholesterol and converts nascent discoidal to mature spherical HDL (2); it binds to the hepatic scavenger receptor and mediates selective uptake of HDL cholesterol esters by the liver (12).

Oxidative stress is thought to be among the key initiating factors in atherogenesis. According to the oxidation modification hypothesis, retention of oxidized low-density lipoproteins (oxLDLs) in the arterial wall triggers a cascade of pro-atherogenic events culminating in foam cell formation, a hallmark of early atherosclerosis (13). In contrast to oxLDL, functional consequences of HDL oxidation remain unclear (14-16). Some studies report that oxidation impairs HDL ability to promote cellular cholesterol efflux (14,17) while others suggest that oxidized HDL may actually enhance cholesterol efflux (18,19). This apparent controversy may result from the differences in the oxidizing agents and their products as well as in the extent of oxidation used in different studies. In fact, Pirillo and colleagues reported that mild oxidation by copper improves HDL ability to accept cell cholesterol, while extensive oxidation inhibits this ability (20). Our recent study is consistent with this report and shows that mild oxidation by copper or hypochlorite destabilizes mature human HDL and accelerates heat-induced protein dissociation and lipoprotein fusion, while extensive oxidation inhibits these reactions (21). To clearly understand these oxidation-induced changes in functions and remodeling of plasma HDL, one has to differentiate between the complex effects of multiple protein and lipid modifications that occur concomitantly at various stages of oxidation. To do so, we use discoidal reconstituted HDLs (rHDLs) of controlled composition, such as complexes of human apoA-I with dimyristoyl phosphatidylcholine (DMPC), as simple experimental models for dissecting the effects of protein oxidative modifications by hypochlorite on HDL structure and stability.

Hypochlorite is a product of myeloperoxidase (MPO), a heme protein expressed by macrophages in atherosclerotic lesions that is thought to be an important *in vivo* mediator of lipoprotein oxidation (15,22,23). The active form of MPO and the proteins modified by MPO-generated HOCl have been found in human atherosclerotic lesions, supporting the notion that MPO mediates HDL oxidation in the artery wall (24,25). Physiological concentrations of HOCl have not been accurately determined, although some *in-vivo*-based reports suggest that, at sites of acute inflammation, the molar ratio of oxidant to HDL may be as high as 30:1 (26 and references therein). At these and lower HOCl levels and near-neutral pH, oxidative modifications in HDL are mainly limited to apolipoproteins (26,27). We use this property of HOCl to test the effects of apolipoprotein oxidation on the structure and stability of model discoidal HDL.

Dose-dependent HOCl-induced modifications in the major HDL proteins, apoA-I and apoA-II, have been well-documented by using limited proteolysis, mass spectrometry and liquid chromatography; the general consensus is that oxidation of Met and Cys occurs first, followed by modifications of Lys and aromatic residues and, eventually, protein cross-linking into homo- or heterodimers and trimers (24,26-30). Specifically, limited proteolysis and mass spectrometry studies of apoA-I in solution or on HDL suggest that, at a 5:1 HOCl:protein molar ratio, Met 86, Met 112 and Met 148 are converted to methionine sulfoxides (MetO); at a 10:1 ratio, Phe 37 and Phe 71 are the primary targets of oxidation; and at a 25:1 ratio, Tyr 192 is chlorinated (25,28). In addition, progressive oxidation of Trp residues in apoA-I has been detected by fluorescence (31). Oxidative modifications in these and other hydrophobic residues, which are located in the apolar faces of the amphipathic apolipoprotein α -helices, are expected to impair lipid binding affinity of apoA-I and affect its functions. In fact, oxidation of Met 112 and Met 148 to MetO reportedly impairs apoA-I ability to activate LCAT (29); it

also leads to partial unfolding and destabilization of the α -helices in lipid-free and in lipid-bound apoA-I and apoA-II, as well as to reduced protein affinity for lipid that is attributed to reduced hydrophobicity of the Met-containing apolar helical faces (29,30,32). Oxidative cross-linking of apoA-I and apoA-II in HDL, which occurs via the tyrosyl radicals, reportedly alters apoA-I conformation, leading to increased solvent exposure of the protein α -helices (33,34) and to enhanced ability of HDL to promote cholesterol efflux from fibroblasts (35). Here, we analyze the effects of these oxidative modifications on thermal stability and remodeling of discoidal rHDLs.

Earlier, we established a kinetic mechanism of lipoprotein stabilization and showed that thermal or chemical denaturation of model discoidal and plasma spherical HDLs leads to protein unfolding and dissociation and to particle fusion (36-40). Importantly, this denaturation reaction resembles metabolic HDL remodeling and fusion by plasma factors such as LCAT (41,42). Thus, our *in vitro* studies of HDL denaturation provide a useful experimental approach for understanding lipoprotein remodeling *in vivo*. Here, we use this approach to analyze the effects of protein oxidation on the structure, stability and remodeling of discoidal rHDLs. The results complement our recent oxidation studies of plasma spherical HDL (21) and help dissect the effects of protein and lipid oxidation on HDL remodeling and functions.

Materials and Methods

Materials

Diethylenetriamine pentaacetic acid (DTPA, >98% purity) and sodium hypochlorite, NaOCl (<5% chlorine) were from Across Organic. Lipids, including DMPC, dipalmitoyl-PC (DPPC), palmitoyl-oleoyl-PC (POPC), and FC were from Avanti Polar Lipids (95+% purity). All chemicals were of highest purity analytical grade.

Protein Preparation and Characterization

ApoA-I and apoA-II (an S-S linked dimer of two 77 a. a. monomers and a second-major HDL protein) were isolated and purified from human plasma HDL and refolded as described (38). Human apoC-I (a minor HDL protein of 57 a. a. that is a structural and functional prototype of larger exchangeable apolipoproteins) was custom-synthesized and purified to 97%+ purity at 21st Century Biochemicals as described (36).

Lipoprotein Reconstitution

DMPC complexes with apolipoproteins were obtained by adding protein stock solution to DMPC suspension (protein:lipid ratio 1:4 mg/mg) followed by overnight incubation at 24 °C. Discoidal complexes of apoA-I with DPPC, POPC, or with PC and FC were prepared by sodium cholate dialysis (43). For DPPC and POPC complexes, PC:A-I:cholate molar ratios of 80:1:80 were used; complexes containing FC were prepared using PC:FC:A-I initial molar ratios of 100:50:1. Lipoprotein formation was confirmed by negative staining electron microscopy (EM) and non-denaturing polyacrylamide gel electrophoresis (PAGE).

Protein Oxidation by Hypochlorite

To oxidize apoA-I to various stages, 1 mg/mL of protein (lipid-free or in complex with lipid) in standard buffer (10 mM Na phosphate, 100 μ M DTPA, pH 7.5) was incubated for 12 h at 24 °C with NaOCl using oxidant:protein molar ratios from 5:1 to 50:1 (44). Concentration of HOCl was determined spectrophotometrically, using extinction coefficient $\epsilon_{290} = 350 \text{ M}^{-1}\text{cm}^{-1}$ (45). The reaction was terminated with 2.5 mM Met; the samples were dialyzed overnight at 4 °C against standard buffer prior to use. Complexes reconstituted from intact apoA-I and DMPC that were oxidized to various stages by HOCl (termed ox(A-I:DMPC)

throughout this work) were compared with similar complexes that were reconstituted from oxidized apoA-I and DMPC (termed (oxA-I):DMPC). The absence of oxidative lipid modifications in ox(A-I):DMPC was confirmed by thin-layer chromatography as described (46).

Gel Electrophoresis

SDS PAGE (12% w/v gel) was performed under non-reducing conditions in a Mini Protean II chamber (Bio-Rad) for 2 h at 120 V; 10 µg protein aliquots were applied per lane. The gel was stained with Imperial protein stain (Pierce).

Non-denaturing PAGE was performed using self-made 4-15% gradient gels. Protein samples (10 µg) were run for 2 h at 120 V. The gel was stained with Imperial protein stain. The particle diameters were determined from comparison with the high molecular-weight standard (GE Healthcare).

Circular Dichroism Spectroscopy

Far-UV CD and 90° light scattering data were collected using an AVIV 215 spectropolarimeter with thermoelectric temperature control as described (38,47). In temperature-jump (T-jump) experiments, the sample temperature was rapidly increased at time $t=0$ from 25 °C to a constant value ranging from 75 to 95 °C and the time course of the protein unfolding was monitored at 222 nm. Helical content was calculated from the measured molar residue ellipticity [Θ_{222}] as described (36).

Electron Microscopy

Protein-lipid complexes were visualized by negative staining EM under low-dose conditions in a CM12 transmission electron microscope (Philips Electron Optics) as described (36,38). The disk diameters were determined from these micrographs using 200-350 particles.

Fluorescence Spectroscopy

Trp emission spectra were recorded at 25 °C with a Fluoromax-2 spectrofluorimeter from apoA-I solutions of 50 µg/mL concentration in standard buffer. The excitation wavelength was 280 nm; the spectra were recorded from 310-500 nm with 5 nm excitation and emission slit widths.

DMPC Clearance

DMPC multilamellar vesicles were prepared by thin film evaporation; DMPC solution in methanol/chloroform (2:1 v/v) was evaporated under nitrogen and the lipid film was hydrated with the standard buffer. Clearance of DMPC multilamellar vesicles by normal or oxidized apoA-I was monitored at 24 °C by turbidity at 325 nm using a Varian Cary-300 UV/vis absorption spectrophotometer equipped with thermoelectric temperature control as described (40).

All experiments in this study were repeated 3-5 times to ensure reproducibility.

Results

First, we analyzed A-I:DMPC complexes at various stages of oxidation corresponding to oxidant:protein ratios from 5:1 to 50:1. Protein cross-linking was assessed by SDS PAGE (Fig. 1). Intact apoA-I ran as a single band corresponding to monomeric protein (~28 kDa, lane 0 in Fig. 1). Oxidation of lipid-free (lanes 1', 2') and DMPC-bound apoA-I (lanes 1, 2) produced two additional bands near 50 and 100 kDa reflecting oxidation-induced dimer and trimer

formation that was confirmed by SDS PAGE (44). Increasing oxidant:protein ratios to 25:1 and beyond resulted in near-complete conversion of monomeric to cross-linked protein on discoidal particles (lanes 3, 4); lipid-free apoA-I (data not shown) undergoes extensive intramolecular cross-linking under these conditions (44). At lower oxidant:protein ratios ($\leq 10:1$) that encompass the range of physiologically relevant conditions (26), lipid-bound and lipid-free apoA-I show comparable population distribution of monomeric, dimeric and trimeric species (lanes 1, 2 and 1', 2' compared); therefore, we used these relatively low ratios in our analysis of (oxA-I):DMPC complexes and their comparison with ox(A-I:DMPC) disks.

Effects of protein oxidation on the particle size at 25 °C were analyzed by native PAGE (Fig. 2). Intact disks formed two distinct populations of about 11 and 13 nm in size (Fig. 2, lane 0). Oxidation of apoA-I:DMPC disks progressively increased their heterogeneity, leading to formation of smaller and larger particles (Fig. 2, lanes 1-4). Complexes reconstituted from oxA-I and DMPC appeared larger than intact or ox(A-I:DMPC) disks and showed an increase in the particle size with increasing degree of protein oxidation (Fig. 2, lanes 1', 2').

EM analysis confirmed these results (Fig. 3). Electron micrographs of intact apoA-I:DMPC complexes showed disks of average diameter $\langle d \rangle = 11$ nm (Fig. 3A). Oxidation of these disks using oxidant:protein ratios of 10:1 and above lead to formation of smaller and larger ox(A-I:DMPC) particles with broad diameter distribution from about 9-22 nm (Fig. 3C-E).

Complexes reconstituted from oxidized apoA-I and DMPC, (oxA-I):DMPC, were larger than intact or ox(A-I:DMPC) disks, and the average diameter of these complexes was about 22 nm (Fig. 3F, G). Thus, the results of native PAGE and EM studies consistently show that oxidation of apoA-I in discoidal particles leads to disk remodeling into larger and smaller particles, while oxidation of lipid-free A-I prior to lipoprotein reconstitution leads to formation of only larger particles.

Oxidation-induced secondary structural changes in apoA-I were analyzed by CD spectroscopy. Far-UV CD spectra in Fig. 4 suggest that intact disks have about 60% α -helical content that is reduced to about 50% at oxidant:protein molar ratios of 5:1 and 10:1, and to about 40% α -helix at oxidant:protein ratios of 25:1 and 50:1. Similarly, lipid-free protein showed up to 20% loss in its helical content under these oxidative conditions (CD data not shown). These results are consistent with the earlier studies reporting partial α -helical unfolding in apoA-I upon oxidation (28).

Oxidation-induced remodeling of A-I:DMPC discs at ambient temperatures (Fig. 3), together with their reduced α -helical content (Fig. 4), suggests that protein oxidation destabilizes the disk assembly. To test this notion, we analyzed thermal denaturation of A-I:DMPC disks that were oxidized to various stages. Intact and ox(A-I:DMPC) complexes were heated and cooled from 10 to 98 °C at a rate of 11 °C/h, and thermal denaturation and reconstitution were monitored at 222 nm by CD and by 90° light scattering for changes in the α -helical structure and in the particle size, respectively (35). The results show that oxidation leads to progressive low-temperature shifts in the apparent temperature of disk denaturation, from $T_m = 82$ °C in intact disks to 60 °C in ox(A-I:DMPC) disks that were modified using 50:1 oxidant:protein ratio (Fig. 5A, B). Thus, the CD and light scattering melting data confirm that oxidation destabilizes A-I:DMPC disks and promotes their fusion.

To test whether this notion applies to (oxA-I):DMPC complexes, we compared the melting data recorded from these complexes by CD (Fig. 5C) and light scattering (not shown) with similar data recorded from ox(A-I:DMPC) disks. This comparison was made using 5:1 and 10:1 oxidant:protein ratios that produce similar cross-linking patterns in lipid-bound and lipid-free protein (Fig. 1, lanes 1, 1' and 2, 2'). The results show that the melting data of ox(A-I:DMPC) and (oxA-I):DMPC complexes largely overlap (Fig. 5C). Thus, despite large

difference in the diameters of these complexes observed at ambient temperatures (Fig. 3B, C and E, F), their apparent thermal stability is similar and is significantly lower than that of intact disks.

To further test this notion and to determine the effect of oxidation on the kinetics of lipoprotein denaturation, we monitored the rate of α -helical unfolding at 222 nm in T-jump experiments. Lipoprotein samples were rapidly heated from 25 °C to high temperature and the protein unfolding was monitored by CD at 222 nm as a function of time. Fig. 6 illustrates the results of such kinetic experiments at 80 °C. Kinetic T-jump data at this and other temperatures clearly show progressive acceleration of protein unfolding upon oxidation, with the slowest unfolding observed in intact disks and fastest in ox(A-I:DMPC) disks that were oxidized using 50:1 HOCl:protein ratio (lines 0-4 in Fig. 6). Furthermore, at similar oxidant:protein ratios of 5:1 or 10:1, the kinetic CD data recorded of (oxA-I):DMPC and ox(A-I:DMPC) complexes fully overlapped, confirming their identical stability (lines 1, 1' in Fig. 6). Although T-jump data of intact A-I:DMPC disks were well-approximated by single exponentials (38), similar data of oxidized complexes could not be fitted with exponential functions, thereby precluding quantitative Arrhenius analysis of the effects of oxidation on the kinetic lipoprotein stability. Nevertheless, our CD and light scattering melting and kinetic data (Figs. 5, 6) clearly show that protein oxidation progressively destabilizes lipoproteins and accelerates protein unfolding and lipoprotein remodeling and fusion.

Fluorescence spectra of lipid-free apoA-I and of A-I:DMPC complexes showed a gradual reduction in the intensity of the Trp emission peak centered at 330 nm with increasing oxidant:protein ratio from 5:1 to 50:1 (Fig. 7A), which is characteristic of Trp oxidation (31). Furthermore, turbidity measurements showed a gradual reduction in the rate of DMPC clearance by apoA-I upon increased protein oxidation (Fig. 7B). These results are consistent with earlier reports (29,32) and with our stability studies (Figs. 4,5) and indicate that oxidation of apoA-I progressively reduces its affinity for lipid. This may result from oxidative modifications of hydrophobic residues (Met, Trp, Tyr, Phe, etc.) located in the lipid-binding helical faces, which reduce the lipophilicity of these helices. In addition, apolipoprotein cross-linking, which reportedly increases solvent exposure of the α -helices (33,34) and is accompanied by partial α -helical unfolding on the disks (Fig. 4), may also contribute to the reduced lipophilicity.

To test whether oxidation-induced lipoprotein destabilization is limited to A-I: DMPC disks or is a general property of discoidal HDLs, we analyzed the effects of oxidation on thermal stability of complexes reconstituted from various proteins and lipids. In addition to apoA-I, we used homologous exchangeable apolipoproteins apoA-II (38,39) and apoC-I; these proteins contain a single Met in the apolar helical face that can form MetO, as well as several Tyr residues that can be cross-linked by HOCl. In addition to DMPC (14:0, 14:0), we reconstituted disks with longer-chain PCs that were fully saturated or mono-unsaturated, such as dipalmitoyl-PC (16:0, 16:0) or palmitoyl:oleoyl:PC (16:0, 18:1); such PCs better approximate lipid composition of plasma HDL. Furthermore, we reconstituted A-I:DMPC:FC disks containing 10% FC, oxidized them to various stages by HOCl, and assessed the ensuing changes in their thermal stability by CD. The CD melting data in Fig 8 illustrate gradual destabilization of A-I:DMPC:FC complexes upon increased degree of protein oxidation. These and other results consistently show that, similar to A-I:DMPC disks, selective protein oxidation in other discoidal rHDLs gradually destabilizes these rHDLs and accelerates protein unfolding and dissociation, as well as lipoprotein remodeling and fusion.

Discussion

The results reported here show that oxidation of the exchangeable apolipoproteins by HOCl progressively destabilizes discoidal rHDLs of various composition and accelerates protein dissociation and lipoprotein remodeling and fusion (Figs. 5, 6, 8). This is consistent with the earlier reports of destabilization of α -helices in lipid-free and lipid-bound apoA-I and apoA-II upon oxidation with hydrogen peroxide (29,32). This destabilization was attributed to reduced protein affinity for lipid upon MetO formation in the apolar lipid-binding faces of the amphipathic apolipoprotein α -helices (29). Such MetO formation occurs at early stages of oxidation and is followed by modifications of other protein groups, such as aromatic residues and Lys, and their cross-linking (24-31). Correlation of rHDL destabilization (Fig. 5,6) with partial α -helical unfolding (Fig. 4), Trp oxidation (Fig. 7a), protein cross-linking (Fig. 1) and decelerated lipid clearance (Fig. 7B) suggests that not only MetO formation but also HOCl-induced modifications of other aromatic groups located in the lipid-binding faces of the protein α -helices, as well as protein cross-linking, may contribute to reduced protein affinity for lipid and to rHDL destabilization. In fact, cross-linking of lipid-bound apoA-I reduces its helical content (Fig. 4) and reportedly increases solvent exposure of specific protein α -helices (16, 33,34), which may weaken protein-lipid interactions and promote protein dissociation and lipoprotein remodeling and fusion.

Our results reveal that oxidative protein modifications in apoA-I:DMPC disks lead to rHDL remodeling into larger and smaller particles. Such a remodeling was observed at or above 10:1 oxidant:protein ratio (Fig. 3C-E), which probably is well in excess of the normal HOCl levels and hence is unlikely to occur *in vivo*. Remodeling of spherical lipoproteins into larger and smaller particles has been well-documented. For example, conversion of plasma HDL into larger and smaller particles occurs upon remodeling by plasma factors, such as phospholipid transfer protein (48), probably due to imbalance between the core and surface of spherical particles. Another example is heat-induced conversion of very low-density lipoproteins into larger and smaller spherical particles resulting in transient formation of small HDL-like lipoproteins (49). To our knowledge, the current study provides the first report of remodeling of discoidal lipoproteins into larger and smaller particles. Larger particles are probably fusion products of intact-size disks; such fusion may result from conformational changes in oxidized apoA-I leading to increased exposure of lipid surface in discoidal complexes. The mechanism of dissociation of smaller particles is not entirely clear. Formation of these particles is concomitant with the disappearance of the monomeric apoA-I at oxidation stages 2-4 (lanes 2-4 in Fig. 1 and panels C-E in Fig. 3), suggesting that the conformation of apoA-I that is altered upon cross-linking or other oxidative modifications favors dissociation of such small particles.

Since apolipoprotein dissociation and lipoprotein fusion occur not only during denaturation of discoidal rHDLs but also during denaturation and metabolic remodeling of plasma spherical HDLs, analysis of the effects of oxidative modifications on these reactions may provide molecular basis for understanding functional consequences of HDL oxidation. Interestingly, in contrast to discoidal rHDLs that are progressively destabilized upon oxidation (Figs. 5, 6), plasma spherical HDLs are destabilized only at early oxidation stages corresponding to low HOCl:protein ratio up to 10:1, but show increased resistance to protein dissociation and particle remodeling at higher oxidant levels (21). We hypothesized that initial destabilization of plasma HDL results from oxidative modifications to its major proteins, apoA-I and apoA-II, while the reversal of this effect upon further oxidation is due to modifications of core and surface lipids (21).

The results of the current work strongly support this hypothesis. They show that, regardless of the initial protein state (lipid-free or in discoidal rHDL), mild oxidation of apoA-I leads to

protein cross-linking into dimers and trimers (Fig. 1) and to destabilization of the rHDL assembly (Figs. 5, 6, 8). This suggests strongly that apoA-I cross-linking into dimers and trimers that occurs at early oxidation stages in plasma spherical HDL is also destabilizing. We propose that such moderate destabilization of plasma HDLs benefits their functions, since it facilitates cholesterol efflux to mildly oxidized HDLs, promotes apolipoprotein exchange among lipoproteins, and accelerates HDL remodeling by plasma factors during RCT (20,21). The results reported here show that this functional advantage results, at least in part, from modifications to HDL proteins that occur at early stages of oxidation, such as oxidation of Met and aromatic groups in the apolar helical faces, that destabilize HDL assembly and promote HDL remodeling.

Acknowledgments

We are indebted to Cheryl England and Michael Gigliotti for apolipoprotein isolation and to Allison Koch for help with particle size analysis.

This work was supported by the National Institutes of Health grants RO1 GM 067260 and HL 026355.

Abbreviations

RCT	reverse cholesterol transport
HDL	high-density lipoprotein
rHDL	reconstituted HDL
LDL	low-density lipoprotein
oxLDL	oxidized LDL
(oxA-I):DMPC	complexes reconstituted using oxidized apoA-I
ox(A-I):DMPC)	complexes that were reconstituted using intact apoA-I and then oxidized
apo	apolipoprotein
PC	phosphatidyl choline
DMPC	1,2-dimyristoyl- <i>sn</i> -glycero-3-phosphocholine
DPPC	1,2-dipalmitoyl- <i>sn</i> -glycero-3-phosphocholine
POPC	1-palmitoyl-2-oleoyl- <i>sn</i> -glycero-3-phosphocholine
FC	free cholesterol
MetO	methionine sulfoxide (S=O double bond)
PAGE	polyacrylamide gel electrophoresis
CD	circular dichroism
T-jump	temperature jump
EM	electron microscopy

References

1. Li L, Chen J, Mishra VK, Kurtz JA, Cao D, Klon AE, Harvey SC, Anantharamaiah GM, Segrest JP. Double belt structure of discoidal high density lipoproteins: Molecular basis for size heterogeneity. *J Mol Biol* 2004;343(5):1293–1311. [PubMed: 15491614]
2. Fielding CJ, Fielding PE. Molecular physiology of reverse cholesterol transport. *J Lipid Res* 1995;36(2):211–228. [PubMed: 7751809]

3. Asztalos BF. High density lipoprotein metabolism and progression of atherosclerosis: New insights from the HDL atherosclerosis treatment study. *Curr Opin Cardiol* 2004;19(4):385–391. [PubMed: 15218401]
4. Barter PJ, Nicholls S, Rye KA, Anantharamaiah GM, Navab M, Fogelman AM. Anti-inflammatory properties of HDL. *Circ Res* 2004;95(8):764–772. [PubMed: 15486323]
5. Parthasarathy S, Barnett J, Fong LG. High-density lipoprotein inhibits the oxidative modification of low-density lipoprotein. *Biochim Biophys Acta* 1990;1044:275–283. [PubMed: 2344447]
6. Kontush A, Chantepie S, Chapman MJ. Small, dense HDL particles exert potent protection of atherogenic LDL against oxidative stress. *Arterioscler Thromb Vasc Biol* 2003;23(10):1881–1888. [PubMed: 12920049]
7. Stocker R, Keaney JF Jr. Role of oxidative modifications in atherosclerosis. *Physiol Rev* 2004;84:1381–1478. [PubMed: 15383655]
8. Navab M, Anantharamaiah GM, Reddy ST, Van Lenten BJ, Ansell BJ, Fonarow GC, Vahabzadeh K, Hama S, Hough G, Kamranpour N, Berliner JA, Lusis AJ, Fogelman AM. The oxidation hypothesis of atherogenesis: the role of oxidized phospholipids and HDL. *J Lipid Res* 2004;45(6):993–1007. [PubMed: 15060092]
9. Witztum JL, Steinberg D. The oxidative modification hypothesis of atherosclerosis: does it hold for humans? *Trends Cardiovasc Med* 2001;11(3-4):93–102. [PubMed: 11686009]
10. Rohrer L, Hersberger M, Von Eckardstein A. High density lipoproteins in the intersection of diabetes mellitus, inflammation and cardiovascular disease. *Curr Opin Lipidol* 2004;15(3):269–278. [PubMed: 15166782]
11. Vedhachalam C, Duong PT, Nickel M, Nguyen D, Dhanasekaran P, Saito H, Rothblat GH, Lund-Katz S, Phillips MC. Mechanism of ATP-binding cassette transporter A1-mediated cellular lipid efflux to apolipoprotein A-I and formation of high density lipoprotein particles. *J Biol Chem* 2007;282(34):25123–25130. [PubMed: 17604270]
12. Xu S, Laccotripe M, Huang X, Rigotti A, Zannis VI, Krieger M. Apolipoproteins of HDL can directly mediate binding to the scavenger receptor, SR-BI, and HDL receptor that mediates selective lipid uptake. *J Lipid Res* 1997;38:1289–1298. [PubMed: 9254056]
13. Heinecke JW. Mechanisms of oxidative damage of low density lipoprotein in human atherosclerosis. *Curr Opin Lipidol* 1997;8:268–274. [PubMed: 9335950]
14. Nagano Y, Ami H, Kita T. High density lipoprotein loses its effect to stimulate efflux of cholesterol from foam cells after oxidative modification. *Proc Natl Acad Sci USA* 1991;88:6457–6461. [PubMed: 1862074]
15. Francis GA. High density lipoprotein oxidation: *In vitro* susceptibility and potential *in vivo* consequences. *Biochim Biophys Acta* 2000;1483:217–235. [PubMed: 10634938]
16. Panzenbock U, Kritharides L, Raftery M, Rye KA, Stocker R. Oxidation of methionine residues to methionine sulfoxides does not decrease potential antiatherogenic properties of apolipoprotein A-I. *J Biol Chem* 2000;275(26):19536–19544. [PubMed: 10751387]
17. Bergt C, Reicher H, Malle E, Sattler W. Hypochlorite modification of high density lipoprotein: effects on cholesterol efflux from J774 macrophages. *FEBS Letters* 1999;452:295–300. [PubMed: 10386609]
18. Francis GA, Mendez AJ, Biennan EL, Heinecke JW. Oxidative tyrosylation of high density lipoprotein by peroxidase enhances cholesterol removal from cultured fibroblasts and macrophage foam cells. *Proc Natl Acad Sci USA* 1993;90:6631–6635. [PubMed: 8341680]
19. Francis GA, Oram JF, Heinecke JW, Bieman EL. Oxidative tyrosylation of HDL enhances the depletion of cellular cholesteryl esters by a mechanism independent of passive sterol desorption. *Biochemistry* 1996;35:15188–15197. [PubMed: 8952466]
20. Pirillo A, Uboldi P, Pappalardo G, Kuhn H, Catapano AL. Modification of HDL₃ by mild oxidative stress increases ATP-binding cassette transporter 1-mediated cholesterol efflux. *Cardiovasc Research* 2007;75(3):566–574.
21. Gao X, Jayaraman S, Gursky O. Mild oxidation promotes and advanced oxidation impairs remodeling of human high-density lipoprotein *in vitro*. *J Mol Biol.* 2008;10.1016/j.jmb.2007.12.030
22. Heinecke JW. Oxidants and antioxidants in the pathogenesis of atherosclerosis: implications for the oxidized low density lipoprotein hypothesis. *Atherosclerosis* 1998;141:1–15. [PubMed: 9863534]

23. Berliner JA, Heinecke JW. The role of oxidized lipoproteins in atherogenesis. *Free Radical Biol Med* 1996;20:707–727. [PubMed: 8721615]
24. Hazen SL, Heinecke LW. 3-Chlorotyrosine, a specific marker of myeloperoxidase-catalyzed oxidation, is markedly elevated in low density lipoprotein isolated from human atherosclerotic intima. *J Clin Invest* 1997;99:2075–2080. [PubMed: 9151778]
25. Bergt C, Pennathur S, Fu X, Byun J, O'Brien K, McDonald TO, Singh P, Anantharamaiah GM, Chait A, Brunzell J, Geary RL, Oram JF, Heinecke JW. The myeloperoxidase product hypochlorous acid oxidizes HDL in the human artery wall and impairs ABCA1-dependent cholesterol transport. *Proc Natl Acad Sci USA* 2004;101(35):13032–13037. [PubMed: 15326314]
26. Panzenbock U, Raitmayer S, Reicher H, Lindner H, Glaner O, Malle E, Sattler W. Effects of reagent and enzymatically generated hypochlorite on physiochemical and metabolic properties of high density lipoproteins. *J Biol Chem* 1997;272:29711–29720. [PubMed: 9368040]
27. Hazen SL, Hsu FF, Duffin K, Heinecke JW. Molecular chlorine generated by the myeloperoxidase–hydrogen peroxide–chloride system of phagocytes converts low density lipoprotein cholesterol into a family of chlorinated sterols. *J Biol Chem* 1996;271:23080–23088. [PubMed: 8798498]
28. Bergt C, Oettl K, Keller W, Andreas F, Leis HJ, Malle E, Sattler W. Reagent or myeloperoxidase-generated hypochlorite affects discrete regions in lipid-free and lipid-associated human apolipoprotein A-I. *Biochem J* 2000;346:345–354. [PubMed: 10677352]
29. Anantharamaiah GM, Hughes TA, Iqbal M, Gawish A, Neame PJ, Medley MF, Segrest JP. Effect of oxidation on the properties of apolipoproteins A-I and A-II. *J Lipid Res* 1988;29:309–318. [PubMed: 3132519]
30. Von Eckardstein A, Walter M, Holz H, Benninghoven A, Assmann G. Site-specific methionine sulfoxide formation is the structural basis of chromatographic heterogeneity of apolipoproteins A-I, C-II, and C-III. *J Lipid Res* 1991;32:1465–1476. [PubMed: 1753217]
31. Peng DQ, Wu Z, Brubaker G, Zheng L, Settle M, Gross E, Kinter M, Hazen SL, Smith JD. Tyrosine modification is not required for myeloperoxidase-induced loss of apolipoprotein A-I functional activities. *J Biol Chem* 2005;280(40):33775–33784. [PubMed: 16091367]
32. Sigalov AB, Stern LJ. Oxidation of methionine residues affects the structure and stability of apolipoprotein A-I in reconstituted high density lipoprotein particles. *Chem Phys Lipids* 2001;113:133–146. [PubMed: 11687233]
33. Artola RL, Conde CB, Bagatolli L, Pécora RP, Fidelio GD, Kivatinitz SC. High-density lipoprotein from hypercholesterolemic animals has peroxidized lipids and oligomeric apolipoprotein A-I: its putative role in atherogenesis. *Biochem Biophys Res Commun* 1997;239(2):570–574. [PubMed: 9344872]
34. Greilberger J, Jurgens G. Oxidation of high-density lipoprotein HDL₃ leads to exposure of apo-AI and apo-AII epitopes and to formation of aldehyde protein adducts, and influences binding of oxidized low-density lipoprotein to type I and type III collagen in vitro. *Biochem J* 1998;331(1):185–191. [PubMed: 9512478]
35. Wang WQ, Merriam DL, Moses AS, Francis GA. Enhanced cholesterol efflux by tyrosyl radical-oxidized high density lipoprotein is mediated by apolipoprotein AI-AII heterodimers. *J Biol Chem* 1998;273(28):17391–17398. [PubMed: 9651324]
36. Gursky O, Mehta R, Gantz DL. Complex of human apolipoprotein C-1 with phospholipid: Thermodynamic or kinetic stability? *Biochemistry* 2002;41:7373–7384. [PubMed: 12044170]
37. Mehta R, Gantz DL, Gursky O. Human plasma high-density lipoproteins are stabilized by kinetic factors. *J Mol Biol* 2003;328(1):183–192. [PubMed: 12684007]
38. Jayaraman S, Gantz DL, Gursky O. Kinetic stabilization and fusion of apolipoprotein A-2:DMPC disks: Comparison with apoA-1 and apoC-1. *Biophys J* 2005;88:2907–2918. [PubMed: 15681655]
39. Jayaraman S, Gantz DL, Gursky O. Effects of salt on the thermal stability of human plasma high-density lipoprotein. *Biochemistry* 2006;45(14):4620–4628. [PubMed: 16584197]
40. Benjwal S, Jayaraman S, Gursky O. Electrostatic effects on the stability of discoidal high-density lipoproteins. *Biochemistry* 2005;44(30):10218–10226. [PubMed: 16042399]
41. Rye KA, Clay MA, Barter PJ. Remodeling of high density lipoproteins by plasma factors. *Atherosclerosis* 1999;145(2):227–238. [PubMed: 10488948]

42. Barter PJ. The regulation and remodeling of HDL by plasma factors. *Atheroscler Suppl* 2002;3(4): 39–47. [PubMed: 12573362]
43. Jonas A. Reconstitution of high-density lipoproteins. *Methods Enzymol* 1986;128:553–582. [PubMed: 3724523]
44. Shao B, Bergt C, Fu X, Green P, Voss CJ, Oda NM, Oram JF, Heinecke JW. Tyrosine 192 in apolipoprotein A-I is the major site of nitration and chlorination by myeloperoxidase, but only chlorination markedly impairs ABCA1-dependent cholesterol transport. *J Biol Chem* 2005;280:5983–5993. [PubMed: 15574409]
45. Arnhold J, Weigel D, Richter O, Hammerschmidt S, Arnold K, Krumbiegel M. Modification of low density lipoproteins by sodium hypochlorite. *Biomed Biochim Acta* 1991;50:967–997. [PubMed: 1723592]
46. Jayaraman S, Gantz DL, Gursky O. Effects of oxidation on the structure and stability of human low-density lipoprotein. *Biochemistry* 2007;46(19):5790–5797. [PubMed: 17444660]
47. Benjwal S, Verma S, Röhm KH, Gursky O. Monitoring protein aggregation during thermal unfolding in circular dichroism experiments. *Protein Sci* 2006;15:635–639. [PubMed: 16452626]
48. Settasatian N, Duong M, Curtiss LK, Ehnholm C, Jauhainen M, Huuskonen J, Rye KA. The mechanism of the remodeling of high density lipoproteins by phospholipid transfer protein. *J Biol Chem* 2001;276(29):26898–26905. [PubMed: 11325961]
49. Guha M, England C, Herscovitz H, Gursky O. Thermal transitions in human very-low density lipoprotein: fusion, rupture and dissociation of HDL-like particles. *Biochemistry* 2007;46(20):6043–6049. [PubMed: 17469851]

Figure 1.

Effects of apoA-I oxidation to various stages on protein cross-linking analyzed by SDS PAGE. Lipid-free apoA-I (lanes 0, 1', 2') or A-I:DMPC disks (lanes 1-4) containing 1 mg/mL protein in standard buffer (10 mM Na phosphate, 100 μ M DTPA, pH 7.5), were incubated with various concentrations of HOCl as described in Methods. Lane numbers correspond to oxidant:protein molar ratios of 0:1 (0), 5:1 (1, 1'), 10:1 (2, 2'), 25:1 (3), 50:1 (4). Gel was stained with Imperial protein stain.

Figure 2.

Effects of protein oxidation on the size of apoA-I:DMPC complexes at 25 °C analyzed by non-denaturing PAGE. Ten μ l of 1 mg/mL apoA-I solution was applied to each lane; at this concentration, lipid-free apoA-I aggregates and runs at about 7 - 8 nm (lane A-I, (1)). Lanes 0-4 show discoidal complexes reconstituted from intact apoA-I and DMPC that were oxidized using oxidant:protein molar ratios of 0:1 (0), 5:1 (1), 10:1 (2), 25:1 (3), or 50:1 (4). Lanes 1' and 2' show complexes reconstituted from oxidized apoA-I and DMPC; the oxidant:protein ratios are 5:1 (1') or 10:1 (2'). Molecular size standards (st) in nm are indicated.

Figure 3.

Effects of protein oxidation on the size and morphology of A-I:DMPC complexes analyzed by negative staining EM. Upper row shows electron micrographs and lower row shows the corresponding particle size distributions. Numbers in panels A-E correspond to those in Fig. 2 and show intact A-I:DMPC disks (0) and disks that were treated with oxidant:protein ratios of 5:1 (1), 10:1 (2), 25:1 (3), 50:1 (4). Panels F, G show disks reconstituted from oxidized apoA-I and DMPC; HOCl:A-I ratios are 5:1 (1') or 10:1 (2'). Average particle diameters with standard deviations are indicated in selected panels.

Figure 4.

Effect of oxidation on the secondary structure of apoA-I in DMPC complexes at 22 °C. Far-UV CD spectra were recorded from the ox(A-I:DMPC) disks (20 $\mu\text{g}/\text{mL}$ protein) that were incubated with HOCl using oxidant:protein ratios of 0:1 (0), 5:1 (1), 10:1 (2), 25:1 (3), 50:1 (4).

Figure 5.

Effects of oxidation on thermal denaturation of A-I:DMPC disks. Disk samples, which were prepared as in Fig. 4, were heated and cooled from 10–98 °C at a constant rate of 11 °C/h, and the CD (A) and light scattering (B) melting data were recorded simultaneously at 222 nm to monitor changes in the protein helical content and in the particle size, respectively; the negative slopes in the light scattering data result from an optical artifact in the CD instrument and from the temperature dependence of the refractive index (47). The heating curves of non-oxidized disks (0, black lines in panels A, B) show a transition centered at 82 °C that corresponds to protein unfolding and concomitant lipoprotein fusion (38). The apparent transition temperature decreases upon oxidation; the heating curves of highly oxidized disks (4, light grey) show a similar transition centered at 60 °C followed by an additional fusion transition observed by light scattering at higher temperatures (panel B). Panel C shows CD melting data recorded at 222 nm, 11 °C/h scan rate of ox(A-I):DMPC (which were reconstituted from intact apoA-1 and then oxidized) and (oxA-I):DMPC (which were reconstituted from oxidized apoA-I); oxidant:protein ratio was 10:1.

Figure 6.

Effects of oxidation on the protein unfolding kinetics in DMPC complexes. Intact A-I:DMPC (0), ox(A-I:DMPC) (1-4), and ox(A-I):DMPC complexes (1', 2'), which were treated using HOCl:A-I molar ratios of 5:1 (1, 1'), 10:1 (2, 2'), 25:1 (3), or 50:1 (4), were subjected to T-jumps from 25 to 80 °C. The time course of the protein unfolding was monitored by CD at 222 nm. The 2, 2' (not shown) and 1, 1' data partially overlap.

Figure 7.

Effects of protein oxidation on the kinetics of DMPC clearance and on Trp modifications in apoA-I:DMPC disks. (A) DMPC multilamellar vesicles in standard buffer were incubated at 24 °C. The clearance was triggered at time $t=0$ by adding the protein; the final concentration was 20 $\mu\text{g/mL}$ apoA-I and 80 $\mu\text{g/mL}$ DMPC. ApoA-I was non-oxidized (0) or oxidized by using HOCl:apoA-I molar ratio of 5:1 (1'), 10:1 (2'), 25:1 (3'), or 50:1 (4'). (B) Trp emission spectra of normal A-I:DMPC disks (0) or of similar disks that were oxidized by using HOCl:A-I molar ratios of 5:1 (1), 10:1 (2), 25:1 (3), or 50:1 (4).

Figure 8.

Effect of oxidation on the thermal denaturation of A-I:DMPC:FC disks. Disk samples containing 10% cholesterol were prepared as described in Materials and Methods. The disks were heated and cooled from 10-98 °C at a constant rate of 11 °C/h. The protein unfolding was monitored by CD at 222 nm. Numbers correspond to those in Figs 4 and 5 and show intact disks (0) and disks that were oxidized using HOCl:apoA-I molar ratio of 5:1 (1), 10:1 (2), 25:1 (3), or 50:1 (4).