Published in final edited form as:

Biochemistry. 2008 April 1; 47(13): 4119–4128. doi:10.1021/bi702082q.

Salt Dependence of IRF1 DBD Binding to DNA Reveals Ions Are Localized Around Protein and DNA

Victoria V. Hargreaves and Robert F. Schleif*

Department of Biology, Johns Hopkins University, Baltimore, MD 21218

Abstract

The equilibrium dissociation constant of the DNA binding domain of interferon regulatory factor 1, IRF1 DBD, for its DNA binding site depends strongly on salt concentration and salt type. These dependencies are consistent with IRF1 DBD binding to DNA resulting in cation release from the DNA and both anion release from the protein and cation uptake by the protein. We demonstrated this by utilizing the fact that fluoride's release from protein upon complex formation does not contribute to the salt concentration dependence of binding and by studying mutants in which charged residues in IRF1 DBD that form salt bridges with DNA phosphates are changed to alanine. The salt concentration dependencies of the dissociation constants of wild type IRF1 DBD and the mutants R64A, D73A, K75A, and D73A K75A were measured in buffer containing NaF, NaCl, or NaBr. The salt concentration and type dependencies of the mutants relative to wild type IRF1 DBD provide evidence of charge neutralization by solution ions for R64 and by a salt bridge between D73 and K75 in buffer containing chloride or bromide salts. These data also allowed us to determine the number, type, and localization of condensed ions around both IRF1 DBD and its DNA binding site.

Counterions are condensed on DNA and reduce its charge density (1-9). These counterions are released from the DNA into the solution when charged protein residues form salt bridges with DNA phosphates. The release gives rise to the salt concentration dependence of binding, which has been widely interpreted to be proportional to the net number of ions released from the DNA (10-14).

It has been assumed (15-19) that because proteins have a low charge density compared to DNA, counterions will not condense on charged residues of proteins, and thus they will not contribute to the salt concentration dependence of binding. In many DNA binding proteins, however, basic residues are clustered at the DNA binding interface, which generates a high local positive charge density. Therefore, it seems possible that condensation of counterions on the DNA binding surface of proteins, which would generate a salt concentration dependence of binding, may occur. In fact, evidence exists suggesting counterions are concentrated in excess of coions around some proteins, and the release of these condensed counterions from the protein upon DNA binding contributes to the salt concentration dependence of binding. This evidence includes sensitivity of the equilibrium binding constant to anion identity (20-22) and non-linearity over large ranges of salt concentration of log-log plots of the binding constant versus salt concentration (22-25). In contrast to local ion effects, global ion redistribution around the protein and DNA has been proposed as a source of the salt concentration dependence of binding (26).

Here we describe the ion effects on DNA binding for the interferon regulatory factor 1 DNA binding domain, IRF1 DBD¹. Charged residues, whose positions in the protein-DNA complex

*Corresponding author Phone: 410-516-5206 Fax: 410-516-5213 schleif@jhu.edu.

allow them to form salt bridges with DNA phosphates, were mutated to alanine, and the salt concentration and type dependencies of binding of the mutant proteins were measured. The salt concentration dependencies of the K_d of the mutants relative to wild type IRF1 DBD revealed whether these charged residues are primarily neutralized by solution ions or primarily form a salt bridge with a neighboring residue of opposite charge. Using this information and the fact that the release of fluoride ions from protein does not contribute to the salt concentration dependence of binding (27), we demonstrated that ions are condensed on IRF1 DBD and its DNA binding site, and we determined the number, type, and average degree of condensation of ions around DNA and three charged residues in IRF1 DBD.

MATERIALS AND METHODS

Purification of IRF1 DBD

A pET-3a plasmid (Novagen) encoding the 113 N-terminal amino acids comprising the DNA binding domain of interferon regulatory factor 1 was provided by P. Privalov (28). The expression and purification of IRF1 DBD has been adapted from Escalante et al. (29). A single colony of cells, BL21(DE3), freshly transformed with pET3a-IRF1 DBD, was grown overnight at 37°C in 100 ml YT medium (30) containing 100 µg/ml ampicillin. The next morning the cells were washed and resuspended in 100 ml fresh YT medium and added to two 2 L baffled flasks, each containing 500 ml YT medium supplemented with 100 µg/ml ampicillin. The cells were grown at 37°C to an apparent OD_{600 nm} of ~0.6 to 0.8. At this point IPTG was added to 0.4 mM and growth was continued for four hours, after which the cells were pelleted and frozen.

The frozen cells were resuspended in three volumes of lysis buffer (50 mM HEPES, pH 7.5, 0.5 M KCl, 10% glycerol, 0.1% Triton X-100, 1 mM EDTA, 2 mM DTE, 0.2 mM PMSF) and lysed by three passages through a French press. 10 µg/mol DNase I were added after lysis. The insoluble fraction, which contains IRF1 DBD, was pelleted and solubilized in 25 ml of lysis buffer containing 6 M guanidine hydrochloride. The 25 ml IRF1 DBD sample was refolded by dialyzing against small volumes of lysis buffer such that when equilibrium was attained, the internal and external concentrations of guanidine hydrochloride were: 2 M, 1 M, 0.5 M, 0.25 M. Each dialysis step was allowed to equilibrate for three hours at 4°C. The final dialysis was against 10 mM sodium phosphate, pH 7.4, 0.3 M NaCl, 5% glycerol, 2 mM DTE. This 25-30 ml protein solution, in which IRF1 DBD was approximately 75 µM and 90% pure, was loaded onto a 5 ml HiTrap Heparin HP column (Pharmacia) pre-equilibrated with the final dialysis buffer. IRF1 DBD eluted at about 640 mM NaCl in a 60 ml gradient from 300 mM to 800 mM NaCl. Heavily overloaded samples run on SDS/14% (w/v) polyacrylamide gels showed no trace of contaminating proteins. Pure IRF1 DBD was stored at -70°C.

Concentrations of IRF1 DBD were determined using an extinction coefficient (ϵ_{280}) of 36,820 M⁻¹ cm⁻¹ as calculated from the amino acid composition by the method of Pace et al. (31). Ten grams of cells typically yielded 20-40 mg of pure IRF1 DBD.

The mutations, R64A, D73A, K75A, or D73A K75A, were introduced into the pET3a-IRF1 DBD plasmid using the QuikChange Site-directed Mutagenesis kit from Stratagene (La Jolla, CA). Plasmid DNA was isolated from cells using the Wizard® Plus Miniprep DNA purification system from Promega (Madison, WI). The mutations were confirmed by DNA sequencing, and the mutant proteins were expressed and purified as described for wild type IRF1 DBD. While WT IRF1 DBD and the R64A mutant protein eluted at 640 mM NaCl, the K75A and D73A K75A mutant proteins eluted at 555 mM NaCl, and the D73A mutant protein eluted at

¹Abbreviations: IRF1 DBD, interferon regulatory factor 1 DNA binding domain; FAM, 6-carboxyfluorescein; PRDI, positive regulatory domain I

725 mM NaCl. The proteins eluted at high concentrations from the column so salt carried over to the experimental buffer was insignificant.

Fluorescence anisotropy measurement of DNA binding

A 13 bp oligonucleotide with the sequence 5'-ACTTTCACCTTCTC-3' labeled at the 5' end with the fluorescent tag 6-carboxyfluorescein (FAM) and its unlabeled complement were purchased from Integrated DNA Technologies, Inc. The positive regulatory domain I (PRD I) site, the binding site of IRF1 DBD, from the human interferon- β enhancer is underlined. The DNA duplex was prepared by mixing the complementary oligonucleotides in equimolar amounts in 10 mM sodium phosphate, pH 7.4, 0.2 M NaCl, heating to 70°C, and cooling overnight.

To measure binding, increasing amounts of pure IRF1 DBD were mixed into a cuvette containing 2.4 ml of 5×10^{-9} M DNA in 10 mM sodium phosphate, pH 7.4, 2 mM DTE, and either NaF, KF, NaCl, KCl, NaBr, or KBr between 200 and 650 mM. Measurements were made on a T-format Photon Technology International (Lawrenceville, NJ) spectrofluorimeter with a 75-watt xenon arc lamp. The FAM label was excited at 490 nm, and its emission at 520 nm was measured. The fluorescence anisotropy was calculated from the measured fluorescence intensities, which were corrected for the G-factor. The K_d of the IRF1 DBD-DNA binding reaction was determined by least-squares fitting the curve of fluorescence anisotropy as a function of total protein concentration to the following binding equation:

$$A = A_0 + \Delta A \frac{(P_t + D_t + K_d) - \sqrt{(P_t + D_t + K_d)^2 - 4P_t \times D_t}}{2D_t}$$

where A_0 is the anisotropy of free DNA, ΔA is the change in anisotropy between the maximum anisotropy of the complex and the anisotropy of free DNA, P_t is the total protein concentration, D_t is the total DNA concentration, and K_d is the protein-DNA dissociation constant.

Measuring slope and standard error

The log of the K_d was plotted versus the log of the salt concentration, and the slope of this function was calculated for wild type IRF1 DBD in six salts and for each of the four mutants in three salts. The standard error of the data points with respect to their least squares fit was calculated using the LINEST statistical function in Microsoft Excel.

Fluorescence intensity of guanidine hydrochloride-induced unfolding

Increasing amounts of guanidine hydrochloride were mixed into a cuvette containing 2.4 ml of 1×10^{-6} M IRF1 DBD or one of the four mutants in 10 mM sodium phosphate, pH 7.4, 0.3 M NaCl, 2 mM DTE. The protein was excited at 280 nm, and the emission at 336 nm was measured. Denaturation curves were plotted of the intrinsic fluorescence normalized to that of native protein versus guanidine hydrochloride concentration and compared for WT IRF1 DBD and the four mutant proteins.

RESULTS

Salt concentration and type dependencies of the K_d for IRF1 DBD interaction with its DNA binding site

To determine the ionic contributions of IRF1 DBD binding to its DNA binding site, we monitored the equilibrium binding of the protein to a 13 bp FAM-labeled DNA duplex by fluorescence anisotropy. The fluorescence anisotropy signal increased with increasing IRF1

DBD concentration as a result of decreasing the tumbling rate of the DNA upon formation of the protein-DNA complex and fixing the fluorescent probe on the protein. From this information, we calculated the equilibrium dissociation constant of IRF1 DBD for DNA. A typical binding curve for WT IRF1 DBD in 300 mM NaCl is shown in Figure 1.

The association of protein, P , and DNA, D , to form protein-DNA complex, PD , in buffer containing salt, MX , may be represented by:

where m represents the number of cations taken up by the protein, n represents the number of anions released from the protein, and q represents the number of cations released from the DNA upon complex formation (24).

The following equation for the salt concentration dependence of binding can be derived from the power law dependence inherent in equation 1 by analogy to Record et al. (10) and Record et al. (32):

$$\partial \log K_d / \partial \log [MX] = q - m + n \quad (2)$$

Equation 2 relates the slope of the plot of $\log K_d$ versus the log of the salt concentration to the effective net number of ions released. Each of the variables in eq. 2 can be partitioned into the number of ion binding sites and the thermodynamic association of ions per site for DNA and protein as shown in the following equation:

$$\partial \log K_d / \partial \log [MX] = b\psi - c\phi + a\delta. \quad (3)$$

In eq. 3, b is the number of DNA phosphates that form salt bridges with protein residues, and ψ is the average degree of condensation or average effective concentration of cations per DNA phosphate. The number of cations released from the DNA, q , is the product of b and ψ . The number of anionic residues of IRF1 DBD that take up cations upon DNA binding is c , and the average degree of condensation of cations per site is ϕ . The number of cations taken up by the protein, m , is the product of c and ϕ . The number of cationic residues that release anions upon DNA binding is a , and the average degree of condensation of anions per site is δ . Different anions are likely to condense to different degrees due to their different sizes so we assign fluoride δ_1 , chloride δ_2 , and bromide δ_3 . The number of anions released from the protein, n , is the product of a and δ . If counterions are completely condensed on DNA and protein, ψ , ϕ , and δ will be one. If counterions are uncondensed, ψ , ϕ , and δ will be zero. In this paper, we determine ψ , ϕ , δ_2 , and δ_3 for three charged residues in IRF1 DBD and its DNA binding site.

Over the range of concentrations used, 200 to 650 mM, of the six salts studied, NaF, KF, NaCl, KCl, NaBr, KBr, the plot of $\log K_d$ versus the log of the salt concentration is linear as shown in Figure 2. Each data point in Figure 2 is derived from one binding curve, an example of which is shown in Figure 1. Aggregation of IRF1 DBD at salt concentrations below 200 mM prevented us from extending the range of salt concentration dependence studies below 200 mM. The increase in K_d with increasing salt concentration between 200 and 650 mM indicates a net release of ions, cations and/or anions, from the protein and/or DNA upon complex formation. We measured the K_d in a variety of salts to extract the relative contributions of cation and anion release to the slopes of the plots in Figure 2. When we state that an experiment was performed in buffer containing a certain salt, that is the only salt the experimental buffer contains. Table

1 shows the salt concentration dependencies of WT IRF1 DBD binding to DNA in buffers containing these salts.

Cation identity affects the K_d of IRF1 DBD for DNA, but it does not affect the salt concentration dependence of the K_d . That is, binding of IRF1 DBD to DNA is tighter in potassium salts than in sodium salts with the same anion, but the salt concentration dependencies of the K_d do not differ significantly from one another (Fig. 2, Table 1). The slopes are approximately 2.6 in potassium and sodium fluoride, 4.0 in potassium and sodium chloride, and 5.5 in potassium and sodium bromide (Table 1). These similar slopes indicate that similar net numbers of potassium and sodium ions are released upon complex formation.

In contrast to cation type, anion type affects both the magnitude of the K_d and the salt concentration dependence of the K_d (Fig. 2, Table 1). That is, when the anion type is changed, and the cation type is kept constant, the dissociation constants differ significantly (Fig. 2). Binding is tightest in fluoride salts, intermediate in chloride salts, and weakest in bromide salts. The salt concentration dependence of the K_d is also sensitive to anion type. The slope of this function is 2.60, 4.02, and 5.53 in buffer containing sodium fluoride, sodium chloride, and sodium bromide, respectively (Table 1). The different slopes demonstrate that different numbers of fluoride, chloride, and bromide ions are displaced from IRF1 DBD upon binding DNA. The fact that the release of chloride and bromide ions from IRF1 DBD contributes to the salt concentration dependence of binding demonstrates that these anions are condensed on free IRF1 DBD.

R64 releases an anion upon IRF1 DBD-DNA binding

To learn more about charge neutralization in IRF1 DBD, we examined the ionic contributions from individual basic residues. In the protein-DNA complex, arginine 64 forms a salt bridge with a DNA phosphate. The epsilon nitrogen of arginine 64 is 3.1 Å from the nearest phosphate oxygen of thymine 215 (PDB ID 1IF1). In free IRF1 DBD, solution ions must neutralize the positive charge of R64 because the protein does not contain a nearby negative residue that could neutralize R64 by forming a salt bridge. To verify that solution ions neutralize the positive charge of R64, we compared the experimentally measured differences between the salt concentration dependencies of binding for the R64A mutant and WT IRF1 DBD to the predicted number of ions that would be released if residue R64 were neutralized by solution ions. First we demonstrated the expected net number of ions are released under conditions in which there was no contribution from anion interaction with the protein. Then we used conditions in which anion release from the protein contributed to the salt concentration dependence. Thus any observed differences must be attributed to condensed anions interacting with the protein, specifically solution anions neutralizing R64.

To simplify the discussion here and in later sections, we partition the ion release and uptake from a residue of IRF1 DBD and a DNA phosphate into the sum of one or more of the five partial reactions shown in Table 2. In this table, we represent the number of ions released as positive values and the number of ions taken up as negative values. Due to the large energy of dehydration, fluoride ions interact weakly with proteins. Larger ions, like chloride and bromide, can more easily become dehydrated, so they interact with proteins stronger (33). Thus, we provide separate columns for fluoride and chloride/bromide.

Consider first the case of fluoride ions. The potential net release of ions upon binding of IRF1 DBD to DNA with respect to residue R64 is described by the sum of cases a and b, which is one ion, in buffer containing NaF (Table 2). When the mutant protein containing the R64A substitution binds DNA, case f describes the situation, and no ions are released with respect to this residue. The result in buffer containing NaF would therefore be a salt concentration dependence of binding approximately one unit smaller for the R64A mutant than for WT. As

expected, experimentally we find the salt concentration dependence of binding is 2.60 ± 0.23 for WT IRF1 DBD and is 1.31 ± 0.14 for R64A in buffer containing NaF (Fig. 3, Table 1). The difference of approximately one in the slopes of the WT protein and the mutant protein must arise from the R64A mutant displacing one less cation from the DNA phosphates compared to WT IRF1 DBD. This difference is also consistent with residue R64 being neutralized by a fluoride ion in the free protein.

Now we consider the cases of chloride and bromide ions. Chloride and bromide ions interact more strongly with protein than fluoride ions interact with protein (33), and the release of these ions from proteins upon DNA binding contributes to the salt concentration dependence of binding (20,21,34). If residue R64 of WT IRF1 DBD is neutralized by chloride, then when the protein binds DNA, the net release of ions with respect to R64 is described by the sum of cases a and b in the M^+ and Cl^- columns of Table 2, or a net of two ions. The R64A mutant protein's net ion release with respect to residue R64 would correspond to case f in Table 2, which is zero. Experimentally we found this to be true; the salt concentration dependence of the K_d is smaller for R64A, 2.37 ± 0.10 , than for WT IRF1 DBD, 4.02 ± 0.19 , in buffer containing NaCl (Fig. 3, Table 1). The difference, 1.6, indicates not only that a sodium ion is condensed on the DNA phosphate, which forms a salt bridge with R64 in the complex, but also that a chloride ion must be condensed on residue R64 in free IRF1 DBD. Similar results were observed in buffer containing NaBr. The salt concentration dependence of the K_d is 5.53 ± 0.38 for WT IRF1 DBD and 3.36 ± 0.36 for the R64A mutant in buffer containing NaBr. Therefore the positive charge of R64 in IRF1 DBD is neutralized by a solution ion in buffer containing NaCl or NaBr.

K75 and D73 are neutralized by ions in fluoride-containing buffer but not in chloride- or bromide-containing buffers

In the IRF1 DBD-DNA complex, lysine 75 forms a salt bridge with a DNA phosphate. The epsilon amino group of lysine 75 is $2.7 \text{ \AA}$ from the nearest phosphate oxygen of adenine 204. Unlike the environment of R64, however, a negatively charged residue, D73, lies near K75. Although a dehydrated salt bridge is usually approximately three Angstroms (35), the charged groups of amino acids D73 and K75 are approximately seven Angstroms apart in the co-crystal structure of IRF1 DBD with DNA (36). It is possible that in the absence of DNA, D73 and K75 possess substantial salt bridge character. To determine how K75 is neutralized, we mutated this residue to an alanine and again measured the salt concentration and type dependencies of binding. We also made the single mutant D73A and the double mutant D73A K75A to further test the hypothesis that D73 and K75 form a salt bridge in the free protein.

Table 3 presents the five situations being considered here and their breakdown into the partial reactions from Table 2, with the resulting prediction of the net ion release for each of the cases in both fluoride buffer and chloride or bromide buffer. Experimentally, the data in fluoride buffers indicate D73 and K75 are neutralized by solution ions. The dependence of K_d on salt concentration in NaF for the D73A mutant is 2.95 ± 0.23 (Fig. 4a), which is approximately equal to that of WT, 2.60 ± 0.23 (Fig. 2), as seen in Table 1. The slopes of these plots for the K75A (Fig. 4b) and D73A K75A (Fig. 4c) mutants in NaF, 1.55 ± 0.11 and 1.17 ± 0.20 respectively, were similar to each other and smaller than those of WT and the D73A mutant, demonstrating that residue K75 is neutralized by a solution ion in buffer containing NaF.

Experimentally, the salt bridge case was observed for residues D73 and K75 in buffer containing chloride or bromide salts. In the presence of NaCl, the slopes are 4.02 ± 0.19 for WT IRF1 DBD (Fig. 2), 4.20 ± 0.16 for the K75A mutant (Fig. 4b), and 4.29 ± 0.13 for the D73A K75A mutant (Fig. 4c). These slopes are approximately equal and less than the slope of the D73A mutant, 6.21 ± 0.51 (Fig. 4a, Table 1). These results demonstrate that residue K75 forms a salt bridge with residue D73 in buffer containing chloride salt (Table 3). Similar data

were collected in bromide salt. In the presence of NaBr, the slopes are 5.53 ± 0.38 for WT IRF1 DBD (Fig. 2), 5.16 ± 0.29 for the K75A mutant (Fig. 4b), and 5.13 ± 0.22 for the D73A K75A mutant (Fig. 4c). These slopes are approximately equal to each other and less than the slope of the D73A mutant, 8.12 ± 0.41 (Fig. 4a, Table 1, Table 3). Therefore, residues D73 and K75 form a salt bridge in chloride and bromide salts.

The mutant proteins are folded

To ensure the mutations do not destabilize the proteins, which could potentially create additional sources of ion effects, we measured the fluorescence intensity upon guanidine hydrochloride-induced unfolding (Fig. 5). The proteins unfolded in a similar fashion, and their calculated Gibbs free energies of unfolding are similar (data not shown). Additional evidence that the mutant proteins are folded in a conformation similar to WT IRF1 DBD is evidenced by their ability to bind DNA.

Global analysis of ion effects

The experiments and analysis described above have shown that for residues 64, 73, and 75, the release of ions from IRF1 DBD and DNA and the uptake of ions by IRF1 DBD contributes to the salt concentration dependence of the protein's binding to DNA. The fact that such effects were observed requires that the degree of condensation be appreciable. In this section, we will estimate the magnitude of the condensation by a global analysis of the binding data.

By analogy to equation 3, the following system of equations describe the salt concentration dependence of binding for WT IRF1 DBD and the mutant proteins R64A, D73A, K75A, and D73A K75A in the presence of NaF, NaCl, or NaBr. The analysis is global because the left-hand sides of the equations express the slopes in terms of the residues, phosphates, mode of neutralization, and degree of condensation. The right-hand sides of the equations are the experimentally determined values of the salt concentration dependence of binding taken from Table 1.

$$\text{WT NaF: } b\psi - (c-1)\phi + a\delta_1 = 2.60 \pm 0.23 \quad (4)$$

$$\text{WT NaCl: } b\psi - c\phi + (a-1)\delta_2 = 4.02 \pm 0.20 \quad (5)$$

$$\text{WT NaBr: } b\psi - c\phi + (a-1)\delta_3 = 5.53 \pm 0.38 \quad (6)$$

$$\text{R64A NaF: } (b-1)\psi - (c-1)\phi + (a-1)\delta_1 = 1.31 \pm 0.14 \quad (7)$$

$$\text{R64A NaCl: } (b-1)\psi - c\phi + (a-2)\delta_2 = 2.37 \pm 0.10 \quad (8)$$

$$\text{R64A NaBr: } (b-1)\psi - c\phi + (a-2)\delta_3 = 3.36 \pm 0.36 \quad (9)$$

$$\text{D73A NaF: } b\psi - (c-1)\phi + a\delta_1 = 2.95 \pm 0.32 \quad (10)$$

$$\text{D73A NaCl: } b\psi - (c-1)\phi + a\delta_2 = 6.21 \pm 0.51 \quad (11)$$

$$\text{D73A NaBr: } b\psi - (c-1)\phi + a\delta_3 = 8.12 \pm 0.41 \quad (12)$$

$$\text{K75A NaF: } (b-1)\psi - (c-1)\phi + (a-1)\delta_1 = 1.55 \pm 0.11 \quad (13)$$

$$\text{K75A NaCl: } (b-1)\psi - (c-1)\phi + (a-1)\delta_2 = 4.20 \pm 0.16 \quad (14)$$

$$\text{K75A NaBr: } (b-1)\psi - (c-1)\phi + (a-1)\delta_3 = 5.16 \pm 0.29 \quad (15)$$

$$\text{D73A K75A NaF: } (b-1)\psi - (c-1)\phi + (a-1)\delta_1 = 1.17 \pm 0.20 \quad (16)$$

$$\text{D73A K75A NaCl: } (b-1)\psi - (c-1)\phi + (a-1)\delta_2 = 4.29 \pm 0.13 \quad (17)$$

$$\text{D73A K75A NaBr: } (b-1)\psi - (c-1)\phi + (a-1)\delta_3 = 5.13 \pm 0.22 \quad (18)$$

Using information from the co-crystal structure of the IRF1 DBD-DNA complex (36), we solved the system of equations for the following unknowns: ψ , the average degree of condensation of cations per DNA phosphate; ϕ , the average degree of condensation of cations per acidic residue, which, in the free protein, forms a salt bridge with a basic residue that forms a salt bridge with a DNA phosphate in the complex; and δ_2 and δ_3 , the average degrees of condensation of chloride and bromide ions respectively per basic residue of IRF1 DBD that forms a salt bridge with a DNA phosphate.

According to the structure, five positively charged residues ($a = 5$) of IRF1 DBD, K43, R64, K75, K78, and R99, form salt bridges with four DNA phosphates ($b = 4$). Residues R99 and K43 contact the same phosphate. It appears from the crystal structure that only residue K75 is capable of forming a salt bridge ($c = 1$) with a negatively charged residue, D73, in the free protein.

Due to the high energy of dehydration, fluoride ions do not interact strongly with protein (33). To simplify our analysis, we assume δ_1 is zero, as has been done previously (20-22,34). Thus subtracting equation 10 from 11 gives $a\delta_2 = 3.2$, and hence δ_2 is 0.64. From the standard errors of the slopes, the error of δ_2 is 0.16. Similar analysis of equations 12 and 10 gives $\delta_3 = 1.02 \pm 0.14$. Substituting these values into equation 4, ψ is calculated to be 0.65 with an error of 0.05. Equation 5 then gives $\phi = 1.16 \pm 0.64$.

The number of cations released from the DNA, q , is approximately 2.6. The number of cations taken up by IRF1 DBD, m , is approximately 1.2. The number of anions released from the protein, n , is approximately 3.2 for chloride. The number of bromides released from IRF1 DBD is approximately 5.1.

We solved for δ_2 and δ_3 using the equations describing the salt concentration dependence of binding of the mutant D73A, equations 10, 11, and 12. The equations describing the salt concentration dependence of binding of the mutant K75A, equations 13, 14, and 15, or those of the mutant D73A K75A, equations 16, 17, and 18, could have been used instead. Using equations 13, 14, and 15, δ_2 is 0.65 ± 0.075 , and δ_3 is 0.9 ± 0.1 . Using equations 16, 17, and 18, δ_2 is 0.78 ± 0.075 , and δ_3 is 0.98 ± 0.1 . Therefore, the values of the unknowns δ_2 and δ_3 do not depend strongly on the equations used to solve for them. This is also true for the determination of ψ and ϕ . When these estimated values for the unknowns are substituted into equations 4 through 18, close agreement of the left-hand terms of the equations with the experimentally determined values on the right-hand side lends reliability to the solutions obtained.

DISCUSSION

In this work we describe the binding of IRF1 DBD to its DNA target sequence over a range of salt concentrations for six salts, NaF, KF, NaCl, KCl, NaBr, and KBr. We mutated charged residues, which form salt bridges with DNA phosphates according to the crystal structure of the IRF1 DBD-DNA complex. The mutations alter the measured stoichiometry of ion release, providing evidence of ion condensation on IRF1 DBD and its DNA binding site. The salt concentration and type dependencies of the K_d of the mutants relative to WT IRF1 DBD also indicate how certain charged residues are neutralized in the free protein, demonstrating whether it is by solution ions or by a salt bridge with a neighboring residue. Using this information, we determined the number, type, and degree of condensation of ions around DNA phosphates and charged protein residues.

Sodium and potassium affect IRF1 DBD-DNA equilibrium

The salt concentration dependencies of IRF1 DBD binding to DNA are similar in the presence of sodium and potassium salts with the same anion (Fig. 2, Table 1). According to eq. 2, this result means that similar net numbers of sodium and potassium ions are released during complex formation. Others have also reported that similar numbers of sodium and potassium ions are released upon DNA binding for Ku (21), single-stranded binding protein (34), and engrailed (20).

While cation type does not significantly affect the salt concentration dependence of the K_d of IRF1 DBD for DNA, cation type does affect the magnitude of the K_d . Binding is tighter in potassium salts than in sodium salts with the same anion (Fig. 2). This difference may be due to cation uptake by the protein and/or cation release from the DNA. Because cations can interact with both protein and DNA, we cannot discern the relative contributions of the interactions responsible for the cation type effect on the K_d . Experiments using X-ray crystallography (37), molecular dynamics simulations (38), and NMR (39) have demonstrated differences in the location but not the number of Na^+ and K^+ ions around DNA, which are consistent with our data that indicate that cation type affects the magnitude of the K_d but not the salt concentration dependence of the K_d . Differences in the location and affinity of cations for protein surfaces have also been observed using molecular dynamics (40,41) and X-ray crystallography (42).

Cation localization around DNA

To determine the degree of condensation or effective concentration of cations around DNA phosphates, Record and colleagues measured the salt concentration dependence of binding of polylysine peptides of varying length to DNA (10). Assuming there was no release of anions from the polylysine peptide upon its binding DNA, Record and colleagues determined the localization of cations around the phosphates of native DNA to be 0.88 (10). Since then many salt concentration dependence studies (11,13-18,21,24,25,34) have used this value for ψ . More recently, grand canonical Monte Carlo simulations were used to predict the local ion effects on binding of an oligocation with eight positive charges to DNA oligomers of varying length (43). The degree of localization or condensation that is measured depends strongly on the DNA length.

The work reported here appears to be the first measurement of the localization of cations around DNA phosphates from the local ion effects on binding of a globular protein without assuming that anions are not released from the protein. We calculate ψ is 0.65 ± 0.05 .

Instead of local ion effects from the neutralization of phosphates, global ion redistribution has been proposed to give rise to the salt concentration dependence of protein-DNA interactions (26). That is, in a protein-DNA complex, the large negative electrostatic potential near the DNA excludes anions from the protein's surface, and the protein's presence excludes cations near the DNA (26). In support of this proposal, Misra and colleagues compared the reported salt concentration dependence of λ cI repressor binding and EcoRI binding with their non-linear Poisson Boltzmann calculations of ion concentrations around the two proteins.

Misra et al. (26) suggest residue E34K in λ cI repressor does not make direct contacts to DNA and hence the difference between the salt concentration dependence of binding of this mutant λ cI repressor and wild type repressor can not be explained by local ion effects. They also propose that the difference between the salt concentration dependence of binding of a deletion mutant of the first 12 residues of the N-terminal region of EcoRI endonuclease relative to that of wild type can not be explained by local ion effects. The papers cited by Misra et al. do not, however, support their proposals. Based on structural evidence, Nelson and Sauer (44) reported that residue E34K in λ cI repressor forms a salt bridge with DNA, and this is consistent with an increased salt concentration dependence of the mutant relative to that of wild type λ cI repressor. Similarly, Jen-Jacobson et al. (45) reported that there is unpublished structural evidence that Arg9 in the N-terminal region of EcoRI endonuclease makes a direct contact to the DNA, which is consistent with a decreased salt concentration dependence of a deletion mutant of the N-terminal region relative to that of the wild type protein.

Our data appear to be well described in terms of local ion effects. For example, the D73A mutant takes up one less sodium ion and releases one more chloride ion relative to WT IRF1 DBD, which explains why the salt concentration dependence of binding of the D73A mutant protein is approximately two units more than that of WT IRF1 DBD. Local ion effects can also simply explain why the salt concentration dependencies for WT IRF1 DBD and the mutant proteins K75A and D73A K75A in buffer containing NaCl are approximately equal. That is, the contribution to the salt concentration dependence from the cation displaced from the phosphate that interacts with K75 in WT IRF1 DBD is compensated for by the cation taken up by D73, and ion interaction with residues 73 and 75 does not contribute to the salt concentration dependence in mutant proteins D73A or D73A K75A.

Anions interact specifically with IRF1 DBD

The salt concentration dependencies of binding differ significantly when the anion is changed (Fig. 2, Table 1). This indicates that when the anion identity is changed, different numbers of

solution anions are released from WT IRF1 DBD upon its binding to DNA. This effect must arise from anion interaction with the protein because anions do not interact with DNA. Evidence for anion release from other DNA binding proteins has been observed previously; specifically, the equilibrium binding constants of single-strand binding protein (34), Ku (21), engrailed (20), and T4 gene 32 protein (22) are sensitive to anion identity.

In addition to affecting the salt concentration dependence of the K_d of IRF1 DBD for DNA, anion type also affects the magnitude of the K_d . DNA binding is tightest in fluoride salts, weaker in chloride salts, and weakest in bromide salts for WT IRF1 DBD (Fig. 2). This order of anions (F^- , Cl^- , Br^-) follows the Hofmeister series and probably reflects the relative strengths of interactions of the ions with IRF1 DBD. Small ions, like fluoride, have organized, tightly held water shells, whereas larger ions, like bromide, hold water less tightly (46). As a result, fluoride interacts weakly with protein, and thus its release from protein does not contribute to the salt concentration dependence of binding (27). Tight binding of IRF1 DBD to DNA in buffer containing fluoride results from a weak interaction between fluoride and the protein. It requires less energy to dehydrate a larger ion, like bromide, therefore it can interact more strongly with protein, and its release from the protein contributes to the salt concentration dependence of binding. That is, the weak binding of IRF1 DBD to DNA in buffer containing bromide results from a strong interaction between bromide and the protein. Our calculated degrees of condensation of chloride and bromide ions per basic residue of IRF1 DBD that forms a salt bridge with a DNA phosphate, 0.64 ± 0.16 and 1.02 ± 0.14 respectively, are consistent with the anion order of the Hofmeister series. The fact that these numbers are less than and close to one also lends credibility to their values.

IRF1 DBD also takes up cations upon binding DNA

The salt concentration dependence of the K_d of the D73A mutant for DNA relative to that of WT IRF1 DBD is consistent with cation uptake by the protein. We predict and provide evidence that D73 and K75 form a salt bridge in IRF1 DBD in buffer containing chloride or bromide salt. The D73A mutant has a greater salt concentration dependence of binding relative to WT IRF1 DBD in buffer containing chloride or bromide salt (Table 1). This indicates that the mutant protein takes up fewer ions than WT IRF1 DBD. An increase in the slope by approximately two in buffer containing NaCl of the D73A mutant relative to WT indicates that one less sodium ion is taken up and one more chloride ion is released from the mutant protein. Hence, we can conclude that in WT IRF1 DBD residue D73 takes up a sodium ion in buffer containing chloride or bromide salt. Data consistent with cation uptake by other DNA binding proteins, non-linearity over large ranges of salt concentration of log-log plots of the binding constant versus salt concentration, have been published for CAP (24), single-strand binding protein (34), papillomavirus E2 proteins BPV-E2/D and HPV16-E2/D (25), AraC (23), and T4 gene 32 protein (22).

Through the use of mutant proteins and fluoride-containing buffers, we were able to calculate the effective concentration of cations around negatively charged residues of IRF1 DBD, ϕ , is 1.16 ± 0.64 . The fact that the most probable value of ϕ is greater than one likely results from the relatively high value of its error.

Aggregation at low ionic strength

In preliminary experiments we observed that the apparent strength of binding of IRF1 DBD to its DNA binding site increased as the salt concentration increased from 25 mM to 200 mM (data not shown). This anomalous behavior, unusually high anisotropy values, and ultracentrifugation data (data not shown) are all consistent with aggregation of IRF1 DBD at protein concentrations above 1 μ M and salt concentrations below 200 mM. The binding strength decreased as the salt concentration increased above 200 mM (Table 1), as predicted

by the polyelectrolyte effect. Therefore, in this paper, we present data collected at salt concentrations only at and above 200 mM. Non-linearity in the salt concentration dependence of binding over large ranges of salt concentration has been observed previously and accounted for by ion association with the protein (23,24). Protein aggregation at low salt concentration is a potential alternative explanation of non-linearity in the salt concentration dependence of binding in such experiments, and data suggest that this is the case in this study.

The role of salt bridges in IRF1 DBD

Residues D73 and K75 form a salt bridge in buffer containing chloride or bromide salt but not in buffer containing fluoride salt. It is unclear why. The fact that fluoride does not interact strongly with proteins can not explain this difference because our experiments are capable of detecting the difference in the salt concentration dependencies of binding for the salt bridge case and the counterion case in buffer containing NaF. A possible explanation for the different forms of neutralization in NaF and NaCl is that it is energetically more costly to condense two ions, Na^+ and Cl^- , on residues D73 and K75 in the free protein than it is to form a salt bridge between the two residues. Alternatively, perhaps in buffer containing NaCl or NaBr, K75 is in a conformation that positions it close to D73 and allows the formation of a salt bridge, but in buffer containing NaF, K75 and D73 are not situated to form a salt bridge so instead these residues are neutralized by counterions. In principle, this hypothesis could be tested by a denaturation experiment in fluoride containing buffer. A different conformation of residues D73 and K75 in different salts supports a global ion redistribution model, but our data are more easily explained by local ion effects.

Our salt concentration dependence of binding data, which are measured between 200 and 650 mM, indicate that D73 and K75 form a salt bridge in free IRF1 DBD in buffer containing chloride or bromide salt (Table 1). This finding opposes the fact that some salt bridges in proteins persist at low salt concentrations (< 0.1 M) and tend to be disrupted at high salt concentrations (47).

According to structural information (35,48), several other DNA binding proteins such as integration host factor, single-strand binding protein, lac repressor, DNA gyrase, the nucleosomal proteins, RNA polymerase core protein, c-Myb, and HU contain cationic residues that participate in salt bridges in the free protein and interact with DNA phosphates in the complex. If two residues have formed a salt bridge, an anion would not be displaced from the positively charged residue upon DNA binding. Also, the negatively charged residue would take up a cation when the positively charged residue interacts with the DNA. Releasing fewer ions and taking up more ions reduces the protein's salt concentration dependence of binding. Mechanisms that reduce the salt concentration dependence of binding are important for a cell to avoid large fluctuations in gene regulation in response to changes in cellular conditions (23,49,50).

In conclusion, the studies presented in this work demonstrate ions condense on free IRF1 DBD in addition to DNA. We measured the number, type, and degree of condensation of ions around this protein and DNA. Upon IRF1 DBD-DNA complex formation, between two and three cations are released from the DNA binding site, one cation is taken up by IRF1 DBD, and three chloride or five bromide ions are released from IRF1 DBD. Specifically, residue D73 takes up a cation, and residues R64 and K75 release anions upon IRF1 DBD binding to DNA in buffer containing fluoride salt. The salt concentration dependencies of four mutant proteins relative to that of the WT protein indicate that in free IRF1 DBD, residue R64 is neutralized by a solution ion and residues D73 and K75 form a salt bridge in buffer containing chloride or bromide salt. We calculated the average degree of condensation of cations per DNA phosphate is 0.65 ± 0.05 ; the average degree of condensation of cations per acidic residue is 1.16 ± 0.64 ; and the average degrees of condensation of chloride and bromide ions per basic residue are

0.64 ± 0.16 and 1.02 ± 0.14 , respectively. We conjecture that, like IRF1 DBD, other DNA binding proteins have a mixture of basic residues that form salt bridges and basic residues that are neutralized by solution ions. We also predict that the degree of condensation of ions around acidic and basic residues in other DNA binding proteins and DNA phosphates will be similar to those we measured for IRF1 DBD and its DNA binding site.

Acknowledgments

The authors thank Dr. David Draper for helpful discussions and the members of Schleif lab, Dr. Michael Rodgers, Katherine Frato, Jennifer Seedorff, and Stephanie Dirla, for insightful comments on the manuscript.

REFERENCES

1. Manning GS. Limiting laws and counterion condensation in polyelectrolyte solutions I. Colligative properties. *J. Chem. Phys* 1969a;51:924–933.
2. Manning GS. Limiting laws and counterion condensation in polyelectrolyte solutions II. Self-diffusion of the small ions. *J. Chem. Phys* 1969b;51:934–938.
3. Manning GS. Limiting laws and counterion condensation in polyelectrolyte solutions III. An analysis based on the Mayer ionic solution theory. *J. Chem. Phys* 1969c;51:3249–3252.
4. Manning GS. Limiting laws and counterion condensation in polyelectrolyte solutions IV. The approach to the limit and the extraordinary stability of the charge fraction. *Biophys. Chem* 1977b;7:95–102. [PubMed: 901909]
5. Manning GS. The molecular theory of polyelectrolyte solutions with applications to the electrostatic properties of polynucleotides. *Q. Rev. Biophys* 1978a;11:179–246. [PubMed: 353876]
6. Le Bret M, Zimm BH. Monte Carlo determination of the distribution of ions about a cylindrical polyelectrolyte. *Biopolymers* 1984;23:287–312.
7. Wilson RW, Bloomfield VA. Counterion-induced condensation of deoxyribonucleic acid. A light-scattering study. *Biochemistry* 1979;18:2192–2196. [PubMed: 444448]
8. Ponomarev SY, Thayer KM, Beveridge DL. Ion motions in molecular dynamics simulations on DNA. *Proc. Natl. Acad. Sci. U.S.A* 2004;101:14771–14775. [PubMed: 15465909]
9. Keyser UF, Koeleman BN, Van Dorp S, Krapf D, Smeets RMM, Lemay SG, Dekker NH, Dekker C. Direct force measurements on DNA in a solid-state nanopore. *Nature Physics* 2006;2:473–477.
10. Record MT Jr, Lohman TM, deHaseth PH. Ion effects on ligand-nucleic acid interactions. *J. Mol. Biol* 1976a;107:145–158. [PubMed: 1003464]
11. Record MT Jr, deHaseth PL, Lohman TM. Interpretation of monovalent and divalent cation effects on the *lac* repressor-operator interaction. *Biochemistry* 1977;16:4791–4796. [PubMed: 911790]
12. Record MT Jr, Anderson CF, Lohman TM. Thermodynamic analysis of ion effects on the binding and conformational equilibria of proteins and nucleic acids: the roles of ion association or release, screening, and ion effects on water activity. *Q. Rev. Biophys* 1978;11:103–178. [PubMed: 353875]
13. deHaseth PL, Lohman TM, Record TM Jr. Nonspecific interaction of *lac* repressor with DNA: an association reaction driven by counterion release. *Biochemistry* 1977a;16:4783–4791. [PubMed: 911789]
14. Barkley MD, Lewis PA, Sullivan GE. Ion effects on the *lac* repressor-operator equilibrium. *Biochemistry* 1981;20:3842–3851. [PubMed: 7272280]
15. Moraitis MI, Xu H, Matthews KS. Ion concentration and temperature dependence of DNA binding: comparison of PurR and LacI repressor proteins. *Biochemistry* 2001;40:8109–8117.
16. Senear DF, Batey R. Comparison of operator-specific and nonspecific DNA binding of the λ cI repressor: [KCl] and pH effects. *Biochemistry* 1991;30:6677–6688. [PubMed: 1829636]
17. Barkley MD. Salt dependence of the kinetics of the *lac* repressor-operator interaction: role of nonoperator deoxyribonucleic acid in the association reaction. *Biochemistry* 1981;20:3833–3842. [PubMed: 7023537]
18. Whitson PA, Olson JS, Matthews KS. Thermodynamic analysis of the lactose repressor-operator DNA interaction. *Biochemistry* 1986;25:3852–3858. [PubMed: 3527258]

19. Manning GS. Is a small number of charge neutralizations sufficient to bend nucleosome core DNA onto its superhelical ramp? *J. Am. Chem. Soc* 2003;125:15087–15092. [PubMed: 14653743]
20. Dragan AI, Li Z, Makeyeva EN, Milgotina EI, Liu Y, Crane-Robinson C, Privalov PL. Forces driving the binding of homeodomains to DNA. *Biochemistry* 2006;45:141–151. [PubMed: 16388589]
21. Arosio D, Costantini TK, Vindigni A. Fluorescence anisotropy studies on the Ku-DNA interaction. *J. Biol. Chem* 2004;279:42826–42835. [PubMed: 15284231]
22. Rouzina I, Pant K, Karpel RL, Williams MC. Theory of electrostatically regulated binding of T4 gene 32 protein to single- and double-stranded DNA. *Biophys. J* 2005;89:1941–1956. [PubMed: 15994897]
23. Martin KJ, Schleif RF. Equilibrium DNA-binding of AraC protein. Compensation for displaced ions. *J. Mol. Biol* 1987;195:741–744. [PubMed: 3656430]
24. Stickle DF, Fried MG. Cation binding linked to a sequence-specific CAP-DNA interaction. *Biophys. Chem* 2007;126:106–116. [PubMed: 16782261]
25. Blakaj DM, Kattamuri C, Khrapunov S, Hegde RS, Brenowitz M. Indirect readout of DNA sequence by Papillomavirus E2 proteins depends upon net cation uptake. *J. Mol. Biol* 2006;358:224–240. [PubMed: 16513133]
26. Misra VK, Hecht JL, Sharp KA, Friedman RA, Honig B. Salt effects on protein-DNA interactions. The λ CI repressor and EcoRI endonuclease. *J. Mol. Biol* 1994;238:264–280. [PubMed: 8158653]
27. von Hippel, PH.; Schleif, T. *Biological Macromolecules*. Timasheff, SN.; Fasman, G., editors. Vol. 2. Marcel Dekker, Inc.; New York: 1969a. p. 417-574.
28. Hargreaves VV, Makeyeva EN, Dragan AI, Privalov PL. Stability and DNA binding ability of the DNA binding domains of interferon regulatory factors 1 and 3. *Biochemistry* 2005;44:14202–14209. [PubMed: 16245936]
29. Escalante CR, Yie J, Thanos D, Aggarwal AK. Expression, purification, and co-crystallization of IRF-1 bound to the interferon- β element PRD I. *FEBS Letts* 1997;414:219–220. [PubMed: 9315689]
30. Schleif, RF.; Wensink, P. *Practical Methods in Molecular Biology*. Springer-Verlag; New York: 1981.
31. Pace CN, Vajdos F, Fee L, Grimsley G, Gray T. How to measure and predict the molar absorption coefficient of a protein. *Protein Sci* 1995;4:2411–2423. [PubMed: 8563639]
32. Record MT Jr, Ha J-H, Fisher M. Analysis of equilibrium and kinetic measurements to determine thermodynamic origins of stability and specificity and mechanism of formation of site-specific complexes between proteins and helical DNA. *Methods in Enzymology* 1991;208:291–343. [PubMed: 1779839]
33. von Hippel PH, Schleif T. Ion effects on the solution structure of biological macromolecules. *Accts. Chem. Res* 1969b;2:257–265.
34. Overman LB, Bujalowski W, Lohman TM. Equilibrium binding of *Escherichia coli* single-strand binding protein to single-stranded nucleic acids in the (SSB)₆₅ binding mode. Cation and anion effects and polynucleotide specificity. *Biochemistry* 1988;27:456–471. [PubMed: 3280021]
35. Saecker RM, Record MT Jr. Protein surface salt bridges and paths for DNA wrapping. *Curr. Opin. Struct. Biol* 2002;12:311–319. [PubMed: 12127449]
36. Escalante CR, Yie J, Thanos D, Aggarwal AK. Structure of IRF-1 with bound DNA reveals determinants of interferon regulation. *Nature* 1998;391:103–106. [PubMed: 9422515]
37. Schneider B, Kabelac M. Stereochemistry of binding of metal cations and water to a phosphate group. *J. Am. Chem. Soc* 1998;120:161–165.
38. Cheng Y, Korolev N, Nordenskiöld L. Similarities and differences in interaction of K⁺ and Na⁺ with condensed ordered DNA. A molecular dynamics computer simulation study. *Nucleic Acid Res* 2006;34:686–696. [PubMed: 16449204]
39. Bleam ML, Anderson CF, Record MT Jr. Relative binding affinities of monovalent cations for double-stranded DNA. *Proc. Natl. Acad. Sci. U.S.A* 1980;77:3085–3089. [PubMed: 16592827]
40. Vrbka L, Jungwirth P, Baudin P, Touraud D, Kunz W. Specific ion effects at protein surfaces: a molecular dynamics study of bovine pancreatic trypsin inhibitor and horseradish peroxidase in selected salt solutions. *J. Phys. Chem. B* 2006a;110:7036–7043. [PubMed: 16571019]

41. Vrbka L, Vondrasek J, Jagoda-Cwiklik B, Vacha R, Jungwirth P. Quantification and rationalization of the higher affinity of sodium over potassium to protein surfaces. *Proc. Natl. Acad. Sci. U. S. A* 2006b;103:15440–15444. [PubMed: 17032760]
42. Chakrabarti P. Systematics in the interaction of metal ions with the main-chain carbonyl group in protein structures. *Biochemistry* 1990;29:651–658. [PubMed: 2337588]
43. Olmstead MC, Bond JP, Anderson CF, Record MT Jr. Grand canonical Monte Carlo molecular and thermodynamic predictions of ion effects on binding of an oligocation (L^{8+}) to the center of DNA oligomers. *Biophys. J* 1995;68:634–647. [PubMed: 7696515]
44. Nelson HC, Sauer RT. Lambda repressor mutations that increase the affinity and specificity of operator binding. *Cell* 1985;42:549–558. [PubMed: 3161621]
45. Jen-Jacobson L, Lesser D, Kurpiewski M. The enfolding arms of EcoRI endonuclease: role in DNA binding and cleavage. *Cell* 1986;45:619–629. [PubMed: 3011275]
46. Collins KD. Sticky ions in biological systems. *Proc. Natl. Acad. Sci. U.S.A* 1995;92:5553–5557. [PubMed: 7539920]
47. Marqusee S, Baldwin RL. Helix stabilization by $Glu^- \dots Lys^+$ salt bridges in short peptides of *de novo* design. *Proc. Natl. Acad. Sci. U.S.A* 1987;84:8898–8902. [PubMed: 3122208]
48. Holbrook JA, Tsodikov OV, Saecker RM, Record MT Jr. Specific and non-specific interactions of integration host factor with DNA: thermodynamic evidence for disruption of multiple IHF surface salt-bridges coupled to DNA binding. *J. Mol. Biol* 2001;310:379–401. [PubMed: 11428896]
49. Fried MG, Stickle DF. Ion-exchange reactions of proteins during DNA binding. *Eur. J. Biochem* 1993;218:469–475. [PubMed: 8269936]
50. Record MT Jr, Anderson CF, Mills P, Mossing M, Roe J. Ions as regulators of protein-nucleic acid interactions *in vitro* and *in vivo*. *Adv. Biophys* 1985;20:109–135. [PubMed: 3914831]

Figure 1.

A representative binding curve of WT IRF1 DBD binding to a FAM-labeled 13 base pair oligonucleotide at 300 mM NaCl. Fluorescence anisotropy is shown as a function of protein concentration. The fluorescent label was excited at 490 nm, and its emission was measured at 520 nm. The solid curve is the non-linear least squares fit of the binding equation to the data.

Figure 2. The dependence of $\log K_d$ of WT IRF1 DBD on the logarithm of the NaF, KF, NaCl, KCl, NaBr, and KBr concentrations. The K_d values were obtained from individual binding experiments.

Figure 3. The dependence of log K_d of the R64A mutant protein on the logarithm of the NaF, NaCl, and NaBr concentrations. The K_d values were obtained from individual binding experiments.

4a

4b

4c

Figure 4.

The dependence of log K_d of D73A (a), K75A (b), or D73A K75A (c) mutant protein on the logarithm of the NaF, NaCl, and NaBr concentrations. The K_d values were obtained from individual binding experiments.

Figure 5.

The dependence of the intrinsic fluorescence intensity, normalized to that of the native protein, on the concentration of GdnHCl for WT IRF1 DBD and the mutant proteins R64A, D73A, K75A, and D73A K75A. The excitation wavelength was 280 nm, and the emission at 336 nm was measured.

Table 1

Salt dependence of binding of WT and mutant IRF1 DBD

Protein	Salt	Slope	Std. Error
WT	NaF	2.60	0.23
	KF	2.65	0.20
	NaCl	4.02	0.19
	KCl	3.95	0.19
	NaBr	5.53	0.38
	KBr	5.51	0.57
R64A	NaF	1.31	0.14
	NaCl	2.37	0.10
	NaBr	3.36	0.36
D73A	NaF	2.95	0.32
	NaCl	6.21	0.51
	NaBr	8.12	0.41
K75A	NaF	1.55	0.11
	NaCl	4.20	0.16
	NaBr	5.16	0.29
D73AK75A	NaF	1.17	0.20
	NaCl	4.29	0.13
	NaBr	5.13	0.22

Table 2
Net numbers of ions released by different structural elements when IRF1 DBD binds DNA

Case	Element	Status when free	Status in complex	Potential number of ions released from element		
				M ⁺ ^a	F ^{-b}	Cl ⁻ or Br ^{-c}
a	DNA phosphate	Neutralized by solution ions	Forms a salt bridge with a basic residue	+1	0	0
b	Basic residue	Neutralized by solution ions	Forms a salt bridge with a DNA phosphate	0	0	+1
c	Acidic residue	Neutralized by solution ions	Neutralized by solution ions	0	0	0
d	Basic residue	Neutralized by a salt bridge with acidic residue	Forms a salt bridge with a DNA phosphate	0	0	0
e	Acidic residue	Neutralized by a salt bridge with basic residue	Neutralized by solution ions	-1	0	0
f	Neutral residue	-	-	0	0	0

^aWhere M⁺ is Na⁺ or K⁺.
^bIf the buffer anion is F⁻, this column applies.
^cIf the buffer anion is Cl⁻ or Br⁻, this column applies.

Table 3

Potential ion contributions to the salt concentration dependence of DNA binding attributable to residues 75 and 73

Free protein	Partial reactions from Table 2	Net number of ions released in FI ⁻ buffer ^a	Net number of ions released in Cl ⁻ or Br ⁻ buffer ^a
WT, neutralization of K75 and D73 by solution ions	a + b + c	1	2
WT, neutralization of K75 and D73 by salt bridge	a + d + e	0	0
K75A	c + f	0	0
D73A	a + b + f	1	2
K75A D73A	f	0	0

^a As calculated from the entries in Table 2.