Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2010 January ; 19(1): 258–264. doi:10.1158/1055-9965.EPI-09-0800.

Population-based study of the association of variants in mismatch repair genes with prostate cancer risk and outcomes

Wendy J. Langeberg^{1,2}, Erika M. Kwon³, Joseph S. Koopmeiners^{1,2}, Elaine A. Ostrander³, and Janet L. Stanford^{1,2}

¹Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA

²School of Public Health, University of Washington, Seattle, WA

³National Human Genome Research Institute, National Institutes of Health, Bethesda, MD

Abstract

Background—Mismatch repair (MMR) gene activity may be associated with prostate cancer (PC) risk and outcomes. This study evaluated whether single nucleotide polymorphisms (SNPs) in key MMR genes are related to PC outcomes.

Methods—Data from two population-based case-control studies of PC among Caucasian and African-American men residing in King County, Washington were combined for this analysis. Cases (n=1,458) were diagnosed with PC in 1993–96 or 2002–05 and identified via the Seattle-Puget Sound SEER cancer registry. Controls (n=1,351) were age-matched to cases and identified via random digit dialing. Logistic regression was used to assess the relationship between haplotype-tagging SNPs and PC risk and disease aggressiveness. Cox proportional hazards regression was used to assess the relationship between SNPs and PC recurrence and PC-specific death.

Results—Nineteen SNPs were evaluated in the key MMR genes: five in *MLH1*, 10 in *MSH2*, and 4 in *PMS2*. Among Caucasian men, one SNP in *MLH1* (rs9852810) was associated with: overall PC risk (OR=1.21, 95% CI=1.02, 1.44; p=0.03), more aggressive PC (OR=1.49, 95% CI=1.15–1.91; p<0.01), and PC recurrence (HR=1.83, 95% CI=1.18, 2.86; p<0.01), but not PC-specific mortality. A non-synonymous coding SNP in *MLH1*, rs1799977 (I219V), was also found to be associated with more aggressive disease. These results did not remain significant after adjusting for multiple comparisons.

Conclusion—This population-based case-control study provides evidence for a possible association with a gene variant in *MLH1* in relation to risk of overall PC, more aggressive disease, and PC recurrence, which warrants replication.

Keywords

prostate cancer; mismatch repair; association study; genetic variant; odds ratio; recurrence; biomarker

Introduction

This year alone, an estimated 30,000 deaths will occur among US men due to prostate cancer [1]. Established risk factors for PC (age, race/ethnicity, and a family history of PC) and features of more aggressive disease (e.g., higher Gleason score, advanced tumor stage, and high

prostate-specific antigen [PSA] levels) are not adequate to predict which cases will become life-threatening; therefore, active investigation is underway to identify biomarkers that will enhance the ability to identify patients at higher risk for adverse PC outcomes [2]. In this analysis, we evaluated the association of variants in key mismatch repair (MMR) genes, *MSH2* (on 2p22–21), *MLH1* (on 3p21), and *PMS2* (on 7p22), in relation to overall PC risk, risk of more aggressive disease, PC recurrence, and PC-specific mortality.

Mutations in MMR genes (*MLH1*, *MSH2*, *MSH3*, *MSH6*, *PMS1*, and *PMS2*) can lead to instability of microsatellites (MSI) and failure to repair DNA damage during DNA replication. This damaged DNA can accumulate and eventually lead to the development of neoplasms, such as hereditary nonpolyposis colon cancer (HNPCC), which is characterized by mutations in five microsatellites [3]. A number of studies have reported more MSI in PC tumor tissue compared to normal prostatic tissue [4–9], but some PC tissue studies have found a low frequency of MSI [10–14]. In addition, reduction or loss of MMR protein expression has been found in human PC cell lines, such as LNCaP, PC-3 and DU145 [15–20]. And some studies, but not all, have correlated *hMSH2* immunohistochemical staining intensity with a higher Gleason score and lower disease-free survival [21–23]. Recently, Norris et al. found elevated levels of PMS2 in the prostate tumor tissue of patients who recurred compared with non-recurrent patients [24].

The non-synonymous coding SNP rs1799977 in *MLH1* (also referred to as Ile-219Val or I219V) has been evaluated in two studies of PC risk, with mixed results. Using 275 PC sibships and 556 unrelated controls, Burmester et al. found the rare allele of the SNP rs1799977 was significantly associated with PC [25]. Fredriksson et al., however, found no difference in allele frequency for rs1799977 between 121 patients with hereditary PC (allele frequency=54.5%), unselected patients with PC (54.0%), 202 patients with benign prostatic hyperplasia (54.0%), and 200 controls (55.0%) [26].

In light of these provocative but inconclusive findings, this study evaluated the association between variants in the key MMR genes and the risk of PC and PC outcomes.

Methods

Study Population

Data were combined for this analysis from two population-based case-control studies of risk factors for PC among Caucasian and African-American men residing in King County, Washington, described previously [27–28]. Both studies ascertained cases from the Seattle-Puget Sound Surveillance Epidemiology and End Results (SEER) cancer registry. The first study included 753 cases diagnosed between January 1, 1993 and December 31, 1996 who were 40 to 64 years of age at diagnosis. The second study included 1,001 cases diagnosed between January 1, 2002 and December 31, 2005 who were 35 to 74 years of age at diagnosis. Controls (n=703 for the first study, n=942 for the second study) were men without a self-reported history of PC, who were recruited via random digit dialing (RDD) during the same ascertainment period and from the same underlying general population as the cases; they were frequency matched to cases by five-year age groups. Among eligible subjects ascertained for the first study, 82% of cases and 75% of controls participated in the study interview, and of these participants, 84% of cases and 80% of controls provided a blood sample. Among eligible subjects ascertained for the second study, 75% of cases and 63% of controls participated in the study interview, and of these participants, 83% of cases and 84% of controls provided a blood sample. After combining these two studies, there were 1,457 PC cases and 1,351 controls with DNA available for the analysis.

Background information was collected from participants at the time of interview and included demographic and lifestyle factors, medical history, PC screening history, and family history of PC. This information was assessed prior to date of diagnosis for cases and prior to a pre-assigned reference date for controls. Clinical information such as Gleason score, tumor stage, serum PSA level at diagnosis, and primary treatment was obtained from the cancer registry. Patient files have been linked to the registry on a regular basis to obtain vital status and primary cause of death of cases; death certificates are requested from the state to confirm underlying cause of death. In 2004, a follow-up survey was sent to 631 of the cases from the first study, 82% of whom responded, to assess secondary treatment(s) and evidence for PC recurrence or progression.

The Institutional Review Board (IRB) of the Fred Hutchinson Cancer Research Center approved study procedures and materials, and written informed consent was obtained from all study participants. Genotyping was approved by the National Human Genome Research Institute's IRB.

TagSNP Selection and Genotyping

DNA samples were genotyped for 20 single nucleotide polymorphisms (SNPs) in the *MLH1*, *MSH2*, and *PMS2* genes. The SNPs were selected using the Genome Variation Server (gvs.gs.washington.edu/gvs) to cover the genes as haplotype-tagging SNPs. The Applied Biosystems (ABI) SNPlex® Genotyping System was used for genotyping and proprietary GeneMapper® software was used for allele assignment (www.appliedbiosystems.com). Discrimination of the specific SNP allele was carried out with the ABI 3730xl DNA Analyzer and is based on the presence of a unique sequence assigned to the original allele-specific oligonucleotide. Quality control included genotyping of 144 blind duplicate samples distributed across all genotyping batches. There was ≥99% agreement between blinded samples for all SNP genotypes. Each batch of DNA aliquots genotyped incorporated similar numbers of case and control samples, and laboratory personnel were blinded to the case-control status of samples. Genotype frequencies in *MLH1*, *MSH2*, and *PMS2* were evaluated among Caucasian and African-American controls separately; all SNPs were consistent with the expected proportions under Hardy-Weinberg, except for rs12112229 among Caucasians, and so this SNP was removed from the analysis.

Statistical Methods

Logistic regression was used to calculate odds ratios (ORs) and 95% confidence intervals (CIs) to estimate the relative risk of PC among cases relative to controls for each SNP genotype. Polytomous logistic regression was used to calculate ORs and 95% CIs to estimate the relative risk of more aggressive and less aggressive PC relative to controls for each SNP genotype. More aggressive PC was defined by a Gleason score of 7(4+3) or 8–10, regional or distant tumor stage, or a diagnostic PSA value ≥20 ng/mL. Codominant and dominant genetic models were considered for each SNP. All models were adjusted for age at reference date, and tested for possible confounding by PC screening history and/or family history of PC. In addition, permuted p-values were calculated to adjust for multiple comparisons, as described previously [29].

Cox proportional hazards regression was used to estimate hazard ratios and 95% CIs to assess the relationship between the SNPs found to be significantly associated with aggressive PC and recurrence or death from PC. The analyses of recurrence were restricted to cases diagnosed with local or regional stage disease and who either subsequently died of PC (prior to the follow-up survey) or completed a follow-up survey, which provided recurrence information and consent to obtain medical records. Recurrence was defined as at least one of the following from self-report and/or medical records: a positive bone scan, CT, MRI, or biopsy showing PC after

primary treatment; use of secondary therapy (androgen deprivation therapy [ADT], external beam radiation therapy, cryotherapy, or chemotherapy); an elevated PSA (≥ 0.2 ng/mL) after radical prostatectomy; an elevated PSA after radiation therapy (nadir PSA +2 ng/mL); a rising PSA while on primary ADT; treatment for evidence of progressive disease that was initiated >12 months after diagnosis in patients on active surveillance; or a self-reported physician's diagnosis of disease recurrence/progression. Time from diagnosis until recurrence was calculated as the difference between the date of diagnosis and the earliest date of evidence of recurrence: date of death from PC, date of recurrence or progression abstracted from medical records, date of recurrence from the follow-up survey, or, for those censored, the end of the year during which the follow-up survey was collected (December 31, 2005). For men who died of PC before December 31, 2005, date of recurrence was imputed to be similar to dates of recurrence for comparable subjects. The analyses of PC death included all cases. The censoring date for members last known to be alive was the date of the last vital status update from the cancer registry (December 1, 2008). The proportional hazards models were adjusted for age and tested for possible confounding by PC screening history or a family history of PC, and recalculated including only cases who received radical prostatectomy as primary therapy.

Most analyses were performed in SAS® version 9.1.3 (SAS Institute, Cary, NC). Hardy-Weinberg equilibrium was calculated in STATA/SE® 10.0 for Windows (StataCorp, College Station, TX).

Results

Among the 1,458 cases and 1,351 controls, a higher proportion of cases than controls were African-American (10.2% vs. 6.3%, respectively; Table 1), had a first-degree relative with PC (21.5% vs. 11.3%), and reported having a PSA or DRE screening test in the five years prior to diagnosis or reference date (89.3% and 86.5%).

Nineteen tagSNPs were evaluated: 5 in *MLH1*, 10 in *MSH2*, and 4 in *PMS2*. Among Caucasian men, one SNP in *MLH1* (rs9852810) was associated with overall PC risk (OR=1.21, 95% CI=1.02, 1.44, $p=0.03$; Table 2 and supplementary data). Rs9852810 and another SNP in *MLH1*, rs1799977, were associated with more aggressive prostate cancer among Caucasian men when aggressive cases were compared with controls (rs9852810: OR, 1.49; 95% CI, 1.15–1.91; $P<0.01$; rs1799977: OR, 1.35; 95% CI, 1.08–1.69; $P=0.03$; Table 2) and when aggressive cases were compared with less aggressive cases (rs9852810: OR, 1.34; 95% CI, 1.03–1.75; $P=0.03$; rs1799977: OR, 1.33; 95% CI, 1.05–1.69; $P=0.02$; data not shown). After adjustment for multiple comparisons using permutation p -values, rs9852810 did not remain significantly associated with overall PC risk ($p_{\text{perm}}=0.22$); in addition the associations between rs9852810 and rs1799977 with more aggressive disease did not attain statistical significance (when compared to controls, $p_{\text{perm}}=0.09$ for both SNPs). The association with overall PC risk and with disease aggressiveness remained similar after adjustment for a first-degree relative with PC or having a PC screening test in the five years prior to reference date. Similar analyses among African-American men revealed no associations between any SNP genotypes and overall prostate cancer risk (Supplementary Data) or disease aggressiveness (data not shown).

Among the 469 Caucasian cases diagnosed with local or regional disease who completed a follow-up survey or died of prostate cancer before December 31, 2005, 143 recurred. Rs9852810, was associated with PC recurrence in Caucasians (110 out of 320 [34.4%] cases with the putative risk genotype and 24 out of 115 [20.9%] cases with the homozygous wild-type genotype recurred; $HR_{GA+AA}=1.83$, 95% CI=1.18, 2.86, $p<0.01$; Table 3). Rs1799977 was not associated with PC recurrence and neither SNP was associated with PC-specific mortality (Table 3).

Discussion

In this population-based case-control study of tagSNPs in key MMR genes (*MLH1*, *MSH2*, and *PMS2*), we found the SNP rs9852810 in *MLH1* to be associated with a modest increase in overall PC risk, risk of more aggressive PC, and PC recurrence. This intronic SNP is in perfect LD with several other SNPs near the start codon of *MLH1* (such as rs11129748). To our knowledge, the association with this variant and PC has not been evaluated previously. We also found an association between the non-synonymous coding SNP rs1799977 in *MLH1* and more aggressive PC. As noted in the introduction, the association between this SNP and PC has been evaluated previously with mixed results [25-26]. This SNP has also recently been reported to be associated with breast cancer risk (OR = 1.87; 95% CI = 1.11, 3.16) [30], and may be associated with susceptibility to childhood acute lymphoblastic leukemia [31].

One limitation to this study is possible type I error due to multiple testing. For each of the 19 SNPs, we calculated 6 significance tests among Caucasians, so one would expect about 6 results might be due solely to chance. The main result (for rs9852810) did not remain significant based on a permuted p-value; however, it was significant in the PC risk analysis, the analysis of aggressive disease, and the analysis of recurrence, which lends strength to the result. If confirmed, this result lends further support for a potential shared susceptibility for PC and colon cancer, which is consistent with prior findings for a SNP in the 8q24 region that confers risk for both cancer types [32-33].

There are several strengths to this study. The data used for this analysis were from two population-based case-control studies, which means men with all grades and stages of disease, and who received a range of initial treatments, were included. In addition, we have over 10 years of patient follow-up to evaluate recurrence and progression, and clinical and patient information was available for evaluation of potential confounders and effect modifiers.

Conclusion

Evidence from previous studies shows that loss of mismatch repair function may be characteristic of prostate carcinogenesis. This population-based study provides evidence for a possible association with a gene variant in *MLH1* in relation to risk of overall PC, more aggressive disease, and PC recurrence, which warrants replication.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the men who participated in these studies; without their help, this work would not be possible. This work was supported by grants RO1 CA056678, RO1 CA082664, RO1 CA092579, and P50 CA097186 from the National Cancer Institute, with additional support from the Fred Hutchinson Cancer Research Center and the Intramural Program of the National Human Genome Research Institute.

References

1. Jemal A, Siegel R, Ward E, et al. Cancer statistics, 2008. *CA Cancer J Clin* Mar-Apr;2008 58(2):71–96. [PubMed: 18287387]
2. Salinas CA, Koopmeiners JS, Kwon EM, et al. Clinical utility of five genetic variants for predicting prostate cancer risk and mortality. *Prostate* Mar 1;2009 69(4):363–72. [PubMed: 19058137]
3. Lynch HT, de la Chapelle A. Hereditary Colorectal Cancer. *New England Journal of Medicine* Mar; 2003 348(10):919–32. [PubMed: 12621137]

4. Colombo P, Patriarca C, Alfano RM, et al. Molecular disorders in transitional vs. peripheral zone prostate adenocarcinoma. *Int J Cancer* 2001;94:383–389. [PubMed: 11745418]
5. Dahiya R, Lee C, McCarville J, Hu W, Kaur G, Deng G. High frequency of genetic instability of microsatellites in human prostatic adenocarcinoma. *Int J Cancer* 1997;72:762–767. [PubMed: 9311591]
6. Egawa S, Uchida T, Suyama K, et al. Genomic instability of microsatellite repeats in prostate cancer: Relationship to clinicopathological variables. *Cancer Res* 1995;55:2418–2421. [PubMed: 7757995]
7. Uchida T, Wada C, Wang C, et al. Microsatellite instability in prostate cancer. *Oncogene* 1995;10:1019–1022. [PubMed: 7898920]
8. Gao X, Wu N, Grignon D, et al. High frequency of mutator phenotype in human prostatic adenocarcinoma. *Oncogene* 1994;9:2999–3003. [PubMed: 8084604]
9. Suzuki H, Komiya A, Aida S, et al. Microsatellite instability and other molecular abnormalities in human prostate cancer. *Jpn J Cancer Res* 1995;86:956–961. [PubMed: 7493915]
10. Cunningham JM, Shan A, Wick MJ, et al. Allelic imbalance and microsatellite instability in prostatic adenocarcinoma. *Cancer Res* 1996;56:4475–4482. [PubMed: 8813143]
11. Zhou XP, Hoang JM, Li YJ, et al. Determination of the replication error phenotype in human tumors without the requirement for matching normal DNA by analysis of mononucleotide repeat microsatellites. *Genes Chromosomes Cancer* 1998;21:101–107. [PubMed: 9491320]
12. Ahman AK, Jonsson BA, Damber JE, Bergh A, Grönberg H. Low frequency of microsatellite instability in hereditary prostate cancer. *BJU Int* Mar;2001 87(4):334–8. [PubMed: 11251526]
13. Terrell RB, Wille AH, Cheville JC, Nystuen AM, Cohen MB, Sheffield VC. Microsatellite instability in adenocarcinoma of the prostate. *Am J Pathol* 1995;147:799–805. [PubMed: 7677191]
14. Rohrbach H, Haas CJ, Baretton GB, et al. Microsatellite instability and loss of heterozygosity in prostatic carcinomas: Comparison of primary tumors, and of corresponding recurrences after androgen deprivation therapy and lymph-node metastases. *Prostate* 1999;40:20–27. [PubMed: 10344720]
15. Boyer JC, Umar A, Risinger JI, et al. Microsatellite instability mismatch repair deficiency, and genetic defects in human cancer cell lines. *Cancer Res* 1995;55:6063–6070. [PubMed: 8521394]
16. Leach FS, Velasco A, Hsieh JT, Sagalowsky AI, McConnell JD. The mismatch repair gene hMSH2 is mutated in the prostate cancer cell line LNCaP. *J Urol* Nov;2000 164(5):1830–3. [PubMed: 11025778]
17. Yeh CC, Lee C, Dahiya R. DNA mismatch repair enzyme activity and gene expression in prostate cancer. *Biochem Biophys Res Commun* 2001;285:409–413. [PubMed: 11444857]
18. Chen Y, Wang J, Fraig MM, et al. Defects of DNA mismatch repair in human prostate cancer. *Cancer Res* May 15;2001 61(10):4112–21. [PubMed: 11358834]
19. Chen Y, Wang J, Fraig MM, et al. Alterations in PMS2, MSH2 and MLH1 expression in human prostate cancer. *Int J Oncol* 2003;22:1033–1043. [PubMed: 12684669]
20. Norris AM, Woodruff RD, D'Agostino RB Jr, Clodfelter JE, Scarpinato KD. Elevated Levels of the Mismatch Repair Protein Pms2 Are Associated With Prostate Cancer. *Prostate* Feb;2007 67(2):214–25. [PubMed: 17044039]
21. Prtalo A, Leach FS, Markwalder R, et al. Tissue microarray analysis of hMSH2 expression predicts outcome in men with prostate cancer. *Journal of Urology* Nov;2005 174:1814–8. [PubMed: 16217293]
22. Velasco A, Albert PS, Rosenberg H, Martinez C, Leach FS. Clinicopathologic implications of hMSH2 gene expression and microsatellite instability in prostate cancer. *Cancer Biol Ther* 2002;1:362–367. [PubMed: 12432247]
23. Velasco A, Hewitt SM, Albert PS, et al. Differential expression of the mismatch repair gene hMSH2 in malignant prostate tissue is associated with cancer recurrence. *Cancer* Feb 1;2002 94(3):690–9. [PubMed: 11857301]
24. Norris AM, Gentry M, Peehl DM, D'Agostino R Jr, Scarpinato KD. The elevated expression of a mismatch repair protein is a predictor for biochemical recurrence after radical prostatectomy. *Cancer Epidemiol Biomarkers Prev* Jan;2009 18(1):57–64. [PubMed: 19124481]
25. Burmester JK, Suarez BK, Lin JH, et al. Analysis of candidate genes for prostate cancer. *Hum Hered* 2004;57(4):172–8. [PubMed: 15583422]

26. Fredriksson H, Ikonen T, Autio V, et al. Identification of germline MLH1 alterations in familial prostate cancer. *Eur J Cancer* Nov;2006 42(16):2802–6. [PubMed: 16963262]
27. Stanford JL, Wicklund KG, McKnight B, Daling JR, Brawer MK. Vasectomy and risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev* Oct;1999 8(10):881–6. [PubMed: 10548316]
28. Agalliu I, Salinas CA, Hansten PD, Ostrander EA, Stanford JL. Statin use and risk of prostate cancer: Results from a population-based epidemiological study. *Am J Epidemiol* 2008;168(3):250–260. [PubMed: 18556686]
29. Fitzgerald LM, Kwon EM, Koopmeiners JS, Salinas CA, Stanford JL, Ostrander EA. Analysis of recently identified prostate cancer susceptibility loci in a population-based study: associations with family history and clinical features. *Clin Cancer Res* May 1;2009 15(9):3231–7. [PubMed: 19366831]
30. Smith TR, Levine EA, Freimanis RI, et al. Polygenic model of DNA repair genetic polymorphisms in human breast cancer risk. *Carcinogenesis* Nov;2008 29(11):2132–8. [PubMed: 18701435]
31. Mathonnet G, Krajcinovic M, Labuda D, Sinnott D. Role of DNA mismatch repair genetic polymorphisms in the risk of childhood acute lymphoblastic leukaemia. *Br J Haematol* Oct;2003 123(1):45–8. [PubMed: 14510941]
32. Amundadottir LT, Sulem P, Gudmundsson J, et al. A Common Variant Associated With Prostate Cancer in European and African Populations. *Nature Genetics* Jun;2006 38(6):652–8. [PubMed: 16682969]
33. Haiman CA, Le Marchand L, Yamamoto J, et al. A common genetic risk factor for colorectal and prostate cancer. *Nat Genet* Aug;2007 39(8):954–6. [PubMed: 17618282]

Table 1
Characteristics of population-based prostate cancer cases and controls

Characteristic	Cases (n=1,458)		Controls (n=1,351)	
	n	(%)	n	(%)
Age at diagnosis/reference date				
35–49	118	(8.1)	126	(9.3)
50–54	215	(14.8)	209	(15.5)
55–59	357	(24.5)	358	(26.5)
60–64	433	(29.7)	348	(25.8)
65–69	177	(12.1)	164	(12.1)
70–74	158	(10.8)	146	(10.8)
Race				
Caucasian	1,309	(89.8)	1,266	(93.7)
African-American	149	(10.2)	85	(6.3)
First-degree relative with prostate cancer				
No	1,145	(78.5)	1,199	(88.8)
Yes	313	(21.5)	152	(11.3)
Screening history ¹				
None	157	(10.8)	182	(13.5)
DRE only	258	(17.7)	519	(38.4)
PSA	1,043	(71.6)	650	(48.1)
PSA value ²				
< 4.0	189	(13.0)	1,253	(92.8)
4.0–9.9	814	(55.8)	80	(5.9)
10.0–19.9	210	(14.4)	16	(1.2)
≥ 20.0	138	(9.5)	2	(0.2)
Missing	107	(7.4)		
Gleason score				
2–4	72	(4.9)	--	--
5–6	741	(50.8)	--	--
7 (3+4)	408	(28.0)	--	--
7 (4+3)	91	(6.2)	--	--
8–10	140	(9.6)	--	--
Missing	6	(0.4)	--	--
Stage at diagnosis				
Local	1,141	(78.3)	--	--
Regional	280	(19.2)	--	--
Distant	37	(2.5)	--	--
Primary treatment				

Characteristic	Cases (n=1,458)		Controls (n=1,351)	
	n	(%)	n	(%)
RP	831	(57.0)	--	--
RT	412	(28.3)	--	--
ADT	72	(4.9)	--	--
Other treatment	5	(0.3)	--	--
Active surveillance	138	(9.5)	--	--

PSA=prostate-specific antigen; RP=radical prostatectomy; RT=radiation therapy; ADT=androgen deprivation therapy

¹ Screening history within five years prior to diagnosis or reference date.

² PSA at diagnosis for cases and measured at interview date for controls.

Table 2
Risk of prostate cancer and disease aggressiveness¹ associated with two SNPs in the *MLH1* gene²

Controls			All cases			Less aggressive cases			More aggressive cases		
(n=1,266) ³			(n=1,309) ³			(n=874) ³			(n=435) ³		
SNP	n	(%)	n	%	OR ⁴	95% CI	p ⁵	n	%	OR ⁴	95% CI
rs9852810, chr7:115949965											
GG	410	(33.1)	364	(28.9)	1.00	Reference		260	(30.9)	1.00	Reference
GA	601	(48.5)	651	(51.8)	1.21	1.02 1.45		427	(50.8)	1.12	0.91 1.36
AA	228	(18.4)	243	(19.3)	1.20	0.96 1.51	0.09	154	(18.3)	1.06	0.82 1.37
GA/AA	829	(66.9)	894	(71.1)	1.21	1.02 1.44	0.03	581	(69.1)	1.10	0.91 1.33
rs1799977, chr7:115969290											
TT	607	(49.1)	578	(46.2)	1.00	Reference		406	(48.5)	1.00	Reference
CT	514	(41.6)	555	(44.4)	1.13	0.96 1.33		357	(42.6)	1.04	0.86 1.25
CC	115	(9.3)	118	(9.4)	1.07	0.81 1.42	0.35	75	(9.0)	0.96	0.7 1.32
CT/CC	629	(50.9)	673	(53.8)	1.12	0.96 1.31	0.16	432	(51.6)	1.02	0.86 1.22

SNP=single nucleotide polymorphism; OR=odds ratio; CI=confidence interval

¹ More aggressive PC is defined by a Gleason score of 7(4+3) or 8–10, regional or distant tumor stage, or a diagnostic PSA value ≥20 ng/ml.

² Among Caucasian cases and controls only.

³ Total number of cases and controls varies by SNP due to missing genotype data.

⁴ Adjusted for age at reference date.

⁵ The first p-value is the test for trend using the co-dominant model; the second p-value is for the dominant model.

Table 3
Risk of prostate cancer recurrence and death associated with two SNPs in the *MLH1* gene¹

Risk of recurrence				Risk of death									
SNP	n ²	no. who recurred ³	(%)	median time to recurrence/ censorship (years)	HR ⁴	95% CI	p	n	no. who died of PC	(%)	HR ⁴	95% CI	p
rs9852810, chr7:1159499650													
GG	115	24	(20.9)	9.0	1.00	Reference		364	13	(3.6)	1.00	Reference	0.42
GA/AA	320	110	(34.4)	8.6	1.83	1.18 2.86	0.007	894	43	(4.8)	1.29	0.69 2.4	
rs1799977, chr7:115969290													
TT	195	55	(28.2)	8.9	1.00	Reference		578	25	(4.3)	1.00	Reference	0.74
CT/CC	238	79	(33.2)	8.7	1.22	0.86 1.72	0.260	673	32	(4.8)	1.09	0.65 1.84	

HR=hazard ratio; CI=confidence interval

¹ Among Caucasian cases only.

² Cases diagnosed with local or regional disease who completed a follow-up survey or died of prostate cancer before Dec. 31, 2005.

³ Recurrence is defined as at least one of the following: positive bone scan, CT, MRI, or biopsy showing PC after primary treatment; biochemical failure after RP as primary treatment (PSA ≥0.2 ng/mL); biochemical failure after RT as primary treatment (nadir PSA +2 ng/mL); ADT as secondary treatment or a rising PSA on ADT; or RT as secondary treatment.

⁴ Risk of recurrence or death, respectively, among PC patients with the at-risk allele relative to PC patients homozygous for the wildtype allele.