Published in final edited form as:

Annu Rev Biophys. 2008 ; 37: 375–397. doi:10.1146/annurev.biophys.37.032807.125817.

Macromolecular crowding and confinement: biochemical, biophysical, and potential physiological consequences*

Huan-Xiang Zhou¹, Germán Rivas², and Allen P. Minton³

Huan-Xiang Zhou: zhou@sb.fsu.edu; Germán Rivas: grivas@cib.csic.es; Allen P. Minton: minton@helix.nih.gov

¹ Department of Physics and Institute of Molecular Biophysics and School of Computational Science, Florida State University, Tallahassee, Florida 32306

² Departamento de Ciencia de Proteínas, Centro de Investigaciones Biológicas, CSIC, Madrid 28040

³ Section on Physical Biochemistry, Laboratory of Biochemistry and Genetics, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, US Department of Health and Human Services, Bethesda, Maryland 20892

Abstract

Expected and observed effects of volume exclusion on the free energy of rigid and flexible macromolecules in crowded and confined systems, and consequent effects of crowding and confinement on macromolecular reaction rates and equilibria are summarized. Findings from relevant theoretical/simulation and experimental literature published from 2004 onward are reviewed. Additional complexity arising from the heterogeneity of local environments in biological media, and the presence of nonspecific interactions between macromolecules over and above steric repulsion are discussed. Theoretical and experimental approaches to the characterization of crowding- and confinement-induced effects in systems approaching the complexity of living organisms are suggested.

Keywords

excluded volume; configurational entropy; free energy; protein-protein interactions; macromolecular associations; protein folding; site-binding

Introduction

It is now widely appreciated that almost all proteins and other biological macromolecules *in vivo* exist, at least transiently, as components of structural and functional complexes (3). The number of published studies of macromolecular interactions has increased almost exponentially for the last 20 years and amounts to several hundred a year at present. However, almost all of these studies are aimed at the characterization of attractive interactions that result in the formation of protein complexes or complexes of protein and other macromolecules (e.g. ribonucleoproteins). In contrast, repulsive interactions between macromolecules, by definition, do not lead to the formation of complexes and thus may not be observed directly. However, the presence and significance of repulsive interactions in fluid media containing a high total concentration of macromolecules and/or structural obstacles to the free motion of macromolecules may be observed indirectly through their effects upon a variety of

*Dedicated to Steven B. Zimmerman, pioneering investigator of excluded volume effects in biological systems, with emphasis on the influence of macromolecular crowding on the structure and function of DNA

macromolecular reactions involving association and conformational isomerization (49,52). Many of these effects may be predicted qualitatively (and sometimes quantitatively) using simple statistical-thermodynamic models, and observed experimentally by measurement of the dependence of thermodynamically-based solution properties and reaction kinetics and equilibria upon the concentration and composition of macromolecular co-solutes that are nominally inert with respect to the reaction of interest. The importance of excluded volume interactions lies in their generality. Such interactions are universal and entirely nonspecific, and have the potential to significantly modulate the kinetics and equilibria of a large number of macromolecular reactions taking place in physiological fluid media.

We shall classify an excluded volume effect according to its origin: *macromolecular crowding* refers to effects of volume exclusion by one soluble macromolecule to another, and *macromolecular confinement* refers to effects of volume exclusion by a fixed (or confining) boundary to a soluble macromolecule. A number of reviews of both crowding and confinement effects have appeared during the past five years (19,27,55,57,71,89,92,93) and it is not our intention to recapitulate published material. Moreover, a separate review of the effect of crowding on macromolecular transport via diffusion appears elsewhere in the present volume. However, in the interest of completeness, in the following section we briefly summarize the various ways that crowding and confinement are expected to influence the equilibria and kinetics of macromolecular associations and isomerizations. Next, results of recently published (2004–2007) theoretical contributions, atomistic simulations, and experiments relating to effects of macromolecular crowding and confinement upon a variety of macromolecular reactions are summarized. (References to work published prior to 2004 may be found in one or more of the reviews cited above.). The present review concludes with discussions of several topics related to the applicability of theoretical predictions or the results of experiments conducted *in vitro* to actual macromolecular processes taking place in living organisms.

A common thermodynamic framework for analysis of macromolecular crowding and confinement

General features of macromolecular crowding and confinement are qualitatively exhibited by their effects upon three prototypical macromolecular reactions:

Bimolecular association

Association of a soluble macromolecular ligand with a specific surface binding site

Two-state folding of a protein

These three reactions are characterized respectively by three standard free energy changes and the corresponding equilibrium constants:

$$\begin{aligned}\Delta F_{AB} &= -RT \ln K_{AB} \\ \Delta F_{LS} &= -RT \ln K_{LS} \\ \Delta F_{UN} &= -RT \ln K_{UN}\end{aligned}$$

where R denotes the molar gas constant and T the absolute temperature. The effect of an environmental perturbation such as crowding or confinement upon the equilibrium state of a particular chemical reaction may be analyzed by constructing a simple thermodynamic cycle as shown in Figures 1A–C and Figures 2A–C. The value of standard free energy changes and equilibrium constants for a particular reaction in bulk dilute solution is denoted by the superscript ‘0’, e.g., $\Delta F_{AB}^0, K_{AB}^0$. Then the free energy of hetero-association, site binding, and unfolding in the bulk and perturbed environments (either crowded or confined) will be related by (57)

$$\Delta\Delta F_{AB} \equiv \Delta F_{AB} - \Delta F_{AB}^0 = \Delta F_{AB}^C - \Delta F_A^C - \Delta F_B^C \quad (1)$$

$$\Delta\Delta F_{LS} \equiv \Delta F_{LS} - \Delta F_{LS}^0 = \Delta F_{LS}^C - \Delta F_L^C \quad (2)$$

and

$$\Delta\Delta F_{UN} \equiv \Delta F_{UN} - \Delta F_{UN}^0 = \Delta F_N^C - \Delta F_U^C \quad (3)$$

where ΔF_X^C denotes the standard free energy change associated with the transfer of X from bulk solution to the crowded or confined environment as indicated in Figures 1 and 2. [Footnote 1] [Footnote 2] Free energy changes (on a per mole basis) are related to changes in the corresponding equilibrium constant by

$$K_X = K_X^0 \exp(-\Delta\Delta F_X / RT) \quad (4)$$

where X denotes AB , LS or UN . The magnitude of the effect of crowding or confinement upon a particular reaction equilibrium may thus be evaluated indirectly by comparing the free energies of transfer of reactants and products from bulk solution to the crowded or confined medium. Examples of such estimates are provided below. [Footnote 3] In these examples, we focus on the excluded volume aspect of crowding and confinement, because this effect appears

¹Equations (1 – 3) apply equally to changes in Helmholtz or Gibbs free energies.

Simple models used to estimate the magnitude of ΔF_X^C generally assume constant volume and hence in the strict sense yield estimates of Helmholtz free energy changes. However, differences between Helmholtz and Gibbs free energy changes associated with reactions in the liquid state are not of qualitative significance.

² ΔF_X^C is formally equivalent to the difference between the equilibrium average free energy of interaction of X with the perturbing co-solutes or boundaries and the equilibrium average free energy of interaction of X with the constituents of bulk solvent that are replaced by co-solute or boundary. Thus ΔF_X^C implicitly takes into account any energetic consequence of de-solvation that may accompany the transfer.

to be universal. Other nonspecific interactions of reactants with crowders and confining walls will be discussed later.

Crowding

Estimation of ΔF_X^{crowd}

The colligative properties of concentrated solutions of globular proteins, under conditions such that long range interactions between protein molecules are damped out, are well described by models in which individual macromolecular species are represented by hard convex particles that resemble the molecules in general size and shape (50,58), justifying the use of hard particle models for quantitative estimation of excluded volume effects. The free energy of transfer of a macromolecule X from a dilute solution to a solution containing an arbitrary concentration of other macromolecules of the same or other species is then equivalent to the free energy of creating a cavity in the solution, free of any part of another molecule, that is large enough to accommodate the newly introduced macromolecule (43). The scaled particle theory (SPT) of hard particle fluids, originally due to Reiss et al (68), was devised specifically to calculate the free energy of cavity formation, and is thus particularly appropriate for numerical estimation of the magnitude of crowding effects. Let us consider as an example a fluid containing a volume fraction ϕ of inert hard spherical particles with radius r_c . According to SPT (14,70) the free energy or work of placing into this fluid a spherocylinder (sc), that is, a right circular cylinder capped on each end by a hemisphere, with a radius $r_{sc} = R_{sc}r_c$ and a cylindrical length of $2Lr_{sc}$, is given by

$$\Delta F_{sc}^{crowd}/RT = -\ln(1-\phi) + A_1Q + A_2Q^2 + A_3Q^3 \quad (5)$$

where

$$\begin{aligned} Q &= \phi/(1-\phi) \\ A_1 &= R_{sc}^3 + 3R_{sc}^2 + 3R_{sc} + 1.5L(R_{sc}^2 + 2R_{sc} + 1) \\ A_2 &= 1.5(2R_{sc}^3 + 3R_{sc}^2) + 4.5L(R_{sc}^2 + R_{sc}) \\ A_3 &= 3R_{sc}^3 + 4.5LR_{sc}^2 \end{aligned}$$

Note that the free energy of transfer of a spherical particle into this fluid is just the special case of equation (5) with $L = 0$.

Association equilibria (Fig 1A)

Consider a bimolecular association reaction (I) taking place in a solution of spherical crowding molecules. For illustration, let monomeric A and B be represented as identical hard spherical particles with radius equal to that of crowder (i.e., $R_{sc} = R_I = 1$, $L = L_I = 0$). The dimer AB is represented as a spherocylindrical particle with volume equal to twice that of monomer, but with $L = L_2$ and $R_{sc} = R_2$ simultaneously adjustable to maintain constant volume, to illustrate the effect of different assumed shapes for dimer. Results obtained using equations (1), (4), and (5) are plotted in Figure 3. It is evident that crowding can substantially enhance the equilibrium tendency of A and B to dimerize when AB is compact ($L_2 \leq 1$), but can also inhibit dimerization

³The treatment presented here may be readily extended to a quasi-equilibrium analysis of the effect of crowding or confinement upon the kinetics of a transition-state limited association or isomerization reaction, in which case one must additionally estimate the free energy change associated with the transfer of the transition state from bulk to the crowded or confined medium (see for example the Appendix of (54)).

when the dimer is so aspherical that it excludes more volume to crowder than two monomers. Many proteins can also form higher order oligomers. The effect of crowding upon the equilibrium constant for concerted association of n monomers to 1 n -mer increases dramatically with the value of n (49).

Association rates

The rate constant of association between two molecules can generally be written as (51,91)

$$k_a = \frac{k_D k_{\text{react}}}{k_D + k_{\text{react}}} \quad (6)$$

where k_D is the rate constant under diffusion control and k_{react} is the rate constant under reaction or transition-state control. Crowding affects the rate constant in the two regimes differently. k_D is proportional to the relative diffusion constant of the two reactants. Increased crowding is expected to monotonically decrease the diffusion constant and thus acts to decrease k_D . At the same time crowding also induces an attractive interaction between the reactants, manifesting itself in the enhanced dimerization noted earlier, which acts to partially compensate for the decrease in k_D due to the decrease in diffusion coefficient (88). However, the overall effect is such that generally k_D is expected to decrease with increased crowding (51).

In the reaction- or transition-state controlled regime, the rate constant is dictated by the energy barrier arising from conformational changes necessary before forming the transition-state complex. Because the transition state for association in solution is generally nearly as compact as product complex, crowding is expected to effectively lower the energy barrier and thus increase the association rate constant by a factor that is close to the factor for increase of the equilibrium association constant. Correspondingly, a small effect of crowding on the dissociation rate constant is expected (54). Since fast associations are typically under diffusion control and slow associations under reaction control (4), crowding is generally expected to slow fast associations and accelerate slow associations (51).

Site-binding equilibria (Fig 1B)

To illustrate the effect of crowding on a site-binding reaction (II), let the free ligand be represented as a hard spherical particle ($L = 0$) with radius R_{sc} times that of a hard spherical crowder. When the bound ligand is buried and thus inaccessible to crowders, crowding affects only the free energy of free ligand (i.e., $\Delta F_{\text{LS}}^{\text{crowd}} = 0$). [Footnote 4]. Results obtained using equations (2), (4), and (5) are plotted in Figure 4, and it is evident that site-binding can be significantly more enhanced by crowding effects than simple bimolecular association in solution. The same approximation may also be used to analyze the effect of crowding on protein solubility, which is modeled by assuming that crowding affects protein in the supernatant solution, but not in the condensed phase (42, 49).

Two-state protein folding equilibria (Fig 1C)

The presence of crowder will influence equilibria between conformational states of a macromolecule in favor of conformations that exclude less volume to crowder. In the case of protein folding, unfolded conformations are more expanded and thus crowding is expected to

⁴The assumption that bound ligand excludes no volume to crowder provides an upper bound estimate of the effect of crowding, which may be less to the extent that bound ligand retains mobility and/or solvent exposure.

favor the native state. However, the use of the hard particle model to estimate ΔF_U^{crowd} is unreliable, as the U state consists of a manifold of conformations instead of a single rigid conformation. Thus the extent to which crowding is predicted to stabilize the native state varies substantially between different models for the unfolded state (see following section for examples).

Two-state folding rates

The rate constant for isomerization can be generally written as

$$k = k_0 \exp(-\Delta F^\ddagger / RT) \quad (7)$$

where $\Delta F^\ddagger$ is the activation free energy (i.e., free energy difference between the transition state and the reactant state) and the pre-factor k_0 depends on the dynamics of the isomerization process. Crowding can in principle affect both $\Delta F^\ddagger$ and k_0 . For example, if the transition state is more compact than the reactant state, as in the case of two-state protein folding, then crowding is expected to reduce $\Delta F^\ddagger$. If intra-chain diffusion plays a significant role, crowding may act to reduce the rate of intra-chain diffusion and hence the value of k_0 as well (see review of relevant literature below). When the transition state is more expanded than the reactant state, as in the case of protein unfolding, crowding is expected to decrease the rate constant.

Confinement

Estimation of $\Delta F_X^{confine}$

Consider a molecule X within a unit volume of bulk solution, which can exist in a finite number of configurational states, each of which is specified by a set of positional coordinates denoted by r and a set of orientational coordinates denoted by θ . The free energy of transferring the molecule X from this volume to an equal volume of solution that may be bounded by hard walls in one, two, or three dimensions is given by the statistical-thermodynamic expression (23)

$$\Delta F_X^{confine} = -RT \ln \left(\frac{\iint_{allowed} d\theta dr}{\iint_{all} d\theta dr} \right) \quad (8)$$

Where the multiple integral in the denominator is taken over all configurational states accessible in bulk solution, and the multiple integral in the numerator is taken over all allowed configurational states in the bounded volume, that is, all states in which no part of X intersects any hard wall boundary.

While both confinement and crowding effects result from the reduction in possible configurations available to a macromolecule due to the presence of a high volume fraction of other macromolecules or static barriers to movement, there is one major qualitative difference between the two phenomena. In contrast to the free energy cost of crowding, the free energy cost of confinement is not necessarily minimal for the molecular conformation that is globally the most compact in the sense of having the smallest radius of gyration. Rather, confinement favors conformations having a shape that is complementary to the shape of the confining volume. For example, while a spherical conformation may be favored in a quasi-spherical cavity, the preferred conformation in a cylindrical pore may be rod-like, and the preferred

conformation in a planar pore (bounded by two parallel hard walls) may be plate-like. Thus numerical estimates of the magnitude of confinement effects are extremely sensitive to the choice of models for the structure of both confining space and confined macromolecular species (52).

Association equilibria (Fig. 2A)

Model calculations of the effect of confinement in differently shaped pores upon the association of two spherical monomers of identical size to form a dimer of twice the volume and varying shapes, suggest that confinement has a small effect on bimolecular association, and is expected to increase the equilibrium constant for bimolecular association by at most a factor of two or three. However, the effect of confinement upon association constants for concerted formation of larger n -mers having a shape compatible with the shape of the confining volume increases strongly as the value of n increases (52).

Site binding equilibria (Fig. 2B)

If the bound ligand is completely immobile, then confinement affects only the free energy of the free ligand (i.e., $\Delta F_{LS}^{confine}=0$). As a simple example, consider a spherical ligand with radius a confined in a spherical cavity with radius R_{cavity} . The change in the site binding constant due to ligand confinement may be calculated from equation (2), (4), and (8) to be

$$K_{LS}/K_{LS}^0 = \left(1 - \frac{a}{R_{cavity}}\right)^{-3} \quad (9)$$

When $a = 0.5R_{cavity}$, the binding constant is increased by a factor of 8.

Two-state protein folding equilibria (Fig 2C)

It is evident upon inspection of Fig 2C that the free energy cost of confining any partially or fully unfolded conformation of a protein will be greater than the cost of confining the native state, and that confinement must stabilize the native state relative to any unfolded state. As pointed out previously when summarizing the effects of crowding on protein folding, numerical estimates of the magnitude of the effect of confinement will be quite sensitive to the nature of approximations made in treating the effect of confinement on the (average) unfolded state. Zhou and Dill (90) presented a simple model in which the unfolded state is modeled as a random walk with a specified radius of gyration. They then estimated the value of $\Delta F_U^{confine}$ associated with the placement of U into a variety of confining geometries. Some of their results are plotted in Figure 5. According to this model, confinement is predicted to decrease the value ΔF_{UN} by as much as 20 – 30 RT in very small cavities with a volume only slightly greater than that of the native state. This result makes intuitive sense. In such small cavities it is essentially impossible for proteins to unfold, as there is no space available for proteins to unfold into. By modeling the transition state for protein folding as a sphere with a somewhat larger radius than that of the folded state, a different but conceptually related model (28) has been used to estimate the change in the energy barrier for folding, and calculate the dependence of folding rate on the size of the confining cavity. A maximum acceleration in folding by tuning the cavity size is predicted.

In conclusion, it should be noted that enhancement of association and site binding equilibria by confinement only occurs when the confined macromolecules are truly confined, i.e., they cannot equilibrate with bulk solution. When protein can equilibrate between a pore and bulk

solution the extent of association resulting in the formation of oligomers in the pore is less than in the bulk, and the extent of site binding in the pore equal to that in the bulk (52).

Review of theoretical/simulation literature since 2004

Most of the effects summarized in the preceding section can be predicted on the basis of simple statistical-thermodynamic models. These models have the advantage that they are focused on specific aspects of crowding and confinement, are intuitive, and their results can often be expressed in the form of reasonably simple analytical expressions.

On the other hand, they do not allow for the exploration of complexities associated with crowding and confinement of real macromolecules by other real macromolecules. A complementary theoretical approach is by atomistic simulations. However, one needs to remain cognizant of the fact that the systems being studied via simulation are still models instead of the “real thing”; not all idiosyncratic details of the model systems are of general value for understanding the real systems. For computational efficiency, in most of the atomistic simulation studies, amino acid residues were modeled by a pseudo-atom representing C_α with or without a second pseudo-atom representing the side-chain, and solvent was not treated explicitly.

Effects of crowding on protein folding

Zhou (94) considered the effect of crowding on the free energy of the unfolded state by treating the unfolded protein as a three-dimensional random walk in the presence of hard spherical obstacles. Calculation of the probability that a random walk consisting of a certain number of steps will not encounter a crowding particle leads to a simple relation:

$$\Delta F_U^{crowd}/RT = -\ln(1 - \varphi) + 3\varphi y^2(1 + 2\pi^{-1/2}y^{-1}) \quad (10)$$

where y is the ratio of the radius of gyration of the unfolded chain in bulk solution to the radius of crowder. Equation (10) takes into account the possibility that an unfolded chain can in principle be accommodated within interstitial voids between spherical crowders that may be too small to accommodate a folded protein modeled as a hard particle. When the folded protein is modeled as a hard sphere, the treatment of Zhou predicts that whereas at low volume fraction of crowder, excluded volume effects stabilize the folded state relative to the unfolded state, at very high volume fractions of crowder, excluded volume stabilizes the unfolded state relative to the folded state. It has also been noted that ΔF_U^{crowd} for a self-avoiding chain can be lower than for an ideal chain with the same radius of gyration (89).

Minton (2005) presented an effective two-state model for protein folding, with the unfolded state modeled as a compressible sphere. Because this model allows for intra-molecular as well as intermolecular excluded volume effects, more compact conformations of the unfolded chain are energetically more costly than in the random walk model, and as a result calculated values of ΔF_U^{crowd} are significantly more positive, and calculated values of $\Delta\Delta F_{UN}$ significantly more negative than in the random walk model. This model also predicts the energetic consequences of neglecting intra-molecular excluded volume, and provides estimates of the extent to which the average radius of gyration of an unfolded polypeptide is reduced by addition of hard particle crowders.

Hu et al. (31) modeled the unfolded state as a chain of small tangent hard spheres. They predicted that smaller crowders stabilize the unfolded state relative to the native state, whereas

larger crowders promote the stability of the folded form. This conclusion stands in contradiction to the results of prior theoretical treatments (56,94), as well as experiment (65).

In a simulation study, Cheung et al (11) found that, at a crowder volume fraction of 0.25, the folding stability of a WW domain was increased by 1.1 kcal/mol. Cheung et al. also found that the folding rate initially increases with crowder concentration but decreases at higher crowder concentrations. The decrease in folding rate was attributed to restriction by crowders of conformational fluctuations necessary for protein folding. A second factor may be an increase in the free energy of the transition state due to the decreased probability of finding voids that can accommodate the protein in an expanded transition state.

Effect of crowding on macromolecular isomerization

Using simple space-filling models for both helix and coil, Snir and Kamien (76) predicted that crowding by hard spheres promotes the formation of helical conformations by random-coil polymers.

Miscellaneous crowding effects

Hall (25,26) has modeled the folding of a tracer protein in the presence of crowders which themselves can undergo folding/unfolding transitions or can self-associate. As the folded or self-associated crowders present less excluded volume, the model predicts that the stabilization of the native state by crowders will be decreased as the crowders fold or self-associate.

Effects of confinement on isomerization and protein folding

By modeling the unfolded state as a polymer chain and the native and transition states as hard spheres, Hayer-Hartl and Minton (28) obtained simple analytical expressions describing the dependence of the rate of two-state folding within a spherical cavity upon the radius of the cavity and the molecular weight of the confined protein. These expressions predicted that folding rates would be maximized at an intermediate cavity size that increased with the molecular weight of the encapsulated protein. They were able to quantitatively rationalize variations of the rate of refolding of several proteins within the central cavity of different mutants of the chaperonin GroEL/GroES that were specifically designed to change the volume of the cavity.

Maximization of folding rate at an intermediate cavity size has also been observed in atomistic simulations (12,80,85). These simulations also indicated that folding rate can be modulated by attractive interactions between the confined protein and the walls of the enclosing cavity. In addition, Cheung and Thirumalai (12) found that the folding yield can be increased by as much as 50% by repeated switching on and switching off of an attractive interaction between cavity wall and the confined protein.

A simulation of Jewett et al (37) suggests that the rate of folding in a confined environment can be increased via an alternative pathway in which a folding intermediate is transiently bound to the cavity wall. In another simulation study, Rathore et al. (66) suggested that although confinement-induced stabilization of the native state is dominated by entropy, stabilizing intramolecular interactions are not as optimal as in bulk solution. Net stabilization is hence sequence-dependent and not as great as expected on the basis of entropic effects alone.

Ziv et al (97) simulated the helix formation of peptides confined in an infinite cylinder and found that the helical state is stabilized relative to the coil state. They attempted to rationalize their results with a simple statistical-thermodynamic model, in which the coil state is modeled as a polymer chain, and the helix is modeled by a stiffer polymer chain. Helix stabilization is

predicted, but the predicted stabilization is independent of peptide length, in contrast to their simulation results.

Pande and co-workers (46,78) simulated helix formation inside a cylindrical pore and folding of the villin headpiece inside a spherical cavity. Unlike previous simulations, solvent (water) was treated explicitly in these studies. Contrary to the result of Ziv et al (97), confinement in the cylindrical pore was reported to disfavor helix formation. Confinement in the spherical cavity was reported to favor folding of the protein when the solvent was allowed to equilibrate with the bulk, but disfavor folding when solvent was trapped within the cavity.

Miscellaneous confinement effects

Elcock (18) simulated the co-translational folding of three proteins. The proteins were fed off the peptidyltransferase active site with the ribosome large subunit, which was represented by atoms and pseudo-atoms. No structure formation was observed while the nascent chain was in the ribosome exit tunnel, and the co-translational folding of two single-domain proteins, CI2 and barnase, were found to follow mechanisms identical to those in bulk water. However, for a two-domain protein, Semliki forest virus protein, co-translational folding was found to follow a mechanism different from that in bulk water. In the latter environment, the two domains first folded independently and then docked together. On the ribosome, the N-terminal domain folded first and the structure of the C-terminal domain then gradually accreted onto the pre-formed domain.

Review of the experimental literature since 2004

Relevant experimental literature published during the past four years has been classified according to one of six categories: (1) Effects of crowding on macromolecular association rates; (2) Effects of crowding on macromolecular association equilibria; (3) Effects of crowding on conformational isomerization; (4) Effects of crowding on protein stability with respect to denaturation; (5) Effects of crowding upon enzyme activity; and (6) Effects of confinement on protein stability with respect to denaturation. Noteworthy findings in the six categories are listed in Tables 1–6.

Effect of crowding upon the competition between protein folding and aggregation

It has been known for some time that many partially and fully unfolded proteins exhibit an increased propensity in vitro to form insoluble aggregates, leading to irreversible denaturation (1,13,35). Reference is made to two types of protein stability, namely stability with respect to unfolding, referred to as ‘thermodynamic stability’, and stability with respect to aggregation, referred to as ‘colloidal stability’. These two types of stability are in fact ordinarily interdependent and may be treated separately only under special conditions, such as in the limit of extreme dilution. The close relationship between the two types of stability is due to the similarity on an atomic scale between the kinds of non-covalent interactions that stabilize the native conformation of a protein and those that stabilize intermolecular non-covalent complexes. The difference in free energy between a natively folded protein and a misfolded and/or aggregated protein may be quite small and crowding can shift the thermodynamic balance between the two forms in either direction depending upon details of the folding and aggregation pathways (19,55).

A cautionary note on the use of polyethylene glycol (PEG) as a crowding agent

The effect of macromolecular crowding on a particular reaction is generally studied experimentally by measuring changes in reaction rates or equilibria in the presence of different concentrations of putatively inert macromolecular cosolutes. One of the most popular cosolutes is the highly water-soluble synthetic polymer polyethylene glycol (or polyethylene oxide), which is actually a polyether with monomeric structure ($-\text{CH}_2 - \text{CH}_2 - \text{O} -$). It has long been evident that PEG fractions with molecular weights in excess of a few thousand have a large and predominantly repulsive interaction with proteins, and tend to induce macromolecular associations and compaction in qualitative accord with crowding theory (see for example (36,67) and works cited in Tables 1–5). However, a number of studies have shown that this interaction cannot be described quantitatively in terms of excluded volume alone, and several independent lines of evidence point to an attractive interaction between PEG and nonpolar or hydrophobic sidechains on the protein surface (5,6,50,83,84). Thus the repulsive excluded volume contribution to PEG-protein interaction is partially compensated to an unknown extent by an attractive interaction, the strength of which can vary significantly between different proteins of approximately equal size. A variety of other water-soluble polymers and proteins (e.g. Dextrans, Ficoll, hemoglobin, defatted BSA) have been shown to lack such an attractive interaction for other proteins, and their interactions with proteins can be described using pure excluded volume models (42,50,69,70,73,84). Since these readily available polysaccharides and proteins have the added advantage of resembling more closely the types of macromolecules encountered in a physiological medium, we recommend them as alternatives to PEG as crowding agents for use in quantitative studies.

From *in vitro* to *in vivo*: Addressing additional complications in biological environments

Beyond excluded volume: effects of other types of nonspecific interactions

Theoretical models for estimating the magnitude of crowding and confinement on macromolecular reactions generally assume that these effects are predominantly entropic in origin, i.e., deriving from the relative reduction in configurational entropy of reactants, transition state, and products due to crowding or confinement. Nevertheless, it has been realized from the outset that other nonspecific interactions such as electrostatic repulsion and attraction and hydrophobic attraction are likely to contribute significantly to overall energetics in highly crowded or confined media (50). It has been found that the effects of such interactions upon the colligative and association properties of highly concentrated solutions of a single protein may be satisfactorily accounted for by a simple semi-empirical model in which the protein molecules are treated as effective hard spheres, the apparent size of which reflects short-ranged “soft” attractions or repulsions in addition to steric exclusion (27,58). However, the “effective hard particle model” cannot be expected to satisfactorily describe a medium containing high concentrations of multiple macromolecular species interacting via qualitatively different potentials, such as a solution containing two concentrated proteins that are oppositely charged at a particular pH (27,50). At the present time it appears that Monte Carlo and/or Brownian dynamics simulations provide the most promising approach to the theoretical study of crowding effects in such solutions, which should also be accompanied by additional experimental studies.

The results of experimental studies of the effect of high concentrations of a single species of inert macromolecule (‘crowder’) upon the associations of dilute test proteins have been interpreted in the context of effective hard particle models that take into account nonspecific repulsive interactions deriving from steric exclusion and electrostatic repulsion in as well as attractive interactions leading to association of the test molecules (69,70). It is assumed in these

models that crowder molecules interact with each other only via volume exclusion. One indication that the situation in physiological media may be more complex is provided by a recently published report (87), where concentrated mixtures of protein and polysaccharide were shown to exhibit non-additive effects upon the refolding of lysozyme. Moreover, concentrated solution mixtures of dextran and polyethylene glycol have been found to spontaneously separate into immiscible phases, between which proteins may partition in accordance with their relative affinity for each phase (45). The physical bases of these phenomena deserve closer study, as one would expect the local environment of most biological macromolecules *in vivo* to consist of more than one volume-excluding species.

We cannot yet adequately assess functional consequences of the nearly ubiquitous proximity of soluble proteins to the surfaces of biological structures (e.g., phospholipid membranes and protein fibers). However some directions for future research have been suggested by relevant *in vitro* studies. A variety of proteins have been shown to associate weakly with actin fibers, microtubules, DNA, and phospholipid membranes in a non site-specific fashion (see (10) for references to specific studies). Proteins are localized at the surfaces of these structures by attractive electrostatic and/or hydrophobic interactions in addition to repulsive volume-exclusion (hard wall) interactions. The reduction in configurational entropy of the protein resulting from this dual mode of localization is greater than that achieved by hard wall confinement alone, and in fact magnifies significantly the consequences of confinement. Simple theoretical models (53) predict that adsorbed macromolecules, like hard-wall confined macromolecules, have a stronger tendency to self- or hetero-associate than they do in bulk solution, and that the tendency to associate increases substantially with the strength of attraction between the soluble macromolecule and the surface. Adsorption may also increase the rate of macromolecular binding to specific surface sites (95). A number of macromolecular associations have been reported to proceed more rapidly or to a greater extent on surfaces than in bulk solution (e.g. (29,40,75,95)), suggesting that the consequences of localization via adsorption may be quite general and of potential importance in heterogeneous physiological media.

Interpretation and significance of in-cell measurements of protein stability and association

Experimental techniques have been developed recently that provide information about either the stability and conformation or the association properties of specific labeled proteins within living cells (reviewed in (24)). The use of some of these techniques to study diffusional transport of labeled macromolecules in cytoplasm and tissue is reviewed elsewhere in this volume. In these studies a labeled protein (or pair of labeled proteins) is introduced into a cell via expression of recombinant proteins or microinjection. Then, a signal that reflects conformation or association of labeled protein, or co-localization (hetero-association) of two labeled proteins is monitored. In certain experiments, the intracellular environment is globally perturbed via addition of denaturant, temperature change, or application of hyper- or hypo-osmotic stress (21,32,33,47). The monitored response of the labeled protein(s) within the cell to the applied perturbation is compared to that of the same protein(s) in dilute solution, and conclusions are drawn regarding the effect of the environment *in vivo* upon the stability of the labeled protein(s).

While each of these techniques does indeed provide information about aspects of the behavior of tracer proteins within a cell, one must be very careful about the interpretation of the results of such experiments. The potential impact of labeling procedures upon the spatial distribution of the tracer species, possible induction of artificial associations, and/or disruption of specific interactions of interest must be assessed. A number of additional questions must be answered: 1) How is the test protein distributed among the different local microenvironments within an

intact cell? If it is found in multiple environments, how does one interpret the overall average signal? 2) If a tracer protein that is not native to a host organism such as *E. coli* is highly over-expressed within that organism, how likely is it that the protein is situated within a microenvironment that closely resembles its native milieu? 3) How does the living cell respond to applied stress? Does this response alter the distribution of the test protein, the composition of the microenvironment(s) of the test protein, and the interactions between the test protein and its surroundings within each microenvironment? 4) Since a living cell is a homeostatic system, within which one cannot vary the composition of individual microenvironments in a systematic and quantifiable fashion, how can one determine the extent to which a particular tracer protein in a particular intracellular microenvironment is either confined or crowded?

Toward a narrowing of the gap between “in vitro” and “in vivo”

We suggest that the influence of crowding and confinement upon macromolecular reactivity *in vivo* may be best explored by means of a “bottom-up” approach, according to which the behavior of test proteins is studied quantitatively within a series of media in which features thought to be essential to a particular microenvironment are incorporated in a systematic fashion, from the most simple to the most complex (see for example (96)). Such a bottom-up approach would ultimately lead to construction of a cytomimetic medium incorporating all of the major elements thought to be present in the selected microenvironment. The ability to control and independently manipulate temperature, pH, salt and osmolyte concentration, the types and concentrations of soluble bystander macromolecules and the types and abundances of structural elements such as membranes and cytoskeletal filaments (if appropriate) in this model system provides a rigorous approach to the characterization of nonspecific interactions influencing the behavior of proteins and other macromolecules within a native-like environment. Clearly this is no simple task, but we believe that if our goal is to understand in quantitative terms the role of nonspecific interactions in biology – a role we believe is absolutely essential to the mechanism of life – we cannot avoid paying attention to the details of these inherently complex systems [Footnote 5].

Summary points

1. Macromolecular crowding nonspecifically enhances reactions leading to the reduction of total excluded volume. In general these include the formation of macromolecular complexes in solution, binding of macromolecules to surface sites, formation of insoluble aggregates, and compaction or folding of proteins. The expected magnitude of the effect is strongly dependent upon the relative sizes and shapes of concentrated crowding species and dilute macromolecular reactants and products.
2. Macromolecular crowding is generally expected to increase the rate of slow, transition-state limited association reactions and to decrease the rate of fast, diffusion-limited association reactions.
3. Simple statistical-thermodynamic theories based upon coarse-grained structural models usually provide reliable predictions of qualitative effects, and in favorable circumstances can provide reasonably good semi-quantitative predictions of the magnitude of an expected effect.
4. Biological fluids are more complex than systems studied theoretically or experimentally *in vitro* due to increased heterogeneity and the probable presence of

⁵“It may not be a dream to imagine that, using reconstituted systems of increasing complexity, the coordinated motility of an artificial cell will eventually be mimicked.... However, in many instances the analysis of molecular events using classical biochemical and structural approaches remains the limiting factor for future progress” (9).

nonspecific attractive and repulsive intermolecular interactions in addition to volume exclusion. Theoretical and experimental exploration of model systems containing well-defined elements of added complexity are strongly encouraged.

Acknowledgments

Research of HXZ is supported by NIH Grant GM058187. Research of GR is supported by Grants BFU2005-04807-C02-01 from the Spanish Ministry of Education and S-BIO-0260/2006 from the Madrid Government. Research of APM is supported by the Intramural Research Program, National Institute of Diabetes and Digestive and Kidney Diseases, NIH.

Terms/definitions

Macromolecular crowding	a term used to connote the influence of mutual volume exclusion upon the energetics and transport properties of macromolecules within a solution containing a high total volume fraction of macromolecules
<i>Macromolecular confinement</i>	a term used to connote the influence of excluded volume upon the free energy and reactivity of a macromolecule situated in a cavity, bounded by impenetrable walls, having a smallest interior dimension only slightly larger than the largest dimension of the macromolecule

Literature cited

1. Acampora G, Hermans JJ. Reversible denaturation of sperm whale myoglobin I. Dependence on temperature, pH and composition. *J Am Chem Soc* 1966;89:1543–7. [PubMed: 6039292]
2. Ai X, Zhou Z, Bai Y, Choy WY. 15N NMR spin relaxation dispersion study of the molecular crowding effects on protein folding under native conditions. *J Am Chem Soc* 2006;128:3916–7. [PubMed: 16551092]
3. Alberts B. The cell as a collection of protein machines: preparing the next generation of molecular biologists. *Cell* 1998;92:291–4. [PubMed: 9476889]
4. Alsallaq R, Zhou HX. Energy landscape and transition state of protein-protein association. *Biophys J* 2007;92:1486–502. [PubMed: 17142273]
5. Bhat R, Timasheff SN. Steric exclusion is the principal source of the preferential hydration of proteins in the presence of polyethylene glycols. *Protein Science* 1992;1:1133–43. [PubMed: 1304392]
6. Bloustine J, Virmani T, Thurston GM, Fraden S. Light scattering and phase behavior of lysozyme-poly(ethylene glycol) mixtures. *Phys Rev Lett* 2006;96:0878031–4.
7. Bolis D, Politou AS, Kelly G, Pastore A, Temussi PA. Protein stability in nanocages: A novel approach for influencing protein stability by molecular confinement. *Journal of Molecular Biology* 2004;336:203–12. [PubMed: 14741216]
8. Campanini B, Bologna S, Cannone F, Chirico G, Mozzarelli A, Bettati S. Unfolding of Green Fluorescent Protein mut2 in wet nanoporous silica gels. *Protein Science* 2005;14:1125–33. [PubMed: 15802645]
9. Carlier M-F, Pantaloni D. Control of actin assembly dynamics in cell motility. *J Biol Chem* 2007;282:23005–9. [PubMed: 17576764]
10. Chatelier RC, Minton AP. Adsorption of globular proteins on locally planar surfaces: models for the effect of excluded surface area and aggregation of adsorbed protein on adsorption equilibria. *Biophysical Journal* 1996;71:2367–74. [PubMed: 8913577]
11. Cheung MS, Klimov D, Thirumalai D. Molecular crowding enhances native state stability and refolding rates of globular proteins. *Proceedings of the National Academy of Sciences of the United States of America* 2005;102:4753–8. [PubMed: 15781864]
12. Cheung MS, Thirumalai D. Nanopore-protein interactions dramatically alter stability and yield of the native state in restricted spaces. *Journal of Molecular Biology* 2006;357:632–43. [PubMed: 16427652]

13. Chiti F, Dobson CM. Protein misfolding, functional amyloid, and human disease. *Ann Rev Biochem* 2006;75:333–66. [PubMed: 16756495]
14. Cotter MA. Hard spherocylinders in an anisotropic mean field: a simple model for a nematic liquid crystal. *J Chem Phys* 1977;66:1098–106.
15. del Alamo M, Rivas G, Mateu MG. Effect of macromolecular crowding agents on human immunodeficiency virus type 1 capsid protein assembly in vitro. *Journal of Virology* 2005;79:14271–81. [PubMed: 16254362]
16. Derham BK, Harding JJ. The effect of the presence of globular proteins and elongated polymers on enzyme activity. *Biochim Biophys Acta* 2006;1764:1000–6. [PubMed: 16720113]
17. Diaz-Lopez T, Davila-Fajardo C, Blasies F, Lillo MP, Giraldo R. Early events in the binding of the pPS10 replication protein RepA to the single iteron and operator DNA sequences. *J Mol Biol* 2006;364:909–20. [PubMed: 17045290]
18. Elcock AH. Molecular simulations of cotranslational protein folding: fragment stabilities, folding cooperativity, and trapping in the ribosome. *PLoS Comput Biol* 2006;2:e98. [PubMed: 16789821]
19. Ellis RJ, Minton AP. Protein aggregation in crowded environments. *Biological Chemistry* 2006;387:485–97. [PubMed: 16740119]
20. Fu CY, Morais MC, Battisti AJ, Rossmann MG, Prevelige PE Jr. Molecular dissection of $\Phi 29$ scaffolding protein function in an in vitro assembly system. *J Mol Biol* 2007;366:1161–73. [PubMed: 17198713]
21. Ghaemmaghami S, Oas TG. Quantitative protein stability measurement in vivo. *Nature Struct Biol* 2001;8:879–82. [PubMed: 11573094]
22. Ghahghaei A, Rekas A, Price WE, Carver JA. The effect of dextran on subunit exchange of the molecular chaperone alpha A-crystallin. *Biochimica Et Biophysica Acta-Proteins and Proteomics* 2007;1774:102–11.
23. Giddings JC, Kucera E, Russell CP, Myers MN. Statistical theory for the equilibrium distribution of rigid molecules in inert porous networks: exclusion chromatography. *J Phys Chem* 1968;72:4397–408.
24. Golemis, EA.; Adams, PD. Protein-protein interactions: a molecular cloning manual. Cold Spring Harbor: Cold Spring Harbor Laboratory Press; 2005. p. 938
25. Hall D. Protein self-association in the cell: a mechanism for fine tuning the level of macromolecular crowding? *European Biophysics Journal with Biophysics Letters* 2006;35:276–80. [PubMed: 16217649]
26. Hall D, Dobson CM. Expanding to fill the gap: A possible role for inert biopolymers in regulating the extent of the ‘macromolecular crowding’ effect. *Febs Letters* 2006;580:2584–90. [PubMed: 16647066]
27. Hall D, Minton AP. Macromolecular crowding: qualitative and semiquantitative successes, quantitative challenges. *Biochim Biophys Acta* 2003;1649:127–39. [PubMed: 12878031]
28. Hayer-Hartl M, Minton AP. A simple semiempirical model for the effect of molecular confinement upon the rate of protein folding. *Biochemistry* 2006;45:13356–60. [PubMed: 17073456]
29. Herrig A, Janke M, Austermann J, Gerke V, Janshoff A, Steinem C. Cooperative adsorption of ezrin on PIP₂-containing membranes. *Biochemistry* 2006;45:13025–34. [PubMed: 17059219]
30. Homchaudhuri L, Sarma N, Swaminathan R. Effect of crowding by dextrans and Ficolls on the rate of alkaline phosphatase-catalyzed hydrolysis: a size-dependent investigation. *Biopolymers* 2006;83:477–86. [PubMed: 16868935]
31. Hu Z, Jiang J, Rajagopalan R. Effects of Macromolecular Crowding on Biochemical Reaction Equilibria: A Molecular Thermodynamic Perspective. *Biophys J*. 2007
32. Ignatova Z, Gierasch LM. Monitoring protein stability and aggregation in vivo by real-time fluorescent labeling. *Proc Natl Acad Sci U S A* 2004;101:523–8. [PubMed: 14701904]
33. Ignatova Z, Krishnan B, Bombardier JP, Marcelino AM, Hong J, Gierasch LM. From the test tube to the cell: exploring the folding and aggregation of a beta-clam protein. *Biopolymers* 2007;88:157–63. [PubMed: 17206628]
34. Ittah V, Kahana E, Amir D, Haas E. Applications of time-resolved resonance energy transfer measurements in studies of the molecular crowding effect. *J Mol Recognit* 2004;17:448–55. [PubMed: 15362104]

35. Jaenicke R. What does protein refolding *in vitro* tell us about protein folding in the cell? Philosophical Transactions of the Royal Society of London Series B 1993;339:287–95. [PubMed: 8098533]
36. Jarvis TC, Ring DM, Daube PH, von Hippel PH. “Macromolecular crowding”: thermodynamic consequences for protein-protein interactions within the T4 DNA replication complex. J Biol Chem 1990;265:15160–7. [PubMed: 2168402]
37. Jewett AI, Baumketner A, Shea JE. Accelerated folding in the weak hydrophobic environment of a chaperonin cavity: creation of an alternate fast folding pathway. Proc Natl Acad Sci U S A 2004;101:13192–7. [PubMed: 15331776]
38. Jiang M, Guo ZH. Effects of macromolecular crowding on the intrinsic catalytic efficiency and structure of enterobactin-specific isochorismate synthase. Journal of the American Chemical Society 2007;129:730–1. [PubMed: 17243787]
39. Klimov DK, Newfield D, Thirumalai D. Simulations of beta-hairpin folding confined to spherical pores using distributed computing. Proc Natl Acad Sci U S A 2002;99:8019–24. [PubMed: 12060748]
40. Knight JD, Miranker AD. Phospholipid catalysis of diabetic amyloid assembly. J Mol Biol 2004;341:1175–87. [PubMed: 15321714]
- 40a. Kozer N, Kuttner YY, Haran G, Schreiber G. Protein-protein association in polymer solutions. Biophys J 2007;92:2139–2149. [PubMed: 17189316]
41. Kozer N, Schreiber G. Effect of crowding on protein-protein association rates: fundamental differences between low and high mass crowding agents. J Mol Biol 2004;336:763–74. [PubMed: 15095986]
- 41a. Kuttner YY, Kozer N, Segal E, Schreiber G, Haran G. Separating the contribution of translational and rotational diffusion to protein association. J Am Chem Soc 2005;127:15138–15144. [PubMed: 16248654]
42. Laurent TC. The Interaction between polysaccharides and other macromolecules. The solubility of proteins in the presence of dextran. Biochem J 1963;89:253–7. [PubMed: 14084609]
43. Lebowitz JL, Helfand E, Praestgaard E. Scaled particle theory of fluid mixtures. J Chem Phys 1965;43:774–9.
44. Lei C, Shin Y, Liu J, Ackerman EJ. Synergetic effects of nonoporous support and urea on enzyme activity. Nano Letters 2007;7:1050–3. [PubMed: 17341123]
45. Long MS, Jones CD, Helfrich MR, Mangeney-Slavin LK, Keating CD. Dynamic microcompartmentation in synthetic cells. Proc Natl Acad Sci U S A 2005;102:5920–5. [PubMed: 15788532]
46. Lucent D, Vishal V, Pande VS. Protein folding under confinement: A role for solvent. Proc Natl Acad Sci USA 2007;104:10430–4. [PubMed: 17563390]
47. McNulty BC, Young GB, Pielak GJ. Macromolecular crowding in the Escherichia coli periplasm maintains alpha-synuclein disorder. Journal of Molecular Biology 2006;355:893–7. [PubMed: 16343531]
48. McPhie P, Ni YS, Minton AP. Macromolecular crowding stabilizes the molten globule form of apomyoglobin with respect to bath cold and heat unfolding. Journal of Molecular Biology 2006;361:7–10. [PubMed: 16824541]
49. Minton AP. Excluded volume as a determinant of macromolecular structure and reactivity. Biopolymers 1981;20:2093–120.
50. Minton AP. The effect of volume occupancy upon the thermodynamic activity of proteins: some biochemical consequences. Molecular and Cellular Biochemistry 1983;55:119–40. [PubMed: 6633513]
51. Minton, AP. Holobiochemistry: an integrated approach to the understanding of biochemical mechanism that emerges from the study of proteins and protein associations in volume-occupied solutions. In: Srere, P.; Jones, ME.; Mathews, C., editors. Structural and Organizational Aspects of Metabolic Regulation. N.Y: Alan R. Liss; 1989. p. 291-306.
52. Minton AP. Confinement as a determinant of macromolecular structure and reactivity. Biophys J 1992;63:1090–100. [PubMed: 1420928]

53. Minton AP. Confinement as a determinant of macromolecular structure and reactivity. II. Effects of weakly attractive interactions between confined macrosolutes and confining structures. *Biophysical Journal* 1995;68:1311–22. [PubMed: 7787020]
54. Minton AP. Effects of excluded surface area and adsorbate clustering on surface adsorption of proteins II. Kinetic models. *Biophys J* 2001;80:1641–8. [PubMed: 11259279]
55. Minton AP. Influence of macromolecular crowding upon the stability and state of association of proteins: Predictions and observations. *Journal of Pharmaceutical Sciences* 2005;94:1668–75. [PubMed: 15986476]
56. Minton AP. Models for excluded volume interaction between an unfolded protein and rigid macromolecular cosolutes: Macromolecular crowding and protein stability revisited. *Biophysical Journal* 2005;88:971–85. [PubMed: 15596487]
57. Minton AP. How can biochemical reactions within cells differ from those in test tubes? *Journal of Cell Science* 2006;119:2863–9. [PubMed: 16825427]
58. Minton AP. The effective hard particle model provides a simple, robust, and broadly applicable description of nonideal behavior in concentrated solutions of bovine serum albumin and other nonassociating proteins. *J Pharmaceutical Sci.* 2007 [epub ahead of print]. 10.1002/jps.20964
59. Moran-Zorzano MT, Viale A, Munoz F, Alonso-Casajus N, Eydollin G, et al. Escherichia coli AspP activity is enhanced by macromolecular crowding and by both glucose-1,6-bisphosphate and nucleotide-sugars. *Febs Letters* 2007;581:1035–40. [PubMed: 17306798]
60. Munishkina LA, Cooper EM, Uversky VN, Fink AL. The effect of macromolecular crowding on protein aggregation and amyloid fibril formation. *Journal of Molecular Recognition* 2004;17:456–64. [PubMed: 15362105]
61. Olsen SN. Applications of isothermal titration calorimetry to measure enzyme kinetics and activity in complex solutions. *Thermochimica Acta* 2006;448:12–8.
62. Olsen SN, Ramlov H, Westh P. Effects of osmolytes on hexokinase kinetics combined with macromolecular crowding. *Comp Biochem & Physiol, part A.* 2007;10.1016/j.cbpa.2007.05.009
63. Peterson RW, Anbalagan K, Tommos C, Wand AJ. Forced folding and structural analysis of metastable proteins. *Journal of the American Chemical Society* 2004;126:9498–9. [PubMed: 15291527]
64. Ping GH, Yang GL, Yuan HM. Depletion force from macromolecular crowding enhances mechanical stability of protein molecules. *Polymer* 2006;47:2564–70.
65. Qu Y, Bolen CL, Bolen DW. Osmolyte-driven contraction of a random coil protein. *Proc Natl Acad Sci U S A* 1998;95:9268–73. [PubMed: 9689069]
66. Rathore N, Knotts TA, de Pablo JJ. Confinement effects on the thermodynamics of protein folding: Monte Carlo simulations. *Biophys J* 2006;90:1767–73. [PubMed: 16361344]
67. Reddy MK, Weitzel SE, Daube SS, Jarvis TC, von Hippel PH. Using macromolecular crowding agents to identify weak interactions within DNA replication complexes. *Methods Enzymol* 1995;262:466–76. [PubMed: 8594371]
68. Reiss H, Frisch HL, Lebowitz JL. Statistical mechanics of rigid spheres. *J Chem Phys* 1959;31:369–80.
69. Rivas G, Fernandez JA, Minton AP. Direct observation of the self-association of dilute proteins in the presence of inert macromolecules at high concentration via tracer sedimentation equilibrium: theory, experiment, and biological significance. *Biochemistry* 1999;38:9379–88. [PubMed: 10413513]
70. Rivas G, Fernandez JA, Minton AP. Direct observation of the enhancement of noncooperative protein self-assembly by macromolecular crowding: indefinite linear self-association of bacterial cell division protein FtsZ. *Proc Natl Acad Sci U S A* 2001;98:3150–5. [PubMed: 11248047]
71. Rivas G, Ferrone F, Herzfeld J. Life in a crowded world. *EMBO reports* 2004;5:23–7. [PubMed: 14710181]
72. Roque A, Ponte I, Suau P. Macromolecular crowding induces a molten globule state in the C-terminal domain of histone H1. *Biophys J* 2007;93:2170–7. [PubMed: 17513371]
73. Sasahara K, McPhie P, Minton AP. Effect of dextran on protein stability and conformation attributed to macromolecular crowding. *J Mol Biol* 2003;326:1227–37. [PubMed: 12589765]

74. Sasaki Y, Miyoshi D, Sugimoto N. Regulation of DNA nucleases by molecular crowding. *Nucleic Acids Res.* 2007
75. Smith MI, Sharp JS, Roberts CJ. Nucleation and growth of insulin fibrils in bulk solution and at hydrophobic polystyrene surfaces. *Biophys J* 2007;93:2143–51. [PubMed: 17496011]
76. Snir Y, Kamien RD. Entropically driven helix formation. *Science* 2005;307:1067. [PubMed: 15718461]
77. Snoussi K, Halle B. Protein self-association induced by macromolecular crowding: A quantitative analysis by magnetic relaxation dispersion. *Biophysical Journal* 2005;88:2855–66. [PubMed: 15665132]
78. Sorin EJ, Pande VS. Nanotube confinement denatures protein helices. *J Am Chem Soc* 2006;128:6316–7. [PubMed: 16683786]
79. Spencer DS, Xu K, Logan TM, Zhou HX. Effects of pH, salt, and macromolecular crowding on the stability of FK506-binding protein: An integrated experimental and theoretical study. *Journal of Molecular Biology* 2005;351:219–32. [PubMed: 15992823]
80. Takagi F, Koga N, Takada S. How protein thermodynamics and folding mechanisms are altered by the chaperonin cage: molecular simulations. *Proc Natl Acad Sci U S A* 2003;100:11367–72. [PubMed: 12947041]
81. Tang YC, Chang HC, Roeben A, Wischnewski D, Wischnewski N, et al. Structural features of the GroEL-GroES nano-cage required for rapid folding of encapsulated protein. *Cell* 2006;125:903–14. [PubMed: 16751100]
82. Tokuriki N, Kinjo M, Negi S, Hoshino M, Goto Y, et al. Protein folding by the effects of macromolecular crowding. *Protein Science* 2004;13:125–33. [PubMed: 14691228]
83. Tubio G, Nerli B, Pico G. Relationship between the protein surface hydrophobicity and its partitioning behaviour in aqueous two-phase systems of polyethyleneglycol-dextran. *J Chromatogr B* 2004;799:293–301.
84. Winzor DJ, Wills PR. Molecular crowding effects of linear polymers in protein solutions. *Biophys Chem* 2006;119:186–95. [PubMed: 16129549]
85. Xu WX, Wang J, Wang W. Folding behavior of chaperonin-mediated substrate protein. *Proteins* 2005;61:777–94. [PubMed: 16231331]
86. Yamin G, Munishkina LA, Karymov MA, Lyubchenko YL, Uversky VN, Fink AL. Forcing nonamyloidogenic beta-synuclein to fibrillate. *Biochemistry* 2005;44:9096–107. [PubMed: 15966733]
87. Zhou BR, Yi L, Fen D, Zheng Z, Jie C. Mixed macromolecular crowding accelerates the oxidative refolding of reduced, denatured lysozyme - Implications for protein folding in intracellular environments. *Journal of Biological Chemistry* 2004;279:55109–16. [PubMed: 15494409]
88. Zhou H-X. Protein folding and binding in confined spaces and in crowded solutions. *J Mol Recognit* 2004;17:368–75. [PubMed: 15362094]
89. Zhou H-X. Protein folding in confined and crowded environments. *Arch Biochem Biophys.* 2007 in press.
90. Zhou H-X, Dill KA. Stabilization of proteins in confined spaces. *Biochemistry* 2001;40:11289–93. [PubMed: 11560476]
91. Zhou H-X, Szabo A. Theory and simulation of the time-dependent rate coefficients of diffusion-influenced reactions. *Biophys J* 1996;71:2440–57. [PubMed: 8913584]
92. Zhou H-X. Loops, linkages, rings, catenanes, cages, and crowders: Entropy-based strategies for stabilizing proteins. *Accounts of Chemical Research* 2004;37:123–30. [PubMed: 14967059]
93. Zhou H-X. Polymer models of protein stability, folding, and interactions. *Biochemistry* 2004;43:2141–54. [PubMed: 14979710]
94. Zhou H-X, Szabo A. Enhancement of association rates by nonspecific binding to DNA and cell membranes. *Phys Rev Lett* 2004;93:1781011–4.
95. Zhu M, Souillac PO, Ionescu-Zanetti C, Carter SA, Fink AL. Surface-catalyzed amyloid fibril formation. *J Biol Chem* 2002;277:50914–22. [PubMed: 12356747]
96. Zimmerman SB, Trach SO. Estimation of macromolecule concentrations and excluded volume effects for the cytoplasm of *Escherichia coli*. *J Mol Biol* 1991;222:599–620. [PubMed: 1748995]

97. Ziv G, Haran G, Thirumalai D. Ribosome exit tunnel can entropically stabilize alpha-helices. *Proc Natl Acad Sci U S A* 2005;102:18956–61. [PubMed: 16357202]
98. Zorrilla S, Rivas G, Acuna AU, Lillo MP. Protein self-association in crowded protein solutions: A time-resolved fluorescence polarization study. *Protein Science* 2004;13:2960–9. [PubMed: 15459331]

Figure 1.

Thermodynamic cycles illustrating linkage between free energy of transfer of reactants and products from dilute solution to crowded medium and standard free energy of (A) association in solution, (B) site-binding, and (C) two-state folding of a protein.

Figure 2.

Thermodynamic cycles illustrating linkage between free energy of transfer of reactants and products from dilute solution to confined volume element and standard free energy of (A) association in solution, (B) site-binding, and (C) two-state folding of a protein.

Figure 3.

Effect of volume fraction ϕ of hard sphere crowders upon equilibrium constant for association of two spherical monomers with $R_1 = 1$ to form a spherocylindrical dimer with twice the volume, and (a) $L_2 = 0$, $R_2 = 1.26$ (spherical dimer); (b) $L_2 = 2/3$, $R_2 = 1$; (c) $L_2 = 1$, $R_2 = 0.928$; (d) $L_2 = 1.5$, $R_2 = 0.851$; and (e) $L_2 = 2$, $R_2 = 0.794$.

Figure 4. Effect of volume fraction ϕ of hard sphere crowders upon equilibrium constant for binding of a spherical macromolecular ligand to an immobile surface site. $MW_{\text{ligand}}/MW_{\text{crowder}} = 2$ (curve a), 1 (curve b), and 0.5 (curve c).

Figure 5.

Effect of confinement in a spherical cavity upon the free energy of two-state folding, calculated as a function of cavity radius according to the theory of Zhou and Dill (90), for proteins containing $N = 100$ residues (a) and 200 residues (b). Vertical lines represent the radii of the hard spherical representation of the folded proteins, calculated according to $r_N (\text{\AA}) = 3.73 N^{1/3}$. The radius of gyration of the unfolded protein is calculated according to $r_g (\text{\AA}) = 3.27 N^{1/2}$.

Table 1

Reported effects of macromolecular crowding on association rates

Addition of proteins, polysaccharides (dextran 138 K, Ficoll 70 K) and PEG (ranging in molar mass from 0.2 to 3.3 K) increases the rate of amyloid fibrillation by alpha-synuclein (60). ~10-fold acceleration in 150 g/L of PEG 3.3 K and ~3-fold in 150 g/L Ficoll. ~5-6-fold acceleration in 50 g/L lysozyme or BSA.
Addition of PEG 10 K induces the fibrillation of beta-synuclein in the presence of Zn2+ (86). Both PEG and Zn2+ are required for fiber formation.
Addition of Ficoll 70 K accelerates HIV capsid protein self-assembly (15). The half-time for protein assembly is decreased ~10-fold in 100 g/L Ficoll.
Addition of PEG 3.3 K induces the co-assembly of the bacteriophage Φ29 monomeric capsid protein and dimeric scaffolding protein to form bonafide capsid particles (20). In the absence of the scaffolding protein, the capsid protein forms amorphous polymers in PEG.
Addition of Ficoll 70 K and PEG (ranging in molar mass from 0.2 to 8 K) reduces the second order rate constant for diffusion-limited bimolecular association of beta-lactamase (TEM) and beta-lactamase inhibitor protein (BLIP) (40a,41,41a). At low crowder concentration, rate scales with rotational diffusion. At higher concentrations, rate decreases more strongly with increasing polymer concentration, possibly due to crowding-induced self-association of reactant species.

Table 2

Reported effects of macromolecular crowding on association equilibria

<i>Addition of the unrelated protein RNase A promotes the dimerization of tracer apomyoglobin (98). Significant effects are observed at > 50 g/L RNase. In contrast, an equivalent mass concentration of HSA did not promote apoMb dimer formation; this differential is much larger than expected from excluded volume models.</i>
<i>Addition of dextran 10 K enhances formation of a decamer of bovine pancreatic trypsin inhibitor (77). ~30-fold increase in decamer fractional abundance and ~5×10⁵-fold increase in the association constant at ~ 200 g/L of dextran.</i>
<i>Addition of BSA or ovalbumin increases the binding affinity of replication protein RepA for specific DNA sequences (17). ~10-fold increase of the equilibrium association constant for RepA-DNA complex formation in 150 g/L BSA.</i>
<i>Addition of dextran 70 K inhibits the exchange of subunits between aggregates of alpha-crystallin. (22) Effect can be attributed to crowding-induced lowering of concentration of monomeric subunit in equilibrium with aggregate.</i>

Table 3

Reported effects of macromolecular crowding on conformational isomerization

<p>Addition of dextran reduces mean distance between residues 169 and 203 in partially unfolded adenylate kinase (34). <i>Inter-residue distance, measured by time resolved FRET, is reduced from 39 Å in 1 M GuHCl to 34 Å in 1 M GuHCl + 270 g/L dextran. Inter-residue distance in native conformation (no GuHCl) is 29 Å.</i></p> <p>Addition of Ficoll 70 K or PEG 20000 shifts near-UV and far-UV CD spectra of RNase A in 2.4 M urea to that characterizing native RNase A (82). <i>Essentially complete transition of spectrum achieved at ca 35 wt-% added polymer</i></p> <p>Addition of Ficoll 70 K or PEG 20K significantly decreases the hydrodynamic radius of partially unfolded RNase A in 2.4 M urea, but does not affect the hydrodynamic radius of native protein (82). <i>Decrease of ca 25% in presence of 35 wt-% polymer</i></p> <p>High concentrations of macromolecules in vitro and in the periplasm of <i>E. coli</i> stabilize compact conformations of α-synuclein relative to expanded conformations (47). <i>Conformational transition from more compact at 10°C to less compact at 35°C in dilute solution is suppressed in 300 g/L BSA or in periplasm of intact E. coli cells.</i></p> <p>Addition of PEG 20K partially restores enzymatic activity of RNase A in 2.4 M urea (82). <i>Enzyme activity increases from 8% of native in the absence of polymer to 26% in the presence of 30 wt-% polymer</i></p> <p>Pulling force required to unfold ubiquitin increases by 20% in the presence of 300 g/L dextran 40K (64). <i>Observed effect is ca 10-fold greater than calculated via excluded volume model</i></p> <p>Addition of 30 wt-% Ficoll 70K or PEG 6K induces conversion of the unstructured C-terminal domain of histone H1 to a molten globule (72).</p>

Table 4
Reported effects of macromolecular crowding on protein stability with respect to denaturation

Addition of Ficoll 70K increases the free energy of unfolding of FK506-binding protein (79). $\Delta G_{N,U}$ increases by ca RT upon addition of 160 g/L dextran. Two-state transition verified. Magnitude consistent with prediction of excluded volume model (88).
Addition of dextran 30K stabilizes the molten globule conformation of apomyoglobin at pH 2 with respect to heat and cold-induced unfolding (48). T_{50} for cold denaturation reduced by $\sim 13^{\circ}\text{C}$ and T_m for heat denaturation raised by $\sim 20^{\circ}\text{C}$ upon addition of 270 g/L dextran. Results in qualitative agreement with prediction of excluded volume theory.
Refolding rate of Rd-apocyt b_{562} increases by 30% at 30° and by 80% at 20° in the presence of 85 g/L PEG 20K (2). Unfolding rate little affected. Measurement via ^{15}N NMR spin-relaxation dispersion
Addition of PEG 4K increases T_m for thermal denaturation of DNase I (74). T_m increases by $> 15^{\circ}$ in 20 wt-% PEG

Table 5

Reported effects of macromolecular crowding on enzyme activity

Enzymatic activity increases, then decreases with increasing concentration of protein crowding agents, but decreases monotonically with increasing concentration of polymeric crowding agents (16). <i>Up to 10-fold increase in specific activity of urease in 30 wt-% hemoglobin</i>
Addition of PEG 6K increases enzyme activity of <i>E. coli</i> AspP (59). <i>50 g/L PEG decreases K_m 4-fold and increases V_{max} 6-fold</i>
Addition of dextran 70K, Ficoll 70K or PEG 6K increases enzyme activity of isochorismate synthase (38). <i>At 25 wt-% of additive, K_m decreases 2–3 fold.</i>
Addition of dextrans (15 – 500K) or Ficoll 70K reduces rate of hydrolysis catalyzed by alkaline phosphatase (30). <i>Larger dextrans have a larger effect per unit w/v concentration. 20 wt-% dextran 500K decreases rate ~7-fold while 20 wt-% dextran 15K decreases rate ~2-fold. Effect attributed largely to reduction in rate of enzyme-substrate encounter.</i>
Specific activity of hexokinase reduced in high concentrations of BSA (61). <i>At 250 g/L BSA, k_M decreases ~33%, k_{cat} decreases ~25%. Reaction rates measured calorimetrically</i>
Effects of small osmolytes and high concentrations of BSA on hexokinase activity are additive (62). <i>Consistent with hypothesis that osmolytes (except urea) interact primarily with proteins via volume exclusion</i>
Addition of PEG 4 – 20K enhances activity of DNase I and S1 nuclease, does not significantly affect activity of exonuclease III, and decreases activity of exonuclease I (74). <i>Polymer does not affect K_m but increases V_{max} of DNase I ~20-fold in 20 wt-% PEG.</i>

Table 6
Reported effects of macromolecular confinement on protein stability and conformation

Confinement in polyacrylamide gels increases T_m of several proteins (7). T_m increases with increasing polyacrylamide concentration in gel. 1.5 – 5° increase in 20% gels.
Confinement of glucose isomerase in functionalized mesoporous silica enhances retention of specific activity at high urea concentrations (44). In ca 300 Å pores, enzyme activity in 8 M urea is equal to that in the absence of urea in bulk solution.
Confinement of GFP-mut2 in silica gel increases free energy of unfolding by GuHCl (8). ~ 3RT increase at 37°C
Structure of mutationally destabilized protein that is disordered in bulk solution reverts to native when protein is encapsulated in reverse micelles (63).
Confinement of glucose isomerase in functionalized mesoporous silica + addition of 0.4 M urea increases specific activity (44). 50% enhancement without covalent crosslinking to silica, 80% enhancement with covalent crosslinking
Confinement of several proteins in the cavity of wild-type and mutant forms of GroEL/GroES accelerates or decelerates rate of refolding in a size-dependent manner (81). Refolding is accelerated in larger cavities and decelerated in smaller cavities, in qualitative or semi-quantitative agreement with predictions of confinement simulations and models (28,39,80)