Case report

Multiple localization of granular cell tumour: a case report

Giovanni Francesco Marangi*, Vito Toto, Igor Poccia, Pierluigi Giglio Fiorito, Beniamino Brunetti and Paolo Persichetti

Address: Department of Plastic and Reconstructive Surgery, University Campus Bio-Medico of Rome, Via Alvaro del Portillo, 00128 Rome, Italy

Email: GFM* - g.marangi@unicampus.it; VT - v.toto@unicampus.it; IP - i.poccia@unicampus.it; PG - p.gigliofiorito@unicampus.it; BB - b.brunetti@unicampus.it; PP - p.persichetti@unicampus.it

* Corresponding author

Received: 23 July 2009 Accepted: 5 August 2009 Published: 8 September 2009

Cases Journal 2009, 2:8751 doi: 10.4076/1757-1626-2-8751

This article is available from: <http://casesjournal.com/casesjournal/article/view/8751>

© 2009 Marangi et al.; licensee Cases Network Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Introduction: Granular cell tumour, also known as Abrikossoff's tumour, is a rare entity occurring in the skin as well as in internal organs, more common among the third to fifth decade of life. It has often been described as a peripheral neuroectodermal tumour and its clinical behaviour is usually benign although malignant and multifocal forms are also known to occur.

Case presentation: We report a case of multiple granular cell tumour in a 17-year-old Caucasian woman who presented with a nodular lesion in the popliteal cave, diagnosed as an Abrikossoff's Tumour, followed 2 years later by another soft tissue mass in the neck.

Conclusion: This case is quite rare: only a few cases were presented in literature with an extended period of time before the clinical presentation of subsequent disease.

Introduction

Granular cell tumour (GCT) is an uncommon soft tissue neoplasm, first described by Abrikossoff in 1926 [1]. Females are slightly more affected than males and it is common in the third to the fifth decade of life [2,3]. It can also occur in childhood but this is rare. These neoplasms arise in different parts of the body, particularly in the tongue, subcutaneous tissue and breast. The head and neck site most likely to be involved is the larynx. Other sites such as the gastrointestinal, urogenital, and respiratory tracts can also be involved. Granular cell tumours usually present as a benign solitary nodule. Malignant granular cell tumours (MGCTs) are extremely rare, high-grade sarcomas of Schwann cell origin. They often metastasize and are associated with short survival. Clinical

manifestations of granular-cell tumours may vary from showing normal, hyperpigmented or ulcerated skin associated with frequent subcutaneous nodules. They can also be either pruritic or painful. Initially, they were thought to arise from skeletal muscle because of the cytologic resemblance to myocytes. Recent studies suggest its derivation from Schwann cells.

We report a rare case of GCT in a young woman which first presented with a nodular lesion in the left popliteal cave and two years later showed a neck subcutaneous nodule.

Case presentation

A 17-year-old Caucasian woman presented to our department with a palpable lesion in the left popliteal cave.

Family history and past medical history were not of significant interest. The lesion appeared 9 months as a 0.5 cm mass, and had slowly enlarged, evolving into a 2 × 1 cm mobile subcutaneous nodule, with clinically distinct margins. The tumour was covered by apparently normal skin. At first the lesion was clinically suspected to be a fibrolipoma and so a simple surgical excision was performed. Histological examination revealed that large polyhedral cells arranged in sheets, with eosinophilic granular cytoplasm. Small oval nuclei were eccentrically placed. Mitotic activity was < 2/10 high power field (HPF). Histological features were suggestive of granular cell tumour. Immunohistochemical study of the nodule was positive for S100 protein, CD68 and p53 in less than 30% of cells. Surgical margins were positive for tumour involvement so a second radical excision including the skin and the subcutaneous tissue was performed. Second histopathological examination assured total excision (Figures 1 and 2).

Approximately 2 years later, a second nodule appeared on her neck. Physical examination revealed a 2 cm subcutaneous, mobile, painless nodule. The lesion gradually enlarged within 6 months. The surrounding skin was normal. Total excision of the nodule was performed. The histopathological examination was identical to the previous specimen showing nests of polygonal cells with abundant eosinophilic granular cytoplasm. There was no cytologic pleomorphism and mitotic activity was >2/10 HPF. The neoplasm was positive for S100 protein and CD68. The patient was discharged on post-operative day 2 with no complication reported. The surgical wound healed in 2 weeks with normal scarring.

Figure 1. Histological section stained with hematoxylin & eosin (X 20).

Figure 2. Histological section stained with hematoxylin & eosin (X 40).

In a two years follow up, the patient is in good health without signs of further tumour development.

Discussion

Granular cell tumours are usually solitary and usually located in the skin or the submucosa of the tongue. The majority of cases involve the head and neck [4,5]. Less frequently involved is the upper respiratory tract [6] and in 10-15% of cases the gastrointestinal tract [7]. Other rare locations include thyroid, breast [8,9], parotid glands, eyes, urinary bladder [10], reproductive organs, abdominal wall [11], cranial and peripheral nerves. It has been also described a correlation with neurofibromatosis, in which multiple subcutaneous nodules are present, especially in children. The neoplasm is typically slow growing, well circumscribed, firm and rounded, with a diameter ranging from 5 to 20 mm, although larger tumours may be seen. The skin surrounding the tumour is usually normal, although it may thicken and rarely may ulcerate. Occasionally tufts of hair may be present on the skin overlying the lesion [12].

The benign tumour consists of large polyhedral cells arranged in sheets with abundant eosinophilic granular cytoplasm. The nuclei are relatively small and mildly pleomorphic with prominent nucleoli. Occasional mitotic figures can be present. The epithelium surrounding the neoplasm often shows pseudoepitheliomatous hyperplasia.

Approximately 1-2 percent of granular-cell tumours are malignant [13,14]. Features suggestive of malignancy include a size greater than 4 cm in diameter, nuclear pleomorphism, lymph nodes metastases, aggressive

clinical behaviour, rapid growth, and ulceration. Histopathological diagnosis of malignancy is based on these six criteria: necrosis, spindling, vesicular nuclei with large nucleoli, increased mitotic activity (> 2 mitoses/10 high-power fields at 200x magnification), high nuclear to cytoplasmic (N:C) ratio, and pleomorphism. Neoplasms that meet three or more of these criteria are classified as malignant, those that meet one or two criteria are classified as atypical, and those that display only focal pleomorphism are classified as benign [14].

Due to low mitotic activity ($< 2/10$ high power field), not significant nuclear to cytoplasmic (N:C) ratio and no pleomorphism detected, we graded both lesions as benign.

Sites of metastases are mainly regional lymphatic nodes, brain tissue, skeleton and lungs.

Initially Abrikosoff suggested that GCT originate from skeletal muscle cells and so referred to them as myoblastomas; this suggestion probably arose from examination of tumours of the tongue in which infiltration between the striated muscle bundles gave the impression of a muscular origin. Recent studies suggest derivation from Schwann cells of the peripheral nerves, and the presence of S100 protein supports this [15]. The KP-1 monoclonal antibody, which recognises the lysosome-associated glycoprotein CD68, reacts positive with schwannomas and granular cell tumours but not with other neuronal neoplasms [16].

Conclusion

The presentation of the second lesion in our patient makes this case report quite distinctive. Few cases were previously described with a second presentation such temporally far from the primary neoplasm.

Even if 10-15% of granular cell tumours present with satellite nodules, in our case clonality was excluded due to the different topographic involvement occurred.

Crawford and De Bakey [17] reported a patient in whom metastasis occurred 14 years after initial treatment. Mullins and Magner [18] reported a case of a woman who first had a right flank subcutaneous mass, diagnosed as granular cell myoblastoma, and then developing 2 years later a mass in the contralateral parotid gland. Our patient was free from clinical disease for approximately 2 years before the development of the second neoplasm. The histopathological examination of the second lesion was identical to the previous specimens with no features suggestive of malignancy.

The treatment of choice for GCT is wide local excision. In the malignant variety, the treatment should include wide

local excision with regional lymph node dissection and radiological evaluation for metastasis. Radiotherapy and chemotherapy are used with variable success, but their effectiveness remains unproven [12].

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

GFM and PP performed the surgical procedures and contributed to the analysis of the clinical data. BB performed the histological examination of the lesion. VT and PG contributed to the interpretation of the data and to the discussion of the manuscript. VT was a major contributor in writing the manuscript together with IP. All authors read and approved the final manuscript.

References

1. Abrikosoff AI: **Über Myome, ausgehend von der quergestreifter willknerlicher Muskulatur.** *Virchows Arch Pathol Anat* 1926, **260**:214-233.
2. Chaudry AP, Jacobs MS, SunderRaj M, Yamane GM, Jain R, Scharlock SE: **A clinico-pathological study of 50 adult oral granular cell tumours.** *J Oral Med* 1984, **39**:97-103.
3. Vuyk HD, Snow GB, Tiwari RM, Van Velzen R, Veldhuizen RW: **Granular cell tumor of the proximal oesophagus.** *Cancer* 1985, **55**:445-449.
4. Strong EW, McDivitt RW, Brasfield RD: **Granular cell myoblastoma.** *Cancer* 1970, **25**:415-422.
5. Peterson LJ: **Granular cell tumour.** *Oral Surg* 1974, **37**:728-735.
6. Sataloff RT, Ressue JC, Portell M, Harris RM, Ossoff R, Merati AL, Zeitels S: **Granular cell tumor of the larynx.** *J Voice* 2000, **14**:119-134.
7. Grande M, Tosato F, De Marchi C, Cassese M, Persichetti P, Paganini A, Paolini A: **Granular cell myoblastoma of the esophagus. Abrikosov's tumor.** *Recenti Prog Med* 1984, **75**:348-352.
8. Chetty R, Kalan MR: **Malignant granular cell tumor of the breast.** *J Surg Oncol* 1992, **49**:135.
9. Patel A, Lefemine V, Yousuf SM, Abou-Samra W: **Granular cell tumour of the pectoral muscle mimicking breast cancer.** *Cases J* 2008, **1**:142.
10. Yoshida T, Hirai S, Horii Y, Yamauchi T: **Granular cell tumor of urinary bladder.** *Int J Urol* 2001, **8**:29-31.
11. Chaudhry A, Griffiths EA, Shah N, Ravi S: **Surgical excision of an abdominal wall granular cell tumour with Permacol mesh reconstruction: a case report.** *Int Semin Surg Oncol* 2008, **5**:4.
12. Qureshi NA, Tahir M, Carmichael AR: **Granular cell tumour of the soft tissues: a case report and literature review.** *International Seminars in Surgical Oncology* 2006, **3**:21.
13. Shimamura K, Osamura RY, Ueyama Y, Hata J, Tamaoki N, Machida N, Fukuda H, Uemura K: **Malignant granular cell tumor of the right sciatic nerve.** *Cancer* 1984, **53**:524-529.
14. Fanburg Smith JC, Meis Kindblom JM, Fante R, Kindblom LG: **Malignant granular cell tumor of soft tissue. Diagnostic criteria and clinicopathologic correlation.** *Am J Surg Pathol* 1998, **22**:779-794.
15. Weber-Cappuis K, Wildmann JJ, Kapanci Y: **Histological and immunohistological profile of benign granular cell tumor. A report of 41 cases.** *Ann Pathol* 1995, **15**:198-202.

16. Kurtin PJ, Bonin DM: **Immunohistochemical demonstration of the lysosome-associated glycoprotein CD68 (KP-1) in granular cell tumors and schwannomas.** *Hum Pathol* 1994, **25**:1172-1178.
17. Crawford ES, De Bakey ME: **Granular cell myoblastoma: two unusual cases.** *Cancer* 1953, **6**:786.
18. Mullins GM, Magner JW: **Multiple granular cell myoblastomas: a case report.** *Irish Medical J* 1984, **77**:286-287.

Do you have a case to share?

Submit your case report today

- Rapid peer review
- Fast publication
- PubMed indexing
- Inclusion in Cases Database

Any patient, any case, can teach us something

**CASES
NETWORK**

www.casesnetwork.com