Published in final edited form as:

Bioorg Med Chem. 2010 January 15; 18(2): 922–929. doi:10.1016/j.bmc.2009.11.034.

Perchlorotriptyl Radical-Fluorophore Conjugates as Dual Fluorescence and EPR Probes for Superoxide Radical Anion

Jinhua Wang^b, Vinh Dang^a, Wei Zhao^d, Dongning Lu^b, Brian K. Rivera^a, Frederick A. Villamena^c, Peng George Wang^b, and Periannan Kuppusamy^a

^aDepartment of Internal Medicine, Davis Heart and Lung Research Institute, The Ohio State University, Columbus, OH 43210, USA

^bDepartment of Chemistry, Davis Heart and Lung Research Institute, The Ohio State University, Columbus, OH 43210, USA

^cDepartment of Pharmacology, Davis Heart and Lung Research Institute, The Ohio State University, Columbus, OH 43210, USA

^dCollege of Pharmacy, Nankai University, Tianjin 300071, China

Abstract

Perchlorotriptyl radicals, monosubstituted with a fluorophore using an amide linker of varying chain length, were synthesized and characterized. Electron paramagnetic resonance (EPR) spectroscopic study indicated free-electron coupling with the aromatic hydrogen nuclei and long-range coupling with the methylene hydrogens of the linker group. Reactivity of the fluorophore-conjugated triptyls with superoxide radical anion showed quenching of EPR signal and enhancement of fluorescence emission spectrum. This work presents the first example of a perchlorotriptyl-fluorophore conjugate that can potentially be employed as a dual probe for the detection of superoxide under oxidative stress-mediated conditions in biological systems.

1. Introduction

Superoxide radical anion ($O_2^{\bullet-}$), a reactive free radical species formed by one-electron reduction of molecular oxygen, has been implicated in a variety of disease conditions such as ischemia-reperfusion injury, diabetes, atherosclerosis, chronic inflammation, viral infection, and malignant diseases.¹ Superoxide produced by mammalian cells serve as a signaling molecule that can modulate events such as enzyme phosphorylation, cell growth, hypertrophy and apoptosis,^{2,3} but increased $O_2^{\bullet-}$ production can lead to cellular dysfunction. Therefore, there is a need for improved techniques for the detection and quantification of $O_2^{\bullet-}$ in order to gain knowledge on its role in the progression of oxidative stress-related diseases and pathophysiological conditions.

A variety of analytical techniques have been developed to detect and quantify $O_2^{\bullet-}$ radicals in biological systems. These include spectrophotometric methods that use reduction of ferricytochrome *c* or nitroblue tetrazolium (NBT) by $O_2^{\bullet-}$, chemiluminescence, fluorescence-

Address for correspondence: Periannan Kuppusamy PhD, 420 West 12th Ave, Room 114, Ohio State University, Columbus, OH 43210. Kuppusamy.1@osu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

based techniques and electron paramagnetic resonance (EPR) spectroscopy using spin-trapping techniques.⁴⁻⁷ However, these methods have certain limitations for measuring $O_2^{\bullet-}$. In the lucigenin-based luminescence detection of $O_2^{\bullet-}$, it has been demonstrated that the probe itself can redox-cycle, resulting in artifactual production of $O_2^{\bullet-}$, whereas the cytochrome c reduction assay is affected by the presence of oxidoreductases in cellular systems.^{8,9} The EPR method requires the use of high concentrations (10 - 200 mM) of spin trap that may perturb the redox balance in the biological system.

Superoxide requires a probe that is sensitive enough to efficiently compete with intracellular components for its detection. This is due to a relatively short half-life of the molecule and the presence of endogenous intracellular antioxidant enzymes that efficiently scavenge $O_2^{\bullet-}$. Additionally, these probes must have target specificity to sub-cellular compartments to reflect radical production at their site of formation. One of the most promising approaches to monitor the redox status or the detection of free radicals in cells is the use of bi-functional fluorogenic spin probes that can be detected by fluorescence and/or EPR spectroscopy. In the past, such dual probes containing a fluorescent chromophore and a paramagnetic nitroxide radical have been reported.¹⁰⁻¹³ In these molecules, the paramagnetic nitroxide moiety acts as a quencher of the fluorescence of the chromophore fragment. Reduction of the nitroxide by free radicals results in the formation of diamagnetic moiety and hence can cause decay of the EPR signal and in turn, will potentiate fluorescence. Hence, the conjugate allows all the nitroxide spin probe features to be retained, but also may gain new advantages by being able to monitor redox and radical production processes by a very sensitive and simpler fluorescence technique. However, the chemical property of nitroxide limits its application for intracellular radical detection due to its susceptibility to be metabolically reduced in the cell. Thus, a highly EPR sensitive (narrow/single line) probe with good biostability against cellular oxidoreductants and high specificity for $O_2^{\bullet-}$ is desirable for radical detection in biological systems.

Recently, triarylmethyl (trityl) radicals have been reported as probes for detecting $O_2^{\bullet-}$ in biological systems.^{14,15} Perchlorotriphenylmethyl radicals are a novel class of trityls whose structures are based on perchlorotriphenyl methyl radical (PTM). Due to the full steric blockage of the central carbon, the PTM radical and its derivatives are highly stable against a variety of reactive chemical agents, and hence are called "inert free radicals".^{16,17} The ability of two derivatives, tris(2,3,5,6-tetrachloro-4-carboxylate-phenyl) methyl radical (PTM-TC) and tris(2,3,5,6-tetrachloro-4-ethyloxycarbonyl-phenyl) methyl radical (PTM-TE), to trap $O_2^{\bullet-}$ have been tested.^{18,19} These species demonstrated enhanced stability in biological systems, excellent water solubility (in the cases of carboxylic acid), and a narrow, single EPR signal. Therefore, the development of trityl-fluorophore conjugate is proposed which may offer improved $O_2^{\bullet-}$ detection by using EPR and/or fluorescence techniques.

2. Results and Discussion

2.1 Synthesis

Due to the steric hindrance and high stability of tris(2,3,5,6-tetrachlorophenyl)methane, **1**, it is difficult to directly derivatize the compound with a chromophore. Instead, our initial efforts involved adding a linker with appropriate functional group between the trityl core structure and the chromophore. Treatment of **1** with *n*-butyl lithium at -78°C followed by nucleophilic addition of the trityl to ethylene oxide, afforded compound **2** with spacer(s) incorporated (Scheme 1). By controlling the molar ratios of **1**, ethylene oxide and base, compounds **2a** or **2b** can be isolated as major products. However, only small amounts of compound **2c** were formed even when six equivalents of ethylene oxide and *n*-butyl lithium were added. It was found that compounds **2a** and **2b** have good solubility in THF and dichloromethane, while compound **2c** was insoluble in any commonly-used solvents. The solubility differences in compounds **2a-2c** can control the equilibrium concentrations of these three compounds in

solution. The primary amine, **4**, was synthesized via Mitsunobu reaction using **2a** and phthalimide followed by hydrazinolysis, with sufficiently high purity for use in the succeeding step according to previously established procedure.²⁰ Two different linker distances were introduced to compounds **4** in order to shed insights into how these distances can affect the chromophore-radical interaction. Amidation reaction of **4** with both 1-pyrenecarboxylic acid *N*-hydroxysuccinimide ester and pyrenebutyric acid *N*-hydroxysuccinimide ester was carried on to form fluorescence chromophore-conjugated compounds **5** and **6**, respectively.²¹ Finally, radicals **7** and **8** were obtained by treating compounds **5** or **6** with excess of $n\text{Bu}_4\text{N}^+\text{OH}^-$ and subsequent oxidation of the resulting anions with *p*-chloranil in a one-pot reaction with near quantitative yield.²²

Two methods were employed for the synthesis of **10** (Scheme 2). In Method A, treatment of **1** with 2.4 equivalents of *n*-BuLi and TMEDA followed by 2.4 equivalents of ethyl chloroformate afforded di-ethoxycarbonyl trityl **9** in 46.7% yield as well as mono-ethoxycarbonyl trityl, which can be reused.¹⁹ However, reaction of **9** with ethylene oxide gave poor yield and a complex mixture that was difficult to characterize. The unsuccessful synthesis of **10** using Method A could be due to the susceptibility of ethoxycarbonyl functionality to nucleophilic addition by ethylene oxide under strong basic condition. Therefore, an alternative method (Method B) was carried out in which **1** was reacted with ethylene oxide first followed by ethyl chloroformate. Although one more hydrolysis step was required, the overall yield increased 2-fold. Following the earlier procedure, radical **15** was synthesized from intermediate **10** (Scheme 3) which was slightly soluble in water, making radical **15** possible to be characterized in aqueous solution.

2.2 EPR Studies

Figure 1 shows the solid state EPR spectra of fluorophore-trityls **7**, **8** and **15**. The peak-to-peak linewidth (ΔB_{pp}) for **15** is the broadest compared to **7** and **8**. In spite of the absence of aromatic protons in **15**, this suggests that the line-broadening mechanism is caused mostly by spin-spin exchange interaction. It can be assumed that the packing interaction for **15** is more efficient due, perhaps, to the short linker group that allows a more ordered lattice structure. Figure 2 shows the EPR spectra of **7**, **8** and **15** in DMSO under anaerobic conditions. Both **7** and **8** exhibited a multiplet (7-line) EPR spectrum arising from the hyperfine coupling of two aromatic and two methylene hydrogen atoms from the linker group, suggesting that the unpaired electron is delocalized over the whole PTM moiety (see Table 1 for the simulated EPR parameters). However, trityl **15** exhibited a triplet spectrum with 1:2:1 ratio due to the methylene protons from the linker group. The observed differences in hyperfine coupling constants of the two aromatic and methylene protons are due to the asymmetry of the molecule as shown in Figure 3.

2.3 Reactivity with superoxide radical anion

EPR Spectroscopy: The addition of KO₂ DMSO solution to trityls **7**, **8** or **15** resulted in the disappearance of their EPR signal (Figure 4) suggesting the formation of a diamagnetic product. Using the analogue perchlorotriphenylmethyl-tricarboxylate, PTM-TC, we postulated that the nature of PTM reactivity to O₂^{•−} is a redox reaction resulting in the formation of the reduced product, perchlorotriphenylmethane and oxidation of O₂^{•−} to molecular oxygen as analyzed by electrochemical and mass spectrometric methods.¹⁸ Competitive kinetics experiments revealed that PTM-TC reacts with O₂^{•−} with an apparent second-order rate constant of $8.3 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$. The high reactivity of PTM-TC with O₂^{•−} may offer a potential opportunity for measurement of superoxide in biological systems. It can, therefore, be assumed that based on these preliminary evidences that O₂^{•−} reactivity with **7**, **8** or **15** can parallel that of PTM-TC. Further, we have previously shown that the perchloro class of trityl radicals, for example

PTM-TC, is not reactive or has no specificity of reaction toward hydroxyl, peroxy, or nitric oxide radicals in aqueous solutions.¹⁸

Fluorescence spectroscopy: The excitation and emission profiles for compounds **7**, **8** and **15** are shown in Figure 5. The spectrum showed two emission bands, at 377 nm and 397 nm, corresponding to excitations at 330 and 346 nm, respectively. Enhanced fluorescence spectra can be observed for all compounds upon addition of $O_2^{\bullet-}$, however, this effect is more pronounced for compounds **7** and **15**. This enhanced effect of $O_2^{\bullet-}$ on the emission spectra is consistent with the trityl radical quenching as demonstrated by the EPR studies mentioned above. This study further demonstrates the potential use of fluorophore-trityl conjugate as both an EPR and fluorescent probe for the detection of $O_2^{\bullet-}$.

3. Summary and Conclusions

Perchlorotrityl-fluorophore conjugates were synthesized and characterized, and EPR spectra show long-range electronic coupling with the methylene protons of the linker group. Reactivity with $O_2^{\bullet-}$ was investigated using EPR and fluorescence spectroscopy and the results show disappearance of the EPR signal of the trityl with a concomitant increase in the emission intensity, respectively, suggesting a strong, perhaps a through-space interaction, between the unpaired spin of the trityl and the excited electronic state of the fluorophore. This work suggests potential application of perchlorotrityl-fluorophore conjugates as dual EPR and fluorescence probe for $O_2^{\bullet-}$ detection.

The new class of bimodal probes with higher sensitivity when compared to existing probes may enable accurate and reliable quantitative determination of superoxide concentration in biological systems. Superoxide determination in cells *in vitro* would require cellular internalization of the probe, while *in vivo* measurements need further considerations including mode of administration of the probe, detection sensitivity, and other complexities in the living system. A further difficulty may be the limitation of the methods for deep tissue measurements. In the case of experimental animal models, one can use excised tissue biopsies following intravenous administration of the probe. The tissue samples could then be immediately flash frozen, preserved, and sectioned for EPR or fluorescence microscopy imaging. However, in the case of clinical situations this may not be readily feasible. Thus the near-term usefulness of the probes is in non-clinical research applications both *in vitro* using cell culture models and *in vivo* using animal models.

4. Experimental Section

Bis[2,3,5,6-tetrachlorophenyl]-[4-(2-hydroxyethyl)-2,3,5,6-tetrachlorophenyl] methane (2a), Bis[4-(2-hydroxyethyl)-2,3,5,6-tetrachlorophenyl]-[2,3,5,6-tetrachlorophenyl] methane (2b), tri[4-(2-hydroxyethyl)-2,3,5,6-tetrachlorophenyl] methane (2c)

To a solution of **1** (1.6 g, 2.46 mmol) and TMEDA (0.44 ml, 2.95 mmol) in anhydrous THF (30 ml) at -78°C , *n*-BuLi (1.84 ml, 1.6 M solution in THF) was added and stirred for 1 hour. Ethylene oxide was added (2.95 ml, 1.0 M solution in Et_2O) and the reaction mixture was warmed to room temperature and stirred overnight. The reaction was quenched by addition of 1.0 ml water. The solvent was removed and the residue was dissolved with CH_2Cl_2 , washed with water and brine, and dried with Na_2SO_4 . Removal of the solvent followed by column chromatography (hexanes: EtOAc = 5:1 to 1:1.5) afforded the product **2a** (0.58 g, 0.48 mmol, 33.7%) **2b** (0.36 g, 0.48 mmol, 19.6%), and **2c** (0.12 g, 0.15 mmol, 6.2%). **2a**: ^1H NMR (400 MHz, CDCl_3) δ 7.61 (s, 2H), 6.97 (s, 1H), 3.87 (t, J = 7.2 Hz, 2H), 3.37 (t, J = 7.2 Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 138.8, 138.7, 137.0, 136.4, 134.8, 134.6, 134.4, 134.4, 133.8, 133.6 (2 carbons), 133.5, 133.3 (2 carbons), 132.4, 132.4, 130.3 (2 carbons), 60.2, 56.3, 37.4; HRMS ($M + \text{Na}$)⁺ calcd for $\text{C}_{21}\text{H}_8\text{Cl}_2\text{ONa}$ 724.6644, found 724.6649. **2b**: ^1H NMR

(400 MHz, CDCl₃) δ 7.61 (s, 1H), 6.97 (s, 1H), 3.87 (t, J = 7.2 Hz, 4H), 3.37 (t, J = 7.2 Hz, 4H); ¹³C NMR (100 MHz, CDCl₃) δ 138.9, 136.9, 136.9, 136.6, 136.5, 134.8, 134.8, 134.6 (2 carbons), 134.4, 133.8, 133.7, 133.5, 133.4, 133.4, 133.3, 132.3, 130.2, 60.2, 56.4, 37.4. HRMS ($M + Na$)⁺ calcd for C₂₃H₁₂Cl₁₂O₂Na 768.6909, found 768.6929. **2c**: ¹H NMR (400 MHz, CDCl₃) δ 7.02 (s, 1H), 3.90 (m, 6H), 3.40 (t, J = 7.2 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 136.8 (6 carbons), 134.8 (3 carbons), 134.5 (3 carbons), 133.8 (3 carbons), 133.4 (3 carbons), 60.3 (3 carbons), 56.7, 37.4 (3 carbons). HRMS ($M + Na$)⁺ calcd for C₂₅H₁₆Cl₁₂O₃Na 812.7171, found 812.7204.

Bis(2,3,5,6-tetrachlorophenyl)-[4-(2-phthalimideethenyl)-2,3,5,6-tetrachlorophenyl] methane (3)

To a solution of **2a** (0.32 g, 0.46 mmol), phthalimide (0.13 g, 0.91 mmol), triphenylphosphine (0.24 g, 0.91 mmol) in anhydrous THF (10 ml), DIAD (0.18 L, 0.91 mmol) was added. The reaction mixture was stirred overnight at room temperature. The solvent was removed *in vacuo* and the residue was dissolved in EtOAc, washed with water and brine, and dried with Na₂SO₄. Removal of the solvent followed by column chromatography (hexanes:EtOAc = 1:0 to 10:1) afforded the product **3** (0.37 g, 0.45 mmol, 97.7%). ¹H NMR (500 MHz, CDCl₃) δ 7.81 (dd, J = 5.4 Hz, 3.0 Hz, 2H), 7.68 (dd, J = 5.4 Hz, 3.0 Hz, 2H), 7.62 (s, 1H), 7.61 (s, 1H), 6.93 (s, 1H), 4.10 (m, 1H), 4.01 (m, 1H), 3.52 (m, 1H), 3.44 (m, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 167.9 (2 carbons), 138.8, 138.6, 136.9, 136.7, 134.8, 134.6, 134.5, 134.3, 134.0 (2 carbons), 133.7, 133.6, 133.5, 133.5, 133.4, 133.2, 132.4, 132.3, 131.9 (2 carbons), 130.2, 130.2, 123.3 (2 carbons), 56.3, 34.7, 33.1; HRMS ($M + Na$)⁺ calcd for C₂₉H₁₁Cl₁₂NO₂Na 851.6888, found 851.6903.

Bis(2,3,5,6-tetrachlorophenyl)-[4-(2-aminoethenyl)-2,3,5,6-tetrachlorophenyl] methane (4)

To a solution of **3** (0.25 g, 0.30 mmol) in absolute EtOH (5 ml) was added hydrazine hydrate. The reaction mixture was refluxed for 4 hours. The mixture was then cooled, diluted with water, and extracted with CH₂Cl₂ (2 $\times$). The organic extracts were washed with brine and dried over MgSO₄. Removal of the solvent afforded the crude product **4** (0.14 g, 0.20 mmol), which was used in the next step without purification. HRMS ($M + H$)⁺ calcd for C₂₁H₁₀Cl₁₂N 701.6987, found 701.6931.

Bis(2,3,5,6-tetrachlorophenyl)-{4-[2-(1-pyrenecarbonyl)aminoethenyl]-2,3,5,6-tetrachlorophenyl} methane (5)

To a solution of **4** (0.14 g, 0.20 mmol), 1-pyrenecarboxyl *N*-hydroxysuccinimide ester (0.08 g, 0.24 mmol) in anhydrous THF (10 ml), Et₃N (0.10 ml, 0.72 mmol) was added. The reaction mixture was stirred at 50°C for 2 hours. The solvent was removed and the residue was purified by column chromatography (hexanes = 100% to hexanes: EtOAc: CH₂Cl₂ = 15/4/10) afforded the product **5** (0.11 g, 0.12 mmol, 59.3%). ¹H NMR (500 MHz, CDCl₃) δ 8.44 (d, J = 10.0 Hz, 1H), 8.16 (d, J = 7.5 Hz, 1H), 8.13 (d, J = 7.5 Hz, 1H), 7.9 - 8.1 (m, 6H), 7.64 (s, 1H), 7.63 (s, 1H), 7.05 (s, 1H), 6.48 (t, J = 5.9 Hz, 1H), 3.8 - 4.0 (m, 2H), 3.54 (t, J = 6.9 Hz, 2H); ¹³C NMR (125 MHz, CDCl₃) δ 170.1, 138.8, 138.7, 137.8, 136.7, 135.0, 134.7, 134.44, 134.39, 133.9, 133.6 (2 carbons), 133.4, 133.3, 132.5, 132.45 (2 carbons), 131.1, 130.6, 130.5, 130.3 (2 carbons), 128.62 (2 carbons), 128.56, 127.0, 126.3, 125.8, 125.7, 124.6, 124.5, 124.35, 124.28, 124.17, 56.5, 38.2, 34.3; HRMS ($M + Na$)⁺ calcd for C₃₈H₁₇Cl₁₂NONa 951.7384, found 951.7368.

Bis(2,3,5,6-tetrachlorophenyl)-{4-[2-(1-pyrenebutylcarboxy)aminoethyl]-2,3,5,6-tetrachlorophenyl} methane (6)

To a solution of **4** (0.14 g, 0.20 mmol), 1-pyrenebutyric acid *N*-hydroxysuccinimide ester (0.09 g, 0.24 mmol) in anhydrous THF (10 ml), Et₃N (0.10 ml, 0.72 mmol) was added. The reaction

mixture was stirred at 50°C for 2 hours. The solvent was removed and the residue was purified by column chromatography (hexanes = 100% to hexanes: EtOAc: CH₂Cl₂ = 1/0.33/1) afforded the product **6** (0.13 g, 0.13 mmol, 67.0%). ¹H NMR (400 MHz, CDCl₃) δ 8.25 (d, *J* = 10.0 Hz, 1H), 8.13 (d, *J* = 7.5 Hz, 2H), 8.0 - 8.1 (m, 2H), 7.9 - 8.0 (m, 3H), 7.79 (d, *J* = 7.8 Hz, 1H), 7.59 (s, 1H), 7.58 (s, 1H), 6.94 (s, 1H), 5.58 (t, *J* = 5.9 Hz, 1H), 3.4 - 3.6 (m, 2H), 3.33 (t, *J* = 7.0 Hz, 2H), 3.23 (t, *J* = 7.0 Hz, 2H), 2.1 - 2.2 (m, 4H); ¹³C NMR (125 MHz, CDCl₃) δ 174.7, 140.67, 140.65, 139.6, 138.5, 137.8, 136.8, 136.5, 136.4, 135.8, 135.5, 135.4, 135.3, 134.4, 133.4, 132.9, 132.3, 132.0, 130.8, 129.5, 129.4, 128.8, 127.9, 127.2, 127.0, 126.96, 126.8, 125.4, 58.3, 39.3, 37.9, 36.2, 34.8, 29.2; HRMS (*M* + Na)⁺ calcd for C₄₁H₂₃Cl₁₂NONa 993.7851, found 993.7851.

Bis(2,3,5,6-tetrachlorophenyl)-{4-[2-(1-pyrenecarbonylamino)ethylethyl]-2,3,5,6-tetrachlorophenyl} methyl radical (PTMH2-PyC) (7)

An aqueous solution of tetra *n*-butyl ammonium hydroxide (84.0 mg, 0.105 mmol) was added to a solution of **5** (65.0 mg, 0.070 mmol) in dry tetrahydrofuran (5 ml) at room temperature. The resulting red solution was stirred overnight, and then *p*-chloranil (51.7 mg, 0.21 mmol) was added. The resulting intensely brown solution was stirred for 1 h. After this time, the solution was evaporated to dryness and the resulting product was purified by chromatography (hexanes = 100% to hexanes: EtOAc: CH₂Cl₂ = 9/1/2) and afforded the product **7** (64.6 mg, 0.070 mmol, 99.3%). HRMS (*M* + Na)⁺ calcd for C₃₈H₁₆Cl₁₂NONa 950.7305, found 950.7291.

Bis(2,3,5,6-tetrachlorophenyl)-{4-[2-(1-pyrenebutylcarboxy)aminoethylethyl]-2,3,5,6-tetrachlorophenyl} methyl radical (PTMH2-PyB), (8)

Aqueous solution of tetra *n*-butyl ammonium hydroxide (73.6 mg, 0.092 mmol) was added to a solution of **6** (60.0 mg, 0.062 mmol) in dry tetrahydrofuran (5 ml) at room temperature. The resulting red solution was stirred overnight and then *p*-chloranil (46.0 mg, 0.186 mmol) was added. The resulting intensely brown solution was stirred for 1 h. After this time, the solution was evaporated to dryness and the resulting product was purified by chromatography (hexanes = 100% to hexanes: EtOAc: CH₂Cl₂ = 1/0.33/1) afforded the product **8** (57.2 mg, 0.059 mmol, 95.4%). HRMS (*M* + Na)⁺ calcd for C₄₁H₂₂Cl₁₂NONa 992.7776, found 992.7775.

Bis[2,3,5,6-tetrachloro-4-ethoxycarbonyl-phenyl]-(2,3,5,6-Tetrachlorophenyl) methane (9)

To a solution of **1** (1.15 g, 1.76 mmol) and TMEDA (0.64 ml, 4.22 mmol) in anhydrous THF (80 ml) at -78°C, *n*-BuLi (2.65 ml, 1.6 M solution in THF) was added and stirred for 1 hour. Ethyl chloroformate was added (0.41 ml, 4.2 mmol) and the reaction mixture was allowed to warm up to room temperature and stirred overnight. The reaction was quenched by addition of 1M HCl to pH = 2. The solvent was removed and the residue was dissolved with CH₂Cl₂, washed with brine, and dried with Na₂SO₄. Removal of the solvent followed by column chromatography (hexanes:CH₂Cl₂:EtOAc = 100:1:1) afforded the product **9** (0.66 g, 0.82 mmol, 46.7%) and mono-substituted compound (0.43 g, 0.59 mmol, 33.4%). ¹H NMR (400 MHz, CDCl₃) δ 7.64 (s, 1H), 6.98 (s, 1H), 4.46 (q, *J* = 7.1 Hz, 4H), 1.40 (t, *J* = 7.1 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 163.2, 138.8 (2 carbons), 138.0, 135.4 (2 carbons), 135.02, 134.99, 134.4, 134.0, 133.9, 133.7, 133.3, 132.6, 130.6, 130.5, 130.4, 129.43, 129.40, 63.1, 56.3, 14.0; HRMS (*M* + Na)⁺ calcd for C₂₅H₂₁Cl₁₂O₄Na 824.6805, found 824.6819.

Bis[2,3,5,6-tetrachloro-4-ethoxycarbonyl-phenyl]-[4-(2-hydroxyethylethyl)-2,3,5,6-tetrachlorophenyl] methane (10)

Method A—To a solution of **9** (0.48 g, 0.60 mmol) and TMEDA (0.23 ml, 1.50 mmol) in anhydrous THF (30 ml) at -78°C, *n*-BuLi (0.94 ml, 1.6 M solution in THF) was added and stirred for 1 hour. Ethylene oxide was added (1.5 ml, 1.0 M solution in Et₂O) and the reaction mixture was stirred allowing the reaction to warm to room temperature. The reaction was

quenched by addition of 1 M HCl to pH = 2. The solvent was removed and the residue was dissolved with CH₂Cl₂, washed with brine, and dried with Na₂SO₄. Removal of the solvent followed by column chromatography (hexanes:EtOAc:CH₂Cl₂ = 6:1:1) afforded the product **10** (0.05 g, 0.059 mmol, 9.9%). *Method B:* To a solution of **11** (0.67 g, 0.73 mmol) in THF:H₂O (1:1, 10 ml) at room temperature, LiOH (0.07 g, 2.92 mmol) was added and stirred overnight. After reaction completed, the pH of reaction mixture was adjusted to 2 by addition of 1M HCl. The reaction mixture was extracted by dichloromethane three times. Removal of the solvent followed by column chromatography (hexanes: EtOAc: CH₂Cl₂ = 6:1:1) afforded the product **10** (0.38 g, 0.45 mmol, 61.5%). ¹H NMR (400 MHz, CDCl₃) δ 6.98 (s, 1H), 4.46 (q, J = 7.1 Hz, 4H), 3.87 (t, J = 7.2 Hz, 2H), 3.37 (t, J = 7.2 Hz, 2H), 1.40 (t, J = 7.1 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 163.3, 139.0, 138.9, 137.4, 135.7, 135.34, 135.31, 135.02, 134.99, 134.75, 134.69, 133.98, 133.92, 133.74, 133.64, 130.43, 130.39, 129.38, 129.36, 63.1, 60.2, 56.5, 37.4, 14.0; HRMS (M + Na)⁺ calcd for C₂₇H₁₆Cl₂O₅Na 868.7068, found 868.7065.

Bis[2,3,5,6-tetrachloro-4-ethoxycarbonyl-phenyl]-[4-(2-ethoxycarbonyloxy-ethyl)-2,3,5,6-tetrachloro-phenyl] methane (11)

To a solution of **2a** (0.73 g, 1.04 mmol) and TMEDA (0.73 ml, 4.84 mmol) in anhydrous THF (60 ml) at -78 °C, *n*-BuLi (3.03 ml, 1.6 M solution in THF) was added and stirred for 1 h. Ethyl chloroformate (0.86 ml, 8.96 mmol) was added and the reaction mixture was stirred allowing the reaction to warm up to -20 °C. The reaction was quenched by addition of 1 M HCl to pH = 2. The solvent was removed and the residue was dissolved with CH₂Cl₂, washed with brine, and dried with Na₂SO₄. Removal of the solvent followed by column chromatography (hexanes:CH₂Cl₂:EtOAc = 20:1:1) afforded the product **11** (0.67 g, 0.73 mmol, 70.2%). ¹H NMR (500 MHz, CDCl₃) δ 7.00 (s, 1H), 4.46 (q, J = 7.1 Hz, 4H), 4.35 (m, 2H), 4.18 (q, J = 7.1 Hz, 2H), 3.48 (t, J = 7.0 Hz, 2H), 1.40 (t, J = 7.1 Hz, 6H), 1.28 (t, J = 7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 163.2, 155.0, 138.92, 138.85, 136.3, 136.2, 135.38, 135.35, 135.01, 134.98, 134.85, 134.83, 133.96, 133.92, 133.81, 133.77, 130.44, 130.40, 129.39, 129.37, 64.2, 63.1, 56.5, 36.6, 33.6, 14.3, 14.0; HRMS (M + Na)⁺ calcd for C₃₀H₂₀Cl₂O₇Na 940.7281, found 940.7281.

Bis[2,3,5,6-tetrachloro-4-ethoxycarbonyl-phenyl]-[4-(2-phthalimide-ethyl)-2,3,5,6-tetrachlorophenyl] methane (12)

To a solution of **10** (110 mg, 0.13 mmol), phthalimide (38 mg, 0.26 mmol), triphenylphosphine (68.4 mg, 0.26 mmol) in anhydrous THF (5 ml), DIAD (51 μL, 0.26 mmol) was added. The reaction mixture was stirred overnight at room temperature. The solvent was removed and the residue was dissolved with EtOAc, washed with water and brine, and dried with Na₂SO₄. Removal of the solvent was followed by column chromatography (hexanes:EtOAc = 8:1 to 3:1) which afforded the product **12** (66.7 mg, 0.068 mmol, 52.3%) with minimal impurity. ¹H NMR (400 MHz, CDCl₃) δ 7.78 (d, J = 5.4 Hz, 3.1 Hz, 2H), 7.67 (d, J = 5.4 Hz, 3.1 Hz, 2H), 6.94 (s, 1H), 4.46 (q, J = 6.9 Hz, 4H), 4.0 - 4.1 (m, 2H), 3.4 - 3.6 (m, 2H), 1.40 (t, J = 6.9 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 167.9, 163.2, 139.0, 138.8, 137.3, 136.1, 135.30, 135.27, 135.1, 134.94, 134.89, 134.6, 134.0, 133.96, 133.85, 133.76, 133.67, 131.9, 130.4, 130.3, 129.34, 129.26, 123.3, 63.0, 56.5, 35.2, 33.1, 14.0; HRMS (M + Na)⁺ calcd for C₃₅H₁₉Cl₂NO₆Na 997.7286, found 997.7291.

Bis[2,3,5,6-tetrachloro-4-ethoxycarbonyl-phenyl]-[4-(2-aminoethyl)-2,3,5,6-tetrachloro phenyl] methane (13)

To a solution of **12** (50 mg, 0.0513 mmol) in absolute EtOH (5 ml) was added hydrazine hydrate. The reaction mixture was heated and refluxed for 2 h. The mixture was cooled while precipitate appeared. The reaction mixture was diluted with water, and extracted with CH₂Cl₂ (2×). The organic extracts were washed with brine, dried over MgSO₄. Removal of

the solvent afforded the crude product **13** (45 mg, 0.053 mmol), which was used in the next step without purification. HRMS (M + H)⁺ calcd for C₂₇H₁₈Cl₁₂NO₄ 845.7410, found 845.7401.

Pyrene-1-carboxylic acid (2-{4-[bis-(2,3,5,6-tetrachloro-4-ethoxycarbonyl-phenyl)-methyl]-2,3,5,6-tetrachloro-phenyl}-ethyl)-amide (14)

To a solution of **13** (45 mg, 0.20 mmol), 1-pyrenecarboxyl *N*-hydroxysuccinimide ester (0.08 g, 0.053 mmol) in anhydrous THF (5 ml), Et₃N (0.033 ml, 0.24 mmol) was added. The reaction mixture was stirred at 50 °C for 2 h. The solvent was removed and the residue was purified by column chromatography (hexanes = 100% to hexanes: EtOAc: CH₂Cl₂ = 4.5/0.8/2) afforded the product **14** (30 mg, 0.03 mmol, 52.5%). ¹H NMR (400 MHz, CDCl₃) δ 8.46 (d, *J* = 10.0 Hz, 1H), 8.19 (d, *J* = 8.5 Hz, 2H), 7.9 - 8.1 (m, 6H), 7.05 (s, 1H), 6.37 (m, 1H), 4.48 (m, 4H), 3.8 - 4.1 (m, 2H), 3.56 (m, 2H), 1.41 (m, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 170.1, 163.3, 139.0, 138.9, 138.2, 136.1, 135.39, 135.37, 135.1, 135.0, 134.9, 134.8, 134.97 (2 carbons), 133.92, 133.7, 132.6, 131.1, 130.7, 130.6, 130.5, 130.4, 129.42, 129.40, 128.7 (2 carbons), 128.6, 127.1, 126.3, 125.8 (2 carbons), 124.8, 124.44, 124.39, 124.35, 124.26, 63.0, 56.5, 38.2, 34.3, 14.0; HRMS (M + Na)⁺ calcd for C₄₄H₂₅Cl₁₂NO₅Na 1095.7809, found 1095.7797.

Bis(2,3,5,6-tetrachlorophenyl)-{4-[2-(1-pyrenecarbonyl)aminoethyl]-2,3,5,6-tetrachlorophenyl} methyl radical (15)

An aqueous solution of tetra *n*-butyl ammonium hydroxide (22.6 μL, 0.028 mmol) was added to a solution of **14** (20.0 mg, 18.6 μmol) in dry tetrahydrofuran (5 ml) at room temperature. The resulting red solution was stirred overnight, and then *p*-chloranil (13.7 mg, 0.056 mmol) was added. The resulting intensely brown solution was stirred for 1 h. After this time, the solution was evaporated to dryness and the resulting product was purified by chromatography (hexanes = 100% to hexanes: EtOAc: CH₂Cl₂ = 6/2/1) afforded the product **15** (19.8 mg, 18.5 μmol, 99.0%); HRMS (M + Na)⁺ calcd for C₄₄H₂₄Cl₁₂NO₅Na 1094.7731, found 1094.7732.

EPR measurements

EPR measurements were performed at room temperature using a Bruker EMX spectrometer equipped with a high-sensitivity resonator. Solutions of the compounds in DMSO were drawn into a 50-ml capillary tube and used for EPR measurements. The acquisition settings were microwave frequency, 9.786 GHz, modulation amplitude, 1G; microwave power, 5 mW; time constant, 40 msec; scan time, 21 sec. Spectral simulations were carried out using a custom-developed software (EPRDAP).

Fluorescence measurements

The fluorescence scan measurements were performed on the compounds in DMSO at room temperature using Aminco-Bowman Series 2 spectrofluorometer (Aminco Spectronic Instruments, New York, USA).

Acknowledgments

This work was supported by NIH grant EB006363.

References

1. Waris G, Ahsan H. J Carcinog 2006;5:14. [PubMed: 16689993]
2. Provinciali M, Donnini A, Argentati K, Di Stasio G, Bartozzi B, Bernardini G. Free Radic Biol Med 2002;32:431. [PubMed: 11864783]
3. Dhanasekaran A, Kotamraju S, Karunakaran C, Kalivendi SV, Thomas S, Joseph J, Kalyanaraman B. Free Radic Biol Med 2005;39:567. [PubMed: 16085176]

4. McCord JM, Fridovich I. *J Biol Chem* 1969;244:6056. [PubMed: 4981789]
5. Auclair C, Torres M, Hakim J. *FEBS Lett* 1978;89:26. [PubMed: 207567]
6. Budd SL, Castilho RF, Nicholls DG. *FEBS Lett* 1997;415:21. [PubMed: 9326361]
7. Roubaud V, Sankarapandi S, Kuppusamy P, Tordo P, Zweier JL. *Anal Biochem* 1997;247:404. [PubMed: 9177705]
8. Fridovich I. *J Biol Chem* 1997;272:18515. [PubMed: 9228011]
9. Thomson L, Trujillo M, Telleri R, Radi R. *Arch Biochem Biophys* 1995;319:491. [PubMed: 7786032]
10. Pou S, Huang YI, Bhan A, Bhadt VS, Hosmane RS, Wu SY, Cao GL, Rosen GM. *Anal Biochem* 1993;212:85. [PubMed: 8396365]
11. Pou S, Bhan A, Bhadt VS, Wu SY, Hosmane RS, Rosen GM. *Faseb J* 1995;9:1085. [PubMed: 7649408]
12. Lozinsky E, Martin VV, Berezina TA, Shames AI, Weis AL, Likhtenshtein GI. *J Biochem Biophys Methods* 1999;38:29. [PubMed: 10078871]
13. Bilski P, Hideg K, Kalai T, Bilska MA, Chignell CF. *Free Radic Biol Med* 2003;34:489. [PubMed: 12566074]
14. Rizzi C, Samouilov A, Kutala VK, Parinandi NL, Zweier JL, Kuppusamy P. *Free Radic Biol Med* 2003;35:1608. [PubMed: 14680684]
15. Kutala VK, Parinandi NL, Zweier JL, Kuppusamy P. *Arch Biochem Biophys* 2004;424:81. [PubMed: 15019839]
16. Ballester M, Miravittles C, Molins E, Carreras C. *J Org Chem* 2003;68:2748. [PubMed: 12662047]
17. Ballester M, Riera J, Castafier J, Badia C, M JM. *J Am Chem Soc* 1971;93:2215.
18. Kutala VK, Villamena FA, Ilangoan G, MasPOCH D, Roques N, Veciana J, Rovira C, Kuppusamy P. *J Phys Chem B* 2008;112:158. [PubMed: 18081340]
19. Dang V, Wang J, Feng S, Buron C, Villamena FA, Wang PG, Kuppusamy P. *Bioorg Med Chem Lett* 2007;17:4062. [PubMed: 17499990]
20. Malkov AV, Gouriou L, Lloyd-Jones GC, Stary I, Langer V, Spoor P, Vinader V, Kocovsky P. *Chemistry* 2006;12:6910. [PubMed: 16807930]
21. Xue L, Charles I, Arya DP. *Chem Commun (Camb)* 2002;70. [PubMed: 12120315]
22. Rovira C, Ruiz-Molina D, Elsner O, Vidal-Gancedo J, Bonvoisin J, Launay JP, Veciana J. *Chemistry* 2001;7:240. [PubMed: 11205016]

Figure 1.

X-band EPR spectra of fluorophore-trityls **7**, **8** and **15** as crystalline powders in air as measured at room temperature. The acquisition settings were: microwave frequency, 9.786 GHz (X-band), modulation amplitude, 1 G; microwave power, 2 mW; time constant, 40 msec; scan time, 15 sec. The first derivative peak-to-peak widths of **7**, **8**, and **15** were 3.26 G, 2.76 G, and 5.14 G, respectively.

Figure 2.

X-band EPR spectra of fluorophore-trityls in anaerobic solution. The EPR spectra of compounds a) **8**; b) **7**; and c) **15** in DMSO (100 μ M) measured at room temperature. (See Experimental Section for instrument settings). The broken red-colored trace superimposed on each spectrum was simulated using the parameters shown in Table 1. The small satellite peaks indicated with * on both sides of the main peaks are due to hyperfine coupling with ^{13}C nuclei (natural abundance).

Figure 3. Preferred conformations for trityls **7** (top) and **8** (bottom) using MMFF MonteCarlo search and geometry optimization at the B3LYP/6-31G* level of theory showing the intramolecular distance between the trityl and fluorophore moieties.

Figure 4.

EPR spectra of (a) **8**, (c) **7** and (d) **15**; and (b), (d) and (f), respectively, are the resulting spectra upon addition of KO_2 in DMSO. The acquisition settings were: microwave frequency, 9.786 GHz, modulation amplitude, 1 G; microwave power, 5 mW; time constant, 40 msec; scan time, 21 sec. The broken red-colored trace superimposed on each spectrum was simulated using the parameters shown in Table 1. The small satellite peaks indicated with * on both sides of the main peaks are due to hyperfine coupling with ^{13}C nuclei (natural abundance).

Figure 5.

Excitation and emission profiles of **8**, **7**, and **15** in the presence and absence of KO_2 in DMSO. Data show two emission bands, at 377 nm and 397 nm, corresponding to excitations at 330 and 346 nm, respectively.

Scheme 1.
Synthesis of compounds **7** and **8**.

Scheme 2.
Methodologies for the synthesis of compound **10**.

Scheme 3.
Synthesis of compound **15**.

Table 1

Experimental EPR hyperfine splitting constants (a_{Hx}) and anaerobic linewidths (ΔB_{pp}) in G for trityls **7**, **8** and **15** in DMSO at 298 K.

Compound	Hyperfine Splitting Constants a_{H} (G)		ΔB_{pp} (G)
	aromatic H's (H_{a} and H_{b})	methylene H's (H_{c} and H_{d})	
7	2.09, 1.91	1.24, 0.95	0.70
8	2.05, 1.96	1.14, 1.02	0.62
15	n/a	1.26, 0.94	0.66