LETTERS

Use of Wikis in Advanced Pharmacy Practice Experiences

To the Editor: Most clinicians have been exposed to, or are aware of, the existence of Wikipedia, the world's most popular wiki project. The online encyclopedia Wikipedia offers the advantage of an open-access library of information that can be authored or edited with little delay by anyone with Internet access.¹ Alternatively, wikis (and their accompanying software programs) may serve as a useful tool in health sciences education and research. A wiki can serve as an "expert pharmacy community" in an online format, fully supported through a collaborative effort by its members. Utilization of wiki technology specifically for pharmacy-related education is limited and its use in pharmacy education has not been well documented. Mirk et al presented information on pharmacy students' prior experience with wikis, but their opinions on their use and impact in a pharmacy education setting are unknown.⁶ At the South Carolina College of Pharmacy on the campus of the University of South Carolina (SCCP-USC), 3 faculty members utilize wiki technology during advanced pharmacy practice experiences (APPE).

Online tools for advancing education and training in health sciences have been used for over a decade, although use of wikis is a more recent development.² Ciesielka described the use of a community wiki in nursing education at Clarion University.³ The students far exceeded course expectations and continued to contribute to the wiki even after grades were assigned. Students were given substantial autonomy and responded positively to the experience, suggesting it greatly facilitated teamwork and allowed them to "see the big picture."³ Incorporation of wikis into medical education has also been documented in a gross anatomy course at the Mayo Medical School.⁴ Given an anatomical outcome, students were instructed to utilize a wiki to design a clinically relevant case. Pre- and post-assessments demonstrated a significant increase in students' confidence in identifying appropriate diagnostic procedures, discerning differential diagnoses and potential treatments, and delivering an oral presentation of the clinical case.⁴

The use of wikis in health sciences expands beyond education into the research arena, as biology databases have begun to utilize wiki technology.⁵ In conjunction with his biologics company, Knewco, Barend Mons has launched a Web site, www.wikiprofessional.org, with the goal of creating a single, interlinked, biomedical database supporting vital research in genetic and protein modeling for many scientists. The use of wikis may be embraced

soon by publishing companies seeking a move to an online format. A wiki format offers the advantage of continuous online group editing done in real time, minimizing inherent delays and costs.

Over the past year, 3 faculty members at SCCP-USC have used wikis to enhance the student experience during APPEs, but student opinions on their use are unknown. These wikis were setup through Google Wikis (<http://www.google.com/sites/>), a free online service. Students participate in a 4- week APPE in an acute care setting focused in internal medicine/infectious diseases, critical care, and/or hematology/oncology. The students create and use wiki entries to answer daily questions that arise during interdisciplinary rounds or stemming from patient-specific and evidence-based care decisions. Additionally, disease-state reviews are created by the students to provide comprehensive information on key topics. In situations where information on the wiki related to a topic already exists, students are asked to review, edit, and expand on the work of their peers. The initial use of wikis required some Web site creation and design, but following that, entries have been exclusively student created. Since using wikis is new to many students, we provide some initial instruction on the types and expectations for content as well as training in "how to" of the Web site. During the first few days of the APPE, students are somewhat reluctant to add entries to the wiki, but reminders to add specific information are well-received. However, once the students are comfortable with the wiki site, they are eager to add new and update existing information.

Thirty students have utilized the wiki over the past 9 months, with 4 students using wikis on multiple APPEs. In order to assess the impact of the wiki on student learning and overall perception, a survey was administered with answers based on a 5-point Likert scale (1=strongly agree; 5=strongly disagree), with additional space for comments.

The survey was administered anonymously online using SurveyMonkey (surveymonkey.com). All students using wikis on APPEs were invited to participate, with 16 students responding to the survey (response rate = 61.5%). Overall, opinions on the survey were positive with greater than 90% of respondents either strongly agreeing or agreeing that wikis helped reinforce concepts and ideas and that they enjoyed working with wikis while on APPEs. Many students (approximately 70%, 11/16) believed that the use of wikis helped create a sense of collegiality and strengthened professional relationships with peers. In addition, a question aimed at gauging students' interest in using wikis in other areas of pharmacy education had a favorable response with approximately 90% of students strongly agreeing or agreeing that they would like to see

wikis and other online tools employed in other areas of pharmacy education. Students gave generally favorable responses to questions gauging their attitudes towards using previously written wiki entries to help review APPE concepts and assessing how the wiki helped them reflect on clerkship concepts.

Student comments regarding the use of wikis detailed how they were enjoyable and helped enhance concept understanding. One student reported that they “enjoyed working on pages for the wiki. [They] felt like they were a good tool to use to help reinforce concepts introduced during the rotation, especially in thinking critically to break the concept down into palatable pieces. . .” Another student stated that, “the wiki forced [them] to put concepts into [their] own words, and as a result enhanced understanding.”

Critics of wikis have commented that open access leads to the possibility of inaccuracies in user contributions. In the medical arena, this could involve potential hazards leading to patient harm. An analysis of online content in Wikipedia compared to Encyclopedia Britannica found little difference in overall accuracy.⁷ At SCCP, faculty members serve as editors for the APPE wikis. Whenever activity occurs on the wiki, a post is made on the announcement page allowing preceptors and students to follow any edits or updates. Currently, our wikis are only available to students as contributors who have been or are currently enrolled in the APPE. This helps to alleviate copyright and liability concerns and controlling for unknown additions to the wiki pages. Our overall experiences with wikis have been positive, and we feel that they provide an opportunity for students to practice critical thinking and writing skills and promote professional collegiality.

We have found the use of wikis to be a useful tool in experiential education, and student responses indicate that they may be helpful in other areas of pharmacy education. Incorporation of wiki technology into didactic elective courses is also being researched and evaluated by the same faculty members. These data will be forthcoming in a future publication.

April D. Miller, PharmD

P. Brandon Bookstaver, PharmD

LeAnn B. Norris, PharmD

South Carolina College of Pharmacy-University of South Carolina

REFERENCES

1. Wiki. <http://en.wikipedia.org/wiki/Wiki>. Accessed May 23, 2009.
2. Boullos MN, Maramba I, Wheeler S. Wikis, blogs and podcasts: a new generation of web-based tools for virtual collaborative clinical practice and education. *BMC Med Educ*. 2006;6:41.

3. Ciesielka D. Using a wiki to meet graduate nursing education competencies in collaboration and community health. *Educ Innovations*. 2008;47:473-476.

4. Philip CT, Unruh KP, Lachman N, Pawlina W. An explorative learning approach to teaching clinical anatomy using student generated content. *Anat Sci Educ*. 2008;1:106-110.

5. Giles J. Key biology databases go wiki. *Nature*. 2007;445(7129):691.

6. Mirk SM, Burkiewicz JS, Fit KE. Student perception of wiki in an elective course. American College of Clinical Pharmacy Phoenix, AZ; 2008.

7. Giles J. Internet encyclopaedias go head to head. *Nature*. 2005;438(7070):900-901.

Mental Health Pharmacy Education at 16 European Universities

To the Editor: Services provided by pharmacists are well-suited for optimizing the use of medications for mental illness.¹ However, lack of mental health education is an important barrier to the provision of pharmacy services.^{2,3} Suboptimal attitudes toward people with mental illness are common among pharmacists and pharmacy students.^{4,5} There are few studies describing pharmacy education about depression, schizophrenia, and substance abuse. A recent study in the *Journal* reported the teaching of psychiatry-related topics at US colleges and schools of pharmacy.⁶ However, no corresponding studies have investigated mental health pharmacy education in Europe. With this in mind, we conducted a preliminary assessment of mental health pharmacy education provided at European universities.

A 20-item online survey instrument was developed using E-lomake software (Eduix Oy, Tampere, Finland). The survey instrument was adapted from survey instruments used to assess mental health pharmacy education at US colleges and schools of pharmacy,⁶ and depression education at medical schools in the Asia-Pacific region.⁷ The survey instrument was divided into 5 sections. Section 1 included 5 items about the respondents and their school of pharmacy. Section 2 included 6 items about the overall course content in relation to mental health. Section 3, 4, and 5 included 3 items each related to education about depression, schizophrenia, and substance use disorders. The survey and functionality of the E-lomake software was pilot tested for face-validity among a convenience sample of native and non-native English speaking educators.

The survey, cover letter, and participant information sheet were emailed to an academic staff member from 30 schools of pharmacy across Europe in December 2008. The pharmacy schools were a convenience sample but were selected to cover a range of sizes and country locations. All

data were analyzed using the Statistical Package for the Social Sciences Version 15.0 (SPSS, Chicago, IL).

Responses were received from 16 pharmacy schools in Belgium, Denmark, Estonia, Finland, Latvia, Lithuania, Norway, Portugal, Slovenia, Spain, Turkey, and the United Kingdom (response rate 53%). Eleven of the respondents were lecturers or associate professors. Fourteen universities were publicly funded and 9 had an undergraduate student enrollment of more than 400 students. All universities taught psychiatry-related topics as part of a therapeutics-based course. Six universities employed or contracted a specialist mental health pharmacist to provide teaching, and 5 universities considered psychiatric pharmacy to be the content focus of their curriculum. All universities provided teaching about depression, schizophrenia, and substance use disorders via didactic lectures (Table 1). Two universities also provided the opportunity for students to participate in a psychiatric pharmacy clerkship to learn about depression and schizophrenia. Depression was taught for an average of 24 contact teaching hours per university degree. Schizophrenia and substance use disorders were taught for an average of 18 and 13 contact teaching hours per degree, respectively.

Our preliminary survey was the first to quantify the content, format, and duration of mental health pharmacy education provided at European universities. Our survey revealed considerable differences in mental health pharmacy education offered in Europe and the United States. Only 2 (13%) European universities that responded offered an elective internship in psychiatric pharmacy compared to 92% of universities in the United States.⁶ Only 6 (38%) European universities that responded employed or contracted a specialist mental health pharmacist to provide teaching, compared to 76% in the United States. These differences may be due to differences in mental health pharmacy practice in Europe and the United States. Pharmacists are able to undergo postgraduate specializa-

tion in mental health pharmacy in both the United States and United Kingdom, but not in most continental European countries. Despite these differences there were also several similarities. All of the responding universities in Europe and the United States offered psychiatric topics as part of a therapeutics-based course. In addition, no universities in Europe and only 4% of universities in the United States required students to undertake a psychiatric pharmacy internship.

Didactic lectures were the most common teaching method in relation to depression, schizophrenia, and substance use disorders. However, didactic lectures alone may not change students' attitudes towards people with mental illness.⁸ Encouragingly, some universities combined didactic lectures with interactive education, including small-group tutorials, role-playing, problem-based learning (PBL), and having actual patients present about their experiences. PBL in medical education has been adopted at different rates across Europe.⁹ Our findings suggest that the same may be true of pharmacy education. Five universities included teaching about sufferer's experience of depression and schizophrenia, and 4 about sufferer's experience of substance use disorders. Having actual patients present about their experience may improve students' attitudes toward people with mental illness.^{8,10} This format of education may also improve pharmacists' confidence to provide services to people with mental illness.¹¹

There are a number of methodological strengths and limitations to our study. We included universities from south, central, and northern Europe. However, the sample size of universities was relatively small. As with the US survey, it was possible that those universities with a research or teaching interest in mental health were more likely to respond. Determining the duration of contact teaching may have been difficult when material about depression, schizophrenia, or substance use disorders

Table 1. Number and Percentage of 16 European Universities Offering Various Formats of Education about Depression, Schizophrenia, and Substance Use Disorders

	Depression, No. (%)	Schizophrenia, No. (%)	Substance Use Disorders, No. (%)
Skills specific to communicating	4 (25)	2 (13)	4 (25)
Sufferer's experience of illness	5 (31)	5 (31)	4 (25)
Didactic lectures	16 (100)	16 (100)	16 (100)
Experimental work (eg, wet laboratories)	2 (13)	1 (6)	0 (0)
Role playing with actors (eg, other students, academics)	4 (25)	3 (19)	2 (13)
Having actual patients present about their experiences	4 (25)	3 (19)	2 (13)
Small group tutorials	9 (56)	8 (50)	6 (38)
Problem based learning	6 (38)	7 (44)	6 (38)
Internships at psychiatric facilities, clinics, pharmacies	2 (13)	2 (13)	1 (6)

was taught as part of general courses on pharmacology, using patient case studies or during internships. In addition, our research did not assess the duration of non-contact teaching, nor the content, format, and duration of education provided at the postgraduate level.

In conclusion, there are considerable differences in the content, format, and duration of education provided in Europe and the United States. There is also considerable variation in mental health pharmacy education across Europe. There is a need to develop core competencies and minimum standards for the provision of mental health pharmacy education at the European and national levels. Our survey highlights the need for a larger European study to determine whether the education provided at European universities adequately prepares pharmacists to provide mental health care services.

Iiro Koski^a

Asko Heikkilä, BSc(Pharm)^a

J Simon Bell, PhD^{a,b,c}

^aFaculty of Pharmacy, University of Helsinki, Finland

^bKuopio Research Centre of Geriatric Care, University of Kuopio, Finland

^cDepartment of Pharmacology and Toxicology, University of Kuopio, Finland

REFERENCES

1. Bell S, McLachlan AJ, Aslani P, Whitehead P, Chen TF. Community pharmacy services to optimise the use of medications for mental illness: a systematic review. *Aust New Zealand Health Policy*. 2005;2:29.
2. Scheerder G, De Coster I, Van Audenhove C. Pharmacists' role in depression care: a survey of attitudes, current practices, and barriers. *Psychiatr Serv*. 2008;59:1155-1160.
3. Scheerder G, De Coster I, Van Audenhove C. Community pharmacists' attitude toward depression: a pilot study. *Res Social Adm Pharm*. 2009;5:Epub 2009 Jan 21.
4. Bell JS, Aaltonen SE, Bronstein E, et al. Attitudes of pharmacy students toward people with mental disorders, a six country study. *Pharm World Sci*. 2008;30:595-599.
5. Phokeo V, Sproule B, Raman-Wilms L. Community pharmacists' attitudes toward and professional interactions with users of psychiatric medication. *Psychiatr Serv*. 2004;55:1434-1436.
6. Cates ME, Monk-Tutor MR, Drummond SO. Mental health and psychiatric pharmacy instruction in US colleges and schools of pharmacy. *Am J Pharm Educ*. 2007;71:4.
7. Lee MS. A preliminary survey of undergraduate education on depression in medical schools in the Asia Pacific region. *Australas Psychiatry*. 2004;12(Suppl 1):S28-S32.
8. Buhler AV, Karimi RM. Peer-level patient presenters decrease pharmacy students' social distance from patients with schizophrenia and clinical depression. *Am J Pharm Educ*. 2008;(5):Article 106.
9. Jippes M, Majoor GD. Influence of national culture on the adoption of integrated and problem-based curricula in Europe. *Med Educ*. 2008;42:279-285.

10. Bell JS, Johns R, Rose G, Chen TF. A comparative study of consumer participation in mental health pharmacy education. *Ann Pharmacother*. 2006;40:1759-1765.

11. Bell JS, Whitehead P, Aslani P, Sacker S, Chen TF. Design and implementation of an educational partnership between community pharmacists and consumer educators in mental health care. *Am J Pharm Educ*. 2006;70:28.

An Academic Perspective on the APPEs' Educational Significance

To the Editor. We question the rationale and evidence that has placed a disproportional emphasis on clinical experience, at the expense of the basic sciences, in doctor of pharmacy (PharmD) programs in the United States. Pharmacy education has evolved over the past 20 years to the point that "clinical experience" may be viewed as paramount by the Accreditation Council for Pharmacy Education (ACPE). Recently, the accreditation requirements for advanced pharmacy practice experiences (APPEs) have been increased from 30 to 36 weeks in the United States and now occupy one-quarter of the typical 4-year PharmD curriculum.¹ During a recent accreditation visit, when ACPE staff evaluators were directly asked for substantial evidence to support the change from 30 to 36 weeks for better education outcomes for pharmacy students, none could be provided. The fact that the accreditation body for US pharmacy programs lacks objective evidence is troubling.

Pharmacy, at least in the past, was viewed in the United States as a science-based profession. Recently, some have commented that the diminishment of science has encouraged a trade school mentality in pharmacy programs and marks a return to apprenticeship training reminiscent of the 19th century rather than the education of pharmacy professionals.^{2,3}

Examining the *Drug Topics* 2008 list of the Top 200 Brand Name Drugs dispensed in the United States, esomeprazole (Nexium), escitalopram (Lexapro), and montelukast (Singulair) were ranked numbers 2, 3, and 4 as the most frequently dispensed drugs.⁴ This indicates that these 3 drugs are considered the standards of clinical practice, at least numerically, for their respective indications.

Esomeprazole (Nexium). Number 2 on the US Top 200 list for 2008 is esomeprazole (Nexium). The Description Section of the package insert states that esomeprazole is the S-isomer of omeprazole.⁵ Omeprazole is available as an over-the-counter (nonprescription) drug in the United States.

In the Clinical Studies section of the package insert, focusing on the 5 trials described for treating Symptomatic Gastroesophageal Reflux Disease (GERD), 2 of the trials were multicenter, randomized, double-blind,

placebo-controlled studies. Esomeprazole was significantly better than placebo. In the remaining 3 GERD trials, esomeprazole 20 mg and 40 mg and omeprazole 20 mg were evaluated. No significant treatment related differences were seen.

There are no claims of superiority in the package insert that esomeprazole is superior to other proton pump inhibitors, including omeprazole, for the treatment of symptomatic GERD.

Escitalopram (Lexapro). Escitalopram is number 3 on the *Drug Topics* list for 2008. Using the Internet and DailyMed to access the package insert, the Description section for the drug notes that escitalopram is the pure S-enantiomer of racemic citalopram. Patients who have been prescribed citalopram since it was first approved in the United States in July 1998 are also getting escitalopram.⁶

The package insert explains that the efficacy of escitalopram was established in 3 placebo controlled trials using the change from baseline to endpoint in the Montgomery Asberg Depression Rating Scale (MADRS) as the primary outcome measure. However, 4 trials are actually described in the package insert. A recent report in the *New England Journal of Medicine* reveals that 4 trials were in fact submitted to the FDA, but only the 3 with statistically positive results were published. The forth trial did not show a statistical difference between escitalopram and placebo and has not been published.⁷ No claims of superiority over any other antidepressant including citalopram appear in the package insert for escitalopram.

Montelukast (Singulair). In the package insert for the number 4 drug on the *Drug Topics* Top 200 list, montelukast, the Clinical Studies section communicates information about 5 seasonal allergic rhinitis clinical trials. In 4 of these 5 trials there was a significant reduction in daytime nasal symptoms scores with montelukast compared to placebo. Three of the 5 trials showed similar results. The fourth trial was not discussed, and it may be assumed that montelukast could not be differentiated from placebo. The last trial used loratadine 10 mg as the active comparator. Loratadine was numerically superior to montelukast in day time nasal symptoms score (the average of individual scores of nasal congestion, rhinorrhea, nasal itching, sneezing) as assessed by patients on a 0-3 categorical scale.⁸

The prescribing of esomeprazole, escitalopram, and montelukast in the US in 2008 suggests the acceptance of these drugs by US prescribers as the standard of practice for their approved uses, despite evidence from randomized controlled trials that these drugs may be no better or not as good as other older medications. This suggests indirectly that the prescribing of these drugs may not be science based.

Pharmaceutical industry promotion may be 1 possible explanation for the high rate of prescribing these 3 drugs; another possible contributing factor is a failure of clinicians' didactic education in the areas of how to critically interpret clinical research; the drug regulatory process; and the uncomfortable history of drug safety disasters.

In the clinical setting, a pharmacy student, who has completed a required APPE at a prestigious medical center, local hospital, or local community pharmacy, remembers that Drug X was widely prescribed and therefore must be the standard of practice everywhere. There may be future negative educational outcomes if the student decides on a career in academics. The new assistant professor brings "clinical experience" misconceptions to the classroom and Drug X, as the standard of practice, is inappropriately perpetuated in the next generation of pharmacists.

A serious question should be addressed by pharmacy educators in and outside of the United States: should pharmacy students attend required APPEs before attaining the scientific education necessary to differentiate between prescribing decisions based on scientific evidence from well-controlled clinical trials and the Hussey-Stetler Test of Time, "clinical experience"? A position that has long been expressed, but apparently ignored by the vast majority of pharmacy educators, is that statistical literacy and the education to critically evaluate clinical research, are basic clinical skills that are as important as proficiency with a sphygmomanometer.⁹

Evidence is lacking that increasing the experiential requirement of Pharm D curricula in the United States from 30 to 36 weeks produces better-educated pharmacy graduates. The public and the profession of pharmacy may be better served if the 36-week requirement for advanced experiential training were cut to 18 weeks of high quality experiences, and the remaining time replaced with 18 weeks of didactic education focusing on the science of drug literature evaluation, the history of drug tragedies, and drug regulatory standards.

Sana R. Sukkari, BSc Pharm. M Phil, PharmD^a

Larry D. Sasich, Pharm.D, MPH, FASHP^a

Mohammed A. Barasain, BSc Pharm, Pharm D (Candidate)^{b,c}

^aLECOM School of Pharmacy

^bCampbell University School of Pharmacy

^cNational Drug and Poison Information Center, Saudi Food and Drug Authority (SFDA), Riyadh, Saudi Arabia

REFERENCES

1. Accreditation Council for Pharmacy Education. Accreditation Standards and Guidelines for the Professional Program in Pharmacy Leading to the Doctor of Pharmacy Degree. American Council for

American Journal of Pharmaceutical Education 2009; 73 (8) Article 139.

Pharmacy Education 2006. http://www.acpe-accredit.org/pdf/ACPE_Revised_PharmD_Standards_Adopted_Jan152006.pdf Accessed June 21, 2009.

2. Dipiro JT. Making the most of pharmacy school. *Am J Pharm Educ.* 2008;72(1):15.

3. Skau K. Pharmacy is a science-based profession. *Am J Pharm Educ.* 2007;71(1):11.

4. Avastar Communications. Top 200 Branded Drugs by Total Prescriptions 2008. SDI/Verispan, VONA, Full year 2008 <http://drugtopics.modernmedicine.com/> Accessed June 21, 2009.

5. AstraZenica LP. Esomeprazole (Nexium) - Professional Product Label. DailyMed.

6. Forest Pharmaceuticals I. Escitalopram (Lexapro) - Professional Product Label. DailyMed <http://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?id=9444> Accessed June 30, 2009.

7. Turner EH, Matthews AM, Linardatos E, Tell RA, Rosenthal R. Selective publication of antidepressant trials and its influence on apparent efficacy. *N Engl J Med.* 2008;358(3):252-260.

8. Merck & Co. I. Montelukast (Singulair) - Professional Product Label. DailyMed <http://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?id=8034> Accessed June 30, 2009.

9. Fletcher RH, Fletcher SW, Wagner EH. Clinical Epidemiology - The Essentials. Baltimore: Williams & Wilkins; 1982.