Published in final edited form as:

J Am Chem Soc. 2010 March 3; 132(8): 2528–2529. doi:10.1021/ja9106572.

Total Synthesis of the Ammosamides

Chambers C. Hughes and William Fenical*

Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92093-0204

Abstract

The ammosamides A-C are chlorinated pyrrolo[4,3,2-*de*]quinoline metabolites isolated from the marine-derived *Streptomyces* strain CNR-698. The natural products, which possess a dense array of heteroatoms, were synthesized in 17–19 steps from 4-chloroisatin. That the five nitrogen atoms were introduced at the appropriate time and in a suitable oxidation state was key to the success of the total synthesis. Compared to synthetic deschloro ammosamide B, natural ammosamide B is much less susceptible to oxidative degradation.

The secondary metabolites of actinomycetes often display exceptional biological activities.¹ These genetically-encoded small molecules, which were subjected to evolutionary pressures over millions of years, generally possess high affinities for their protein targets. Our studies of marine actinomycetes have focused on those strains isolated from deep-ocean sediments.²

The ammosamides A-B (**1–2**) are chlorinated alkaloids characterized by a dense array of heteroatoms (N,O,S) and a relative paucity of hydrogen atoms (Chart 1). The molecules are related to the pyrroloiminoquinone class of natural products,³ although they exhibit several distinct features. The isolation and structural elucidation of ammosamides A and B (**1, 2**) from a marine sediment-derived *Streptomyces* strain CNR-698 was described in a previous communication by our group.⁴ In a back to back communication, we also showed that the metabolites target the cellular cytokinetic protein myosin.⁵ Few natural products have been demonstrated to target this protein.⁶

The pronounced bioactivity of the ammosamides, derived from a disruption in cellular cytokinesis, encouraged us to undertake a total synthesis of this unique class of metabolites. An examination of synthetic analogues with different functional groups on the pyrroloquinoline core, we thought, might provide a molecule with increased potency and/or overall improved properties. Although interest in related pyrroloiminoquinones culminated in several total syntheses,⁷ none of the reported routes was directly amenable to the ammosamide class.

wfenical@ucsd.edu.

Supporting Information Available: Details of the isolation and spectral data for **14**. Crystallographic data for *N*-methyl **4** (CCDC 756061) in CIF format. HRMS data, proton, and carbon NMR spectra for selected intermediates. The material is available free of charge via the Internet at <http://pubs.acs.org>.

We envisioned a total synthesis commencing from 4-chloroisatin (**3**), itself derived from reaction of 3-chloroaniline with chloral hydrate (Scheme 1).⁸ The synthesis was designed with several key principles in mind, namely that (1) the aromatic amines, to which we ascribed much of the natural products' hydrophilicity and lack of solubility, would be disguised as nitro groups; (2) the C-4 chlorine substituent in **3** would be displaced by ammonia (or its equivalent) in a nucleophilic aromatic substitution reaction; (3) the C-3 carbonyl of **3** would be elongated at a later stage through Wittig olefination; (4) the C-7 chlorine substituent would be installed late in the synthesis. A late-stage chlorination would mean that the synthesis could be adapted to provide deschloro derivatives or derivatives with bromine or iodine at C-7. Since comparable activity was observed for ammosamides A (**1**) and B (**2**), we pursued a synthesis specifically of **2** to avoid manipulations with the thiolactam in **1**. We had shown earlier that **2** could be converted to **1** using Lawesson's reagent.⁴

Isatin **3** was first mono-nitrated at 0 °C to give **4**. The position of the nitro group at C-5 was determined by X-ray crystallographic analysis of *N*-methyl **4**.⁹ Nitroisatin **4** was treated with *t*-butylamine in dioxane at elevated temperatures to produce **5** in 87% yield. *N*-methylation of this material was followed by acidic deprotection of the C-4 amine to give **6**. The quinoline ring was then constructed via olefination with *t*-butyl (triphenylphosphoranylidene) acetate and acidic deprotection/condensation. The Wittig reagent was added over a 3 h period using a syringe pump, minimizing the formation of phosphorus-containing by-products that are derived from conjugate addition of excess reagent to the intermediate enone.^{10,11} The quinoline scaffold was selectively nitrated at C-7, and the C-2 alcohol was then converted to the chloride to give **7** in 34% yield from **3**.

Treatment of **7** with hydroiodic acid led to reduction of the nitro groups to amines and substitution at C-4 with iodide (Scheme 2). This compound (not shown), and subsequent diamino intermediates, adopted physical characteristics similar to the ammosamides. Its solubility in organic solvents was severely limited, purification with silica gel chromatography became unmanageable, and the material assumed a dark purple color ($\lambda_{\text{max}} = 520 \text{ nm}$). Thus, the iodide was directly converted to **8** upon treatment with trifluoroacetic anhydride. Iodide **8** was then transformed into the corresponding C-4 nitrile employing copper (I) cyanide. Conversion of the aromatic amines to trifluoroacetamides improved yields and allowed for purification by silica gel chromatography. The amines were subsequently deprotected under acidic conditions. Ester **10** was obtained by conversion of the nitrile to the imide with methanolic KOH and subsequent acidic esterification.

Deschloro ammosamide B (**12**) was prepared after installation of the final nitrogen atom. First, ester **10** was converted to **11** to facilitate silica gel purification. Deprotection and conversion to the C-4 amide resulted from treatment with magnesium nitride.¹² The ¹H NMR spectrum of **12** was similar to that of ammosamide B (**2**) except for the additional C-7 signal at 6.16 ppm.

Ammosamide B (**2**) was also synthesized from ester **10** after chlorination in neat thionyl chloride under UV light (254 nm).¹³ The reaction represents the first example of the chlorination of an unsubstituted 1,3-dianiline. Synthetic **2**, indistinguishable from natural ammosamide B (LC/MS, NMR, HRMS), was obtained after trifluoroacetylation to **13** and amidation.

Though their biological properties are comparable, there is a dramatic difference in the stability of natural ammosamide B (**2**) and synthetic deschloro ammosamide B (**12**). Exposed to air and light, **12** readily degraded within a week via a nitroso intermediate.¹⁴ Natural **2**, however, could be stored in this way for much longer (degradation $t_{1/2} \sim 1 \text{ month}$). The exact role of the biosynthetic chlorination may be to confer stability toward oxidative degradation.

In an effort to generate additional analogues for structure-activity studies, several semisynthetic derivatives of the ammosamides were prepared. In one case, conversion of ammosamide A (**1**) to quinolinium **14** was accomplished via *S*-methylation and reduction with sodium borohydride (see Scheme 1). The conspicuous UV profile of this derivative was recognized later, in very low titer, in the LC/MS chromatogram of the crude CNR-698 extract. The structure of a third member of the ammosamide class, ammosamide C (**14**), was thus elucidated. Metabolite **14** is likely a biosynthetic precursor to **1** and **2**, the latter species arising from nucleophilic oxidation or sulfuration at C-2.

The ammosamides are one of the first natural products known to target myosin.⁶ Our synthesis of the ammosamides has provided new analogues with which to assess the role of each functional group on the pyrroloquinoline core. Based on observations of the deschloro compound, halogenation of the 1,3-diamino moiety helps protect the natural product from oxidative degradation. Studies detailing the bioactivity of each synthetic compound will soon be disclosed.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support was provided by the National Cancer Institute, NIH (under grant R37 CA44848). We thank Phil S. Baran (The Scripps Research Institute) for indispensable advice, and Drs. Arnold L. Rheingold and Antonio G. D'Pasquale (UCSD) for assistance with X-ray diffraction experiments.

References

1. Cragg GM, Grothaus PG, Newman DJ. *Chem Rev* 2009;109:3012–3043. [PubMed: 19422222]
2. (a) Fenical W, Jensen PR. *Nat Chem Biol* 2006;2:666–673. and references therein. [PubMed: 17108984] (b) Hughes CC, Prieto-Davó A, Jensen PR, Fenical W. *Org Lett* 2008;10:629–631. [PubMed: 18205372] (c) Kwon HC, Kauffman CA, Jensen PR, Fenical W. *J Org Chem* 2009;74:675–684. [PubMed: 19132943]
3. For a review: Antunes EM, Copp BR, Davies-Coleman MT, Samaai T. *Nat Prod Rep* 2005;22:62–72. [PubMed: 15692617]
4. Hughes CC, MacMillan JB, Gaudêncio SP, Jensen PR, Fenical W. *Angew Chem Int Ed* 2009;48:725–727.
5. Hughes CC, MacMillan JB, Gaudêncio SP, Fenical W, La Clair JJ. *Angew Chem Int Ed* 2009;48:728–732.
6. The marine natural product pentabromopseudilin also inhibits myosin motor activity: Fedorov R, Böhl M, Tsiavalieris G, Hartmann FK, Taft MH, Baruch P, Brenner B, Martin R, Knölker HJ, Gutzeit HO, Manstein DJ. *Nat Struct Biol* 2009;16:80–88.
7. For example: (a) Peat AJ, Buchwald SL. *J Am Chem Soc* 1996;118:1028–1030. (b) Roberts D, Joule JA, Bros MA, Alvarez M. *J Org Chem* 1997;62:568–577. [PubMed: 11671451] For the first and only synthesis of lymphostin: (c) Tatsuta K, Imamura K, Itoh S, Kasai S. *Tetrahedron Lett* 2004;45:2847–2850.
8. Sandmeyer T. *Helv Chim Acta* 1919;2:234–242.
9. Material for X-ray analysis was prepared via nitration of *N*-methyl-4-chloroisatin. This isatin derivative was identical (¹H NMR, LC/MS) to the product resulting from methylation of **4**.
10. This reactivity has been observed with other electron-deficient enones: (a) Faber K, Stueckler H, Kappe T. *J Heterocycl Chem* 1984;21:1177–1181. (b) Augustin M, Jeschke P. *J Prakt Chem* 1987;329:637–648.
11. With 4-amino-1-methyl-5,7-dinitroisatin, formation of the undesirable phosphocycle [LRESIMS: *m/z* (MH⁺) = 741] was predominant.

12. Veitch GE, Bridgwood KL, Ley SV. *Org Lett* 2008;10:3623–3625. [PubMed: 18642822] Concerning the potential for magnesium nitride to explode: Crane S. *Chem and Engineer News* 2009;87:2–4.
13. Thionyl chloride is reduced to sulfur monoxide. For similar examples: (a) Kim DH, Santilli AA. *Tetrahedron Lett* 1971;12:2441–2442. (b) Soidinsalo O, Wähälä K. *Phosphorus, Sulfur, and Silicon* 2007;182:2761–2767.
14. The difference in stability between a 2-halo-1,3-nitrosoanilines and their nonhalogenated analogues has been noted: Frumkin AE, Churakov AM, Strelenko YA, Smirnov, Tartakovsky VA. *Russ Chem Bull* 1999;48:2103–2107.

Scheme 1.
Synthesis of the ammosamide core.

Scheme 2.

Total synthesis of ammosamides A (1), B (2), C (14) and deschloro ammosamide B (12).

- 1:** ammosamide A (X = S)
2: ammosamide B (X = O)

Chart 1.
Ammosamides A (1) and B (2).