Published in final edited form as:

Biochem Biophys Res Commun. 2010 January 29; 392(1): 1–3. doi:10.1016/j.bbrc.2009.10.072.

Loss of ovarian function in mice results in abrogated skeletal muscle PPAR δ and FoxO1-mediated gene expression

Nicole H Rogers^{a,1}, James W Perfield II^{a,2}, Katherine J Strissel^a, Martin S Obin^a, and Andrew S Greenberg^a

^a Obesity and Metabolism, JM-USDA HNRCA at Tufts University, Boston, MA, USA, 02111

Abstract

Menopause, the age-related loss of ovarian hormone production, promotes increased adiposity and associated metabolic pathology, but molecular mechanisms remain unclear. We previously reported that estrogen increases skeletal muscle PPAR δ expression *in vivo*, and transgenic mice overexpressing muscle-specific PPAR δ are reportedly protected from diet-induced obesity. We thus hypothesized that obesity observed in ovariectomized mice, a model of menopause, may result in part from abrogated expression of muscle PPAR δ and/or downstream mediators such as FoxO1. To test this hypothesis, we ovariectomized (OVX) or sham-ovariectomized (SHM) 10-week old female C57Bl6/J mice, and subsequently harvested quadriceps muscles 12 weeks later for gene expression studies. Compared to SHM, muscle from OVX mice displayed significantly decreased expression of PPAR δ (3.4-fold), FoxO1 (4.5-fold), PDK-4 (2.3-fold), and UCP-2 (1.8-fold). Consistent with studies indicating PPAR δ and FoxO1 regulate muscle fiber type, we observed dramatic OVX-specific decreases in slow isoforms of the contractile proteins myosin light chain (11.1-fold) and troponin C (11.8-fold). In addition, muscles from OVX mice expressed 57% less myogenin (drives type I fiber formation), 2-fold more MyoD (drives type II fiber formation), and 1.6-fold less musclin (produced exclusively by type II fibers) than SHM, collectively suggesting a shift towards less type I oxidative fibers. Finally, and consistent with changes in PPAR δ and FoxO1 activity, we observed decreased expression of atrogen-1 (2.3-fold) and MuRF-1 (1.9-fold) in OVX mice. In conclusion, muscles from ovariectomized mice display decreased PPAR δ and FoxO1 expression, abrogated expression of downstream targets involved in lipid and protein metabolism, and gene expression profiles indicating less type I oxidative fibers.

Keywords

Estrogen; menopause; Atrogen-1/MAFbx; MuRF-1; Musclin; myogenin

INTRODUCTION

Menopause promotes increased visceral adiposity and type 2 diabetes [1], but mechanisms remain unclear. Skeletal muscle, responsible for greater than 70% of whole body insulin

For correspondence please contact: Andrew S Greenberg, MD, HNRCA Room 603, 711 Washington St, Boston, MA 02111, 617-556-3144 (office), 617-556-3224 (fax), Andrew.greenberg@tufts.edu.

¹Present Address: Department of Metabolism and Aging, The Scripps Research Institute, Jupiter, FL, USA, 33458; nrogers@scripps.edu

²Present Address: Departments of Nutrition & Exercise Physiology and Food Science, University of Missouri, Columbia, MO, USA 65211; perfieldj@missouri.edu

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

stimulated glucose uptake, expresses estrogen receptors and is thus a likely target of estrogenic actions.

Skeletal muscle is comprised of both type I “slow” oxidative fibers and type II “fast” glycolytic fibers. Peroxisome proliferators-activated receptor- δ (PPAR δ) is a transcription factor that potently drives skeletal muscle fat oxidation [2] and type I slow oxidative fiber development [3]. Importantly, obesity is inversely associated with the proportion of oxidative slow-twitch fibers [4]. We have previously shown that exogenous estrogen increases skeletal muscle PPAR δ expression in ovariectomized mice [5], suggesting ovariectomy promotes obesity in part by decreasing skeletal muscle PPAR δ expression. We demonstrate here that ovariectomized mice, as a model of menopause, display abrogated PPAR δ and FoxO1 related gene expression in skeletal muscle, with evidence of a shift towards less type I oxidative fibers. These gene expression changes could help explain the decreased energy expenditure and increased adiposity we previously observed in this model [6].

MATERIALS AND METHODS

Animals

Experiments were conducted in a viral pathogen free facility at the Jean Mayer-U.S. Department of Agriculture Human Nutrition Research Center on Aging at Tufts University in accordance with institutional animal care and use committee guidelines. SHM-ovariectomized (SHM, n=10) and ovariectomized (OVX, n=11) C57B/6J mice were purchased from Charles River, with operations performed at 10 weeks of age. Animals were individually housed with 12-h light/dark cycles and given free access to water and food (phytoestrogen free chow, Harlan Teklad).

Quantitative Polymerase Chain Reaction (QPCR)

After 12 weeks, mice were euthanized, and quadriceps muscles dissected and immediately snap frozen in liquid nitrogen and stored at -80°C . RNA was isolated using commercial spin kits (Ambion, Austin, TX) according to manufacturer’s instructions. RNA was quantified and checked for purity using the Nanodrop spectrophotometer (Nanodrop 1000, Wilmington, DE). cDNA was generated from 1 μg of RNA using AMV Reverse Transcriptase (Promega, Madison, WI), and QPCR performed using SYBR Green (Applied Biosystems 7300, Foster City, CA). Fold changes were calculated as $2^{-\Delta\Delta\text{CT}}$ compared to the endogenous control gene cyclophilin B. Primer sequences available upon request.

Statistical analyses

Comparisons were made using student’s T-tests and data presented are means $\pm$ SEM. Significance was defined as $p \leq 0.05$.

RESULTS

Quadriceps from OVX mice displayed significantly decreased PPAR δ expression (3.4-fold). In addition, expression of downstream targets of PPAR δ that mediate lipid oxidation were decreased in muscle from OVX mice. These genes include pyruvate dehydrogenase kinase-4 (PDK-4; 2.3-fold, $p \leq 0.001$), uncoupling protein-2 (UCP-2; 1.8-fold, $p \leq 0.001$) and the forkhead box transcription factor family member FoxO1 (4.5-fold, $p \leq 0.001$) (Fig. 1).

In addition to altering fuel utilization, PPAR δ and FoxO1 promote type I (slow oxidative) fiber development by increasing the expression of slow isoforms of contractile proteins (i.e. troponins, myosins) [7]. OVX resulted in dramatically decreased expression of the myosin light chain slow isoform (11.1-fold, $p \leq 0.001$), and troponin C slow isoform (11.8-fold, $p \leq$

0.05). Moreover, compared to SHM mice, quadriceps from OVX mice expressed 50% less myogenin ($p \leq 0.05$) and twice as much myoD ($p \leq 0.01$). These changes are consistent with a fiber type shift, as myogenin specifically promotes the development of type I oxidative fibers [8], while myoD promotes the development of type II glycolytic fibers [9]. Finally, the expression of musclin, which is negatively regulated by FoxO1 [10] and made exclusively by type II fibers [11], was increased 1.6 fold ($p \leq 0.05$) in muscle from OVX mice. These data, presented in Fig. 1, collectively suggest that OVX in mice results in a fiber type switch toward a less type I oxidative muscle.

PPAR δ [12] and FoxO1 [13] also regulate expression of atrophy gene-1/muscle atrophy F-box (Atrogin-1/MAFbx) and muscle ring-finger protein 1 (MuRF-1), which are ubiquitin ligases involved in muscle atrophy. OVX muscle displayed 2.3-fold less Atrogin-1/MAFbx ($p \leq 0.001$) and 1.9-fold less MuRF-1 ($p \leq 0.05$) expression (Fig. 1). Decreases in these genes are consistent with changes in PPAR δ and FoxO1 expression, and suggest a novel role for ovarian hormones in protein metabolism.

DISCUSSION

Estrogen receptor expression [14] makes muscle a likely target tissue for ovarian hormone action, but studies investigating the impact of ovarian hormones on skeletal muscle are limited. We show here that PPAR δ and downstream mediators of oxidative metabolism, FoxO1, PDK-4 and UCP-2, are decreased in muscle from OVX mice. Moreover, expression of slow isoforms of contractile proteins were decreased in quadriceps from OVX and SHM mice, consistent with other models demonstrating fiber type switching via ablated PPAR δ [15] or FoxO1 [7]. Also, similar to changes reported with FoxO1-ablation induced fiber-type switching in skeletal muscle [7], OVX mice displayed decreased myogenin and increased myoD [9] expression. Finally, and again consistent with less type I oxidative fibers, musclin, which is produced exclusively by type II fibers, was increased in OVX mouse muscle. Collectively, these data (summarized in Fig. 2) suggest that ovariectomy-induced changes in PPAR δ and FoxO1 may lead to an abrogated capacity for fatty acid oxidation and detrimental shifts in fiber type that are likely to contribute to ovariectomy-induced adiposity [6].

PPAR δ promotes catabolism of not only fatty acids, but also proteins, the latter mediated via upregulated atrogin-1/MAFbx and MuRF1 expression (muscle specific ubiquitin ligases involved in atrophy). Consistent with the observed decreased PPAR δ expression, we found atrogin-1 and MuRF1 expression to be attenuated in skeletal muscle from ovariectomized mice. These changes are consistent with observed links between estrogen status and protein turnover [16], as atrogin-1 and MuRF1 have been recently implicated as important regulators of healthy protein turnover [17].

We conclude that ovariectomy alters PPAR δ and FoxO1 related gene expression in murine skeletal muscle. These observations provide a potential explanation for the observed increase in adiposity and co-morbidities with menopause. Future detailed studies are warranted.

Acknowledgments

The authors would like to thank Dr. Richard Karas and Dr. Jennifer Sacke for their thoughtful insight during manuscript preparation.

References

1. Carr MC. The emergence of the metabolic syndrome with menopause. *J Clin Endocrinol Metab* 2003;88:2404–11. [PubMed: 12788835]

2. Wang YX, Lee CH, Tiep S, Yu RT, Ham J, Kang H, Evans RM. Peroxisome-proliferator-activated receptor delta activates fat metabolism to prevent obesity. *Cell* 2003;113:159–70. [PubMed: 12705865]
3. Gaudel C, Schwartz C, Giordano C, Abumrad NA, Grimaldi P. Pharmacological activation of PPAR {beta} promotes rapid and calcineurin-dependent fiber remodeling and angiogenesis in mouse skeletal muscle. *Am J Physiol Endocrinol Metab*. 2008
4. Karjalainen J, Tikkanen H, Hernelahti M, Kujala UM. Muscle fiber-type distribution predicts weight gain and unfavorable left ventricular geometry: a 19 year follow-up study. *BMC Cardiovasc Disord* 2006;6:2. [PubMed: 16403232]
5. D'Eon TM, Souza SC, Aronovitz M, Obin MS, Fried SK, Greenberg AS. Estrogen regulation of adiposity and fuel partitioning. Evidence of genomic and non-genomic regulation of lipogenic and oxidative pathways. *J Biol Chem* 2005;280:35983–91. [PubMed: 16109719]
6. Rogers NH, Perfield JW 2nd, Strissel KJ, Obin MS, Greenberg AS. Reduced energy expenditure and increased inflammation are early events in the development of ovariectomy-induced obesity. *Endocrinology*. 2009
7. Kitamura T, Kitamura YI, Funahashi Y, Shawber CJ, Castrillon DH, Kollipara R, DePinho RA, Kitajewski J, Accili D. A Foxo/Notch pathway controls myogenic differentiation and fiber type specification. *J Clin Invest* 2007;117:2477–85. [PubMed: 17717603]
8. Ekmark M, Gronevik E, Schjerling P, Gundersen K. Myogenin induces higher oxidative capacity in pre-existing mouse muscle fibres after somatic DNA transfer. *J Physiol* 2003;548:259–69. [PubMed: 12598590]
9. Hughes SM, Taylor JM, Tapscott SJ, Gurley CM, Carter WJ, Peterson CA. Selective accumulation of MyoD and myogenin mRNAs in fast and slow adult skeletal muscle is controlled by innervation and hormones. *Development* 1993;118:1137–47. [PubMed: 8269844]
10. Yasui A, Nishizawa H, Okuno Y, Morita K, Kobayashi H, Kawai K, Matsuda M, Kishida K, Kihara S, Kamei Y, Ogawa Y, Funahashi T, Shimomura I. Foxo1 represses expression of myosin, a skeletal muscle-derived secretory factor. *Biochem Biophys Res Commun* 2007;364:358–65. [PubMed: 17950246]
11. Banzet S, Koulmann N, Sanchez H, Serrurier B, Peinnequin A, Bigard AX. Myosin gene expression is strongly related to fast-glycolytic phenotype. *Biochem Biophys Res Commun* 2007;353:713–8. [PubMed: 17189616]
12. Constantin D, Constantin-Teodosiu D, Layfield R, Tsintzas K, Bennett AJ, Greenhaff PL. PPARdelta agonism induces a change in fuel metabolism and activation of an atrophy programme, but does not impair mitochondrial function in rat skeletal muscle. *J Physiol* 2007;583:381–90. [PubMed: 17540700]
13. Stitt TN, Drujan D, Clarke BA, Panaro F, Timofeyeva Y, Kline WO, Gonzalez M, Yancopoulos GD, Glass DJ. The IGF-1/PI3K/Akt pathway prevents expression of muscle atrophy-induced ubiquitin ligases by inhibiting FOXO transcription factors. *Mol Cell* 2004;14:395–403. [PubMed: 15125842]
14. Wiik A, Glenmark B, Ekman M, Esbjornsson-Liljedahl M, Johansson O, Bodin K, Enmark E, Jansson E. Oestrogen receptor beta is expressed in adult human skeletal muscle both at the mRNA and protein level. *Acta Physiol Scand* 2003;179:381–7. [PubMed: 14656376]
15. Schuler M, Ali F, Chambon C, Duteil D, Bornert JM, Tardivel A, Desvergne B, Wahli W, Chambon P, Metzger D. PGC1alpha expression is controlled in skeletal muscles by PPARbeta, whose ablation results in fiber-type switching, obesity, and type 2 diabetes. *Cell Metab* 2006;4:407–14. [PubMed: 17084713]
16. Toth MJ, Sites CK, Matthews DE, Casson PR. Ovarian suppression with gonadotropin-releasing hormone agonist reduces whole body protein turnover in women. *Am J Physiol Endocrinol Metab* 2006;291:E483–90. [PubMed: 16621894]
17. Fielitz J, Kim MS, Shelton JM, Latif S, Spencer JA, Glass DJ, Richardson JA, Bassel-Duby R, Olson EN. Myosin accumulation and striated muscle myopathy result from the loss of muscle RING finger 1 and 3. *J Clin Invest* 2007;117:2486–95. [PubMed: 17786241]

Figure 1. OVX mice have altered PPAR δ and FoxO1 mediated oxidative gene expression

Real-time PCR was used to determine expression of genes (relative to the endogenous control cyclophilin B) in quadriceps from sham-ovariectomized (SHAM, white bars) and ovariectomized (OVX, black bars) mice. MLC = myosin light chain. Error bars indicate SE of the mean. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 2.

Summary of OVX-induced gene expression changes in mouse quadriceps: OVX decreases PPAR δ expression, leading to decreased UCP-2, PDK4 and FoxO1 expression. Attenuated FoxO1 expression is concomitant with decreased myogenin and increased myoD expression, promoting less type I fiber development. Consistent with a fiber type shift, as only type II fibers produce musclin, OVX mice display increased musclin expression. Expression of PPAR δ and FoxO1 downstream targets involved in protein metabolism (e.g. MuRF-1, atrogin-1) are similarly decreased.