Published in final edited form as:

Environ Sci Technol. 2010 January 1; 44(1): 216–221. doi:10.1021/es802392u.

Exposures to Particulate Air Pollution and Nitro-Polycyclic Aromatic Hydrocarbons amongst Taxi Drivers in Shenyang, China

Justin P. Miller-Schulze[#], Michael Paulsen[@], Akira Toriba[&], Ning Tang[&], Kazuichi Hayakawa[&], Kenji Tamura^{\$}, Lijun Dong[†], Xuemei Zhang[†], and Christopher D. Simpson^{@,*}

[#] Department of Chemistry, University of Washington, Seattle, WA 98195, USA

[&] Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-Machi, Kanazawa 920-1192, Japan

^{\$} National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba 305-8506, Japan

[†] Shenyang Center for Disease Control and Prevention, NO. 37 Qishan Zhong Road Huanggu District, Shenyang, 110031, China

[@] Department of Environmental and Occupational Health Sciences, School of Public Health and Community Medicine, University of Washington, Box 357234, Seattle, WA 98195, USA

Abstract

Exposures to particulate matter (PM) of both 10–2.5 μm ($\text{PM}_{10-2.5}$) and below 2.5 μm ($\text{PM}_{2.5}$) were measured for a cohort of taxi drivers in Shenyang, China, during August of 2007. PM samples were collected inside and outside the taxi during the drivers' workshifts, and also inside the drivers' homes when they were off-shift. Ambient PM samples were also collected at a stationary location in Shenyang. Elemental carbon (EC) and organic carbon (OC) were also measured in PM collected on quartz filters inside the taxis as well as at the stationary site. Concentrations of three nitro-polycyclic aromatic hydrocarbons (NPAHs): 1-nitropyrene (1NP), 2-nitropyrene (2NP), and 2-nitrofluoranthene (2NFI) were determined in extracts of the PM samples by using a 2D-HPLC-MS/MS method. The 2NP and 2NFI concentrations did not change substantially with sampling location, but the 1NP concentrations were much higher in samples collected inside and outside the taxis as compared with sampling locations that were more removed from traffic. Concentration ratios of specific NPAHs were used to assess the atmospheric conditions in Shenyang during the sampling period. The relatively high ratios of 2NFI/1NP (~8–50) indicate an important contribution from secondary NPAH formation to ambient NPAH levels, especially for the non-taxi samples. The ratios of 2NFI/2NP (2.5–4.3) indicate that 2NFI is primarily formed via the hydroxyl-initiated reaction.

Introduction

Particulate matter (PM) of both anthropogenic and natural origin is a well-chronicled potential hazard to human health (1–4). The EPA has concluded that there are likely causal associations between both coarse ($\text{PM}_{10-2.5}$) and fine ($\text{PM}_{2.5}$) PM and cardiovascular and respiratory morbidity and mortality (2). A variety of the components of anthropogenically generated PM have been associated with these health hazards in epidemiology and toxicology studies, including metals, polycyclic aromatic hydrocarbons (PAHs) and their nitrated analogs, nitro-PAHs (NPAHs), as well as numerous other organic compounds (5). Therefore, determining

*Corresponding Author. Tel.: +1 206-543-3222; fax: +1 206 616 2687, simpson1@u.washington.edu (CD Simpson).

Supporting Information Available

Supporting information is available for this manuscript. This information is free of charge via the Internet at <http://pubs.acs.org>.

the contribution of specific sources to ambient PM levels is an important step toward more effective management of particulate air pollution and to reduced public health risk associated with PM exposure.

NPAH species are a particularly important constituent of ambient PM, due to their carcinogenicity and mutagenicity (5) as well as their utility in source apportionment (6). Distinct NPAH are formed via different routes of production. For example, 1-nitropyrene (1NP) has been proposed as a molecular marker for diesel particulate matter (DPM) since it is by far the most abundant NPAH in DPM while being much less abundant in PM derived from other sources (7). Other NPAH isomers, such as 2-nitrofluoranthene (2NFI) and 2-nitropyrene (2NP) have been shown to be formed exclusively via gas-phase atmospheric reactions (8,9). The concentration ratio of 2NFI to 1NP allows estimation of the contribution to ambient NPAH concentrations of atmospheric reaction formation routes as compared with direct emission from primary PM sources (5,10). In addition, the ratio of the concentration of 2NFI to 2NP has been used to assess the relative contribution of two distinct production routes for 2NFI (11,12). For the reasons discussed above, measurements of ambient levels of NPAHs can yield valuable information about the atmospheric reaction conditions, sources of PM emissions, and processes contributing to NPAH formation for a particular geographical region.

In this paper, exposures to the three NPAH species listed above are reported for a cohort of taxi drivers from Shenyang, China. Shenyang is a city of 7.2 million people (as of the 2000 census) in Liaoning Province, the political and economic center of northeast China (13). Shenyang and its surrounding areas are home to numerous heavy industries, including smelters, machinery and chemical plants, coke plants, and power plants (14). In addition, 8 million tons of coal is burned each year in Shenyang, 20% of which is burned for domestic heating and cooking (14). We chose to study taxi drivers in Shenyang because it was anticipated that their exposures to vehicle exhaust (and hence, to 1NP) would be relatively high. However, the data presented herein indicate that exposures to PM derived from secondary sources are also important in this population.

Materials and Methods

Sampling

Twenty-four taxi drivers were recruited for this study, and sample collection took place during August 2007. PM_{10-2.5} and PM_{2.5} were collected inside and outside the subjects' taxis during their workshift (designated as "Inside Car" and "Outside Car") and personal PM samples were collected following the work-shift until the start of the next workshift the following day (samples designated as "Home"). Each driver was monitored over one 24 hour period, and either four or five drivers were monitored simultaneously on a particular day. PM_{10-2.5} and PM_{2.5} were also collected at a stationary location in Shenyang, the Center for Disease Control building (designated "CDC"). The vehicles driven by the subjects were a mix of diesel vehicles and gasoline-natural gas hybrid vehicles. The samplers for the Inside Car and Outside Car samples were placed on the passenger seat and roof of the drivers' taxis, respectively. All windows were open during the drivers' workshifts. The Home samples were collected by placing samplers in drivers' homes, in their bedrooms or kitchen, for example. The drivers' house windows were expected to be left open during this sampling. The CDC samples were obtained outside the CDC building, but samplers were placed at locations away from streets. The PM_{10-2.5} and PM_{2.5} samples were collected using a Sibata ATPS-20H dual-stage impactor connected to a portable Sibata MP-Σ3 pump (Sibata Sci Tech, Tokyo, Japan). The pumps were operated at a flow rate of 1.5 L/min. Particles greater than 10 µm in diameter were removed on a metal impaction plate coated with grease positioned immediately downstream of the sampler inlet. Particles 10-2.5 µm passed around the impaction plate and were collected on a 10 mm glass fiber filter that was placed on the second impaction stage. Particles less than 2.5

µm were collected on a 20 mm Teflon filter located in the final stage of the sampler. The PM_{10-2.5} and PM_{2.5} mass concentrations were obtained by weighing the filters before and after the sampling, always after a storage period (24 h) in a temperature- and humidity-controlled room (ambient temperature, 23±0.2 °C, relative humidity, 35±5%), by an ultra-microbalance (sensitivity 0.1 µg, UMX-2, Mettler-Toledo, Inc., Columbus, OH, USA). In addition, PM_{10-2.5} and PM_{2.5} were collected on quartz filters at the Inside Car and CDC locations. These quartz filters were subsequently analyzed for elemental carbon (EC) and organic carbon (OC) by a commercial laboratory using NIOSH method 5040. Sampling was conducted for between 8 and 10 hours per filter. The Inside Car and Outside Car filter samples were collected during the day and the Home filter samples were collected during nighttime hours. Separate CDC filter samples were collected both during the day and at night. For the purposes of this sampling, “Day” was classified as the time period from 7–8 AM to 5–6 PM, and “Night” was classified as 6–7 PM to 7–8 AM.

Materials

Quartz (2500QAT-UP) and Teflon Fiberfilm™ (a borosilicate glass fiber coated with fluorocarbon, TX60A20) filters were obtained from Pall Life Sciences, Ann Arbor, MI, USA. The 2NF1, 4-nitropyrene (4NP), 3-nitrofluoranthene (3NFl) and deuterated (*d*₉) 1NP (1dNP) were obtained from Midwest Research Institute, Kansas City, MO, USA. 1NP used in the calibration solutions and positive control samples was purchased from Sigma-Aldrich, St. Louis, MO, USA, while 2NP was obtained from RT Corp, Laramie, WY, USA. All solvents used in the extraction procedure and 2D-HPLC-MS/MS analysis were of high purity, HPLC grade quality, and were obtained from Sigma-Aldrich or JT Baker, Mallinckrodt Baker, Inc., Phillipsburg, NJ, USA. Ascorbic Acid was also obtained from JT Baker. The low-volume Turbovap evaporative concentrator was made by Zymark Corp., Hopkinton, MA, USA. Syringe filters were purchased from PALL Corporation, East Hills, NY, USA. The column used for the separation of the NPAHs (AC1) was obtained from Waters Corporation (Milford, MA USA), whereas the column used for the separation of the reduced NPAHs (AC2) was from Agilent Technologies, Inc., Santa Clara, CA, USA.

Sample Preparation

The PM samples were extracted and analyzed over a period of 3 months. Approximately 15 PM extracts were analyzed per analysis sequence, along with sets of quality control samples and calibrants. The extraction procedure employed for these filter samples was adapted from an ultrasonic extraction procedure that has been reported previously (15). Filters were deposited in silanized 7.5 ml screw top glass test tubes. Filters were spiked with 60 pg internal standard (1dNP) and allowed to age for 30 min. 7 ml extraction solvent (CH₂Cl₂) was added to each test tube and the tubes were then sonicated for 60 min. at full power. The CH₂Cl₂ was then decanted into a second silanized test tube and evaporated to dryness in a low volume Turbovap evaporative concentrator. The extracts were reconstituted in 150 µl 75% ethanol / 25% 0.02 M sodium acetate/acetic acid buffer, pH=5.5 sonicated for 10 min. at full power, and then vortexed for 10 min. in a sample vortexer. The reconstituted extracts were then filtered through PTFE 0.45 µm syringe filters into autosampler vials with silanized vial inserts, crimp capped, and analyzed as described below.

Analytical Method for the Determination of NPAHs—The filter extracts were analyzed with a two-dimensional high performance liquid chromatography tandem mass spectrometry method (2D-HPLC-MS/MS). The method has been adapted from Miller-Schulze et al. (15), and is described in detail in the supporting information.

The 2D-HPLC-MS/MS method involves an injection of 40 µl of PM extract, and initial separation on the first analytical column (AC-1) followed by online reduction in a column

packed with a Pt/Rh catalyst. Passage through the reduction column causes the NPAH species and the deuterated internal standard to be fully reduced into the corresponding amino-PAH derivatives. The two-dimensional method uses a switching valve to toggle between trapping and eluting configurations: in the trapping configuration, the mobile phase carrying the reduced NPAH species is diluted 1:5 with water, causing the NPAH determinants to be retained on the front of the trapping column. In the eluting configuration, a higher strength mobile phase, described below, is pumped through the trapping column in the reverse direction. The reduced NPAH determinants and 1dNP are then separated from persistent interfering compounds on the 2nd analytical column (AC2) and detected using an electrospray ionization triple quadrupole mass spectrometer, operated in the multiple reaction monitoring (MRM) mode.

Quantification of Determinants

All three NPAH species were quantified using the same internal standard (1dNP) which was added at a constant level (400 fg/ μ l) to calibration solutions containing all three NPAHs at concentrations ranging from 2.5 to 5000 fg/ μ l. This set of calibration solutions was analyzed at the start of each analysis sequence, and selected calibration solutions were re-analyzed periodically throughout each sequence to monitor instrument performance. Calibration curves were constructed by plotting the ratio of the NPAH response to internal standard response versus NPAH concentration. All peaks, including those in the calibrants and for PM sample extracts, were integrated manually within the Agilent Masshunter software. The NPAH concentration in each sample was calculated from the appropriate regression equation, which employed 1/X weighting for all determinants.

Quality control (QC) samples were included with each batch of filters to give insight on batch-to-batch variations in assay performance, and blank contamination. The relevant results of these QC analyses are summarized here. The average NPAH concentrations calculated to be present in the field blanks were 2NP = 0.32 pg/m³, 2NFl = 0.55 pg/m³, and 1NP = 0.20 pg/m³. Comparison of these blank NPAH concentrations with subsequent data for the NPAH concentrations in Shenyang shows that the blank contamination is negligible. The average NPAH recovery of the extraction and analysis, as measured by dividing the 1dNP internal standard peak area for each sample in a particular analysis batch by the average of the two 1dNP “control” samples analyzed in that batch (vial inserts spiked with an equivalent amount of 1dNP and diluted appropriately with extract matrix), was 92%. Calculated concentrations of NPAH species in spiked filters were within 30% of the expected values: 2NP = 73% $\pm$ 8%, 2NFl = 79% $\pm$ 8%, 1NP = 84% $\pm$ 7%.

Results and Discussion

Samples were collected over a 5 day period between August 16 and August 20, 2007. During this period the meteorological conditions were as follows: ambient temperature = 22–27 °C; mean visibility = 2.6–9.7 miles; average wind speed = 1.9–6.9 knots; precipitation: 0–1 inches/24 hr.

The concentrations of the air contaminants measured on the filter samples for each sampling location are given in Table 1. With the exception of the PM metrics, all concentrations given represent the total determinant concentration for each location, i.e., the sum of the concentrations in both the PM_{10-2.5} and PM_{2.5} filters.

Figure 1 shows the multiple reaction monitoring (MRM) chromatograms for the NPAH analysis of a representative PM_{2.5} sample collected at a traffic-based sampling location. As the chromatogram in Figure 1 shows, all peaks are well resolved. The chromatograms for PM extracts from all sampling locations show similar peaks, albeit with different relative abundances. 1NP, 2NP and 2NFl were readily detected in all samples at concentrations well

above the detection limit of the assay. Two additional compounds are observed in the 2nd (bottom) m/z 218 $\Rightarrow$ 201 transition. Based on the analysis of additional nitro-PAH standard compounds during method development efforts, we speculated that these peaks were 4-nitropyrene (4NP) and 3-nitrofluoranthene (3NFl). We confirmed this with standard addition experiments, i.e., spiking previously analyzed extracts with 3NFl and 4NP and assessing the increase in response of the unidentified peaks. These experiments confirmed that the additional peaks are 4NP and 3NFl. 4NP has been shown to be a product of the same atmospheric processes that generate 2NFl (16), and 3NFl has been shown to be present in the primary emissions of diesel engines, second in concentration only to 1NP (17).

NPAH Concentrations

As the NPAH data in Table 1 shows, while the average concentrations of 2NP and 2NFl are similar across the different sampling locations, the 1NP concentration is dramatically higher for the traffic-based samples (Inside Car, Outside Car) compared to the sampling locations at least somewhat removed from traffic (Home, CDC). This observation is consistent with the fact that 1NP is present exclusively in primary emissions – predominantly in diesel exhaust. In contrast, 2NP and 2NFl are produced via atmospheric reactions with gas-phase PAH precursors, and, as such, would be expected to vary less between traffic and non-traffic sampling locations, as compared to 1NP.

Table 2 gives some concentrations of 1NP, 2NP, and 2NFl at locations around the world for purposes of comparison with the concentrations of these species found in Shenyang.

As Tables 1 and 2 show, the 1NP concentrations in Shenyang are towards the upper end of the range of values reported previously at other locations around the world. The traffic locations in Shenyang have values of 1NP concentrations in-line with the downtown/high traffic locations in Ho Chi Minh City, Vietnam and Denmark, while the non-traffic locations in Shenyang are closer to the non-traffic locations in Ho Chi Minh City and Baltimore, MD, USA in the summer.

The 2NP and 2NFl concentrations in Shenyang are substantially greater than those found in the locations listed in Table 2, although the levels of 2NFl in Shenyang are closest to those found in Ho Chi Minh City, as well as Torrance, CA, USA. This comparison suggests that the atmospheric reaction conditions, including the abundance of gas phase precursors, are more favorable for 2NP and 2NFl formation in Shenyang during the sampling period than at the locations studied in the works summarized in Table 2.

Another observation regarding the data reported in Table 1 is that the coefficients of variation (CVs) are consistently higher for 2NFl and 2NP (CVs ~100%) compared to 1NP (CVs ~30–50%). A major determinant of the high CVs for 2NFl and 2NP (and also for PM) appears to be substantial day-to-day variation in the concentrations of these contaminants (illustrated for 2NFl in Figure S2 of supporting information). In contrast, there is relatively little day-to-day variation in the 1NP concentrations-especially at the traffic locations (Figure S3, supporting information).

The two-stage impactor utilized in the current study permitted separate analysis of NPAHs in the fine and coarse PM fractions. For all locations approximately 90% of the mass of all NPAH species was found in the PM_{2.5} fraction, and the variation in this distribution between locations was quite low (CV <5%; see Table S1 in supporting information). Over 84% of EC was found in the fine PM fractions and PM_{2.5} constituted 57% of the PM₁₀ (range = 47–67%).

Two distinct pathways have been described for the atmospheric formation of 2NFl and 2NP: the hydroxyl radical-initiated pathway, which yields both 2NFl and 2NP, and the NO₃ initiated

pathway, which yields primarily 2NFI (22). Because the -NO_3 radical (and associated NO_x species, NO_2 and N_2O_5) can play an important role in atmospheric chemistry (11), particularly at night when these NO_x species exist at higher concentrations, the ratio of 2NFI to 2NP is of interest not only for assessment of NPAH formation, but also because it provides insight into the overall atmospheric chemistry of a specific location or airshed.

Figures 2a and 2b show the median values for the concentration ratios of 2NFI to 2NP and 2NFI and 1NP, respectively, for each sampling location.

The median values for the 2NFI/2NP concentration ratio range from 2.5–4.3 across all four locations. Studies by Arey, Zielinska, and colleagues (11) concluded that ratios of 2NFI/2NP close to 10 indicated 2NFI formation mostly via the hydroxyl-initiated pathway, whereas ratios closer to 100 indicated mostly -NO_3 initiated 2NFI formation. Therefore, our data indicate that formation of 2NFI and 2NP occurs primarily via the hydroxyl-initiated reaction in this airshed. It is important to note that the concentration of -NO_3 radical is episodic, being more abundant at night. Thus, one would expect that the impact of the -NO_3 initiated reaction and hence the ratio of 2NFI/2NP would be greater at night. However, examination of data from the CDC location where we collected separate, co-located daytime and nighttime filters, revealed that the 2NFI/2NP ratio was less than 10 in all samples, and on only one occasion was the nighttime 2NFI/2NP ratio significantly greater than the preceding daytime value. This observation supports our conclusion that – at least during the period of the current study – the hydroxyl-initiated reaction is the dominant source of 2NP and 2NFI. 2NFI/2NP ratios significantly below 10 have been observed in numerous areas around the world, with averages of 2.1 in Athens, Greece (23), 3.8 in Milan, 6.3 in Rome, 1.6 in Naples, 4.5 in Madrid, and 1.9 in Brazil (10), to list a few published values. The relative consistency of the 2NFI/2NP ratio in our study indicates that the formation process and reaction conditions that generate these two compounds are the same at all four locations in Shenyang during the sampling period.

There is also a positive linear correlation between the 2NFI and 2NP concentrations for all four locations, $r^2 = 0.631$ ($r^2 = 0.709$ if a single potential “outlier” is excluded). While this positive correlation does not necessarily implicate the hydroxyl-initiated pathway specifically for 2NFI and 2NP formation, it does suggest that the same process or combination of processes is prevalent over all sampling locations.

The ratio of the concentrations of 2NFI to 1NP has also been employed to assess the relative influence of primary sources as opposed to atmospheric formation for ambient NPAH levels (5,10,25). Some studies have implied that a 2NFI/1NP concentration ratio of 5 or greater is indicative of NPAH levels dominated by atmospheric reactions whereas a ratio less than 5 indicated a dominance of primary emissions (5,10,25). Notably, these benchmark ratios have been based on ambient measurements as opposed to the chamber studies of Arey and Atkinson described cited earlier. The concentration ratio of 2NFI to 1NP, grouped by sampling location, is illustrated in Figure 2b. As expected given differing formation mechanisms for these two compounds, the sampling locations close to vehicle traffic have lower ratios of 2NFI to 1NP, indicating that ambient NPAH levels are more strongly influenced by vehicle exhaust, specifically diesel exhaust. The median value for the ratio of 2NFI/1NP is below the benchmark ratio of 5 described above for the Inside Car samples (median 2NFI/1NP = 2.3) – indicating a dominance of primary emissions. However, the median 2NFI/1NP ratio is above the benchmark for the Outside Car samples (median 2NFI/1NP = 6.5) as are the mean values for both sample types (Outside Car: 9.10 ± 8.70 , Inside Car: 7.97 ± 7.89). This suggests that even for filter samples collected inside and outside of the taxis, the NPAH concentrations are still significantly influenced by secondary NPAH formation mechanisms in the atmosphere. The sampling locations removed from vehicle traffic have higher ratios of 2NFI to 1NP (median of 2NFI/1NP ratios-CDC: 49.7, Home: 19.4), indicating that the ambient NPAH concentrations at these

locations are dominated by photochemically derived NPAHs. In addition, the ratio of 2NFl to 1NP for the CDC location, which is presumably the most removed from traffic, is greater than the corresponding mean ratio for the home locations.

Associations between 1NP and EC

Elemental Carbon (EC) is currently used as a marker for diesel exhaust (7). However, use of EC as a surrogate measure for diesel exhaust exposure may be confounded by combustion sources other than diesel exhaust that emit EC (26). In the current study, EC was quantified for the Inside Car and CDC sampling locations contemporaneously with the other PM samples, and associations between EC and 1NP were explored.

The correlations between 1NP and EC for the Inside Car location and CDC locations are shown in Figures 3a and 3b. We observed a high correlation between 1NP and EC concentrations in the PM samples collected at the CDC location. In contrast, the 1NP and EC concentrations at the Inside Car sampling location show no correlation. The lack of correlation between the 1NP and EC concentrations at the Inside Car location is not consistent with the expectation that both EC and 1NP are markers of primary diesel emissions. Both the EC and 1NP concentrations are noticeably higher at the Inside Car location as compared with the CDC location, as would be expected for markers of diesel exhaust. However, in the case of 1NP, the difference is approximately seven-fold, whereas in the case of the EC concentrations, the difference is approximately two-fold.

The observation of a lack of correlation between the 1NP and EC concentrations at the Inside Car location (Spearman $r = -0.11$, $p=0.62$) combined with the strong correlation of these concentrations at the CDC site (Spearman $r = 0.905$, $p=0.002$) warrants some discussion. The ambient air in the city of Shenyang is heavily influenced by both industrial coal burning and residential coal burning, although the residential coal burning may not have been significant during the period of this study, as sampling was done in mid-August (14). Since EC is a significant constituent of coal combustion emissions (26), it is likely that coal combustion emissions confounded the expected association between EC and 1NP at the Inside Car sampling location. At the CDC location we found that all the air pollutants we measured were strongly correlated with EC (see supplemental information, Figure S4a–4d). The CDC location is the least influenced by direct traffic emissions, and in the current study it appears to reflect urban Shenyang PM derived from a consistent mixture of sources.

In conclusion, our data showed that for this cohort of taxi drivers, exposures to NPAH, particularly to 2NFl, were high compared to most of the ambient NPAH concentrations that have been previously reported in the literature. 1NP concentrations were ~4–5 fold higher during the drivers' workshift compared to the concentrations measured inside their homes. Therefore, we conclude that the major portion of the drivers' 1NP exposure occurred during their workshift, and, further, we anticipate that 1NP exposures are also high for individuals whose occupations place them in close proximity to traffic. In contrast, the high concentrations we observed for 2NFl exhibited little spatial variation, which indicates that a large fraction of the population of Shenyang is likely to have high exposures to 2NFl. The predominance of 2NFl compared to 1NP indicates a major contribution to ambient NPAH concentrations of secondary aerosol formed via atmospheric reactions with the parent PAHs, as opposed to primary emissions. The significance of atmospheric reactions for ambient NPAH concentrations is important as NPAH species in PM are responsible for at least some of the health risk associated with PM of anthropogenic origin.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported in part by grant number R21-ES014917 from the National Institute of Environmental Health Sciences (NIEHS), NIH USA. The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of NIEHS or NIH.

References

- Davidson CI, Phalen RF, Solomon PA. Airborne Particulate Matter and Human Health: A Review. *Aerosol Sci and Technol* 2005;39:737–749.
- U.S. Environmental Protection Agency, National Center for Environmental Assessment. Air Quality Criteria for Particulate Matter. Vol. I. U.S. EPA, National Center of Environmental Assessment-RTP Office, Office of Research and Development; Research Triangle Park, NC, USA: 2004.
- Liu LJS, Box M, Kalman D, Kaufman J, Koenig J, Larson T, Lumley T, Sheppard L, Wallace L. Exposure Assessment of Particulate Matter for Susceptible Populations in Seattle. *Environ Health Perspect* 2003;111:909–918. [PubMed: 12782491]
- Lena TS, Ochieng V, Carter M, Holguin-Veras J, Kinney PL. Elemental Carbon and PM_{2.5} Levels in an Urban Community Heavily Impacted by Truck Traffic. *Environ Health Perspect* 2002;110:1009–1015. [PubMed: 12361926]
- Hien TT, Thanh LT, Kameda T, Takenaka N, Bandow H. Nitro-polycyclic hydrocarbons and polycyclic hydrocarbons in particulate matter in an urban area of a tropical region: Ho Chi Minh City, Vietnam. *Atmos Environ* 2007;41:7715–7725.
- Cecinato A, Marino F, Di Filippo P, Lepore L, Possanzini M. Distribution of *n*-alkanes, polynuclear aromatic hydrocarbons and nitrated polynuclear hydrocarbons between the fine and coarse fractions of inhalable atmospheric particulates. *J Chromatogr A* 1999;849:255–264. [PubMed: 10420616]
- Scheepers PTJ, Martens MHJ, Velders DD, Fijneman P, van Kerkhoven M, Noordehoek J, Bos RP. 1-Nitropyrene as a Marker for the Mutagenicity of Diesel Exhaust-Derived Particulate Matter in Workplace Atmospheres. *Environ Mol Mutagen* 1995;25:134–147. [PubMed: 7535227]
- Arey J, Zielinska B, Atkinson R, Winer AM, Ramdahl T, Pitts JN Jr. The Formation of Nitro-PAH from the Gas-Phase Reactions of Fluoranthene and Pyrene with the OH Radical in the Presence of NO_x. *Atmos Environ* 1986;20(12):2339–2345.
- Sweetman JA, Zielinska B, Atkinson R, Ramdahl T, Winer AM, Pitts JN Jr. A Possible Formation Pathway for the 2-Nitrofluoranthene Observed in Ambient Particulate Organic Matter. *Atmos Environ* 1986;20(1):235–238.
- Ciccioli P, et al. Formation and transport of 2-nitrofluoranthene and 2-nitropyrene of photochemical origin in the troposphere. *J Geophys Res* 1996;101:19567–19581.
- Arey J, Zielinska B, Atkinson R, Aschmann SA. Nitroarene Products from the Gas-Phase Reactions of Volatile Polycyclic Aromatic Hydrocarbons with the OH Radical and N₂O₅. *Int J of Chem Kinet* 1989;21:775–799.
- Feilberg A, Poulsen MWB, Nielsen T, Skov H. Occurrence and sources of particulate nitro-polycyclic aromatic hydrocarbons in ambient air in Denmark. *Atmos Environ* 2001;35:353–366.
- Li LK, Clifford GM, Yao WQ, Arslan A, Li N, Shi JF, Snijders PJF, Meijer CJLM, Qiao YL, Franceschi S. Human papillomavirus infection in Shenyang City, People's Republic of China: a population-based study. *British Journal of Cancer* 2006;95:1593–1597. [PubMed: 17088908]
- Xu Z, Yu D, Jing L, Xu X. Air Pollution and daily Mortality in Shenyang, China. *Archives of Environmental Health* 2000;55(2):116–120.
- Miller-Schulze JP, Paulsen M, Toriba A, Hayakawa K, Simpson CD. Analysis of 1-nitropyrene in air particulate matter standard reference materials by using two-dimensional high performance liquid chromatography with online reduction and tandem mass spectrometry detection. *J Chromatogr A* 2007;1167:154–160. [PubMed: 17826788]
- Zielinska B, Arey J, Atkinson R, Winer AM. The Nitroarenes of Molecular Weight 247 in Ambient Particulate Samples Collected in Southern California. *Atmos Environ* 1989;1:223–229.
- Bamford HA, Bezabeh DZ, Schantz MM, Wise SA, Baker JE. Determination and Comparison of Nitrated-Polycyclic Aromatic Hydrocarbons Measure in Air and Diesel Particulate Reference Materials. *Chemosphere* 2003;50:575–587. [PubMed: 12685733]

18. Arey J, Zielinska B, Atkinson R, Winer AM. Polycyclic aromatic hydrocarbon and nitroarene concentrations in ambient air during a wintertime high-NO_x episode in the Los Angeles basin. *Atmos Environ* 1987;21(6):1437–1444.
19. Reisen F, Arey J. Atmospheric Reactions Influence Seasonal PAH and the Nitro-PAH Concentrations in the Los Angeles Basin. *Environ Sci and Technol* 2005;39:64–73. [PubMed: 15667076]
20. Bamford HA, Baker JE. Nitro-polycyclic hydrocarbon concentrations and sources in urban and suburban atmospheres of the Mid-Atlantic region. *Atmos Environ* 2003;37:2077–2091.
21. Murahashi T, Hayakawa K. A sensitive method for the determination of 6-nitrochrysene, 2-nitro-fluoranthene and 1-,2-, and 4-nitropyrenes in airborne particulates using high performance liquid chromatography with chemiluminescence detection. *Anal Chim Acta* 1997;343:251–257.
22. Atkinson R, Arey J, Zielinska B, Aschmann SA. Kinetics and Nitro-Products of the Gas-Phase OH and NO₃ Radical-Initiated Reactions of Naphthalene-*d*₈, Fluoranthene-*d*₁₀, and Pyrene. *Int J Chem Kinet* 1990;22:999–1014.
23. Marino F, Cecinato A, Siskos PA. Nitro-PAH in ambient particulate matter in the atmosphere of Athens. *Chemosphere* 2000;533–557. [PubMed: 10665391]
24. Tang N, Hattori T, Taga R, Igarashi K, Yang X, Tamura K, Kakimoto H, Mishukov VF, Toriba A, Kizu R, Hayakawa K. Polycyclic aromatic hydrocarbons and nitropolycyclic aromatic hydrocarbons in urban air particulates and their relationship to emission sources in the Pan-Japan Sea countries. *Atmos Environ* 2005;39:5817–5826.
25. Albinet A, Leoz-Garziandia E, Budzinski H, Villenave E, Jaffrezo J-L. Nitrated and oxygenated derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleys Part I: Concentrations, sources and gas/particle partitioning. *Atmos Environ* 2008;42:43–54.
26. Schauer JJ. Evaluation of elemental carbon as a marker for diesel particulate matter. *J of Exposure Anal Environ Epidemiol* 2003;13:443–453.

Figure 1.
2D-HPLC-MS/MS Chromatograms for a traffic based sampling location. Top pane represents the 1st time segment and the m/z transition from m/z 218 $\Rightarrow$ 201. Middle pane represents the m/z transition from m/z 227 $\Rightarrow$ 210 in the 2nd time segment. Lower pane represents the m/z transition from m/z 218 $\Rightarrow$ 201, also in the 2nd time segment. Peaks annotated in figure, see text for abbreviations.

Figure 2.

Figure 2a: Median concentration ratios of 2NFI to 2NP grouped by sampling location. Inside and outside car samples were collected during daytime. Home samples were collected at night and CDC samples were collected both during the day and at night.

Figure 2b: Median concentration ratios of 2NFI to 1NP grouped by sampling location.

Figure 3. Figures 3a and 3b: Relationship of 1NP to EC for (A) Inside Car and (B) CDC sampling locations.

Table 1

Air contaminants measured at various sample locations in Shenyang. NPAH, EC and OC concentrations are the total of the determinant concentration in both the PM_{10-2.5} and PM_{2.5} size fractions.

Location	2NP (pg/m ³)	2NFl (pg/m ³)	INP (pg/m ³)	EC (µg/m ³)	OC (µg/m ³)	PM _{2.5} (µg/m ³)	PM _{10-2.5} (µg/m ³)
Inside Car (n=23)	Mean ± Std Dev	192±199	552±477	80.8±26.0	25.7±5.6	53.4±7.1	101±44.5
	Median	55.0	195	79.3	25.0	52.8	69.9
	Range	(15.2–555)	(73.9–1270)	(35.1–150)	(17.2–40.2)	(41.7–67.5)	(51.6–173)
Outside Car (n=24)	Mean ± Std Dev	232±231	753±614	97.1±31.1	NA**	119±62.4	112±50.4
	Median	157	690	95.6	NA**	92.6	108
	Range	(14.9–614.)	(89.9–1820)	(38.7–187)	(58.7–322)	(44.4–244)	
Home (n=24)	Mean ± Std Dev	210±280	533±485	22.4±11.4	75.2±33.0	51.0±36.7	
	Median	99.9	368	18.7	NA**	73.2	40.3
	Range	(14.8–999)	(29.0–1510)	(7.9–49.7)	(25.1–130)	(6.2–124)	
CDC (n=8)	Mean ± Std Dev	369±546	767±723	11.2±6.0	28.5±12.1	81.5±54.4	50.6±45.4
	Median	110	723	12.0	32.0	69.7	38.7
	Range	(8.4–1550)	(30.5–1720)	(3.3–18.0)	(12.4–47.1)	(24.6–158)	(7.6–124)

Plus/Minus values represent 1 standard deviation.

** EC and OC values for Outside Car and Home locations not available as quartz filters were not deployed at these locations

Table 2

Summary of selected NPAH concentrations in literature

Sampling Location (City)	Sampling Location (environment type)	Sampling Date	2NP pg/m ³	2NF1 pg/m ³	INP pg/m ³	Reference Number
Ho Chi Minh City, Vietnam	Urban	1/05-3/06	n/a	165±65	8.1±3.9	5
	Urban	1/05-3/06	n/a	190±97	9.1±3.9	
	Traffic Site	1/05-3/06	n/a	191±64	73±40	
Denmark	Urban/High Traffic	12/96-6/96,	20±5	91±54	127±44	12
	Rural/Open-Land	2/98-2/99	8±4	60±22	30±15	
Torrance, CA	Urban	2/86	30	400	30	18
Los Angeles (LA) and Riverside (Riv), CA	Traffic (LA)	8/02	6.2	62.8	6.5	19
	Traffic (LA)	1/03	7.8	74.2	21	
	Downwind (Riv)	8/02	3.1	154	6.5	
	Downwind (Riv)	1/03	4.3	79.2	12.2	
Baltimore, MD.	Urban (Baltimore)	1/01	6.5	60	27	20
Fort Meade, MD.	Urban (Baltimore)	7/01	2.7	99	8.1	
	Rural (Ft. Meade)	1/01	4	49	20	
	Rural (Ft. Meade)	7/01	0.76	28	1.4	
Kanazawa, Japan.	Urban/Downtown	10/94	4.45	59.34	32.14	21
	Suburban	10/94	1.56	16.56	1.68	