Published in final edited form as:

Vulnerable Child Youth Stud. 2009 March ; 4(1): 67–82. doi:10.1080/17450120802385729.

Maternal HIV/AIDS and adolescent depression: A covariance structure analysis of the “Parents and Adolescents Coping Together” (PACT) model

Debra A. Murphy, Ph.D.¹, William D. Marelich, Ph.D.², and Hortensia Amaro, Ph.D.³

¹ Health Risk Reduction Projects, Integrated Substance Abuse Programs, Department of Psychiatry, University of California at Los Angeles

² Department of Psychology, California State University, Fullerton

³ Bouve College of Health Sciences, Northeastern University

Abstract

The current study assessed the efficacy of selected variables from the Parents and children Coping Together (PACT) model, which was designed to predict maternal HIV effects on child/adolescent outcomes. Data from two longitudinal studies applying PACT measures were utilized, encompassing a seven-year assessment timespan for HIV-infected mothers and their children. Both maternal and child-based measures were evaluated, and a sequential longitudinal design was adopted. Structural equation modeling using FIML was performed to assess the proposed model. Results show the PACT model was viable in predicting child/adolescent outcomes of self-concept and depression. Study implications are discussed, including the influences of maternal factors on child's self-concept and depression, and a reconsideration of the affect of family cohesion on child/adolescent outcomes.

Keywords

Maternal HIV; Child self-concept; Child depression; Structural equation modeling

The advent of new antiretroviral medication regimens has shifted HIV from a terminal illness to a chronic disease (Valdiserri, Holtgrave, & West, 1999). Therefore, children are living longer with chronically ill parents who have a highly stigmatized disease. Chronic illness is a major stressor for family members, and has been linked to increased psychological distress in adults (Derogatis et al., 1983; Woods, Haberman, & Packard, 1993) and their children (Armistead, Klein, & Forehand, 1995; Worsham, Compas, & Ey, 1997). Only recently have the effects of parental HIV on early and middle age adolescents begun to be examined (Murphy, Marelich, Hoffman, & Schuster, 2006).

Framework for Maternal HIV Effects on Child Outcomes

The literature predicts that both parent and child background and situational factors will affect long-term child outcomes (i.e., behavioral adjustment, mental health, and social adjustment) in response to parental illness. A model of maternal HIV effects on child outcomes has been developed for a long-term study of mothers living with HIV/AIDS and their well children (cf. Murphy, Marelich et al., 2006), the Parents and children Coping Together (PACT) study. The

five-year PACT study was designed to longitudinally assess mothers living with HIV and their young, well children age 6 to 11 years. A subsequent study, Parents and Adolescents Coping Together (PACT II), followed up the majority of these families through child transition to early and middle adolescence. The PACT model (see Figure 1) was adapted from work by Sandler (Sandler, Reynolds, Kliwer, & Ramirez, 1992; Sandler, Tein, & West, 1994) describing child and maternal situational and background factors anticipated to affect child adjustment in response to disruptive events (e.g., divorce and parental death). In the PACT model, background and intermediate outcomes of parents and children are anticipated to impact child adjustment and long-term child outcomes. In this study, analysis of a longitudinal model was conducted to determine the relationship between selected background and intermediate factors with child depression (see bolded variables in Figure 1). The support from the literature for each of the variables in the model selected for this study, and their links and hypothesized paths to other variables in the model, will be briefly reviewed.

Maternal and Child/Adolescent Background Factors

Ethnicity

Ethnicity is a powerful variable in family responses to HIV, shaping perceptions of illness, health care use, and attitudes toward providers (Chavez, Hubbell, McMullin, Martinez, & Mishra, 1995; Clark, 1998; Sargent & Brettell, 1996). Among HIV-positive adults, ethnicity is a factor in treatment status, with ethnic minorities, primarily African Americans, more likely to be untreated (Kalichman, Graham, Luke, & Austin, 2002). Further, rates of disclosure to family and friends are significantly lower for African Americans than for whites (Petrak, Doyle, Smith, Skinner, & Hedge, 2001).

Maternal Health

The effects of maternal illness on child outcomes among families affected by HIV have been mixed. Some studies suggest that as illness progresses, mothers with HIV are likely to exhibit a number of maladaptive behaviors that disrupt relationships with their children (Cates, Graham, Boeglin, & Tiekler, 1990; Lamping et al., 1991). For example, when HIV-infected mothers remain healthy, their children are less likely to exhibit depressive symptoms (Murphy, Marelich, & Hoffman, 2002). Dorsey et al. (1999) found a linear increase in children's report of externalizing and internalizing difficulties as their mothers progressed through stages of HIV infection and AIDS. However, in some studies maternal HIV was not found to significantly add to the risk for child problem behaviors (Mellins, Brackis-Cott, Dolezal, & Meyer-Bahlburg, 2005).

Maternal and Child/Adolescent Intermediate Factors

Disclosure

Maternal disclosure of HIV can affect child adjustment, parent-child relationships, and the ability of parents to access resources. Many HIV-positive mothers do not disclose to their children because they believe the child might be too young to understand, or are concerned the child will disclose to others or be angry or fearful (Murphy, Steers, & Dello Stritto, 2001). However, overall, research indicates support for disclosure to children: (1) clinicians advise parents to disclose (Armistead & Forehand, 1995; Zayas & Romano, 1994); (2) mothers who have disclosed report significantly lower levels of depression (Wiener, Battles, & Heilman, 1998), and stronger family cohesion following disclosure; and (3) mothers typically report that their children are emotional at first, but exhibit no lasting negative impact (Kennedy et al., 2007; Murphy, Roberts, & Hoffman, 2006; Schrimshaw & Seigel, 2002). Moreover, children who have been disclosed to appear to be doing better on a number of measures compared to

children who have not (e.g., Murphy, 2008; Murphy, Marelich, & Hoffman, 2002; Murphy et al., 2001).

Family Functioning

Parental illness and family functioning have been linked in numerous research studies. For example, Dura and Beck (1988) found that families with maternal diabetes or chronic pain reported lower cohesiveness than families with no illness. Parental perceptions of the extent to which illness interferes with daily life has been inversely associated with perceptions of family cohesiveness (Mikail & von Baeyer, 1990). Moreover, family functioning has long been associated with child outcomes. In particular, family cohesion is associated with child resilience (Carbonell, Reinherz, & Giaconia, 1998), and has been found to mediate the relationship of negative life events to child depression (Roosa, Dumka, & Tein, 1996). In a large study HIV-positive parents in a European multi-center study (Nostlinger, Bartoli, Gordillo, Roberfroid, & Colebunders, 2006), with respect to influencing factors on children's emotional and behavioral symptoms, healthy general family functioning emerged as the best predictor.

Child/Adolescent Self-concept

Identity develops through processes of individuation and connectedness within the family (Hamilton, 1996). Having an HIV-infected mother is likely to affect developmental processes related to identity formation occurring in early and middle adolescence, and thus influence self-concept. Children of women with breast cancer and of women with diabetes tend to score significantly lower on self-concept than children in a comparative control sample (Armsden & Lewis, 1994). Reyland, McMahon, Higgins-Delessandro, & Luthar (2002) found that children of HIV-seropositive mothers had lower self-esteem than children attending public school in the same community. But other factors may mediate parental illness effects on child self-concept. Murphy et al. (2001) found that among children affected by maternal HIV, for those children whose mothers had disclosed their HIV-positive serostatus, lower levels of negative self-esteem were found. Finally, behavioral and emotional problems may impact self-concept. Adolescents with depression have been shown to have a worse self-concept (Kolaitis & Liakopoulou, 2005; McGrath & Repetti, 2002).

Child Mental Health, Behavioral, and Social Outcomes

As can be seen by the previous review of background factors and intermediate factors that may affect child outcomes, children affected by parental HIV/AIDS are at risk for poorer mental health, behavioral, and social outcomes. Somatic illness in a parent has long been considered a risk factor for psychological problems in children. Among families where the mother had breast cancer, one-third of the patients reported an increase in behavioral disorders in their younger children (Buckley, 1977). In a review of 15 years of literature on children of somatically ill parents, Romer, Barkmann, Schulte-Markwort, Thomalla, and Riedesser (2002) found that children of seriously ill parents had higher scores than controls on symptom scales, with a tendency towards internalizing symptomatology. Murphy, Greenwell, Mouttapa, Brecht, & Schuster (2006) have found that stability in HIV-infected mothers' health appears to be associated with a more rapid improvement in children's mental health over time.

Summary

In the present study, selected variables from the PACT model (see Figure 1) were analyzed. While some of these variables and associations have been evaluated in the PACT sample in many studies, this study takes a longitudinal perspective and includes more variables from the model to establish a "bigger picture" of these families. Within the constraints of our limited

sample size and statistical power, we included as many of the PACT variables as possible for formal model assessment. Maternal background factors of ethnicity and health status were included in the analysis. Maternal intermediate factors included: disclosure of HIV status; family functioning; and maternal coping (specifically, physical functioning). The child intermediate factor included in the analysis was self-valuing (measured as self-concept). The main child outcome was mental health, specifically child depression.

Method

Participants

One hundred thirty-five mothers living with HIV/AIDS and their young, well children 6 to 11 years old were recruited from 14 sites in Los Angeles County (11 clinical primary care sites and 3 AIDS service organizations) from November 1997 to June 1999 into the “Parents and children Coping Together” (PACT) study. “Parents and Adolescents Coping Together” (PACT II), followed the majority of these families when the children were transitioning to early and middle adolescence.

Of the 135 mothers recruited for PACT I, 81 remained in the study across all 30 months, re-consented to participate in PACT II, and provided data at the PACT II baseline (36 additional families were recruited from June 2003 through October 2004 using the same procedures outlined above [with children age 11 – 14 ½ years], making the total sample of 118 for PACT II – however, for the current study only those available from both PACT studies were retained). PACT II baseline was conducted approximately three years after the last PACT I follow-up, and assessments were conducted every six months. Four time points from the PACT studies combined are included in the current analysis: PACT I baseline interview and follow-ups at 30 months, 66 months, and 84 months (numbered in 6-month increments from the first assessment – month total includes break between PACT I and PACT II grants). Some minor differences between the 81 mother/child pairs who reconsented for PACT II and the 54 who did not reconsent have previously been noted (see Murphy, Greenwell et al., 2006, for attrition analysis). No child differences were found, and only two maternal differences were evident; mothers who could not participate in PACT II due to losing custody of the child reported higher family conflict scores, and mothers who had died (and thus could not participate) showed lower CD4 counts than those who reconsented.

At PACT I baseline for the 81 participants, mothers’ mean age was 35.0 years ($SD = 5.96$; ranging from 23 – 52 years); 34.6% African American, 30.9% Latina, 25.9% white, and the remainder other/mixed ethnicity. Those married comprised 19% of the sample, while 36% reported never being married, 23% widowed, 14% separated, and 9% divorced. Approximately 26% had completed high school, while 51% had not completed high school--the remainder had completed some college, had a college degree, or had attended a technical school. The majority (81.5%) were unemployed.

The mean age of children in the study at PACT I baseline was 8.43 ($SD = 1.83$; ranging from 5 – 12 years using mothers’ report). At 66 months (PACT II baseline), mean age was 13.44 ($SD = 1.93$; ranging from 10 – 17 years). By 84 months, mean age was 14.83 years old ($SD = 1.90$; ranging from 12 – 19 years). Overall, 53.1% of the children were male.

Inclusion criteria for PACT I were: mother had AIDS or was HIV symptomatic; she had a well child age 6 – 11; and she was English or Spanish speaking. HIV symptomatic was defined using the CDC Guidelines for CD4+T Lymphocyte Category 2 and Clinical Category B, including a CD4 count between 200 – 400 and the occurrence of a specified opportunistic disease or the occurrence of diseases for which treatment was complicated by HIV. Medical

chart abstraction was conducted to verify eligibility. Of the 214 mothers contacted for study participation, 24% were ineligible based on the above criteria, and 13% chose not to participate.

Procedures

Clinic staff at recruitment sites referred eligible families, and flyers and brochures allowed patients/clients to contact study staff directly. Interviewers only discussed the mother's diagnosis with children when prior child knowledge was confirmed through both parent report and child confirmation of that knowledge through an indirect screening. After receiving a description of the study, mothers who agreed to participate signed the IRB-approved Informed Consent forms and children signed the assent form. Trained bilingual interviewers conducted face-to-face interviews in the family's home.

For both PACT I and PACT II, interviews of mothers and children/adolescents were conducted separately using a computer-assisted interviewing program (CAPI) on laptop computers. Interviews with mothers lasted 1.5 hours and interviews with children/adolescents lasted 1 hour. Mothers received an incentive of \$35.00 and children/adolescents received \$25.00.

Measures

Background Factors

Maternal ethnicity (baseline): Mothers were asked to report their ethnicity (Asian/Pacific Islander; Black or African-American; Native American/American Indian; White; Hispanic [Spanish] or Latino Origin, or Other or Mixed Ethnicity).

Maternal disclosure (baseline): Mothers were asked how much their child knew about their HIV status. If the child knew the mother had HIV or AIDS, disclosure status was coded with a "1", otherwise disclosure was coded "0".

Maternal viral load and CD4 cell count (baseline): Mothers' health status was assessed using viral load and CD4 cell counts from medical records abstraction. Missing medical abstract CD4 cell count and viral load values were replaced with available self-report measures (a Pearson correlation of medical abstract CD4 and self-report CD4 was .77; viral load correlation between medical abstract/self-report was .74). Eight cases were missing baseline CD4 and were replaced with self-report. Seven cases were missing viral load, and four were replaced with self-report. Viral load was highly skewed, and a log transformation was applied (a "1" constant was added prior to transformation due to zero values) to adjust the distribution.

Child gender and age (baseline): Mother's report of child's age and gender was collected.

Maternal physical functioning (66 months): The Medical Outcome Short Form 36 was administered to mothers (Ware & Sherbourne, 1992), and the 10-item physical functioning subscale was chosen because it is based on a mother's report of her activity limitations rather than how she felt. Higher scores indicate better functioning. Internal consistency reliability for the current sample is .91.

Intermediate Factors

Family functioning (30 months): The sub-scale measuring family cohesion from the Family Functioning Scale (Bloom & Naar, 1994) was administered to mothers. Higher scores indicate more family cohesion. Internal consistency reliability for the current sample is .66.

Self-concept (66 months): The Piers-Harris Children's Self-Concept Scale (Piers, 1993) was administered. Subscales for this study and alphas are: Physical Appearance and Attribution, .84, Happiness and Satisfaction, .63, Popularity, .71, and Intellectual and School Status, .77.

Child Outcomes

Child depression (84 months): The Children's Depression Inventory (CDI; Kovacs, 1992) was administered. The scale consists of five subscales: Negative Mood, Interpersonal Problems, Ineffectiveness, Anhedonia, and Negative Self-Esteem. Total score reliability for the current sample is .91.

Analysis

A sequential longitudinal design (MacCallum & Austin, 2000) was adopted to test the PACT model. Model evaluation was performed using EQS (Bentler, 2006) applying full information maximum likelihood (FIML). Beyond direct associations, indirect effects of exogenous variables using a Sobel test extension were evaluated (Bentler, 2006; see James, Muliak, & Brett, 2006, for SEM mediation interpretation). Stability of the model fit statistics and parameter estimates were assessed using a model-based bootstrap procedure, with resampling fixed to our sample N , and requesting 1000 bootstrap samples. Observed means and standard deviations for indicator variables are presented in Table 1.

Results

To assess the model, due to limited sample size, select measurement variables were used representing different points of time. Final predictor variables include baseline measures from PACT I (mother's ethnicity, illness status [CD4, viral load], serostatus disclosure), a 30-month follow-up measure from PACT I (mothers' perception of family cohesion), and baseline measures (66-months) from PACT II (mothers' physical functioning; the Piers-Harris subscales representing the latent construct of child's self-concept). Child depression (our main outcome) was taken at 84-month follow-up from PACT II, consisting of the five CDI subscales representing the latent construct of child depression. See Table 2 for observed pair-wise correlations between the 15 measured variables. For the Piers-Harris and CDI subscales, a latent construct approach was adopted since the subscales function as multiple indicators around the construct centroid (Little, Lindberger, & Nesselrode, 1999) which may lead to better construct measurement as opposed to global indicators.

Two proposed background factors (gender, age) are not included in the final proposed model. Early exploratory runs yielded poor associations between these variables and adolescent depression, and between those variables and other model variables. Since we had to be selective in the measured variables, both gender and age were removed as exogenous variables.

To assess the proposed model (see Figure 2), disturbance terms for the constructs with multiple indicators (e.g., self-concept, child depression) were allowed to vary and provided start values of 5, while one indicator path was set to 1.0 for identification purposes. Error variances for all endogenous variables were allowed to vary. Regarding assumptions, two cases were possible multivariate outliers, but were retained due to small sample size considerations. Multicollinearity was not evident based on bivariate correlations of model variables (see Table 2), multicollinearity diagnostics on the observed data, or tolerance defaults in EQS. The assumption of multivariate kurtosis (assessed using the Bonett-Woodward-Randall test; Bentler, 2006) was violated. Thus, "robust" model fit indices were requested, including the Yuan-Bentler Scaled χ^2 , the comparative fit index (CFI), the incremental fit index (IFI), the non-normed fit index (NNFI), and the root mean squared error of the approximation (RMSEA). Robust standard errors were used to assess model parameter significance.

A test of Figure 2 yielded a Yuan-Bentler Scaled χ^2 of 253.03 ($df = 84$), with a CFI of .94, an IFI of .94, an NNFI of .92, and a RMSEA of .111 (90% C.I. = .085 to .135). Since these indices suggest ill fit, the standardized residuals from the model were evaluated to improve fit. One indicator, the log of viral load, exhibited a large standardized residual and was removed from the model. Nonsignificant paths ($p > .10$) were also removed. A new model (Figure 3) yielded a Yuan-Bentler Scaled χ^2 of 184.86 ($df = 76$), with a CFI of .98, an IFI of .98, an NNFI of .98, and a RMSEA of .059 (90% C.I. = .000 to .091), suggesting good fit. Total variance accounted for in child self-concept by the predictors was 21%, and 45% was accounted for in child depression.¹ Model-based bootstrap verified the stability of these findings. Of the 1000 samples drawn, 995 converged with a mean average Yuan-Bentler Scaled χ^2 of 165.23 (mean $SD = 29.87$), placing the final chi-square test statistic within one standard deviation of the 995 replications (similar validation was found for the CFI and RMSEA – no statistics are available for the other measures). Parameter estimates were assessed calculating new z-statistics based on the bootstrap standard error means – all maintained the same level of significance as noted in our final model.

Final model parameter estimates are in Figure 3. All paths are significant at .05 or better except from family cohesion to child self-concept ($p < .10$). As hypothesized, disclosure was positively associated with family cohesion and self-concept; those mothers who had disclosed their HIV status to their families reported higher levels of physical functioning. Higher CD4 counts at baseline were also associated with higher reports of mothers' physical functioning. However, counter to hypotheses, higher family cohesion was associated with lower levels of physical functioning, lower levels of self-concept, and higher levels of child depression. As hypothesized, higher self-concept was associated with lower levels of child depression.

A number of indirect associations were noted ($p < .05$) on child depression utilizing a Sobel test extension. Although these indirect associations are suggestive of mediation effects, a definitive assessment cannot be made since the complete complement of PACT model variables were not available. However, we have retained the mediation labels suggested by James et al. (2006) since they are descriptive of the variable relationships. Associations resembling complete mediation (indirect effects only are noted for a predictor on an endogenous outcome) were noted for both CD4 count (through physical functioning and child self-concept) and physical functioning (through child self-concept). Higher CD4 counts were indirectly associated with lower child depression (standardized indirect coefficient [IC] = $-.04$), and higher physical functioning was indirectly associated with lower child depression (IC = $-.15$). Associations resembling partial mediation (both indirect and direct effects are noted for a predictor on an endogenous outcome) were noted for family cohesion (through child self-concept) on child depression; higher family cohesion was indirectly associated with higher child depression (IC = $.08$).

Discussion

This was a preliminary study of the model for the "Parents and children Coping Together" (PACT) project. Overall, partial support was found for the PACT model (Murphy, Marelich et al., 2006), evidencing associations through which maternal illness level affects

¹A second model was evaluated replacing child's self-concept with child's coping to assess further the effects of a child intermediate factor on depression. Coping was measured using four items from the Child General Coping Efficacy scale (Sandler et al., 1994); the degree to which coping strategies employed during the past month were effective in making them feel better; the degree to which they were satisfied with the strategies they employed; how well they felt they handled their problems compared to other kids; and how well they thought they would cope in the future. Fitting the same paths from the Figure 3, fit statistics showed a Yuan-Bentler Scaled χ^2 of 193.48 ($df = 76$), with a CFI of .98, an IFI of .98, an NNFI of .98, and a RMSEA of .060 (90% C.I. = .000 to .092). Although model fit is good, a major difference between paths in this model and those in Figure 3 was in the prediction child's self-concept; mothers' ethnicity was the only significant direct path. Other outcomes were essentially the same.

child depression, which are also influenced by child and parent background and situational factors.

Maternal physical functioning as measured by CD4 counts at baseline is associated with later maternal physical functioning, which in turn is related to child self-concept (with higher maternal physical functioning associated with better child concept), which subsequently influences child depression (higher self-concept is related to lower levels of depression). Maternal physical functioning has an indirect effect on child depression through child self-concept, and maternal disclosure (of her serostatus to the child) has an indirect effect on child depression. Maternal disclosure is also associated directly with self-concept: when disclosure has occurred, child self-concept is better. This is consistent with studies indicating that children aware of their mother's HIV status are doing well compared to children unaware of their mothers' status (Murphy, Marelich, & Hoffman, 2002; Murphy et al., 2001; Schrimshaw & Seigel, 2002).

Interestingly, maternal ethnicity was negatively associated with child self-concept for white mothers. This is consistent with previous literature indicating that minority youth report higher self-esteem than white youth (e.g., Gray-Little & Hafdahl, 2000). However, it should be noted that we had only a rough analysis of ethnicity given the fact that the majority of our mothers were either Latina or African-American. Future studies will need to further investigate ethnicity findings.

One unanticipated finding was that higher family cohesion was associated with lower levels of maternal physical functioning, poorer child self-concept, and higher child depression. There are several studies of families affected by HIV and cancer that have also reported this association. Among families affected by maternal HIV, Bauman, Camacho, Silver, Hudis, & Draimin (2002) reported that family cohesion was a risk factor for poorer child depression controlling for other protective factors. Similarly, among parents of children with cancer, Sloper (2000) found that family cohesion predicted distress at a later time point. The explanation postulated was Dolgin and Phipps' (1996) suggestion that the construction of family is related to the idea of centripetal and centrifugal forces, which operate around events in the normal family life cycle to draw families together or pull them apart. That is, certain challenges serve to pull a family together (centripetal orientation, such as adding a child to a family), whereas some challenges pull a family apart (centrifugal, such as an adolescent preparing to leave home). Serious illness in a family member is generally thought to produce centripetal forces (Moore, Stambrook, & Peters, 1993), which may explain increased cohesion reported in families with a member with cancer (Dolgin & Phipps, 1996). When a family is in centripetal orientation, the normative stresses combined with illness-related stresses can be additive and family cohesion may become excessive, leading to enmeshment. And although serious illness is generally seen as producing centripetal forces, if it occurs in families who are already undergoing centrifugal forces, greater family disengagement may result.

Houtzager et al. (2004), in a study of coping and family functioning of siblings of childhood cancer patients, found that adjustment problems were associated with high family adaptation and cohesion. The authors noted that their finding was congruent with the so-called circumplex model of family functioning (Olson, Sprenkle, & Russell, 1979), where family cohesion is defined as the amount of closeness and mutual involvement experienced in the family system. Moderate levels of cohesion are considered to be related to the most favorable adjustment outcomes in families faced with stress, whereas extreme levels of cohesion may be related to less adaptive functioning. Our previous research found cohesion to be in the direction expected here (e.g., conflict between mothers and their partners associated with lower family cohesion, higher maternal depression associated with lower family cohesion; Murphy, Marelich, Dello Stritto, Swendeman, & Witkin, 2002). However, findings in this study were unexpected and

are consistent with the literature reviewed in this section regarding enmeshment. Future research utilizing cohesion in similar populations may wish to consider both the positive and negative effects of cohesion when specifying hypotheses. For example, it is plausible that cohesion could have a curvilinear relationship with the stated outcomes, although in the current paper this was investigated posthoc but not found tenable. Alternatively, a clearer picture of cohesion could emerge using repeated measures or growth curve analyses.

One major limitation to the current study regards the level of child depression in our sample. Although self-concept was directly associated with depression (lower levels of self-concept predicting higher levels of depression, agreeing with past research between depression and self-esteem; Alfeld-Liro & Sigelman, 1998; Battle, 1990; Bhatti, Derezotes, Kim, & Specht, 1989; Orvaschell, Beeferman, & Kabacoff, 1997), only 11% of children had CDI total depression scores indicating clinical depression (cutoff of 12; American Psychiatric Publishing, Inc., 2008; Nelson, Politano, Finch, Wendel, & Mayhall, 1987), with another 13% borderline (scores from 9–11). Hence, a majority of our sample children were not clinically depressed. However, other studies indicate the CDI is better suited as a continuous measure of mood (Matthey & Petrovski, 2002), and the current results should be viewed with this in mind. It may be more accurate to state for the current study that greater negative mood is a function of lower levels of self-concept.

Another limitation is sample size. To address this, we took a number of steps. We limited the number of measured variables and parameters estimated since ratios between sample N and parameters can affect modeling outcomes (Jackson, 2003). We also utilized a bootstrap procedure to assess fit on “like” samples and found our final model results tenable. Finally, an alternative model using coping¹ derived similar findings to the main model. Thus, although we had less power, effects were indeed found with good model fit. Another caveat regards the reported indirect effects. Although these are suggestive of mediation, we did not have data reflecting all the PACT model variables that could be influencing factors. The reported indirect findings should not be viewed as evidence of mediation. Additionally, as with other covariance structural analyses, alternative arrangements of the variables/paths could yield plausible models. Therefore, our results may reflect statistical artifact.

A final limitation is the static as opposed to dynamic nature of the current statistical analysis. The model predicts child depression at 84 months, but does not account for previous levels of depression (this applies as well to the other non-baseline measures). A superior analysis would be to evaluate the model with measurements of the variables at each time-point, and adopt a latent curve approach with these measurements as time-varying covariates (Bollen & Curran, 2006) to account for prior levels. This approach could also account for continued change in CD4 cell counts, maternal disclosure, and family cohesion. Such an analysis was considered but deemed impossible due to sample size. Whether the model could be confirmed adjusting for prior levels and/or continued change in variables cannot be ascertained at this time, nor can causality be intimated given the correlational nature of data.

There are a number of important implications from this study. First, as Bauman et al. (2002) have noted, if instead of very high family cohesiveness being a protective factor it is a risk factor for poorer child mental health, then these families are enmeshed. Very highly cohesive families may need to be targeted with specific interventions to assist children coping with parental illness (Bauman et al., pg. 50). Among these PACT families, higher family cohesion was associated with poorer maternal physical functioning. For example, perhaps family members do too much for the mother in an effort to help, thus impinging on her exercise and thus negatively affecting physical health outcomes.

The second implication from this study focuses on improving depression levels of children affected by maternal HIV. Child concept was found to be directly associated with child depression among children affected by maternal HIV/AIDS. Self-concept has been postulated to be a main foundation for all motivated behavior, and that it gives rise to possible selves (Franken, 1994). Moreover, self-concept is constructed by an individual through interactions with the environment and through considering those interactions (e.g., Huitt, 2004), indicating it can be modified. Numerous studies on interventions to improve the self-concept of children with learning problems have shown that self-concept can be improved (Elbaum & Vaughn, 2001). Thus, one implication is that interventions focused on improving self-concept among children affected by maternal HIV who have depressive symptoms could assist those children in decreasing their depression. To our knowledge, no such studies have been conducted. A limited number of intervention studies with children of parents with somatic illness have been reported, and most have had no systematic evaluation (Diareme et al, 2007). Gunther, Crandles, Williams and Swain (1998) conducted a group psychotherapy program for children of parents in all phases of AIDS that focused on healthy coping and developing peer support, and found the intervention assisted children in terms of their feelings of isolation and depression, but the outcomes were based only on clinical report.

We have speculated that targeting self-concept may improve depression; however the directionality may be reversed, in that targeting depression may improve self-esteem. Interventions may need to target the constructs that affect self-concept, meaning that interventions would also need to target healthy disclosure and child depressive symptoms, as well as family cohesion with an emphasis on positive cohesion rather than enmeshment. Future interventions for adolescents of HIV-positive mothers should investigate strategies designed to directly improve these constructs.

Acknowledgments

This research was supported by Grant #R01 MH 57207 from the National Institute of Mental Health to the first author.

We would like to thank the mothers and young adolescents who participated in this study, as well as the research interviewers, Sonia Ruiz and Claudia Perdomo.

References

- Alfeld-Liro C, Sigelman CK. Sex differences in self-concept and symptoms of depression during the transition to college. *Journal of Youth and Adolescence* 1998;27:219–244.
- American Psychiatric Publishing, Inc. *Handbook of psychiatric measures*. 2. Washington, DC: Author; 2008.
- Armistead L, Forehand R. For whom the bell tolls: Parenting decisions and challenges faced by mothers who are HIV seropositive. *Clinical Psychology: Science and Practice* 1995;2:239–250.
- Armistead L, Klein K, Forehand R. Parental physical illness and child functioning. Special Issue: The impact of the family on child adjustment and psychopathology. *Clinical Psychology Review* 1995;15:409–422.
- Armsden GC, Lewis FM. Behavioral adjustment and self-esteem of school-age children of women with breast cancer. *Oncology Nursing Forum* 1994;21:39–45. [PubMed: 8140000]
- Battle, J. *Self-esteem: The new revolution*. Edmonton, Alberta, Canada: James Battle & Associates; 1990.
- Bauman LJ, Camacho S, Silver EJ, Hudis J, Draimin B. Behavioral problems in school-aged children of mothers with HIV/AIDS. *Clinical Child Psychology & Psychiatry* 2002;7:39–54.
- Bentler, PM. *EQS 6 Structural equations modeling program manual*. Encino, CA: Multivariate Software, Inc; 2006.
- Bhatti, B.; Derezotes, D.; Kim, S-O.; Specht, H. The association between child maltreatment and self-esteem. In: Mecca, A.; Smelser, NJ.; Vasconcellos, J., editors. *Importance of self-esteem*. Berkeley, CA: University of California Press; 1989. p. 24-71.

- Bloom BL, Naar S. Self-report measures of family functioning: Extensions of a factorial analysis. *Family Process* 1994;33:203–216. [PubMed: 7925929]
- Bollen, KA.; Curran, PJ. *Latent curve models: A structural equation approach*. Hoboken, NJ: Wiley; 2006.
- Buckley, IE. *Listen to the children: Impact on the mental health of children of a parent's catastrophic illness*. New York: Cancer Care, Inc; 1977.
- Carbonell DM, Reinherz HZ, Giaconia RM. Risk and resilience in late adolescence. *Child & Adolescent Social Work Journal* 1998;15:251–272.
- Cates JA, Graham LL, Boeglin D, Tiekler S. The effect of AIDS on the family system. *Families in Society* 1990;71:195–201.
- Chavez LR, Hubbell FA, McMullin JM, Martinez RG, Mishra SI. Structure and meaning in models of breast and cervical cancer risk factors: A comparison of perceptions among Latinas, Anglo women, and physicians. *Medical Anthropology Quarterly* 1995;9:40–74. [PubMed: 7697550]
- Clark, L. Gender and generation in poor women's household health production experiences. In: Brown, PJ., editor. *Understanding and applying medical anthropology*. Mountain View, CA: Mayfield Publishing Company; 1998. p. 158-168.
- Derogatis LR, Morrow GR, Fetting J, Penman D, Piasetski S, Schmale AM, Henrichs M, Carnicke CLM Jr. The prevalence of psychiatric disorders among cancer patients. *Journal of the American Medical Association* 1983;249:751–757. [PubMed: 6823028]
- Diareme S, Tsiantis J, Romer G, Tsalamani E, Anasontzi S, Paliokosta E, Kolaitis G. Mental health support for children of parents with somatic illness: A review of the theory and intervention concepts. *Families, Systems, & Health* 2007;25:98–118.
- Dolgin, MJ.; Phipps, S. Reciprocal influences in family adjustment to childhood cancer. In: Baider, L.; Cooper, CL.; Kaplan, A., editors. *Cancer and the family*. Chichester, UK: John Wiley; 1996. p. 79-92.
- Dorsey S, Forehand R, Armistead LP, Morse E, Morse P, Stock M. Mother knows best? Mother and child report of behavioral difficulties of children of HIV-infected mothers. *Journal of Psychopathology & Behavioral Assessment* 1999;21:191–206.
- Dura JR, Beck SJ. A comparison of family functioning when mothers have chronic pain. *Pain* 1988;35:79–89. [PubMed: 3200601]
- Elbaum B, Vaughn S. School-based interventions to enhance the self-concept of students with learning disabilities: A meta-analysis. *Elementary School Journal. Special Issue: Instructional interventions for students with learning disabilities* 2001;101:303–329.
- Franken, R. *Human motivation*. 3. Pacific Grove, CA: Brooks/Cole Publishing; 1994.
- Gray-Little B, Hafidahl AR. Factors influencing racial comparisons of self-esteem: A quantitative review. *Psychological Bulletin* 2000;126:26–54. [PubMed: 10668349]
- Gunther M, Crandles S, Williams G, Swain M. A place called HOPE: Group psychotherapy for adolescents of parents with HIV/AIDS. *Child Welfare* 1998;77:251–271. [PubMed: 9514001]
- Hamilton, LA. *Dissertation Abstracts International*. Vol. 57. 1996. Dyadic family relationships and gender in adolescent identity formation: A social relations analysis; p. 4056 (University Microfilms No. AAM96-33181)
- Houtzager BA, Oort FJ, Hoekstra-Weebers JEHM, Caron HN, Grootenhuis MA, Last BF. Coping and family functioning predict longitudinal psychological adaptation of siblings of childhood cancer patients. *Journal of Pediatric Psychology* 2004;29:591–605. [PubMed: 15491981]
- Huitt, WG. *Becoming a brilliant star: An introduction*. Presentation at the Forum for Integrated Education and Educational Reform sponsored by the Council for Global Integrative Education; Santa Cruz, CA. 2004 Oct 29. Retrieved from <http://chiron.valdosta.edu/whuitt/brilstar/chapters/BrilStarintro.pdf>
- Jackson DL. Revisiting sample size and number of parameter estimates: Some support for the N:q hypothesis. *Structural Equation Modeling* 2003;10:128–141.
- James LR, Muliak SA, Brett JM. A tale of two methods. *Organizational Research Methods* 2006;9:233–244.
- Kalichman SC, Graham J, Luke W, Austin J. Perceptions of health care among persons living with HIV/AIDS who are not receiving antiretroviral medications. *AIDS Patient Care & STD's* 2002;16:233–240.

- Kennedy DP, Cowgill BO, Bogart LM, Corona R, Ryan G, Murphy DA, Schuster MA. Parents and children discuss the disclosure of a parent's HIV-status. 2007 Manuscript submitted for publication.
- Kolaitis G, Liakopoulou M. Adolescent behavioural and emotional problems in relation to their self-concept and family functioning. *Psychiatriki* 2005;16:351–362.
- Kovacs, M. *Children's Depression Inventory Manual*. Los Angeles: Western Psychological Services, Multi-Health Systems, Inc; 1992.
- Lamping, DL.; Sewitch, M.; Clark, E.; Ryan, B.; Gilmore, N.; Grover, SA.; Williams, JL.; Meister, C.; Hamel, M.; Di Meco, P. HIV-related mental health distress in persons with HIV infection, caregivers, and family members/significant others: Results of a cross-Canada survey. *International Conference on AIDS*; Florence, Italy. 1991 Jun.
- Little TD, Lindberger U, Nesselroade JR. On selecting indicators of multivariate measurement and modeling with latent variables: When "Good" indicators are bad and "Bad" indicators are good. *Psychological Methods* 1999;4:192–211.
- MacCallum RC, Austin JT. Applications of structural equation modeling in psychological research. *Annual Review of Psychology* 2000;51:201–226.
- Matthey S, Petrovski P. The Children's Depression Inventory: Error in cutoff scores for screening purposes. *Psychological Assessment* 2002;14:146–149. [PubMed: 12056076]
- McGrath EP, Repetti RL. A longitudinal study of children's depressive symptoms, self-perceptions, and cognitive distortions about the self. *Journal of Abnormal Psychology* 2002;111:77–87. [PubMed: 11866181]
- Mellins CA, Brackis-Cott E, Dolezal C, Meyer-Bahlburg HFL. Behavioral risk in early adolescents with HIV + mothers. *Journal of Adolescent Health* 2005;36:342–351. [PubMed: 15780790]
- Mikail SF, von Baeyer CL. Pain, somatic focus, and emotional adjustment in children of chronic headache sufferers and controls. *Social Science & Medicine* 1990;31:51–59. [PubMed: 2367882]
- Moore A, Stambrook M, Peters L. Centripetal and centrifugal family life cycle factors in long-term outcome following traumatic brain injury. *Brain Injury* 1993;7:247–255. [PubMed: 8508181]
- Murphy DA. HIV-positive mothers' disclosure of their serostatus to their young children: A review. *Clinical Child Psychology and Psychiatry* 2008;13:105–122. [PubMed: 18411869]
- Murphy DA, Greenwell L, Mouttapa M, Brecht ML, Schuster MA. Physical health of mothers with HIV/AIDS and the mental health of their children. *Journal of Developmental and Behavioral Pediatrics* 2006;27:386–395. [PubMed: 17041275]
- Murphy DA, Marelich WD, Dello Stritto ME, Swendeman D, Witkin A. Mothers living with HIV/AIDS: Mental, physical, and family functioning. *AIDS Care* 2002;14:633–644. [PubMed: 12419113]
- Murphy DA, Marelich WD, Hoffman D. A longitudinal study of the impact on young children of maternal HIV serostatus disclosure. *Clinical Child Psychology and Psychiatry* 2002;7:55–70.
- Murphy, DA.; Marelich, WD.; Hoffman, DA.; Schuster, MA. Parental HIV/AIDS: An empirical model of the impact on children in the United States. In: Crane, DR.; Marshall, ES., editors. *Handbook of families and health: Interdisciplinary perspectives*. Thousand Oaks, CA: Sage; 2006. p. 178-194.
- Murphy DA, Roberts KJ, Hoffman D. Young children's reactions to mothers' disclosure of maternal HIV + sersostatus. *Journal of Child and Family Studies* 2006;15:39–56.
- Murphy DA, Steers WN, Dello Stritto ME. Maternal disclosure of mother's HIV serostatus to their young children. *Journal of Family Psychology* 2001;15:441–450. [PubMed: 11584794]
- Nelson WM, Politano PM, Finch AJ, Wendel N, Mayhall CA. Children's Depression Inventory: Normative data and utility with emotionally disturbed children. *Journal of the American Academy of Child & Adolescent Psychiatry* 1987;26:43–48. [PubMed: 3583999]
- Nostlinger C, Bartoli G, Gordillo V, Roberfroid D, Colebunders R. Children and adolescents living with HIV positive parents: Emotional and behavioural problems. *Vulnerable Children and Youth Studies* 2006;1:29–43.
- Olson DH, Sprenkle DH, Russell CS. Circumplex model of marital and family systems: I. cohesion and adaptability dimensions, family types, and clinical applications. *Family Process* 1979;18:3–28. [PubMed: 437067]
- Orvaschel H, Beeferman D, Kabacoff R. Depression, self-esteem, sex, and age in a child and adolescent clinical sample. *Journal of Clinical Child Psychology* 1997;26:285–289. [PubMed: 9292386]

- Petrak JA, Doyle AM, Smith A, Skinner C, Hedge B. Factors associated with self-disclosure of HIV serostatus to significant others. *British Journal of Health Psychology* 2001;6:69–79. [PubMed: 14596739]
- Piers, EV. Piers-Harris Children's Self-Concept Scale: Revised Manual 1984. Los Angeles: Western Psychological Services; 1993.
- Reyland SA, McMahon TJ, Higgins-Delessandro A, Luthar SS. Inner-city children living with an HIV-Seropositive mother: Parent-child relationships, perception of social support, and psychological disturbance. *Journal of Child & Family Studies* 2002;11:313–329.
- Romer G, Barkmann C, Schulte-Markwort M, Thomalla G, Riedesser P. Children of somatically ill parents: A methodological review. *Clinical Child Psychology & Psychiatry* 2002;7:17–38.
- Roosa MW, Dumka L, Tein J. Family characteristics as mediators of the influence of problem drinking and multiple risk status on child mental health. *American Journal of Community Psychology* 1996;24:607–624. [PubMed: 9145493]
- Sandler IN, Reynolds KD, Kliewer W, Ramirez R. Specificity of the relation between life events and psychological symptomatology. *Journal of Clinical Child Psychology* 1992;21:240–248.
- Sandler IN, Tein J, West SG. Coping, stress, and the psychological symptoms of children of divorce: A cross-sectional and longitudinal study. *Child Development* 1994;65:1744–1763. [PubMed: 7859552]
- Sargent, C.; Brettell, C. Introduction: Gender, medicine, and health. In: Sargent, C.; Brettell, C., editors. *Gender and health: An international perspective*. Englewood Cliffs, NJ: Prentice Hall; 1996. p. 1-21.
- Schrimshaw EW, Siegel K. HIV-infected mothers' disclosure to their uninfected children: Rates, reasons, and reactions. *Journal of Social and Personal Relationships*. Special Issue: Personal and social relationships of individuals living with HIV and/or AIDS 2002;19:19–44.
- Sloper P. Predictors of distress in parents of children with cancer: A prospective study. *Journal of Pediatric Psychology* 2000;25:79–91. [PubMed: 10820946]
- Valdiserri RO, Holtgrave DR, West GR. Promoting early HIV diagnosis and entry into care. *AIDS* 1999;13:2317–2330. [PubMed: 10597773]
- Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36): I. Conceptual framework and item selection. *Medical Care* 1992;30:473–483. [PubMed: 1593914]
- Wiener LS, Battles HB, Heilman NE. Factors associated with parents' decision to disclose their HIV diagnosis to their children. *Child Welfare*. Special HIV/AIDS and children, youth, and families: Lessons learned 1998;77:115–135.
- Woods NF, Haberman MR, Packard NJ. Demands of illness and individual, dyadic, and family adaptation in chronic illness. *Western Journal of Nursing Research* 1993;15:10–25. [PubMed: 8421914]
- Worsham, NL.; Compas, BE.; Ey, S. Children's coping with parental illness. In: Wolchik, SA.; Sandler, IN., editors. *Handbook of children's coping: Linking theory and intervention*. New York: Plenum Press; 1997. p. 195-213.
- Zayas, LH.; Romano, K. Adolescents and parental death from AIDS. In: Dane, BO.; Levine, C., editors. *AIDS and the new orphans*. Westport, CT: Auburn House; 1994. p. 59-76.

Figure 1. PACT Model

Note: Selected variables for this study are in bold

Figure 2. Hypothesized measurement model for PACT over 83 months. Observed variables are shown within rectangles, and latent constructs are shown within ellipses

Note: Estimated parameters are noted with a “*”, and directional relationships are indicated with +/-.

Note: Error variance terms for measured variables are not shown.

Figure 3. Final standardized parameter estimates for PACT measurement model after modifications

Note: Error variance terms for measured variables not shown.

** $p < .01$; * $p < .05$, + $p < .10$

Table 1

Observed means and standard deviations of proposed PACT model variables

Variables	<i>M</i>	<i>SD</i>	<i>Min, Max</i>
Ethnicity (0 = Non-white, 1 = White) ¹	0.26	0.44	0, 1
Disclosure (0 = No, 1 = Yes) ¹	0.36	0.48	0, 1
CD4 Cell Count	388.12	280.99	13, 1664
Viral Load (log with +1 constant)	2.77	1.51	0, 5.44
Family Cohesion	15.97	2.38	10, 20
Physical Functioning (MOS 36)	71.30	26.53	0, 100
Self-Concept (Piers-Harris)			
Intellectual and School Status	13.41	2.59	7, 17
Physical Appearance	10.73	2.81	2, 13
Popularity	10.11	2.03	4, 12
Happiness and Satisfaction	9.30	1.20	4, 10
Depression (CDI)			
Negative Mood	1.10	1.41	0, 7
Interpersonal Problems	0.52	1.03	0, 5
Ineffectiveness	1.32	1.65	0, 7
Anhedonia	1.94	2.16	0, 9
Negative Self-Esteem	0.46	0.94	0, 5

¹ Percentages for ethnicity (26% nonwhite) and disclosure (36% nondisclosure)

Table 2
Observed pairwise Pearson correlations and sample size of proposed model variables: r (N)

Measures	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)
<i>Baseline</i>															
(1) CD4 Baseline	--														
(2) Viral Load Baseline	-.37 (78)	--													
(3) Disclosure Baseline	-.06 (81)	-.05 (78)	--												
(4) Mothers' Race/Ethnicity	.16 (81)	-.08 (78)	.03 (81)	--											
<i>30-Month FU</i>															
(5) Family Cohesion	.13 (69)	-.04 (66)	.20 (69)	.01 (69)	--										
<i>6-Month FU</i>															
(6) Physical Functioning	.13 (81)	-.03 (78)	-.11 (81)	.19 (81)	-.20 (69)	--									
<i>Child Self-Concept</i>															
(7) Intellectual and School Status	-.02 (81)	-.07 (78)	.20 (81)	-.21 (81)	.00 (69)	.18 (81)	--								
(8) Physical Appearance and Attributes	-.22 (77)	-.24 (74)	.16 (77)	-.09 (77)	-.07 (65)	.05 (77)	.68 (77)	--							
(9) Popularity	.06 (80)	-.10 (77)	.09 (80)	-.02 (80)	-.11 (68)	.20 (80)	.60 (80)	.54 (77)	--						
(10) Happiness and Satisfaction	-.20 (79)	-.05 (76)	.14 (79)	-.11 (79)	.08 (67)	.12 (79)	.53 (79)	.56 (77)	.44 (78)	--					
<i>84-Month FU</i>															
<i>Child Depression</i>															
(11) Negative Mood	.01 (71)	-.16 (68)	-.07 (71)	.07 (71)	.09 (60)	-.21 (71)	-.17 (71)	-.30 (67)	-.29 (70)	-.25 (69)	--				
(12) Interpersonal Problems	.01 (71)	-.02 (68)	.03 (71)	-.03 (71)	.05 (60)	-.04 (71)	-.17 (71)	-.12 (67)	-.01 (70)	-.13 (69)	.24 (71)	--			
(13) Ineffectiveness	.01 (71)	.14 (68)	-.17 (71)	.08 (71)	.23 (60)	-.12 (71)	-.57 (71)	-.55 (67)	-.40 (70)	-.43 (69)	.30 (71)	.22 (71)	--		
(14) Anhedonia	.00 (71)	-.10 (68)	-.04 (71)	.19 (71)	.12 (60)	-.09 (71)	-.31 (71)	-.23 (67)	-.28 (70)	-.12 (69)	.49 (71)	.21 (71)	.42 (71)	--	
(15) Negative Self-Esteem	.12 (71)	.13 (68)	-.28 (71)	.25 (71)	-.03 (60)	.05 (71)	-.34 (71)	-.48 (67)	-.23 (70)	-.23 (69)	.44 (71)	.32 (71)	.40 (71)	.52 (71)	--