Published in final edited form as:

Infant Child Dev. 2010 ; 19(1): 27–38. doi:10.1002/icd.651.

Psychological and psychophysiological considerations regarding the maternal-fetal relationship

Janet A. DiPietro

Department of Population, Family and Reproductive Health, Johns Hopkins University, 615 N. Wolfe St, E4531, Baltimore MD 21205.

Abstract

The earliest relationship does not begin with birth. Pregnant women construct mental representations of the fetus, and feelings of affiliation or “maternal-fetal attachment” generally increase over the course of gestation. While there is a fairly substantial literature on the development and moderation of psychological features of the maternal-fetal relationship, including the role of ultrasound imaging, relatively little is known about the manner in which maternal psychological functioning influences the fetus. Dispositional levels of maternal stress and anxiety are modestly associated with aspects of fetal heart rate and motor activity. Both induced maternal arousal and relaxation generate fairly immediate alterations to fetal neurobehaviors; the most consistently observed fetal response to changes in maternal psychological state involves suppression of motor activity. These effects may be mediated, in part, by an orienting response of the fetus to changes in the intrauterine environment. Conversely, there is evidence that fetal behaviors elicit maternal physiological responses. Integration of this finding into a more dynamic model of the maternal-fetal dyad, and implications for the postnatal relationship are discussed. Research on the period before birth affords tremendous opportunity for developmental scientists to advance understanding of the origins of human attachment.

Introduction

The connection between the pregnant woman and her developing fetus is perhaps the most profound but enigmatic of all human relationships. Speculation about its nature has been largely relegated to the realm of folklore, literature, and cultural beliefs. The initial academic consideration of this topic was principally psychodynamic in orientation. Empirical interest in the maternal psychological orientation towards the pregnancy and fetus began to emerge in earnest during the 1960s. In part, this research area reflects a downward application of constructs developed to characterize maternal-infant relationships, including attachment, to the period before birth. However, this period of human development is unlike any other because the physiological relationship between the pair is paramount and inextricable. This raises the possibility that variation in one may alter the trajectory of development in the other, with implications that extend beyond parturition.

Technological advances have opened a window on the intrauterine environment, affording the opportunity to characterize the interplay between maternal emotions, maternal and fetal physiology, and fetal behavior in a manner that didn't previously exist. The emerging knowledge base demonstrates the intricate nature of this relationship, including evidence of a bidirectionality. This review will focus on both the psychological and psychophysiological

nature of the maternal-fetal relationship and offer speculation on promising avenues for continued examination of this relatively new field of study.

Psychological considerations

There is a fairly sizable body of literature examining maternal, and occasionally paternal, feelings of affiliation or affection towards the fetus. Research findings in this area, generally labeled “maternal-fetal attachment” are summarized in two reviews that detail 63 articles published prior to 2003 (Cannella, 2005) and from 2000 to 2007 (Alhusen, 2008). Several instruments have been developed to explicitly measure the construct (Condon, 1993; Cranley, 1981; Muller, 1993), although related items are often contained within other scales that focus on pregnancy stress or anxiety. Commonly measured components of maternal-fetal attachment scales include preoccupation with thoughts about the fetus, maternal interaction with the fetus (i.e., talking to or palpating the abdomen to elicit a fetal response), affective valence of feelings towards either the fetus or pregnancy, and the degree to which women report that they modulate their behavior or diet to foster fetal well-being. In general, maternal to fetal attachment increases as gestational age advances (Lerum & LoBiondo-Wood, 1989; Reading, Cox, Sledmere, & Campbell, 1984), and this is often suggested to be linked to increased perception of fetal movements. Randomization into an intervention consisting of asking women to engage in routine periods of fetal movement counting using standard obstetric methods resulted in increased reporting of maternal-fetal attachment when compared to women who were not instructed in these techniques (Mikhail et al., 1991), suggesting that focused awareness of the fetus as a separate individual may indeed be contributory. Attempts to link the strength of maternal-fetal attachment to specific maternal psychological or demographic characteristics have been generally unsuccessful (Cannella, 2005) and there is significant inconsistency in research on factors that may influence the psychological relationship between mother and fetus (Van den Bergh & Simons, 2008).

Fetal imaging and maternal-fetal attachment—The manner in which the introduction of real-time ultrasound to obstetrics in the late 1970s changed the psychological landscape of pregnancy cannot be overstated. The potential impact of ultrasound on maternal to fetal attachment was discussed shortly after ultrasound was introduced to clinical obstetrics with the description of two case reports published in the *New England Journal of Medicine* (Fletcher & Evans, 1983). A flurry of letters followed arguing the advantages and disadvantages of this phenomenon. Twenty-five years later, the medical and diagnostic use of ultrasound has been disassociated from the simple viewing capability as facilities offer imaging services as a way for parents to “connect” with their fetus. The ability to visualize, photograph, and videotape the fetus, initially with 2-dimensional ultrasound and now with 3-dimensional ultrasound with a time component (i.e., 4- dimensional), has allowed parents to extend the documentation of a child's development back from birth to the second post-conceptional month. Sonogram photos routinely regale refrigerators and are toted in wallets and purses. The consequences of the pervasiveness of fetal imaging has altered the way that the fetus is perceived both within the maternal-fetal relationship as well as in medicine and the broader society (Stormer, 2003).

A number of studies have indicated that viewing a fetus on ultrasound confers at least a short-term maternal psychological benefit including reduction in anxiety following a reassuring sonogram (Zlotogorski, Tadmor, Rabinovitz, & Diamant, 1997) and enhanced ability to mentally image the fetus (Ji et al., 2005). There is some evidence that sonograms can also facilitate maternal-fetal attachment, at least in the short term (Pretorius et al., 2006). However, there is no evidence for a shift in levels of maternal-fetal attachment when studies that collected maternal report measures of their perceptions of their fetuses as described above are compared before and after ultrasound became a routine component of obstetric care (Levine, Zagoory-Sharon, Feldman, & Weller, 2007). It is also clear that maternal feelings of attachment increase

over the course of gestation regardless of whether an ultrasound was done (Kemp & Page, 1987; Phipps & Zinn, 1986). A number of recent studies have been directed at determining whether 3D/4D ultrasound generates a greater psychological impact than 2D ultrasound. Results have been conflicting with some reporting that 3D/4D images have greater psychological impact (Ji et al., 2005) but others have not (Lapaire et al., 2007; Rustico et al., 2005; Sedgmen et al., 2006). Discrepancies may be attributed, in part, to differences in whether the emphasis on outcomes is focused on maternal feelings of affiliation towards the fetus or simply being able to construct a clearer mental representation.

Although ultrasound provides an opportunity to visualize the physical features and behaviors of the fetus, parents also develop representational models of more intrinsic characteristics of fetal personality (Zeanah, Zeanah, & Stewart, 1990). Data collected using temperament scales developed for infancy but modified for administration to expectant parents shows stability in both maternal and paternal projections of fetal temperament during the third trimester and these extend into the newborn period (Zeanah, Keener, Stewart, & Anders, 1985). An interview-based assessment, the Working Model of the Child, provides more comprehensive information about the underlying characteristics and affective tone of the parental representation of the fetus. Pregnant women who provided a balanced view of the fetus, as opposed to a disengaged or distorted perception, were significantly more likely to have infants classified as securely attached at 12 months (Benoit, Parker, & Zeanah, 1997).

The prenatal hormonal milieu and maternal-fetal attachment—Very little is known about the contributory influence of the shifting prenatal neuroendocrine environment on maternal psychological adaptation to pregnancy or early maternal behavior. A prime candidate for a potential mediating role is oxytocin, a neuropeptide with a well-established link to affiliative and parental behaviors in a number of animal models (Insel, 1993). A prospective study found no relation between maternal-fetal attachment and either first or third trimester levels of oxytocin in human pregnancies, but women who displayed a pattern of increasing oxytocin from the first trimester through the first month postpartum had reported higher levels of maternal-fetal attachment in pregnancy (Levine et al., 2007). In contrast, no significant associations between maternal-fetal attachment and other steroids, including estradiol, progesterone, testosterone and cortisol, during pregnancy were detected in another study (Fleming, Ruble, Krieger, & Wong, 1997). However, first trimester levels of oxytocin and the rise in the estradiol/progesterone ratio during pregnancy were positively related to stronger early postpartum maternal attachment behaviors and feelings, respectively, in those cohorts (Feldman, Weller, Zagoory-Sharon, & Levine, 2007; Fleming et al., 1997). It is clear that there are significant opportunities to better understand the linkage between the neuroendocrine milieu of pregnancy, the growth of affiliative feelings toward the fetus during pregnancy, and the ultimate impact of both on early parenting behaviors and attachment.

Psychophysiological considerations

In the 1930's, Sontag and colleagues at the Fels Institute began perhaps the earliest systematic scientific inquiry into factors that influence the development of the human fetus, including those provided by the maternal psychological and physical environment (Sontag & Richards, 1938). Although their methods of access to the fetus were fairly primitive, many of the questions they posed then remain active areas of research today. Technological advances in our ability to view and monitor the fetus since that time allows greater reach into the uterus. The traditional obstetrical literature, and the sub-speciality of maternal-fetal medicine in particular, has detailed the complex interface that links the developing fetus to the pregnant woman with emphasis on the pathophysiology of pregnancy. However, study of whether and how normal variation in maternal psychological state may be transduced to the fetus has

traditionally been investigated within developmental, psychophysiological, or psychobiological frameworks.

Fetal responsivity to maternal psychological state—There have been naturalistic observations of fetal heart rate or motor responsivity to episodes of maternal alarm, including maternal distress following a fall (Hepper, 1990), an earthquake (Ianniruberto & Tajani, 1981), and sounding of an air raid alarm during the Gulf War (Yoles, Hod, Kaplan, & Ovadia, 1993). Note that these anecdotal reports persisted through the 1990s, illustrating the lack of systematic scientific study in this area until fairly recently. Research methodology in this area tends to rely on either observation of correspondence between measures of maternal and fetal functioning under undisturbed, baseline conditions, or observation of changes in fetal neurobehavioral functioning following experimental manipulation of maternal psychological state. Fetal neurobehaviors refer to measurable activities of the fetus that are known to be linked to neural development. This essentially reflects a downward extension of the neurobehavioral constructs first developed and applied to the newborn. Fetal neurobehavioral parameters include heart rate and heart rate variability, fetal motor activity, and fetal behavioral state, which is a manifestation of the developing interrelationship between fetal heart rate and motor activity. More detailed information on fetal neurobehavioral development and its putative nervous systems origins can be found elsewhere (DiPietro, 2005). Unlike psychological assessments of maternal-fetal interaction which can be collected at any time during pregnancy, assessments of maternal-fetal physiological interaction has technological constraints related to the ability to adequately monitor the fetal heart and behavior. In addition, fetal neurobehavioral development is relatively immature through at least the first half of pregnancy. Thus most of the studies described in the remainder of this section occurred after 24 weeks gestation.

Maternal dispositional indicators of anxiety or perceived stress, measured through self-report, have been linked to higher levels of fetal motor activity (DiPietro et al., 2002; Field et al., 2003; Van den Bergh et al., 1989), greater variability in fetal heart rate (DiPietro et al., 2002), and state-specific alterations to both (Groome et al., 1995; Sjöstrom, Valentin, Thelin, & Marsal, 2002) commencing in the second half of gestation. However, self-reported psychological trait or state parameters have not been closely linked to variation in maternal physiological parameters, such as cortisol (e.g., Petraglia et al., 2001; Urizar et al., 2004). As a result, studies that rely solely on maternal report provide limited opportunities to understand the potential mechanisms that may mediate these observations.

In one of the first studies to deliberately manipulate maternal psychological state to observe a fetal response, fetal tachycardia was elicited by a series of experimental conditions ranging from a loud, unexpected sound to deceiving women that their fetus was inadequately oxygenated (Copher & Huber, 1967). A less threatening stimulus - a tape recording of a crying infant - generated a decelerative fetal heart rate response in anxious, but not in non-anxious or depressed women (Benson et al., 1987). To date, there are three reports of fetal responses to maternal exposure to the Stroop Color Word task, an experimental cognitive challenge that reliably evokes a robust sympathetic response. These include increased variability in heart rate concomitant with suppression of motor activity (DiPietro, Costigan, & Gurewitsch, 2003) and increased fetal heart rate in fetuses of women with high trait anxiety or depression (Monk et al., 2000; Monk et al., 2004) at or near 36 weeks gestation. Maternal viewing of a labor and delivery documentary at 32 weeks gestation by 136 pregnant participants was associated with a decrease in fetal heart rate variability but also fetal motor suppression (DiPietro, Ghera, & Costigan, 2008b). However, induced maternal relaxation, also evaluated at 32 weeks gestation ($n = 100$) and effected through the use of guided imagery audiotape, yields similar fetal consequences, including increased fetal heart rate variability and suppressed motor activity, although decreased fetal heart rate was also observed (DiPietro et al., 2008a). An earlier, small

study ($n = 10$) found no effect on fetal motor activity in response to maternal viewing of a labor and delivery film (Van den Bergh et al., 1989).

Potential mechanisms of maternal to fetal transduction—Understanding of how changes in maternal psychological state generate a fetal response remains limited. Because there are no neural connections between pregnant woman and fetus, all observed associations must be mediated by other pathways. Baseline maternal and fetal heart rate are strongly correlated, whether averaged over 24-hour (Patrick, Campbell, Carmichael, & Probert, 1982) or 50-minute intervals (DiPietro, Irizarry, Costigan, & Gurewitsch, 2004). There is evidence of a circadian rhythm in fetal heart rate such that intervals of slowest heart rate occur in the middle of the night (Patrick et al., 1982), but this does not appear to be affected by maternal sleep states per se (Hoppenbrouwers et al., 1981). However, there is no evidence that maternal heart rate directly influences fetal heart rate when correspondence is examined using time-dependent analytic methods (DiPietro et al., 2004; van Leeuwen et al., 2003). The most commonly proposed explanatory mechanisms for observations of maternal influences on fetal neurobehaviors involve single or joint activation of either the sympathetic arm of the autonomic nervous system and/or the hypothalamic-pituitary-adrenal axis (HPA). There is some evidence to support each of these possibilities. For example, maternal skin conductance reactivity to both the Stroop and a labor and delivery film has been mildly associated with the degree of fetal motor responsivity to each (DiPietro et al., 2003; DiPietro et al., 2008b). The degree of fetal heart rate reactivity has also been associated with the maternal blood pressure response to the Stroop (Monk et al., 2004). With respect to the HPA axis, higher levels of maternal salivary cortisol have been significantly correlated with higher levels of fetal motor activity in two studies (DiPietro, Kivlighan, Costigan, & Laudenslager, submitted; Field et al., 2005).

Nonetheless, the literature exploring the manner in which maternal psychophysiological alterations affect the fetus is small and leaves many unanswered questions. In general, limited correspondence has been shown between the magnitude of a maternal autonomic response and the magnitude of the fetal neurobehavioral response. When significant correlations have been detected, these tend to be small in magnitude and can explain only a small proportion of the shared variance. In addition, the pervasive fetal response of suppression of motor activity to a number of different maternal manipulations, including both induced arousal and relaxation, is curious. Since the fetal response tends to be observed soon, sometimes almost immediately, after onset of a maternal response to the manipulation, this excludes full expression of an HPA axis response. Fetal responses have also been observed within seconds of the maternal stimulation in studies of sensory capacities, including maternal postural changes (Lecaneut & Jacquet, 2002) and auditory stimuli (Groome et al., 2000) at term. Together, this pattern of findings has resulted in speculation concerning indirect mediation of physiological processes. Specifically, we suspect that at least the initiation, if not the maintenance, of the fetal response to some maternal manipulations may be mediated by fetal perceptual detection of changes in the intrauterine milieu. Sounds generated by maternal heart rate, vasculature, and the digestive tract are prominent in the uterine auditory environment (Querleu, Renard, Boutteville, & Crepin, 1989). The concept of a fetal orienting response to intrauterine changes has also been proposed in non-human primate models (Novak, 2004). It is possible that sudden maternal psychophysiological shifts elicit phasic responses that include a rapid sensory-mediated component as the fetus detects a change, followed by a secondary response with more complex neuroendocrine or autonomic input that may account for more chronic activation. A certain degree of fetal neurologic maturation would be requisite for such a response; based on observations of the trajectory of neurobehavioral development across a number of domains, such a response pattern would be most likely to attain a mature level of function between the 28th and 32nd week of gestation.

Maternal responsivity to fetal behavior—The psychophysiological studies reviewed so far have approached the maternal-fetal relationship as a unidirectional phenomenon: that is, the fetus is a passive recipient of maternal direction. This stage of inquiry into the maternal-fetal relationship parallels the era in developmental science when the infant and child were viewed as the malleable targets of parental care. In 1968, a seminal paper by Bell (Bell, 1968) introduced the now well-established construct that the relationship between parent and infant is bidirectional and that constitutional variation in infants elicits differential behavior in parents. Animal models in which offspring help shape the development of the maternal nervous system have been offered in rodents, such that the reproductive experience fosters growth in maternal neural structures that foster offspring survival, with applicability to functions that extend beyond maternal care (Kinsley, 2008).

Over 70 years ago, Fels Institute investigators speculated that the increased intensity of fetal motor activity with advancing gestation stimulated the coincidentally observed increase in maternal basal metabolic rate (Richards, Newbery & Fallgatter, 1938). More recently, our work has identified a fetal effect on the maternal nervous system by applying time series analytical methods to examine the temporal relations between maternal physiological parameters and fetal neurobehavioral measures in 1-sec intervals for periods of 50 minutes at monthly intervals from 24 weeks gestation through term. The analysis revealed a significant relation between maternal skin conductance, an indicator of sympathetic activation, and fetal motor activity. While this finding was not unexpected, its directionality was certainly unpredicted. Specifically, fetal movements, which most frequently occur in the absence of maternal perception, stimulate a small rise in maternal skin conductance following a temporal lag of 2-3 s (DiPietro et al., 2004). Neither the magnitude nor the temporal characteristics of the fetal motor activity-maternal skin conductance association changed during gestation. Moreover, there was significant stability within individual pairs in the magnitude of this relationship, indicating that this aspect of fetal activation of the maternal nervous system may be a stable characteristic of individual maternal-fetal pairs, with potential implications for postnatal interaction. These findings were replicated on a second sample of maternal-fetal pairs drawn from a South American population of different sociodemographic and ethnic background than the original U.S.-based sample, suggesting that fetal stimulation of the maternal sympathetic nervous system may be an intrinsic feature of human gestation (DiPietro et al., 2006). Fetuses move, on average, approximately once per minute during the second half of pregnancy (Manning, Platt, & Sipos, 1979; Nasello-Paterson, Natale, & Connors, 1988). Thus, despite the ubiquity of the stimulus, the lack of change in the nature of this association during gestation suggests that women do not physiologically habituate or become desensitized to fetal movement.

Towards an integrative model

The fetus is increasingly regarded as an active agent in both its own ontogeny and pregnancy outcomes (Challis et al., 2001; Visser, 2004). Fetal behaviors can maximize intrauterine environmental adaptation, generate epigenetic shaping of the sensory, skeletal, and nervous systems, and provide preparation for successful transition to the postnatal environment (Challis et al., 2001; Prechtl, 1984; Smotherman & Robinson, 1996; Visser, 2004). In turn, there is mounting evidence that features of maternal psychophysiological functioning contribute to the developmental context of the fetus. Although the reports of maternal to fetal signal transduction in response to an experimental manipulation described above are fairly recent and novel findings, all subscribe to a relatively simple unidirectional orientation in which influence flows from pregnant woman to fetus. Thus, research into the dynamic processes that direct dyadic interaction within the maternal-fetal pair remains in its “infancy”, and understanding of these processes lags well behind those of postnatal interaction. Part of this is related to the challenges inherent in conducting fetal research which restricts the pool of investigators who conduct this

work. However, developmental science has historically seen a downward extension of theories, constructs and research questions from later periods of development to earlier ones. Journals devoted to infant research did not emerge until the late 1970s. Over the past decade there has been increasing empirical documentation of dyadic coherence or synchrony in both physiologic and behavioral processes in early infancy (Feldman, 2006; Harrist & Waugh, 2002; Thompson & Trevathan, 2008). Comparative models have begun to serve as a basis for forming a more integrative theory of early attachment (Lickliter, 2008).

Although there is much basic empirical work remaining before understanding of the complex interaction between pregnant woman and fetus is complete, it is not too early to consider the broader context of human development in which this emerging information is integrated. For example, the periodic sympathetic surges women experience in response to fetal motor activity, as described above, may contribute to the observed decrements in cognitive performance observed during pregnancy (Buckwalter et al., 1999; de Groot, Adam, & Hornstra, 2003) by interfering with parasympathetic processes that are required for maintenance of attention. In turn, the coincident dampening of maternal responsivity to external physical and mental challenges during pregnancy (Barron et al., 1986; Kammerer, Adams, von Castelberg, & Glover, 2002; Matthews & Rodin, 1992) may benefit increased awareness and responsivity to internally-generated signals. Thus, periodic sympathetic surges instigated by fetal activity may serve as a signaling function to entrain maternal arousal patterns to the behavior of the fetus in preparation for the consuming demands of newborn care. Among individuals, those maternal-fetal pairs with relatively low levels of this aspect of synchrony, either because the fetus is less active or the mother less sensitive, may experience less early postnatal interactional success than pairs with moderate or higher levels of coherence. This variation, in turn, may have implications for subsequent maternal and infant behavior and affiliation. Thus, relatively small levels of variation in features of the maternal-fetal relationship may be magnified as development proceeds.

Conclusions

It is clear that the earliest relationship does not begin with birth. Psychological maternal-fetal attachment commences fairly early in pregnancy and increases over gestation, culminating in the birth of an infant. There are indications that changes in pregnancy hormones may correspond to maternal psychological adaptation to pregnancy and newborn care, but there is much to be learned in this regard. Maternal psychological functioning can be transduced to the fetus, although the mechanisms that mediate this are unclear. However, it is not accurate to regard the effects of induced maternal psychological stress as damaging to fetal neurobehaviors, since both maternal relaxation and arousal produce similar consequences. Moreover, the consequences of either suppressing or increasing fetal motor activity within the larger context of a dyadic model are unknown. Extending these experimental findings to features of daily life, the end result might even be that pregnant women who express greater psychophysiological reactivity to events or circumstances in their daily environment provide greater experiential conditioning of the fetal nervous system with positive consequences for development. In addition, while the uterus is the developmental niche of the fetus, it has become clear that the fetus is also an active inhabitant of that niche. There is tremendous opportunity for developmental scientists to characterize the manner in which the fetus affects the pregnant woman, and the nature and contribution of maternal-fetal physiological synchrony to both individual and dyadic development of mothers and infants after birth.

Acknowledgments

Preparation of this manuscript was supported by R01 HD27592, National Institute of Child Health and Human Development.

References

- Alhusen JL. A literature update on maternal-fetal attachment. *Journal of Obstetric, Gynecological, and Neonatal Nursing* 2008;37:315–328.
- Barron W, Mujais S, Zinaman M, Bravo E, Lindheimer M. Plasma catecholamine responses to physiologic stimuli in normal human pregnancy. *American Journal of Obstetrics and Gynecology* 1986;154:80–84. [PubMed: 3511708]
- Bell RQ. A reinterpretation of the direction of effects in studies of socialization. *Psychological Review* 1968;75:81–95. [PubMed: 4870552]
- Benoit D, Parker K, Zeanah C. Mothers' representation of their infants assessed prenatally: stability and association with infants' attachment classifications. *Journal of Child Psychology and Psychiatry* 1997;38:307–313. [PubMed: 9232477]
- Benson P, Little BC, Talbert DG, Dewhurst J, Priest RG. Foetal heart rate and maternal emotional state. *British Journal of Medical Psychology* 1987;60:151–154. [PubMed: 3620391]
- Buckwalter J, Stanczyk F, McCleary C, Bluestein B, Buckwalter D, Randin K, et al. Pregnancy, the postpartum, and steroid hormones: effects on cognition and mood. *Psychoneuroendocrinology* 1999;24:69–84. [PubMed: 10098220]
- Cannella BL. Maternal-fetal attachment: an integrative review. *Journal of Advanced Nursing* 2005;50:60–68. [PubMed: 15788066]
- Challis J, Lye S, Gibb W, Whittle W, Patel F, Alfaidy N. Understanding preterm labor. *Annals of the New York Academy of Sciences* 2001;943:225–234. [PubMed: 11594542]
- Condon J. The assessment of antenatal emotional attachment: development of a questionnaire instrument. *British Journal of Medical Psychology* 1993;66:167–183. [PubMed: 8353110]
- Copher DE, Huber C. Heart rate response of the human fetus to induced maternal hypoxia. *American Journal of Obstetrics and Gynecology* 1967;98:320–335. [PubMed: 5621450]
- Cranley M. Development of a tool for the measurement of maternal attachment during pregnancy. *Nursing Research* 1981;30:281–284. [PubMed: 6912989]
- de Groot R, Adam J, Hornstra G. Selective attention deficits during human pregnancy. *Neuroscience Letters* 2003;340:21–24. [PubMed: 12648749]
- DiPietro JA. Neurobehavioral assessment before birth. *Mental Retardation and Developmental Disabilities Research* 2005;11:4–13.
- DiPietro JA, Costigan KA, Nelson P, Gurewitsch E, Laudenslager M. Maternal and fetal responses to induced relaxation during pregnancy. *Biological Psychology* 2008a;77:11–19. [PubMed: 17919804]
- DiPietro JA, Costigan KA, Gurewitsch E. Fetal response to induced maternal stress. *Early Human Development* 2003;74:125–138. [PubMed: 14580752]
- DiPietro JA, Ghera M, Costigan KA. Prenatal origins of temperamental reactivity in infancy. *Early Human Development* 2008b;84:569–575. [PubMed: 18289801]
- DiPietro JA, Kivlighan KT, Costigan KA, Laudenslager MM. Fetal motor activity is associated with maternal cortisol. submitted.
- DiPietro JA, Caulfield LE, Irizarry RA, Chen P, Merialdi M, Zavaleta N. Prenatal development of intrafetal and maternal-fetal synchrony. *Behavioral Neuroscience* 2006;120:687–701. [PubMed: 16768620]
- DiPietro JA, Hilton SC, Hawkins M, Costigan KA, Pressman EK. Maternal stress and affect influence fetal neurobehavioral development. *Developmental Psychology* 2002;38:659–668. [PubMed: 12220045]
- DiPietro JA, Irizarry RA, Costigan KA, Gurewitsch ED. The psychophysiology of the maternal-fetal relationship. *Psychophysiology* 2004;41:510–520. [PubMed: 15189474]
- Feldman R. From biological rhythms to social rhythms: Physiological precursors of mother-infant synchrony. *Developmental Psychology* 2006;42:175–188. [PubMed: 16420127]
- Feldman R, Weller A, Zagoory-Sharon O, Levine A. Evidence for a neurological foundation of human affiliation. *Psychological Science* 2007;18:965–970. [PubMed: 17958710]
- Field T, Diego M, Hernandez-Reif M, Gil K, Vera Y. Prenatal maternal cortisol, fetal activity, and growth. *International Journal of Neuroscience* 2005;115:423–429. [PubMed: 15825247]

- Field T, Diego M, Hernandez-Reif M, Schanberg S, Kuhn C, Yando R, et al. Pregnancy anxiety and comorbid depression and anger: Effects on the fetus and neonate. *Depression and Anxiety* 2003;17:140–151. [PubMed: 12768648]
- Fleming A, Ruble D, Krieger H, Wong P. Hormonal and experiential correlates of maternal responsiveness during pregnancy and the puerperium in human mothers. *Hormones and Behavior* 1997;31:145–158. [PubMed: 9154435]
- Fletcher J, Evans M. Maternal bonding in early fetal ultrasound evaluations. *New England Journal of Medicine* 1983;309:392–393. [PubMed: 6823243]
- Groome LJ, Mooney DM, Holland SB, Smith YD, Atterbury JL, Dykman RA. Temporal pattern and spectral complexity as stimulus parameters for eliciting a cardiac orienting reflex in human fetuses. *Perception and Psychophysics* 2000;62:313–320. [PubMed: 10723210]
- Groome LJ, Swiber MJ, Bentz LS, Holland SB, Atterbury JL. Maternal anxiety during pregnancy: Effect on fetal behavior at 38 and 40 weeks of gestation. *Journal of Developmental and Behavioral Pediatrics* 1995;16:391–396. [PubMed: 8746547]
- Harrist A, Waugh R. Dyadic synchrony: Its structure and function in children's development. *Developmental Review* 2002;22:555–592.
- Hepper PG. Fetal response to maternal shock. *The Lancet* 1990:1068.
- Hoppenbrouwers T, Combs D, Ugartechea JC, Hodgman J, Sterman MB, Harper RM. Fetal heart rates during maternal wakefulness and sleep. *Obstetrics & Gynecology* 1981;57(3):301–309. [PubMed: 7465144]
- Ianniruberto A, Tajani E. Ultrasonographic study of fetal movement. *Seminars in Perinatology* 1981;5:175–181. [PubMed: 7323822]
- Insel T. Oxytocin-a neuropeptide for affiliation: evidence from behavioral, receptor autoradiographic, and comparative studies. *Psychoneuroendocrinology* 1993;17:3–35. [PubMed: 1319071]
- Ji E, Pretorius D, Newton R, Uyan K, Hull A, Hollenbach K, et al. Effects of ultrasound on maternal-fetal bonding: a comparison of two- and three-dimensional ultrasound. *Ultrasound in Obstetrics and Gynecology* 2005;25:473–477. [PubMed: 15846757]
- Kammerer M, Adams D, von Castelberg B, Glover V. Pregnant women become insensitive to cold stress. *BMC Pregnancy and Childbirth* 2002;2:8. [PubMed: 12437774]
- Kemp V, Page C. Maternal prenatal attachment in normal and high-risk pregnancies. *Journal of Obstetric, Gynecologic, and Neonatal Nursing* 1987;16:179–184.
- Kinsley CH. The neuroplastic maternal brain. *Hormones and Behavior* 2008;54:1–4. [PubMed: 18329032]
- Lapaire O, Alder J, Peukert R, Holzgreve W, Tercanli S. Two- versus three-dimensional ultrasound in the second and third trimester of pregnancy: Impact on recognition and maternal-fetal bonding. A prospective pilot study. *Archives of Gynecology and Obstetrics* 2007;276:475–479. [PubMed: 17457600]
- Lecaneut JP, Jacquet AY. Fetal responsiveness to maternal passive swinging in low heart rate variability state: effects of stimulation direction and duration. *Developmental Psychobiology* 2002;40:57–67. [PubMed: 11835151]
- Lerum C, LoBiondo-Wood G. The relationship of maternal age, quickening, and physical symptoms of pregnancy to the development of maternal-fetal attachment. *Birth* 1989;16:13–17. [PubMed: 2742632]
- Levine A, Zagoory-Sharon O, Feldman R, Weller A. Oxytocin during pregnancy and early postpartum: individual patterns and maternal-fetal attachment. *Peptides* 2007;28:1162–1169. [PubMed: 17513013]
- Lickliter R. Theories of attachment: the long and winding road to an integrative developmental science. *Integrative Psychological and Behavioral Science*. 2008 epub ahead of print.
- Manning FA, Platt LD, Sapos L. Fetal movements in human pregnancies in the third trimester. *Obstetrics and Gynecology* 1979;54:699–702. [PubMed: 117410]
- Matthews KA, Rodin J. Pregnancy alters blood pressure responses to psychological and physical challenge. *Psychophysiology* 1992;29:232–240. [PubMed: 1635966]

- Mikhail M, Freda M, Merkatz R, Polizzotto E, Merkatz I. The effect of fetal movement counting on maternal attachment to fetus. *American Journal of Obstetrics and Gynecology* 1991;165:988–991. [PubMed: 1951568]
- Monk C, Fifer W, Myers M, Sloan R, Trien L, Hurtado A. Maternal stress responses and anxiety during pregnancy: Effects on fetal heart rate. *Developmental Psychobiology* 2000;36:67–77. [PubMed: 10607362]
- Monk C, Sloan RP, Myers MM, Ellman L, Werner E, Jeon J, et al. Fetal heart rate reactivity differs by women's psychiatric status: an early marker for developmental risk? *Journal of the American Academy of Child and Adolescent Psychiatry* 2004;43:283–290. [PubMed: 15076261]
- Muller M. Development of the prenatal attachment inventory. *Western Journal of Nursing Research* 1993;15:199–215. [PubMed: 8470375]
- Nasello-Paterson C, Natale R, Connors G. Ultrasonic evaluation of fetal body movements over twenty-four hours in the human fetus at twenty-four to twenty-eight weeks' gestation. *American Journal of Obstetrics and Gynecology* 1988;158:312–316. [PubMed: 3277429]
- Novak MFS. Fetal-maternal interactions: prenatal psychobiological precursors to adaptive infant development. *Current Topics in Developmental Biology* 2004;59:37–60. [PubMed: 14975246]
- Patrick J, Campbell K, Carmichael L, Probert C. Influence of maternal heart rate and gross fetal body movements on the daily pattern of fetal heart rate near term. *American Journal of Obstetrics and Gynecology* 1982;144:533–538. [PubMed: 7137240]
- Petraglia F, Hatch M, Lapinski R, Stomati M, Reis F, Cobellis L, et al. Lack of effect of psychosocial stress on maternal corticotropin-releasing factor and catecholamine levels at 28 weeks gestation. *Journal of the Society for Gynecologic Investigation* 2001;8:83–88. [PubMed: 11336878]
- Phipps S, Zinn A. Psychological response to amniocentesis I: Mood state and adaptation to pregnancy. *American Journal of Medical Genetics* 1986;25:131–1148. [PubMed: 3799712]
- Prechtl, HFR. Continuity and change in early neural development. In: Prechtl, H., editor. *Continuity in neural functions from prenatal to postnatal life*. J.B. Lippincott Co; Philadelphia, PA: 1984. p. 1-15. Vol. Clinics in Developmental Medicine No. 94
- Pretorius D, Gattu S, Ji E, Hollenbach K, Newton R, Hull A, et al. Preexamination and postexamination assessment of parental-fetal bonding in patients undergoing 3-/4-dimensional obstetric ultrasonography. *Journal of Ultrasound in Medicine* 2006;25:1411–1421. [PubMed: 17060427]
- Querleu D, Renard X, Boutteville C, Crepin G. Hearing by the human fetus? *Seminars in Perinatology* 1989;13:409–420. [PubMed: 2683112]
- Reading AE, Cox DN, Sledmere CM, Campbell S. Psychological changes over the course of pregnancy: a study of attitudes toward the fetus/neonate. *Health Psychology* 1984;3(3):211–221. [PubMed: 6399247]
- Richards TW, Newbery H, Fallgatter R. Studies in fetal behavior: II. Activity of the human fetus in utero and its relation to other prenatal conditions, particularly the mother's basal metabolic rate. *Child Development* 1938;9:69–78.
- Rustico M, Mastromatteo C, Grigio M, Maggioni C, Gregori D, Nicolini U. Two-dimensional vs. Two-plus four-dimensional ultrasound in pregnancy and the effect on maternal emotional status: a randomized study. *Ultrasound in Obstetrics and Gynecology* 2005;25:468–472. [PubMed: 15849804]
- Sedgmen B, McMahon C, Cairns D, Benzie R, Woodfield R. The impact of two-dimensional versus three-dimensional ultrasound exposure on maternal-fetal attachment and maternal health behavior in pregnancy. *Ultrasound in Obstetrics and Gynecology* 2006;27:245–251. [PubMed: 16482614]
- Sjostrom K, Valentin L, Thelin T, Marsal K. Maternal anxiety in late pregnancy: Effect on fetal movements and fetal heart rate. *Early Human Development* 2002;67:87–100. [PubMed: 11893440]
- Smotherman WP, Robinson SR. The development of behavior before birth. *Developmental Psychology* 1996;32:425–434.
- Sontag LW, Richards TW. Studies in fetal behavior: I. Fetal heart rate as a behavioral indicator. *Monographs of the Society for Research in Child Development* 1938;3(4):1–67. Serial No. 17
- Stormer N. Seeing the fetus: the role of technology and image in the maternal-fetal relationship. *Journal of the American Medical Association* 2003;289:1700. [PubMed: 12672742]

- Thompson LA, Trevathan WR. Cortisol reactivity, maternal sensitivity, and learning in 3-month-old infants. *Infant Behavior & Development* 2008;31:92–106. [PubMed: 17716739]
- Urizar GG, Milazzon M, Le HN, Delucchi K, Sotelo R, Munoz RF. Impact of stress reduction instructions on stress and cortisol levels during pregnancy. *Biological Psychology* 2004;67:275–282. [PubMed: 15294386]
- Van den Bergh BRH, Mulder EJH, Visser GHA, Poelmann-Weesjes G, Bekedam DJ, Prechtl HFR. The effect of (induced) maternal emotions on fetal behaviour: a controlled study. *Early Human Development* 1989;19:9–19. [PubMed: 2721422]
- Van den Bergh B, Simons A. A review of scales to measure the mother-foetus relationship. *Journal of Reproductive and Infant Psychology* 2008;26:1–13.
- van Leeuwen P, Geue D, Lange S, Cysarz D, Bettermann H, Gronemeyer D. Is there evidence of fetal-maternal heart rate synchronization? *BMC Physiology* 2003;3
- Visser GHA. Fetal behaviour: A commentary. *Neurobiology of Aging* 2004;24:S47–S49.
- Yoles I, Hod M, Kaplan B, Ovadia J. Fetal ‘fright-bradycardia’ brought on by air-raid alarm in israel. *International Journal of Gynaecology and Obstetrics* 1993;40:157–160. [PubMed: 8094687]
- Zeanah C, Keener M, Stewart L, Anders A. Prenatal perception of infant personality: a preliminary investigation. *Journal of the American Academy of Child Psychiatry* 1985;24:204–210. [PubMed: 3989164]
- Zeanah C, Zeanah P, Stewart L. Parents constructions of their infants' personalities before and after birth: A descriptive study. *Child Psychiatry and Human Development* 1990;20:191–206. [PubMed: 2347251]
- Zlotogorski Z, Tadmor O, Rabinovitz R, Diamant Y. Parental attitudes toward obstetric ultrasound examination. *Journal of Obstetric and Gynaecological Research* 1997;23:25–28.