Published in final edited form as:

Adv Synth Catal. 2005 October 1; 347(11-13): 1614–1620. doi:10.1002/adsc.200505172.

Synthesis of *N*-Aryl-2-allyl Pyrrolidines via Palladium-catalyzed Carboamination Reactions of γ -(*N*-Arylamino)alkenes with Vinyl Bromides

Joshua E. Ney, Michael B. Hay, Qifei Yang, and John P. Wolfe^{*,[a]}

^[a]University of Michigan, Department of Chemistry 930 N. University Ave, Ann Arbor, Michigan, 48109-1055, USA

Abstract

A palladium-catalyzed carboamination reaction of γ -*N*-arylamino alkenes with vinyl bromides that affords *N*-aryl-2-allyl pyrrolidines is described. These reactions proceed with high diastereoselectivity for the formation of *trans*-2,3- and *cis*-2,5-disubstituted pyrrolidines. Conditions for a tandem *N*-arylation/carboamination sequence that leads to the formation of an *N*-aryl-2-allyl pyrrolidine or indoline via the coupling of a primary γ -amino alkene, an aryl bromide, and a vinyl bromide are also reported. The mechanism of the carboamination reactions and the origin of unexpected products that formally derive from rearrangement of the vinyl bromide are discussed.

Keywords

Heterocycles; Homogeneous Catalysis; Pyrrolidines; Palladium; Tandem Reactions; Vinyl Halides

Introduction

The pyrrolidine ring is a common motif found in a number of biologically active small molecules.[1] As a result, a great deal of effort has been devoted to the development of synthetic methods for the construction of this core.[2] Some of the most interesting and useful of these methods have utilized palladium catalysis to effect ring closure of a β - or γ -amino alkene or alkyne with concomitant carbon—carbon bond formation.[3]–[5] For example, Larock and Weinreb have described the synthesis of 2-vinyl pyrrolidines via the coupling and cyclization of β -*N*-tosylamino alkenes with vinyl halides.[3] Tamaru has also described a palladium-catalyzed synthesis of pyrrolidines, which employs a Wacker-type carbonylative cyclization to afford products bearing ester substituents.[4]

We have recently reported a new method for the stereoselective synthesis of *N*-aryl-2-benzyl pyrrolidines via the carboamination of γ -*N*-arylamino alkenes with aryl bromides in the presence of a palladium(0) catalyst and a base (eq 1).[6]–[9] These reactions proceed in good yields and are effective with a variety of aryl bromide coupling partners. High diastereoselectivity for the formation of *trans*-2,3- and *cis*-2,5-disubstituted pyrrolidines is observed for most substrate combinations (dr >20:1).

* Fax: (734)-763-2307 jpwolfe@umich.edu .

Supporting Information for this article is available on the WWW under <http://asc.wiley-vch.de>

(1)

(2)

A significant extension of this methodology would involve the use of vinyl bromides as coupling partners to afford *N*-aryl-2-allyl pyrrolidines. However, initial attempts to couple **1** with β -bromostyrene using conditions identical to those employed in the analogous reactions of aryl bromides resulted in low yields of the desired pyrrolidine **3b** due to competing *N*-vinylation of the substrate (eq 2). We reasoned that this problem could be circumvented by employing a palladium/phosphine catalyst system that would disfavor carbon—nitrogen bond-forming reductive elimination,[10] as the side product presumably results from relatively facile reductive elimination of an intermediate vinylpalladium complex.[11] Herein we report the use of a $\text{Pd}_2(\text{dba})_3$ /tri-2-furylphosphine catalyst system to effect the diastereoselective synthesis of *N*-aryl-2-allyl pyrrolidines via the palladium-catalyzed carboamination of γ -*N*-arylamino alkenes with vinyl bromides.

Results and Discussion

Catalyst optimization

In our preliminary experiments we elected to optimize the palladium-catalyzed reaction of amine **1** with vinyl bromide **2a**, which was easily prepared in isomerically pure form (eq 3, Table 1). Unfortunately, the $\text{Pd}_2(\text{dba})_3$ /dppb catalyst system, which provided high yields for the synthesis of 2-benzylpyrrolidines from γ -*N*-arylamino olefins and aryl bromides, afforded an inseparable mixture of the desired product **3a**, regioisomer **4a**, and unexpected isomer **5a** (Table 1, entry 1). Curiously, *N*-vinylation was less problematic with this substrate combination than in the reaction of **1** with β -bromostyrene described above. After some optimization we found that **3a** was obtained in 81 % isolated yield when the reaction was conducted with tri-2-furylphosphine as the ligand (Table 1, entry 4). These conditions provided good selectivity for the formation of **3a** (versus **4a** and **5a**), and did not afford detectable amounts of *N*-vinylated side products.

Scope and diastereoselectivity

With optimized conditions in hand, we proceeded to examine the reaction of **1** with a variety of vinyl bromides. As shown in Table 2, *N*-aryl-2-allylpyrrolidines **3a–3h** were obtained in good to excellent yields with 13:1 to >20:1 selectivity for **3** versus regioisomer **4**. Isomers **5**, which derive from formal rearrangement of the vinyl bromide, were observed in only two reactions, both of which employed *E*-substituted vinyl bromides (Table 2, entries 1 and 4). Notably, an analogous rearranged product was not observed in the reaction of the corresponding

Z-substituted vinyl bromide **2e** (Table 2, entry 5). Side products resulting from *N*-vinylation or Heck vinylation of the alkene were generally not observed in reactions catalyzed by Pd₂(dba)₃/tri-2-furylphosphine, even with the previously problematic β-bromostyrene (**2b**). All products were obtained as a single olefin stereoisomer,[12] and reactions of Z-substituted vinyl bromides afforded the desired pyrrolidine products with retention of the vinyl bromide geometry (Table 2, entries 3 and 5). However, although the reaction of **1** with (2-bromovinyl)trimethylsilane (**2h**) produced a single pyrrolidine product (Table 2, entry 8), the analogous reaction of **1** with the isomeric (1-bromovinyl)trimethylsilane (**2i**) afforded an inseparable 7:4 mixture of pyrrolidines **3i** and **3h** (Table 2, entry 9).

To probe the diastereoselectivity of these reactions, several γ-*N*-arylamino alkenes with substituents at the 1- or 3-position were prepared and subjected to our optimized reaction conditions. As shown in Table 3, reactions of amines **6**–**9** with various vinyl bromides afforded 2,3- and 2,5-disubstituted pyrrolidines in good yields with good to excellent levels of diastereoselectivity (dr 7:1 to >20:1). These reactions were effective with substrates bearing both electron-rich (Table 3, entries 1–2 and 4–8) and electron-poor (Table 3, entry 3) *N*-aryl substituents. The observed preference for the formation of *trans*-2,3- and *cis*-2,5-disubstituted pyrrolidines is analogous to that observed in the previously described reactions of related substrates with aryl bromides, and is complementary to the selectivity for the formation of *cis*-2,3-disubstituted *N*-tosyl pyrrolidines described by Larock and Weinreb.[3]

The size of the vinyl bromide had a notable effect on the stereoselectivity of the carboamination reactions. In reactions of 1-substituted amines **6**–**8** the highest selectivities (dr >20:1) were observed in cases where the vinyl bromide and/or the substituent on the amine were small (Table 3, entries 2–5). In contrast, the reaction of amine **6**, which bears a 1-(*p*-biphenyl) substituent, with hindered vinyl bromide **2g** provided **10g** in good yield but with somewhat lower (dr 7:1) diastereoselectivity (Table 3, entry 1). The reactions of 3-substituted amine **9** provided the highest selectivity (dr >20:1) when the relatively small β-bromostyrene (**2b**) was employed (Table 3, entry 8). Use of α-bromostyrene (**2f**) or tetrasubstituted vinyl bromide **2g** afforded products with lower (dr 10:1) diastereoselectivity (Table 3, entries 6–7).

Tandem reactions

We have recently reported the development of palladium-catalyzed tandem *N*-arylation/carboamination reactions for the synthesis of *N*-aryl-2-benzyl pyrrolidines[13] and indolines [14] from appropriate primary amine precursors (eq 4, R = aryl). These reactions enable the modular assembly of heterocyclic compounds from very simple precursors and could be amenable to library synthesis. In these reactions, the primary amine substrate undergoes *N*-arylation in the presence of a monophosphine-supported palladium(0) catalyst.[15] After the first transformation is complete, an *in situ* ligand exchange process is effected via the addition of a bisphosphine to the reaction mixture, and then a second aryl bromide is added to provide the desired heterocycle in a one-pot process. An extension of this methodology to allow for the use of vinyl bromides as the second coupling partner would allow for the synthesis of an even broader array of compounds.

As shown in Table 4, the conditions reported previously[13],[14] for the synthesis of *N*-aryl-2-benzyl pyrrolidines and indolines via this tandem reaction were also effective for the synthesis of *N*-aryl-2-allyl pyrrolidines and indolines (eq 4, R = vinyl). Use of 2-(di-*tert*-butylphosphino) biphenyl as the ligand for the first step of the sequence was optimal with both the aliphatic and the aromatic amine substrates. In the second step, however, the ligand dppe provided the best results for the pyrrolidine forming reaction (Table 4, entries 1–2), whereas DPEphos was optimal in the reaction that afforded indoline products (Table 4, entries 3–4).

Mechanism

The carboamination reactions of γ -*N*-arylamino alkenes **1** and **6–9** with vinyl bromides likely proceed through a mechanism similar to that described previously for the analogous transformation involving aryl bromides (Scheme 1).[6] The catalytic cycle presumably commences with oxidative addition of the vinyl bromide to palladium(0), which affords vinylpalladium bromide intermediate **I**. The reaction of **I** with the amine substrate and NaOt-Bu yields vinylpalladium amido intermediate **II**, which undergoes intramolecular amidopalladation to afford **III**. Carbon—carbon bond-forming reductive elimination from **III** provides the observed pyrrolidine **3** and regenerates the palladium(0) catalyst. The *N*-vinylated side product that is observed under the non-optimized reaction conditions presumably derives from carbon—nitrogen bond-forming reductive elimination of intermediate **II**.^[11] This side reaction is effectively suppressed by the use of the relatively small tri-2-furylphosphine ligand. Interestingly, the fact that *N*-vinylation is more problematic than *N*-arylation with the Pd₂(dba)₃/dppb catalyst system suggests that vinyl sp² carbon—nitrogen bond-forming reductive elimination is a more facile transformation than aryl sp² carbon—nitrogen bond-forming reductive elimination in cases where the aryl and vinyl bromides have similar electronic properties (e.g., β -bromostyrene and 2-bromonaphthalene).^[16]

As previously suggested,[6],[7] regioisomer **4** likely originates from β -hydride elimination of **III** to afford π -olefin complex **IV** (Scheme 2). Reinsertion of the olefin into the Pd—H bond with reversal of regiochemistry would provide **V**, which can undergo a second β -hydride elimination/reinsertion process to afford **VI**. Carbon—carbon bond-forming reductive elimination would generate regioisomer **4** with concomitant regeneration of the palladium(0) catalyst.

The formation of isomer **5** as a side product in reactions involving *E*-substituted vinyl bromides **2a** and **2d** (Table 2, entries 1 and 4) and the formation of pyrrolidine **3h** as a side product in the reaction of **1** with **2i** (Table 2, entry 9) both appear to involve a formal rearrangement of

the vinyl bromide coupling partner.[17],[18] One possible origin of these products involves an unusual β -hydride elimination reaction of vinylpalladium bromide intermediate **I** to afford π -alkynylpalladium hydride complex **VII**. Reinsertion of the alkyne into the Pd—H bond with reversal of regiochemistry would afford *iso*-**I**,[19] which could re-enter the catalytic cycle shown in Scheme 1 to afford **5** (Scheme 3). This mechanism has previously been invoked to explain the origin of similar rearranged products observed in cross-coupling reactions of (1-bromovinyl)trimethylsilane (**2i**).[17a—b,d] The fact that side products **5** were observed in reactions of amine **1** with *E*-substituted vinyl bromides **2a** and **2d** but not in reactions with *Z*-substituted vinyl bromide **2e** (Table 2, entries 1, 4, and 5) also supports the mechanism shown in Scheme 3, as β -hydride elimination reactions of alkylpalladium complexes require a *syn* relationship between palladium and the involved β -hydrogen atom.[20] The relatively large amount of rearranged product **3h** observed in the reaction of amine **1** with vinyl bromide **2i** may be due to a β -silyl effect, which would stabilize the developing positive charge on the β -carbon atom in the transition state between **I** and **VII**. [21]

Conclusion

In conclusion, we have developed conditions for the synthesis of *N*-aryl-2-allyl pyrrolidines via the palladium-catalyzed carboamination of γ -*N*-arylamino alkenes with vinyl bromides. These reactions display broad scope with respect to vinyl halide structure and afford *trans*-2,3- and *cis*-2,5-disubstituted pyrrolidines with good to excellent levels of diastereoselectivity. Additionally, we have shown that tandem *N*-arylation/carboamination reactions of unsaturated primary amines are effective for the synthesis of *N*-aryl-2-allyl pyrrolidines and indolines. Efforts to apply this methodology to the total synthesis of structurally interesting and biologically active targets are currently underway.

Experimental Section

General remarks

All reactions were carried out under an argon atmosphere in flame-dried glassware. Tris (dibenzylideneacetone)dipalladium(0) and all phosphine ligands were purchased from Strem Chemical Co. and used without further purification. Toluene was purified using a GlassContour solvent purification system. Unless otherwise noted, yields refer to isolated yields of compounds estimated to be $\geq 95\%$ pure as determined by ^1H NMR and combustion analysis. The yields and product ratios reported in the experimental section and supporting information describe the result of a single experiment, whereas those reported in Tables 1–4 are averages of two or more experiments. Thus, the yields and ratios reported in the experimental section and supporting information may differ from those shown Tables 1–4.

General procedure for palladium-catalyzed carboamination reactions with vinyl bromides

A flame-dried Schlenk tube was charged with $\text{Pd}_2(\text{dba})_3$ (1 mol % complex, 2 mol % Pd), tri-2-furylphosphine (4 mol %), and sodium *tert*-butoxide (1.2 equiv). The tube was purged with argon and toluene (4 mL/mmol amine substrate), the amine substrate (1.0 equiv), and the vinyl bromide (1.1–2.0 equiv) were added via syringe. The mixture was heated to 60–110 °C with stirring until the starting material had been consumed as judged by GC or ^1H NMR analysis. The reaction mixture was cooled to room temperature, quenched with saturated aqueous ammonium chloride (2 mL), and diluted with ethyl acetate (10 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2 $\times$ 10 mL). The combined organic extracts were dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo*. The crude product was then purified by flash chromatography on silica gel.

(E)-1-Phenyl-2-[undec-2-enyl]pyrrolidine (3a)

Reaction of amine **1** (40 mg, 0.25 mmol) with vinyl bromide **2a** (60 mg, 0.275 mmol, >20:1 *E/Z*) and sodium *tert*-butoxide (29 mg, 0.3 mmol) at 60 °C following the general procedure afforded the title compound (**3a**; 66 mg, 88 %, >20:1 *E/Z*) as a colorless oil. This material was obtained as a 14:1 mixture of inseparable regioisomers and contained ~6 % **5a**, as judged by ¹H NMR analysis; data are for the major isomer. ¹H NMR (400 MHz, CDCl₃) δ 7.28–7.22 (m, 2 H), 6.67 (t, *J* = 7.4 Hz, 1 H), 6.6 (d, *J* = 7.6 Hz, 2 H), 5.57–5.48 (m, 1 H), 5.47–5.38 (m, 1 H), 3.75–3.69 (m, 1 H), 3.47–3.40 (m, 1 H), 3.22–3.14 (m, 1 H), 2.48–2.41 (m, 1 H), 2.07–1.87 (m, 7 H), 1.44–1.25 (m, 12 H), 0.91 (t, *J* = 6.6 Hz, 3 H); ¹³C NMR (100 MHz, CDCl₃) δ 147.4, 133.4, 129.4, 126.9, 115.4, 112.0, 58.7, 48.5, 36.3, 33.0, 32.1, 29.9, 29.7, 29.5, 29.4, 23.5, 22.9, 14.3 (two aliphatic signals are incidentally equivalent); IR (film) 1598, 1505 cm⁻¹. Anal calcd for C₂₁H₃₃N: C, 84.22; H, 11.11; N, 4.68. Found: C, 84.47; H, 11.29; N, 4.70.

General procedure for palladium-catalyzed tandem *N*-arylation/carboamination reactions

A flame-dried Schlenk tube was charged with Pd₂(dba)₃ (1 mol % complex, 2 mol % Pd), 2-(di-*tert*-butylphosphino)biphenyl (2 mol %) and sodium *tert*-butoxide (2.4 equiv). The tube was purged with argon and toluene (1 mL), the amine substrate (1.0 equiv) and the aryl bromide (1.0 equiv) were added via syringe. The mixture was placed in a pre-heated oil bath at 60 °C with stirring until the starting materials had been consumed as judged by GC analysis. A solution of dppe (2 mol %) in toluene (1 mL) was added, and the temperature was increased to 110 °C. After 15 min of stirring at 110 °C the vinyl bromide (1.2–2.0 equiv) was added via syringe. Heating was continued until the intermediate arylamine had been consumed as judged by GC analysis. The reaction mixture was then cooled to room temperature, quenched with saturated aqueous ammonium chloride (2 mL) and diluted with ethyl acetate (10 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2 × 10 mL). The combined organic extracts were dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo*. The crude product was then purified by flash chromatography on silica gel.

(E)-4-{2-[Undec-2-enyl]pyrrolidin-1-yl}benzonitrile (17)

Reaction of amine **14** (21 mg, 0.25 mmol) with 4-bromobenzonitrile (46 mg, 0.25 mmol), sodium *tert*-butoxide (58 mg, 0.6 mmol) and vinyl bromide **2a** (110 mg, 0.5 mmol, >20:1 *E/Z*) following the general procedure afforded the title compound (**17**; 55 mg, 68 %, >20:1 *E/Z*) as a pale yellow oil. This material was obtained as a 30:1 mixture of inseparable regioisomers and contained ~7 % 4-[2-(2-methylenedecyl)pyrrolidin-1-yl]benzonitrile as judged by ¹H NMR analysis; data are for the major isomer. ¹H NMR (400 MHz, CDCl₃) δ 7.44 (d, *J* = 9.2 Hz, 2 H), 6.52 (d, *J* = 8.4 Hz, 2 H), 5.56–5.47 (m, 1 H), 5.40–5.31 (m, 1 H), 3.80–3.73 (m, 1 H), 3.45–3.39 (m, 1 H), 3.25–3.17 (m, 1 H), 2.39–2.31 (m, 1 H), 2.12–1.91 (m, 7 H), 1.40–1.22 (m, 12 H), 0.88 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (100 MHz, CDCl₃) δ 149.6, 134.3, 133.7, 125.9, 121.2, 112.0, 96.8, 58.7, 48.3, 35.6, 32.9, 32.1, 29.7, 29.7, 29.6, 29.5, 29.4, 23.2, 22.9, 14.3; IR (film) 2213, 1607, 1520 cm⁻¹. Anal calcd for C₂₂H₃₂N₂: C, 81.43; H, 9.94; N, 8.63. Found: C, 81.51; H, 10.00; N, 8.72.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank the University of Michigan and the National Institutes of Health—National Institute of General Medical Science (GM-071650) for financial support of this work. JPW thanks the Camille and Henry Dreyfus Foundation for a new faculty award and Research Corporation for an innovation award. JEN acknowledges the

University of Michigan for a Regents Fellowship and Pfizer for a Graduate Research Fellowship. Additional unrestricted support was provided by Amgen, Eli Lilly, and 3M.

References and Notes

- [1] (a). Massiot G, Delaude C. *Alkaloids* 1986;27:269–322. (b) O'Hagan D. *Nat. Prod. Rep* 2000;17:435–446. [PubMed: 11072891] (c) Lewis JR. *Nat. Prod. Rep* 2001;18:95–128. [PubMed: 11245403]
- [2] (a). Pichon M, Figadere B. *Tetrahedron: Asymmetry* 1996;7:927–964. (b) Mitchinson A, Nadin A. *J. Chem. Soc., Perkin Trans. 1* 2000:2862–2892.
- [3] (a). Harris GD Jr, Herr RJ, Weinreb SM. *J. Org. Chem* 1992;57:2528–2530. (b) Larock RC, Yang H, Weinreb SM, Herr RJ. *J. Org. Chem* 1994;59:4172–4178.
- [4] (a). Tamaru Y, Kimura M. *Synlett* 1997:749–757. (b) Tamaru Y, Hojo M, Higashimura H, Yoshida Z. -i. *J. Am. Chem. Soc* 1988;110:3994–4002. (c) Tamaru Y, Hojo M, Yoshida Z. —i. *J. Org. Chem* 1988;53:5731–5741. (d) Harayama H, Abe A, Sakado T, Kimura M, Fugami K, Tanaka S, Tamaru Y. *J. Org. Chem* 1997;62:2113–2122. [PubMed: 11671516]
- [5]. Jacobi PA, Liu H. *Org. Lett* 1999;1:341–344. [PubMed: 10822573]
- [6] (a). Ney JE, Wolfe JP. *Angew. Chem* 2004;116:3689–3692. *Angew. Chem. Int. Ed* 2004;43:3605–3608. (b) Ney JE, Wolfe JP. *J. Am. Chem. Soc* 2005;127:8644–8651. [PubMed: 15954769]
- [7]. For a related synthesis of *N*-acyl- and *N*-boc-2-benzyl pyrrolidines, see: Bertrand MB, Wolfe JP. *Tetrahedron* 2005;61:6447–6459.
- [8] (a). For a related synthesis of tetrahydrofurans, see: Wolfe JP, Rossi MA. *J. Am. Chem. Soc* 2004;126:1620–1621. [PubMed: 14871078] (b) Hay MB, Hardin AR, Wolfe JP. *J. Org. Chem* 2005;70:3099–3107. [PubMed: 15822970]
- [9]. For a recent report of an intramolecular carboamination reaction of *N*-(arylsulfonyl)-2-allylanilines in the presence of excess Cu(OAc)₂, see: Sherman ES, Chemler SR, Tan TB, Gerlits O. *Org. Lett* 2004;6:1573–1575. [PubMed: 15128239]
- [10] (a). See reference ^{6b}. For a discussion of ligand effects on the rate of carbon—nitrogen bond-forming reductive elimination, see: (b) Hamann BC, Hartwig JF. *J. Am. Chem. Soc* 1998;120:3694–3703. For a discussion of ligand effects on the rate of carbon—carbon bond-forming reductive elimination, see: (c) Gillie A, Stille JK. *J. Am. Chem. Soc* 1980;102:4933–4941.
- [11] (a). For recent reports of palladium-catalyzed *N*-vinylation reactions, see: Dehli JR, Legros J, Bolm C. *Chem. Commun* 2005:973–986. and references cited therein. (b) Barluenga J, Fernandez MA, Aznar F, Valdes C. *Chem. Eur. J* 2004;10:494–507.
- [12]. Pyrrolidines **3b** and **3h** were obtained with >20:1 *E/Z* ratios even though they were derived from geometrically impure vinyl bromides **2b** and **2h**. These vinyl bromides were present in excess, and (*E*)-vinyl bromides have been shown to undergo oxidative addition faster than the corresponding (*Z*)-vinyl bromides. See: Jutand A, Negri S. *Organometallics* 2003;22:4229–4237.
- [13]. Yang Q, Ney JE, Wolfe JP. *Org Lett* 2005;7:2575–2578. [PubMed: 15957894]
- [14]. Lira R, Wolfe JP. *J. Am. Chem. Soc* 2004;126:13906–13907. [PubMed: 15506735]
- [15] (a). For recent reviews on palladium-catalyzed *N*-arylation reactions, see: Muci AR, Buchwald SL. 2002;219:131–209. (b) Hartwig, JF. *Modern Arene Chemistry*. Astruc, D., editor. Wiley-VCH; Weinheim, Germany: 2002. p. 107–168. (c) Schlummer B, Scholz U. *Adv. Synth. Catal* 2004;346:1599–1626.
- [16]. Previous competition experiments have shown that *N*-vinylation occurs preferentially to *N*-arylation when amines are treated with 1:1 mixtures of aryl and vinyl halides under catalytic amination conditions. However, the rate of oxidative addition of vinyl halides is faster than that of aryl halides. Thus, the competition experiments reported do not give a direct measure of the relative rates of reductive elimination. For details of these previous experiments, and references concerning the relative rates of oxidative addition of aryl and vinyl halides, see reference ^{11b}.
- [17] (a). For reports of rearranged products in palladium-catalyzed coupling reactions of (1-bromovinyl) trimethylsilane (**2i**), see: Cazes B, Colovray V, Gore J. *Tetrahedron Lett* 1988;29:627–630. (b) Carpita A, Rossi R, Scamuzzi B. *Tetrahedron Lett* 1989;30:2699–2702. (c) Ennis DS, Gilchrist TL. *Tetrahedron* 1990;46:2623–2632. (d) Sasmal PK, Maier ME. *Org. Lett* 2002;4:1271–1274. [PubMed: 11950340]

- [18]. To the best of our knowledge, the formation of a side product analogous to **5** in a palladium-catalyzed coupling reaction of an *E*-substituted 1-bromo-1-alkene has been reported only once. See: Flamme EM, Roush WR. *Org. Lett* 2005;7:1411–1414. [PubMed: 15787519]
- [19] (a). This reinsertion step fundamentally involves the hydropalladation of an alkyne. For examples of other reactions involving alkyne hydropalladation, see: Grushin VV. *Chem. Rev* 1996;96:2011–2034. [PubMed: 11848819] and references cited therein. (b) Trost BM. *Acc. Chem. Res* 1990;23:34–42. (c) Ji J, Wang Z, Lu X. *Organometallics* 1996;15:2821–2828. (d) Kadota I, Lutete LM, Shibuya A, Yamamoto Y. *Tetrahedron Lett* 2001;42:6207–6210.
- [20]. Beletskaya IP, Cheprakov AV. *Chem. Rev* 2000;100:3009–3066. [PubMed: 11749313]
- [21] (a). β -hydride elimination reactions have been shown to involve the buildup of positive charge at the β -carbon atom in the transition state. See: Mueller JA, Goller CP, Sigman MS. *J. Am. Chem. Soc* 2004;126:9724–9734. [PubMed: 15291576] (b) Mueller JA, Sigman MS. *J. Am. Chem. Soc* 2003;125:7005–7013. [PubMed: 12783555]

Scheme 1.
Proposed mechanism of carboamination reaction.

Scheme 2.
Proposed mechanism for formation of regioisomer **4**.

Scheme 3.
Proposed mechanism for formation of **5**.

Table 1Catalyst optimization.^[a]

(3)

Entry	Ligand	3a/4a ^[b]	3a/5a ^[b]	Isolated yield ^[c]
1	dppb	20:1	3:1	60
2	dppe ^[d]	15:1	5:1	
3	PCy ₃ ^[d]	13:1	4:1	
4	P(2-furyl) ₃	21:1	19:1	81

^[a] Conditions: amine **1** (1.0 equiv), vinyl bromide **2a** (1.1 equiv), NaOt-Bu (1.2 equiv), $\text{Pd}_2(\text{dba})_3$ (1 mol%), Ligand (2 mol% of bidentate ligand or 4 mol% of monodentate ligand), toluene (0.25 M), 60 °C.

^[b] Ratios determined by GC analysis of crude reaction mixtures.

^[c] Isolated yields refer to the total yield of an inseparable mixture of products **3a**, **4a**, and **5a**.

^[d] This reaction was conducted at 110 °C.

Table 2Palladium-catalyzed synthesis of *N*-phenyl-2-allylpyrrolidines.^[a]

Entry	Vinyl bromide	Product (<i>E/Z</i>)
1		2a (>20:1 <i>E/Z</i>)
2		2b (9:1 <i>E/Z</i>)

Entry	Vinyl bromide	Product (<i>E/Z</i>)
-------	---------------	------------------------

Entry	Vinyl bromide	Product (E/Z)
-------	---------------	---------------

6

7

Entry	Vinyl bromide	Product (<i>E/Z</i>)
-------	---------------	------------------------

8

2h (7:1 *E/Z*)

Entry	Vinyl bromide	Product (E/Z)
-------	---------------	---------------

9

[a] Conditions: amine **1** (1.0 equiv), vinyl bromide **2** (1.1–2.0 equiv), NaO *t*-Bu (1.2 equiv), Pd₂(dba)₃ (1 mol%), P(2-furyl)₃ (4 mol %), toluene (0.25 M), 60–110 °C.

[b] Product ratios refer to the isolated material, as judged by ¹H NMR analysis. Product ratios and yields refer to the average of two or more experiments.

[c] This material contained ~5 % of **5a** as an inseparable impurity.

[d] This material contained ~10 % of **5d** as an inseparable impurity.

Table 3

Product (<i>E/Z</i> ratio) ^[b]	Regio ^[b]	Yield [%] ^[b]
--	----------------------	--------------------------

>20:1
72
(dr 7:1)

Product (E/Z ratio) ^[b]	Regio ^[b]	Yield [%] ^[b]
10n (>20:1 E/Z)	20:1	82 (dr >20:1)
11n (>20:1 E/Z)	>20:1	71 (dr >20:1)
12g	>20:1	84 (dr >20:1)

Product (E/Z ratio) ^[b]	Regio ^[b]	Yield [%] ^[b]
------------------------------------	----------------------	--------------------------

7:1
79
(dr >20:1)

TMS
7:1 E/Z

Product (E/Z ratio) ^[b]	Regio ^[b]	Yield [%] ^[b]
------------------------------------	----------------------	--------------------------

>20:1
55
(dr 10:1)

Product (E/Z ratio) ^[b]	Regio ^[b]	Yield [%] ^[b]
13g	>20:1	64 ^[c] (dr 10:1)
	3.6:1	61 ^[c] (dr >20:1)

l-Dppe (2 mol %) used as ligand in this reaction. PMP = *p*-methoxyphenyl.

Yield [%] ^[b]	Product (E/Z ratio) ^[b]
<div data-bbox="454 1533 909 2079"> <p>17 (E/Z ratio > 20:1)</p> </div>	<div data-bbox="454 315 909 1470"> <p>17 (>20:1 E/Z)</p> </div>

Yield [%] ^[b]	Product (E/Z ratio) ^[b]
50[c], 4e], [f] (>20:1 dr)	18 (>20:1 E/Z)

Yield [%] ^[a] / ^[b]	Product (E/Z ratio) ^[a] / ^[b]
60 ^[d] , 77 ^[f]	19 (>20:1 E/Z)

Yield [%]^[b]Product (E/Z ratio)^[b]