Published in final edited form as:

Clin Gastroenterol Hepatol. 2009 May ; 7(5): 549–553. doi:10.1016/j.cgh.2008.10.007.

Suboptimal Rates of Cervical Testing Among Women with Inflammatory Bowel Disease

Millie D Long, MD, MPH^{1,2}, Carol Q Porter³, Robert S Sandler, MD, MPH^{1,2}, and Michael D Kappelman, MD, MPH⁴

¹ University of North Carolina – Chapel Hill, Division of Gastroenterology and Hepatology

² University of North Carolina – Chapel Hill, Department of Epidemiology

³ Cecil G. Sheps Center for Health Services Research, University of North Carolina -Chapel Hill

⁴ University of North Carolina – Chapel Hill, Division of Pediatric Gastroenterology and Hepatology

Abstract

BACKGROUND & AIMS—Women with inflammatory bowel disease (IBD) have a high incidence of abnormal cervical cytology. However, little is known about how frequently women with IBD are tested for cervical abnormalities. We aimed to determine cervical testing rates among women with IBD, specifically those on immunosuppressant medications, and identify risk factors associated with low incidence of screening.

METHODS—Using the PharMetrics Patient-Centric Database from 1996 to 2005, we identified cases of IBD and matched controls via a validated algorithm. Using logistic regression, we compared utilization of cervical testing with IBD case status, patients' age, use of immunosuppressive medications, Medicaid insurance status, and use of primary care services.

RESULTS—Only 70.4% of women with IBD (n=9356) and 65.2% of matched controls (n=25849) received cervical testing (at least once every 3 years). Women with IBD who used primary care services had increased odds of cervical testing (OR 1.37, 95% CI 1.19–1.59). Factors associated with reduced testing included Medicaid insurance (OR 0.28, 95% CI 0.19–0.41), immunosuppressant medication use (OR 0.81, 95% CI 0.74–0.88) and increased age (p for trend <0.01). Among women on immunosuppressive medications (n=7415), 50.1% were tested over a 15-month period. Women on immunosuppressive medications who used primary care services has improved odds of cervical testing (OR 1.28 95% CI 1.14–1.45) whereas those with Medicaid insurance had reduced odds (OR 0.54 95% CI 0.39–0.74).

CONCLUSIONS—Women with IBD are tested for cervical abnormalities at suboptimal rates. Quality improvement initiatives are needed to improve disease prevention services for women with IBD.

Correspondence: Millie Long MD, MPH, CB #7080, University of North Carolina – Chapel Hill, Chapel Hill, NC 27599-7080, Phone: 919-966-4131, Fax: 919-966-6482.

Financial Disclosures: The authors have no financial disclosures.

Writing Assistance: None

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

INTRODUCTION

It is estimated that 11,070 women will be diagnosed with cervical cancer and 3,870 women will die of cancer of the cervix in 2008.¹ A much higher percentage of women will develop abnormal cervical pathology, a precursor to cervical cancer. The American College of Obstetrics and Gynecology (ACOG) and the American Cancer Society (ACS) recommend screening with Pap smear at least every 3 years for all women, and annual screening for women <30 years of age and women who are immunosuppressed.^{2, 3} This is due to the risk of human papilloma virus (HPV) infection leading to dysplasia among immunosuppressed women. Although cervical malignancy is largely preventable with proper screening, it is estimated that 50% of women who receive diagnoses of cervical cancer have never been screened.²

Women with inflammatory bowel disease (IBD) are a population that commonly use immunosuppressant medications and may thus be at higher risk for cervical abnormalities. Currently, it is unclear whether there is also an increased risk of cervical cancer in this population. Two recent studies found a higher proportion of abnormal Pap smears in women with IBD compared to matched controls.^{4, 5} One of these studies found that this increased risk was associated with immunosuppression.⁴ These findings suggest that although screening for cervical dysplasia is recommended for all women, this is a particularly important component of healthcare maintenance and prevention for women with IBD.

Despite the importance of performing recommended Pap smear screening in women with IBD, a growing body of literature suggests that women with IBD⁶ and other chronic illnesses⁷ do not receive optimal screening and preventive care. If women with IBD have an increased risk of cervical abnormalities and a reduced rate of screening, this represents an area to intervene with quality improvement initiatives and potential guidelines for vaccination. Therefore, using a large administrative database, we sought to compare the actual rate of cervical testing in women with and without IBD. We also sought to determine whether the high risk sub-group of women with IBD who were on immunosuppressant medications were more likely to be screened. Our secondary aim was to identify factors associated with reduced Pap smear testing rates, such as age, insurance status and utilization of primary care services.

METHODS

Study Design and Data Source

In this cross-sectional study, we analyzed the medical, surgical, and pharmaceutical insurance claims contained in the PharMetrics Patient-Centric Database (IMS Health, Watertown, MA) for the period August 1, 1996 through June 30, 2005. This longitudinal, patient-level database has been used in previous epidemiological studies of inflammatory bowel disease,⁸ and at the time of this study included claims from 87 health plans in 33 states. Prior studies have reported PharMetrics to be representative of the national commercially-insured population on a variety of demographic measures, including geographic region, age, gender, and health plan type.⁹

Patient selection

All female patients aged 20–64 with at least 36 months of continuous health plan enrollment were eligible for inclusion in this analysis. We chose an initial age of 20 because guidelines recommend initiation of Pap smears three years after the onset of sexual activity (median age of first intercourse for women is 17.4)¹⁰ or age 21. We chose 64 as the upper limit for evaluation since the United States Preventive Services Task Force (USPSTF) recommends discontinuing routine Pap smear screening at ages >65¹¹ and because Medicare eligibility

begins at age 65. We identified cases of Crohn's disease (CD) and ulcerative colitis (UC) using a previously reported administrative definition.⁸ This definition included patients with at least 3 health care contacts, on different days, associated with an International Classification of Diseases, 9th Revision, Clinical Modification diagnosis code for CD (555.xx) or UC (556.xx), or patients with at least one claim for CD or UC and at least 1 pharmacy claim for any of the following medications: mesalamine, olsalazine, balsalazide, sulfasalazine, 6-mercaptopurine, azathioprine, infliximab, adalimumab, and enteral budesonide. For patients who had claims for both CD and UC, disease assignment was made according to the majority of the last 9 claims. For each case, we randomly selected up to 3 non-IBD controls, matched for age, gender, health plan, geographic region, and Medicaid exposure (patients managed by a managed Medicaid plan).

Assessment of outcome (cervical testing)

We ascertained whether or not cervical testing was performed in each patient and control via a previously validated claims algorithm reported to have greater than 95% sensitivity and 95% specificity when compared to the medical record.¹² The algorithm heavily weighs pathology and lab codes for cervical specimen analysis as evidence of cervical testing. The dependent variable for cervical testing was coded in a binary fashion (yes/no).

Assessment of factors associated with cervical testing

We analyzed whether the following potential factors were associated with increased or decreased use of cervical testing: utilization of primary care services/provider (PCP), use of immunosuppressant medications, Medicaid insurance status (a proxy for socioeconomic status), and age. Utilization of PCP was defined as any visit to a general practitioner, internist, family practitioner or geriatrician over the analysis window (binary yes/no). We also performed separate analyses including OB/Gyn visits in the definition of primary care utilization. Immunosuppressant medication use was defined as 2 or more filled prescriptions of at least one of the following medications: prednisone, azathioprine, 6-mercaptopurine, methotrexate, infliximab or adalimumab (binary yes/no). Medicaid insurance status was specified within the dataset (binary yes/no). Age was analyzed in ten year categories.

Additional Analyses

Analyses were then repeated for the second population of women with IBD on chronic immunosuppressant medications over a 15 month analysis window (this interval was chosen to approximate the American College of Obstetrics and Gynecology's recommended 12 month interval for cervical testing for immunosuppressed women). Chronic use of immunosuppressant medications was defined as at least 2 claims for any of the following medications: prednisone, azathioprine, 6-mercaptopurine, methotrexate, infliximab or adalimumab.

Statistical analysis

Bivariate analyses using Pearson's chi squared test statistic were performed to evaluate the frequency of cervical testing over the 36 month analysis window by IBD case status, and over the 15 month analysis window for IBD patients treated with chronic immunosuppression. Similar bivariate analyses were used to evaluate the association between performance of cervical testing and the following independent variables: visit to a primary care physician, immunosuppressant medication use, Medicaid insurance status, and age. Next, odds ratios (OR) and 95% confidence intervals for each of these independent variables were estimated from unconditional logistic regression models.

For all analyses, *p*-values were two-sided, and a *p*-value of 0.05 or less was considered statistically significant. All statistical analyses were performed using Stata version 9.0 (Texas Station, TX). Analyses were performed for the overall population of women with IBD and also stratified by ulcerative colitis and Crohn's disease diagnosis. The study protocol was granted exemption from review by the Institutional Review Board at University of North Carolina because it involved the use of existing, de-identified data.

RESULTS

A total of 9356 cases of women with IBD with at least 36 months of continuous health plan enrollment meeting met our a priori criteria as did 25,849 age and geographically matched women without IBD. The characteristics of the population are shown in Table 1. The mean age of the IBD patients and matched controls was 44 (s.d. 10). Each of the 4 major U.S. census regions were adequately represented in our patient population. More women with IBD had a primary care visit over the 36 month window ($p < 0.01$).

Only 70.4% of women with IBD and 65.2% of their non-IBD matched controls received the recommended Pap smear screening over the three year observation period. Among those women with IBD, 71.1% who utilized primary care services versus 64.0% that did not utilize primary care services underwent appropriate testing ($p < 0.01$). Significantly fewer women with IBD who had Medicaid insurance, as compared to commercial insurance, were screened (43.9% versus 70.7%, $p < 0.01$). Slightly fewer women with IBD on immunosuppressant medications underwent cervical testing (68.3% versus 71.7%, $p < 0.01$). Multivariate analyses confirmed these findings: those with a PCP had significantly increased odds of Pap smear screening, and those on Medicaid insurance, on immunosuppressive medications, and of increasing age had lower rates (Table 2). Additional analyses were performed including OB/Gyn visits in the definition of PCP visit, and showed similar results of improved cervical testing with access to primary care. Women with IBD who visited a PCP or OB/Gyn were over twice as likely to obtain a Pap smear compared to women without a PCP or OB/Gyn visit (OR 2.28 95% CI 1.89–2.75).

To evaluate the possibility that women prescribed immunosuppressant medications have more severe disease and seek GI specialty care at the expense of primary care, we analyzed the bivariate relationship between immunosuppressive use and PCP care. We found that a slightly greater percentage of women with IBD on immunosuppressive medications had primary care visits over the study period (91.8% versus 89.3%, $p < 0.001$).

These analyses were repeated within strata of Crohn's disease and ulcerative colitis and there were no differences (data not shown).

Next we analyzed whether women with IBD who may be at greater risk for cervical dysplasia as a result of immunosuppression received annual screening with Pap smears as recommended by ACOG guidelines. A total of 7415 women with IBD and 2 or more prescriptions for immunosuppressant medications were identified. The demographic characteristics of this subgroup are shown in Table 3 and in general were similar to the overall population of women with IBD. For these high-risk women, only 50.1% were screened with a pap smear over a 15 month window. Those who visited a PCP had a higher rate of Pap smear testing (51.0% versus 46%, $p < 0.01$), and those with Medicaid insurance had a significantly lower rate of Pap smear testing as compared to those with commercial insurance (37.8% versus 50.4%, $p < 0.01$). On multivariate analyses, those with a PCP had higher odds of Pap smear testing (OR 1.28 95% CI 1.14–1.45), this was again seen when Ob/Gyn was included in the definition of PCP (OR 2.95 95% CI 2.51–3.46) and those with Medicaid insurance had a lower odds of Pap smear testing (OR 0.54 95% CI 0.39–0.74).

Older women were also less likely to undergo cervical testing when analyzed in 10 year increments. (Table 4) Analyses were repeated using a more conservative definition of immunosuppressive medication use (azathioprine, 6-mercaptopurine, methotrexate, infliximab or adalimumab but not corticosteroids). Similar to the initial analysis, women with a PCP visit had increased odds of cervical testing (adjusted OR 1.39 (95% CI 1.20–1.61)) and women with Medicaid insurance had reduced odds of cervical testing (adjusted OR 0.72 (95% CI 0.46 – 1.12)).

DISCUSSION

In this large, cross sectional study we found that only approximately 2/3 of women with IBD receive screening for cervical dysplasia and/or cancer as recommended by USPSTF, ACS and ACOG guidelines. We also found that those women without utilization of primary care services, those who were insured by a Medicaid plan, and those at an older age were the least likely to receive this important preventive service. This suboptimal preventive care is particularly alarming, given the abundant evidence that Pap smear screening can reduce the incidence of and mortality from cervical cancer.¹¹ Indeed, although this malignancy is largely preventable with proper screening, it is estimated that 50% of women who receive diagnoses of cervical cancer have never been screened.²

These results are both timely and significant as two recent studies have demonstrated an increased risk of abnormal Pap smears among women with IBD. Bhatia et al found a higher prevalence of abnormal Pap smears among women with IBD as compared to age matched controls (18% versus 5%).⁵ Similarly, Kane et al⁴ also demonstrated an increased risk of abnormal Pap smear among women with IBD at a tertiary care referral center as compared to non-IBD controls. In this small study, women with IBD who used immunosuppressant medications had an even greater risk of having an abnormal Pap smear. With evidence of increased abnormal cervical pathology among women with IBD, it is essential to 1) determine whether US women with IBD are receiving appropriate screening and 2) to identify the factors associated with reduced adherence to cervical screening recommendations.

To our knowledge, we are the first to examine utilization patterns of cervical testing in a large, geographically diverse, population-based sample of women with IBD. A smaller study of IBD patients followed at two midwestern academic medical centers found that approximately 90% of patients received a pap smear every three years.⁶ We believe that the lower rate of cervical screening observed here (70.4%) is more likely to reflect the care delivered to women in the broader US community. Indeed our results mirror those observed in other chronic disease processes where immunosuppression is used, such as rheumatoid arthritis. Kremer et al evaluated utilization of preventive services among women with rheumatoid arthritis from 1987–1995 via medical record review of inpatient and outpatient records and found that only 77% complied with recommended cervical testing every 3 years.⁷

Understanding factors related to utilization of preventive services is paramount in improving screening rates. Our finding that women who visited a primary care physician were more likely to receive a screening Pap smear suggests that women with IBD ought to be co-managed by a primary care provider along with a gastroenterologist. Nevertheless, the suboptimal cervical screening even among IBD patients who visited a primary care provider indicates that “missed opportunities” still exist. We also found that Medicaid insurance (as a proxy for poor socioeconomic status) was associated with poor cervical screening. While there were small numbers of Medicaid patients within our dataset, this is an important, albeit exploratory, observation. To our knowledge socioeconomic disparities in the screening of

patients with IBD have not seen previously described; however, similar social disparities in cancer screening have been reported in non-IBD populations¹³ and highlight the fact that our health system must strive for more equitable care, even among IBD patients.

The fact that we found immunosuppressive medication use to be inversely associated with cervical screening is quite concerning. If the prior studies from Kane et al⁴ and other studies from the rheumatologic literature⁷ are correct, then it appears that the women at highest risk for cervical pathology are the ones who, paradoxically, are the least likely to be screened. This high risk yet under-screened population should be a high priority for quality improvement initiatives.

The strengths of this study include both the size and diversity of the study population. By drawing from a large number of health plans of varying size, type, and location, we believe that the results presented here are broadly generalized to the commercially insured population of the United States. Another strength of this study was the ability to use pharmacy claims in order to delineate use of immunosuppressive medications

An inherent limitation to using administrative data for epidemiological studies is the lack of clinical detail, resulting in the possibility misclassification of our included patients. We used a stringent case definition that required either multiple IBD-related health contacts or IBD-specific pharmaceutical claims to establish a diagnosis of CD or UC. Similar administrative definitions have been previously reported by our group⁸ and others.^{14, 15} Similarly, we were unable to detect Pap smears that were performed and not billed which would result in an underestimation of the actual cervical testing in this population. However, the algorithm used to identify Pap smear utilization has been previously validated against chart review with very high sensitivity and specificity¹² and thus we feel that we had near complete capture of the Pap smears that were performed. Due to the lack of clinical data, we were unable to assess HPV status and smoking, which are known risk factors for cervical cancer. To the extent that these are associated with the performance of pap smears, they may represent unmeasured confounders.

Another limitation to this study is that although our data source is broadly generalizable to the commercially insured population of the United States, several populations were notably excluded from this analysis such as the uninsured and the elderly. The uninsured may receive Pap smears through charity care, and these would not be captured in our database. However, we expect that the uninsured population would be less likely to receive preventive services such as Pap smears, and thus, the overall cervical screening provided to IBD patients in the US might be even worse than that reported here. We also did not have data on patients over the age of 65; however given that the USPSTF only recommends cervical screening for those less than 65 years of age this does not represent a significant limitation.

Part of the effect of reduced cervical screening with advancing age that we observed in our study may have been related to increased rates of hysterectomy with advancing age. Approximately 23.3% of women over 18 years of age in the United States have had a hysterectomy.¹⁶ However, the number of these procedures done over the past decade has been decreasing. The prevalence reaches approximately 18% during the reproductive years, and increases to about 48% after this (peaking at approximately age 75).¹⁶ We were unable to account for prior hysterectomy in our analysis, owing to the fact that we only had claims for a 3 year time period. Instead, we repeated our analyses exclusively in women < 44 years of age (hysterectomy rate peaks between ages 45–64), and the main findings did not change (data not shown).

In summary, this study provides convincing evidence that women with IBD do not receive recommended screening for cervical dysplasia/cancer. We also found that women who are

on immunosuppressive medications, those who are older, insured by Medicaid, and who do not utilize primary care services are the least likely to receive this important preventive service. Given recent data that women with IBD may be at increased risk for abnormal Pap smears⁴, quality improvement initiatives are needed to improve screening rates, particularly in these vulnerable populations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported in part by funding from the National Institutes of Health for the Digestive Disease Epidemiology Training Program, Grant # T32 DK007634, and the Center for Gastrointestinal Biology and Disease, Grant # P30 DK3497.

Abbreviations

ACOG	American College of Obstetrics and Gynecology
ACS	American Cancer Society
CD	Crohn's disease
HPV	Human Papilloma Virus
IBD	Inflammatory Bowel Disease
Ob/Gyn	Obstetrics and Gynecology Provider
PCP	Primary care provider
UC	Ulcerative colitis
USPSTF	United States Preventive Services Task Force

References

1. SEER Data. 2007
2. ACOG practice bulletin. Clinical management guidelines for obstetrician-gynecologists. Number 44, July 2003. (Replaces Committee Opinion Number 252, March 2001). *Obstet Gynecol* 2003;102:203–13. [PubMed: 12850637]
3. Smith RA, Cokkinides V, Brawley OW. Cancer screening in the United States, 2008: a review of current american cancer society guidelines and cancer screening issues. *CA Cancer J Clin* 2008;58:161–79. [PubMed: 18443206]
4. Kane S, Khatibi B, Reddy D. Higher incidence of abnormal pap smears in women with inflammatory bowel disease. *Am J Gastroenterol* 2008;103:631–6. [PubMed: 17941962]
5. Bhatia J, Bratcher J, Korelitz B, Vakher K, Mannor S, Shevchuk M, Panagopoulos G, Ofer A, Tamas E, Kotsali P, Vele O. Abnormalities of uterine cervix in women with inflammatory bowel disease. *World J Gastroenterol* 2006;12:6167–71. [PubMed: 17036389]
6. Selby L, Kane S, Wilson J, Balla P, Riff B, Bingcan C, Hoellein A, Pande S, de Villiers WJ. Receipt of preventive health services by IBD patients is significantly lower than by primary care patients. *Inflamm Bowel Dis* 2008;14:253–8. [PubMed: 17932966]
7. Kremers HM, Bidaut-Russell M, Scott CG, Reinalda MS, Zinsmeister AR, Gabriel SE. Preventive medical services among patients with rheumatoid arthritis. *J Rheumatol* 2003;30:1940–7. [PubMed: 12966594]

8. Kappelman MD, Rifas-Shiman SL, Kleinman K, Ollendorf D, Bousvaros A, Grand RJ, Finkelstein JA. The prevalence and geographic distribution of Crohn's disease and ulcerative colitis in the United States. *Clin Gastroenterol Hepatol* 2007;5:1424–9. [PubMed: 17904915]
9. Stempel DA, Mauskopf J, McLaughlin T, Yazdani C, Stanford RH. Comparison of asthma costs in patients starting fluticasone propionate compared to patients starting montelukast. *Respir Med* 2001;95:227–34. [PubMed: 11266241]
10. <http://www.kff.org/youthhivstds/upload/U-S-Teen-Sexual-Activity-Fact-Sheet.pdf>
11. <http://www.ahrq.gov/clinic/uspstf/uspscerv.htm>
12. Fowles JB, Fowler E, Craft C, McCoy CE. Comparing claims data and self-reported data with the medical record for Pap smear rates. *Eval Health Prof* 1997;20:324–42. [PubMed: 10183327]
13. Cabeza E, Esteve M, Pujol A, Thomas V, Sanchez-Contador C. Social disparities in breast and cervical cancer preventive practices. *Eur J Cancer Prev* 2007;16:372–9. [PubMed: 17554211]
14. Herrinton LJ, Liu L, Lafata JE, Allison JE, Andrade SE, Korner EJ, Chan KA, Platt R, Hiatt D, O'Connor S. Estimation of the period prevalence of inflammatory bowel disease among nine health plans using computerized diagnoses and outpatient pharmacy dispensings. *Inflamm Bowel Dis* 2007;13:451–61. [PubMed: 17219403]
15. Bernstein CN, Blanchard JF, Rawsthorne P, Wajda A. Epidemiology of Crohn's disease and ulcerative colitis in a central Canadian province: a population-based study. *Am J Epidemiol* 1999;149:916–24. [PubMed: 10342800]
16. Merrill RM. Hysterectomy surveillance in the United States, 1997 through 2005. *Med Sci Monit* 2008;14:CR24–31. [PubMed: 18160941]

Table 1

Characteristics of the population (women with at least 36 months of continuous enrollment) by IBD case status^{*}.

Characteristic	IBD (n=9356)		No IBD (n=25849)	
Age	9356	44.1 (10.4)	25849	44.2 (10.3)
Region				
East	3034	32.4%	8705	33.7%
Midwest	3292	35.2%	9473	36.7%
South	742	7.9%	1301	5.0%
West	2288	24.5%	6370	24.6%
PCP ^{**}	8446	90.3%	21495	80.2%
Medicaid	107	1.1%	350	1.4%
Pap smear [†]	6583	70.4%	16851	65.2%

^{*} Mean (standard deviation) for continuous variables by student's t-test, percent for categorical variables by Pearson's chi square test statistic

^{**} Defined as any visit to a primary care provider (internist, family practitioner, general practitioner or geriatrician) over the 36 month period

[†] See appendix 1 for claims algorithm for pap smear over the 36 month study period

Table 2

Multivariate adjusted odds ratios of factors associated with Pap smear screening in US women with IBD over the 36 month study period*

Cervical Testing		
	OR	95% CI
No PCP	1.0 (referent)	
PCP	1.37	1.19–1.59
Commercial insurance	1.0 (referent)	
Medicaid insurance	0.28	0.19–0.41
No immunosuppressive medication use	1.0 (referent)	
Immunosuppressive medication use	0.81	0.74, 0.88
20–29	1.0 (referent)	
30–39	0.93	0.78–1.11
40–49	0.76	0.64–0.90
50–59	0.55	0.46–0.65
60–64	0.36	0.28–0.47

* adjusted for region of the country, medicaid insurance, primary care visit and age as appropriate

Table 3

Characteristics of the population of women with IBD, overall and stratified by Ulcerative Colitis and Crohn's disease, with at least 15 months of continuous enrollment and use of immunosuppressant medications (defined as at least 2 prescriptions filled for immunosuppressant medication over this time period)*

Characteristic	IBD (n=7415)	CD (n=4626)	UC (n=2789)
Age	7415 43.2 (11.2)	4626 42.6 (11.2)	2789 44.0 (11.1)
Region			
East	1646 22.2%	990 21.4%	656 23.5%
Midwest	2788 37.6%	1829 39.5%	959 34.4%
South	1688 22.8%	1051 22.7%	637 22.8%
West	1293 17.4%	756 16.3%	537 19.3%
PCP**	6068 81.8%	3786 81.8%	2282 81.8%
Medicaid	172 2.3%	133 2.9%	39 1.4%
Pap smear†	3717 50.1%	2263 48.9%	1454 52.1%

* Mean (standard deviation) for continuous variables, percent for categorical variables

** Defined as any visit to a primary care provider (internist, family practitioner, general practitioner or geriatrician) over the 15 month period

† See appendix 1 for claims algorithm for Pap smear over the 15 month period of continuous enrollment

Table 4

Multivariate adjusted odds ratios of factors associated with Pap smear screening in US women with IBD on immunosuppressant medications over a 15 month study period*

Cervical Testing		
	OR	95% CI
No PCP	1.0 (referent)	
PCP	1.28	1.14–1.45
Commercial insurance	1.0 (referent)	
Medicaid insurance	0.54	0.39–0.74
20–29	1.0 (referent)	
30–39	0.78	0.67–0.91
40–49	0.67	0.58–0.78
50–59	0.56	0.48–0.65
≥ 60	0.40	0.32–0.50

* adjusted for region of the country, medicaid insurance, primary care visit and age as appropriate