Published in final edited form as:

J Allergy Clin Immunol. 2010 March ; 125(3): 757–760. doi:10.1016/j.jaci.2009.10.054.

Evidence for Neuronal Expression of Functional Fc (epsilon and gamma) Receptors

Hanneke van der Kleij, PhD^a, Nicolas Charles, PhD^b, Khalil Karimi^a, Yu-Kang Mao^a, Jane Foster^a, Luke Janssen^a, Ping Chang Yang^a, Wolfgang Kunze^a, Juan Rivera, PhD^b, and John Bienenstock, MD^a

^a Brain-Body Institute, St. Joseph's Healthcare Hamilton and Department of Pathology & Molecular Medicine, McMaster University, Hamilton, Ontario, Canada L8N4A6

^b Laboratory of Immune Cell Signaling, National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of Health, Bethesda, Maryland 20892, USA

Short Summary

We report that neurons express functional Fc receptors (including FcεR1), that can be activated by antigen, transmit signals along nerve fibers *in vitro* and *in vivo*. These results open new avenues of investigation in neurogenic inflammation and allergic diseases.

Keywords

Neuron; IgE; allergy; neurogenic inflammation; Fc receptors

To the Editor:

The Fc receptor family plays a key role in adaptive immunity through the binding of immunoglobulin antibodies that recognize an immune insult and elicit an inflammatory response leading to its clearance. Dysregulation of this receptor family may have untoward consequences that result in autoimmune and allergic diseases. Many of these diseases seem to involve the nervous system and are exacerbated by stress or other neurological challenges. Recently, the presence of Fc receptors was uncovered on dorsal root ganglion neurons and suggested an IgG and possibly IgE-mediated activation of neurons ^{1–3}. We set out to more extensively explore which Fc receptors might be expressed in neurons, and whether they were functional and able to transmit signals to interconnected neurites *in vitro* and *in vivo*

Messenger RNA was isolated from a highly pure culture of mouse superior cervical ganglion (SCG) neurons ⁴ and expression of Fc receptor transcripts assessed by reverse transcriptase-polymerase chain reactions (RT-PCR) using specific primers for Fcγ and Fcε family members. Fig. 1*a* demonstrates the presence of transcripts for the immunoglobulin-binding α chain of FcγRI, II, III, and IV in three individual SCG neuron mRNA preparations. A small amount of the transcript for the low affinity IgE receptor (FcεRII or CD23) was also detected relative to

Correspondence: John Bienenstock, The Brain-Body Institute, St. Joseph's Healthcare Hamilton, Department of Pathology & Molecular Medicine, McMaster University, 50 Charlton Avenue East, Juravinski Innovation Tower, Room 3304, Hamilton, Ontario, Canada L8N4A6. bienens@mcmaster.ca. tel: 905 522 1155 ext 35203, Fax: 905 540 6593.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

that seen in B cells, known to express this receptor. Fc γ RI transcripts were detected in bone marrow derived mouse mast cells but these levels were less than seen in the neurons. This observation and the inability to detect mMCP-6mRNA in either Balb/c or B16 mice, together with the absence of CD23 transcripts in both neurons and mast cells provided confidence that the observed Fc receptor transcripts in neurons was not a result of mast cell contamination of cultures.

We also unexpectedly observed the presence of transcripts for the α , β , and γ chains of the high affinity IgE receptor (Fc ϵ RI) (Fig. 1,*b*). While the trimeric form ($\alpha\gamma_2$) of this receptor has been described in cells other than mast cells or basophils (such as in human Langerhans cells 5), the expression of the tetrameric form ($\alpha\beta\gamma_2$) was previously thought to be limited to these pro-inflammatory cells. The trimeric Fc ϵ RI shows weak calcium signals when compared to the tetrameric form due to the absence of the Fc ϵ RI β in the former ⁶. To determine if the Fc ϵ RI was expressed on the cell surface of SCG neurons, cells were incubated with IgE and with an antibody to the neuronal specific protein gene product (PGP) 9.5 (which encodes a neuronal ubiquitin C-terminal hydrolase not found on glia) and binding visualized with a fluorescent secondary antibody. IgE was detected on the neuronal cell surface (Fig. 2,*a*). While the presence of PGP9.5 was most evident in the cell body, IgE binding was detected in plasma membrane of both the cell body and neurite extensions. As shown in Figs. 2,*b–d*, the expression of Fc ϵ RI was further confirmed by detection of the α , β , and γ chains of this receptor.

Scorpion venom is a known potent selective activator of neurons and elicits a rapid rise in intracellular Ca²⁺ (Fig. 3,*a*). To test the functionality of Fc ϵ RI expressed on neurons, SCG neurons sensitized with DNP-specific IgE were challenged with DNP-HSA (Ag) and a rapid rise in intracellular Ca²⁺ was observed (Fig. 3,*a*). No changes in intracellular Ca²⁺ were observed when serum albumin alone was used as Ag or when cells were not sensitized with IgE (data not shown). Fc γ RIV was recently described to bind the IgE^b allotype but does not recognize the IgE^a allotype ⁷, whereas Fc ϵ RI binds both allotypes. We excluded that IgE/Ag-mediated calcium responses might occur through Fc γ RIV, by use of both a IgE^a and IgE^b allotypes. Both IgE allotypes similarly elicited calcium responses (Fig. 3,*a*). Addition of Ag elicited a relatively uniform Ca²⁺ responses in the stimulated cell population (Fig. 3,*b*). Based on the high affinity binding of monomeric IgE (was not removed by washing the cells) and the dose response to Ag (Fig. 3, *D*), we could also exclude the involvement of CD23.

Thus, the findings demonstrate the presence of functional high affinity Fc ϵ receptors on SCG neurons. Since Fc γ RIII is known to activate mast cells, its functionality was also tested. SCG neurons sensitized with DNP-specific IgG1, which preferentially binds Fc γ RII and III but weakly to Fc γ RI and not to Fc γ RIV ⁸, showed modest increases in intracellular Ca²⁺ that increased with a large dose of Ag (Fig. 3,*c*). Both IgE and IgG-mediated responses were concentration dependent and a Ca²⁺ response (Fig. 3,*d*) was not elicited in all challenged neurons. Moreover, as expected (given the weak binding of monomeric IgG1 to Fc γ receptors), increased responsiveness via IgG required much higher concentrations of Ag than for IgE.

To test if the Ca²⁺ signals elicited by Fc ϵ RI stimulation could be transferred to interconnected neurites, we explored whether Ca²⁺ rises might be elicited in neighboring neurites after Ag challenge of an IgE sensitized cell body or neurite. Using a spritzer micro-pipette, antigen was puffed directly onto a neuronal cell body, causing an instantaneous (<5 sec) increased fluorescence in that cell body which moved from there to the connected neurites and propagated to the neighboring cell bodies and neurites (Fig. 4). These findings showed that Fc ϵ RI stimulation causes communication among interconnected neurites. To extend these findings to a more physiological setting, we explored whether neurons from the highly innervated intact jejunum ⁹ would respond to an Fc ϵ RI stimulus. After placement of a micropipette spritzer (dotted lines shown in Fig. 5,*a* and *g*) on a large myenteric plexus ganglion neuron (plain lines

shown in Fig. 5,a and g), the anti-DNP IgE sensitized plexus was challenged with Ag. Challenge with a spritz of 1 µg of Ag gave robust calcium responses in adjacent neurons along the nerve fiber in sensitized (2/2) but not in non-sensitized mice (0/3). Repeated spritzes of non-conjugated HSA at this concentration elicited no responses (0/2) (data not shown). To exclude possible mast cell involvement in the transmission of these robust signals, we conducted similar experiments in mast cell deficient *W/W^v* mice and in their wild type control *WBB6F1*. In *W/W^v* mice (3/3) detectable intracellular calcium increases were observed upon Ag challenge (Fig. 5,a–c). No calcium signal was seen when the same ganglion was first challenged with HSA alone (Fig. 5,d–f). The wild type littermates (*WBB6F1*) responded positively (3/3), upon challenge with specific Ag (Fig. 5,g–i) but gave no response to HSA alone. These findings confirm that the observed signal transmission by FcεRI was not likely caused by mast cells and demonstrate the *in vivo* presence of functional FcεRI on jejunal neurons, since sham sensitization *in vivo* prior to an *ex vivo* challenge yielded no response to Ag challenge.

It is well known that sensory nerves may participate in hypersensitivity reactions, a process known as neurogenic inflammation and several lines of evidence support the notion that sensory nerves may play an important role in cutaneous, lung, GI and joint inflammatory diseases. Here we now demonstrate that functionally active FcεRI is expressed on SCG and myenteric plexus neurons. The discovery of functional Fcε and Fcγ receptors on nerves clearly shows that this biological compartment is able to respond to the direct stimulus of antibody-antigen interactions. Our findings define an independent neuronal (non-mast cell/non-basophil) compartment with probable involvement in allergic and possibly other diseases.

Acknowledgments

The work performed in this study was supported by a grant from the McMaster Brain-Body Institute, St Joseph's Healthcare, Hamilton. The work of NC and JR was supported by the intramural research program of NIAMS, NIH. The authors have no financial relationships or interests in the subject of this paper to disclose.

References

1. Andoh T, Kuraishi Y. Expression of Fc epsilon receptor I on primary sensory neurons in mice. *Neuroreport* 2004;15:2029–31. [PubMed: 15486476]
2. Andoh T, Kuraishi Y. Direct action of immunoglobulin G on primary sensory neurons through Fc gamma receptor I. *FASEB J* 2004;18:182–4. [PubMed: 14630707]
3. Rijnierse A, Kroese AB, Redegeld FA, Blokhuis BR, van der Heijden MW, Koster AS, et al. Immunoglobulin-free light chains mediate antigen-specific responses of murine dorsal root ganglion neurons. *J Neuroimmunol* 2009;208:80–6. [PubMed: 19232443]
4. Furuno T, Ma D, van der Kleij HP, Nakanishi M, Bienenstock J. Bone marrow-derived mast cells in mice respond in co-culture to scorpion venom activation of superior cervical ganglion neurites according to level of expression of NK-1 receptors. *Neurosci Lett* 2004;372:185–9. [PubMed: 15542237]
5. Bieber T. Fc epsilon RI on antigen-presenting cells. *Curr Opin Immunol* 1996;8:773–7. [PubMed: 8994854]
6. Lin S, Cicala C, Scharenberg AM, Kinet JP. The Fc(epsilon)RIbeta subunit functions as an amplifier of Fc(epsilon)RIgamma-mediated cell activation signals. *Cell* 1996;85:985–95. [PubMed: 8674126]
7. Hirano M, Davis RS, Fine WD, Nakamura S, Shimizu K, Yagi H, et al. IgEb immune complexes activate macrophages through FcgammaRIV binding. *Nat Immunol* 2007;8:762–71. [PubMed: 17558411]
8. Nimmerjahn F, Bruhns P, Horiuchi K, Ravetch JV. FcgammaRIV: a novel FcR with distinct IgG subclass specificity. *Immunity* 2005;23:41–51. [PubMed: 16039578]
9. Mao Y, Wang B, Kunze W. Characterization of myenteric sensory neurons in the mouse small intestine. *J Neurophysiol* 2006;96:998–1010. [PubMed: 16899648]

Figure 1.

mRNA expression of the subunits of FcεRI, FcγRI-IV, and CD23 in SCG neurons. mRNA was collected from three (1–3) SCG neuronal cultures and expression of the indicated mRNA was measured by RT-PCR. **(a)** Expression of the α subunit of the low affinity receptors for IgG (FcγRI-IV) and for IgE (CD23). **(b, c)** Expression of the subunits of FcεRI and of mMCP-1 **(c)** in SCG neurons derived from Balb/c **(a–c)** or C57BL/6 **(c)** mice. Spleen cDNA and mRNA from bone marrow derived mouse mast cells or from B cells were used as positive controls. The mRNA of the mouse IMCD-3 kidney epithelial cell line from the inner medulla collective duct, was a negative control for FcεRI, mMCP-1 and FcγRI, III, and IV expression. A control for PCR contamination was performed using H₂O.

Figure 2.

Neurons were sensitized with IgE and incubated overnight with antibody against IgE (**a**, left panel) or with an antibody of unknown specificity (**a**, right panel). Staining with the neuron specific marker PGP9.5 (**b–c**, left panel in red) and FcεRIα (**b**), β (**c**) or γ-chain (**d**) (middle panel in green) is shown. Confocal images were overlaid (**b–d**, right panel); yellow represents merging of red and green colours.

Figure 3.

(a) Kinetics of $[Ca^{2+}]_i$ increase upon stimulation with scorpion venom (SV, Control) or with anti-DNP IgE^{a/b} sensitized neurons following Ag stimulation (10 ng/ml). The arrows indicate the time of addition of the stimulus. (b) Confocal image of calcium responses of IgE-sensitized neurons prior to and after Ag addition. (c) Kinetics of $[Ca^{2+}]_i$ increase in anti-DNP IgG 1 sensitized neurons after Ag stimulation. (d) Dose response of IgE and IgG-mediated neuron activation.

Figure 4.

Activation of interconnected neurites by FcεRI stimulation of a single neuronal cell body. **(a)** Bright field image of neuron network showing relation of Ag-containing spritzer to neuronal cell body sensitized with IgE anti-DNP. **(b)** At time zero, beginning of 500ms spritz with Ag (DNP-HSA). **(c)** 0.5s and **(d)** 1.8s after onset of spritz.

Figure 5.

(A–I) Myenteric ganglion calcium imaging. Spritzer (internal bore, 40 μm) is indicated by dotted lines and myenteric plexus by solid lines. (a–c) Anti-DNP IgE sensitized myenteric neurons were imaged in mast cell-deficient W/W^v mice. **a**, Resting condition. **b**, Fluorescent image captured 0.16s after 20ms spritz. **c**, 4s after spritz. (d–f) No increase in calcium fluorescence was observed in non-haptenated HSA spritz. (g–i) Positive calcium increases in WBB6F1 control littermates. Time sequence same as previous.