CORRESPONDENCE

Dangerous Errors in the Diagnosis and Treatment of Bony Tuberculosis

by Dr. med. Alexander Herzog in volume 36/20090

Polymerase Chain Reaction

The article very well written and clinically important, but why did the author not mention the option of using polymerase chain reaction (PCR) as a diagnostic tool? PCR is not a universal diagnostic miracle weapon, but when it works it does so within just a few days and is highly specific. Another advantage is that it can be performed using minimal volumes of aspirates, etc. Even if the procedure was not used in the specific case presented in the article it should have been evaluated in the discussion. The same holds true for both the disease and the test: They need to be borne in mind.

DOI: 10.3238/arztebl.2010.0147a

REFERENCES

1. Herzog A: Case report: dangerous errors in the diagnosis and treatment of bony tuberculosis. Dtsch Arztebl Int 2009; 106 (36): 573–7.

Dr. med. Christof Birkenmaier

Klinikum Großhadern der Ludwig-Maximilians-Universität
Marchionistr. 15
81377 München, Germany
christof.birkenmaier@med.uni-muenchen.de

Non-tuberculous Mycobacteria

The author reported a case of mycobacterial infection, which was diagnosed after a lengthy process. However, the case is actually not one of tuberculosis in the stricter sense: the confirmed species—*Mycobacterium avium intracellulare complex*—belongs to the group of non-tuberculous mycobacteria. These are often found in the environment and, in cultures, are possibly only contaminants. However, in this case the diagnosis was firmly established following the criteria of the American Thoracic Society (ATS) – the specimens came from otherwise sterile body fluids (1). Microscopy using different staining methods, as mentioned in the article, is less relevant than cultures and PCR for confirming a diagnosis, since microscopy has lower sensitivity and, in contrast to the other methods, cannot differentiate between species (2). It therefore needs to be emphasized that the diagnostic mistake in the case report consisted mainly in not doing culture and PCR.

Non-tuberculous mycobacteria—such as the described species—display sensitivities to anti-mycobacterial chemotherapy different from *Mycobacterium tuberculosis complex*. After resistance testing, recommended initial treatment should therefore consist of a combination of clarithromycin, ethambutol, and, if required,

rifabutin. By contrast to the treatment described in the article, however, isoniazid is not the treatment of choice (3).

DOI: 10.3238/arztebl.2010.0147b

REFERENCES

1. Griffith DE, Aksamit T, Brown-Elliott BA, et al.: An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. Am J Respir Crit Care Med 2007; 175(4): 367–416.
2. Schaaf SH, Zumla AI: Tuberculosis. London: Elsevier; 2009; p. 1014.
3. Kaplan JE, Benson C, Holmes KH, et al.: Guidelines for prevention and treatment of opportunistic infections in HIV-infected adults and adolescents: recommendations from CDC, the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America. MMWR Recomm Rep 2009; 58(RR-4): 1–207; quiz CE201–204.
4. Herzog A: Case report: dangerous errors in the diagnosis and treatment of bony tuberculosis. Dtsch Arztebl Int 2009; 106 (36): 573–7.

Dr. med. Norbert Heinrich

PD Dr. med. Michael Hoelscher

Abteilung für Infektions- und Tropenmedizin
Klinikum der LMU
Leopoldstr. 5
80802 München, Germany
heinrich@lrz.uni-muenchen.de

PD Dr. rer. nat. Elvira Richter

Forschungszentrum Borstel
Nationales Referenzzentrum für
Mycobakterien
Parkallee 18
23845 Borstel, Germany

Rare Pathologies

We wish to point out that in this case, no actual tuberculosis was present; rather, the patient had non-tuberculous mycobacteriosis (NTM), caused by *Mycobacterium avium intracellulare*. The infection was finally successfully treated over a period of 2 years with antibiotic combination therapy including clarithromycin (1, 2).

Such pathologies are rare and often affect immunocompromised persons, especially patients with HIV infection.

In the described case, PCR of the microscopically positive direct specimen would have been advantageous as this allows for rapid distinction between tuberculosis and NTM.

Nowadays, an Interferon-Gamma-Release Assay (IGRA) may also facilitate this distinction, in particular since the tuberculin skin test was of limited diagnostic value in the present patient, who had previously undergone BCG vaccination (3).

DOI: 10.3238/arztebl.2010.0147c

REFERENCES

1. Griffith DE, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, et al.: An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. Am J Respir Crit Care Med 2007; 175: 367–416.

2. Abraham E: Erratum: ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med* 2007; 175: 744–5.
3. Hauer B, Loddenkemper R, Diel R: Interferon-Gamma-Tests (IGRAs) – neue diagnostische Möglichkeiten in der Tuberkulosekontrolle. *Epidemiologisches Bulletin (Robert Koch-Institut)* 2009; 11: 98–101.
4. Herzog A: Case report: dangerous errors in the diagnosis and treatment of bony tuberculosis. *Dtsch Arztebl Int* 2009; 106 (36): 573–7.

Dr. med. Barbara Hauer

Prof. Dr. med. Robert Loddenkemper

Deutsches Zentralkomitee zur Bekämpfung der Tuberkulose
Stralauer Platz 34
10243 Berlin, Germany
rloddenkemper@dzk-tuberkulose.de

Chronic Recurrent Multifocal Osteomyelitis

Herzog's article is of particularly topical interest, as the author has drawn renewed attention to tuberculosis (TB), which over the past 50 years seems to have disappeared and has therefore been forgotten as a differential diagnosis (1). TB—including TB of the bone—is threatening to make a comeback and be included, as a pathomorphosis, in a confusing array of differential diagnoses, such as was shown in this case report of bony tuberculosis. The range of differential diagnoses should also include the generally little known "chronic recurrent multifocal (also monofocal) osteomyelitis" (CRMO) as an important possible diagnosis. However, this diagnosis is hampered by the erroneous interpretation of results. Consequently, the rare "primary chronic osteomyelitis" might remain unrecognized as a result of such an indirectly didactic approach.

CRMO does not manifest in sterile abscesses—which by definition do not actually exist—but in larger areas of sometimes "migrating" sterile inflammation of the bone, the localization being age dependent. These inflammations develop in three stages and are primarily lymphoplasmacellular, not "purulent"—that is, never granular leukocyte related. Magnetic resonance tomography and biopsy specimens usually provide unequivocal findings that morphologically depend on which stage of development the pathological process has reached. Further, the localization of the focus does not correspond to any of the 6 CRMO types, where none of the foci reside in the skullcap of CRMO patients. Corticosteroids are not indicated therapeutically.

A recently discovered common radiological feature may, however, erroneously link TB and CRMO in terms of the differential diagnosis: a potential parosseous, especially paravertebral, inflammation of the soft tissues with a tendency to ossification, which may result in vertebral CRMO being mistaken for TB.

However, this specific differential diagnosis is not likely to be common. In our SAPHO/CRMO project in Mainz, there was only one case among 183 patients.

DOI: 10.3238/arztebl.2010.0148a

REFERENCES

1. Herzog A: Case report: dangerous errors in the diagnosis and treatment of bony tuberculosis [Gefährliche Fehler in Diagnostik und

Therapie einer Knochentuberkulose]. *Dtsch Arztebl Int* 2009; 106 (36): 573–7.

2. Schilling F: Abschiedsvorlesung „CRMO“, Univ.-Klinik Mainz 2003, www.sapho-syndrom-crm.de

Prof. Dr. med. Fritz Schilling

Hebbelstr. 20
55127 Mainz, Germany

Lymphoblastic Neoplasia

In my opinion, the main error lies in the initial, inadequate diagnosis in the context of CT guided biopsy. Further information about the exact histological findings would have been desirable.

To base lymphatic leukemia (ALL) therapy merely on a diagnosis of suspected differentiated T-cell lymphoma seems questionable, to say the least. On the one hand, the term "differentiated T-cell lymphoma" is not a widely used medical term; on the other hand, ALL is an immature, lymphoblastic neoplasia that can be confirmed by markers such as terminal desoxyadenylate transferase. In view of the consequences of such a diagnosis, statements such as "consistent with ALL" should not have been used. I don't see any scope for different interpretations on the basis of the patient's history. In case of doubt, a fresh and bigger biopsy specimen should be taken, and if there is still doubt, molecular genetic testing should be used to confirm the clone (T-cell receptor rearrangement).

DOI: 10.3238/arztebl.2010.0148b

REFERENCES

1. Herzog A: Case report: dangerous errors in the diagnosis and treatment of bony tuberculosis. *Dtsch Arztebl Int* 2009; 106 (36): 573–7.

Dr. med. Lars Fischer

Charité Campus Benjamin Franklin
Hindenburgdamm 30
12200 Berlin, Germany
lars.fischer@charite.de

Differences to Tuberculosis

The confirmed pathogen *Mycobacterium avium intracellulare* is in the group of non-tuberculous mycobacteria—also known as ubiquitous mycobacteria or "mycobacteria other than tuberculosis" (MOTT). In contrast to tuberculosis, the strain cannot be transmitted from person to person and the disease is not notifiable under the infection protection law. The described case is thus not one of tuberculosis of the bone—the title is erroneous and incorrect.

Where non-tuberculous mycobacteria are found, the case may be one of saprophytic colonization without clinical significance or a manifest infection that requires therapy. *Mycobacterium avium* complex is the most common non-tuberculous bacteria species that is associated with pathologies in humans.

The first diagnostic puncture did not provide any histological indication of tuberculosis. Because of the raised inflammatory markers and the positive Tine test, tuberculosis was then considered as a possible differential diagnosis. In the abscess specimen obtained in

February 2001, no acid-fast bacilli were found on microscopy—I presume this refers to direct microscopy. The result of a TB culture is not reported. Using a positive culture from this specimen would have enabled the correct diagnosis earlier on, as is mentioned in the discussion section. No mention is made of whether an antibiogram was available. To select the medication on the basis of a positive culture and a confirmed pathogen alone is not consistent with latest clinical practice because therapy without an antibiogram is equivalent to “blind” therapy and entails the risk of—potentially fatal—resistance developing. Especially the treatment of non-tuberculous mycobacteriosis is difficult and takes a long time—18–24 months as a rule.

DOI: 10.3238/arztebl.2010.0148c

REFERENCES

1. An Official American Thoracic Society/Infectious Diseases Society of America Statement: diagnosis, treatment, and prevention of non-tuberculous mycobacterial diseases. *Am J Respir Crit Care Med* 2007; 175: 367–416.
2. Wagner D, Young LS: Nontuberculous mycobacterial infections: a clinical review. *Infection* 2003; 31: 257–70.
3. Karlsdorf B, et al.: Klinik und Therapie der Tuberkulose. *Pneumologie* 2008; 62: 284–94.
4. Herzog A: Case report: dangerous errors in the diagnosis and treatment of bony tuberculosis. *Dtsch Arztebl Int* 2009; 106 (36): 573–7.

Dr. med. Klaus-Detlef Schneider
Hägerweg 2
30659 Hannover, Germany

Disseminated Non-tuberculous Mycobacteriosis

The impressive case report of a pathology that manifested primarily as osteolyses in different skeletal areas cannot be assigned to a diagnosis of tuberculosis of the bone.

Tuberculosis is the exclusive name for all pathologies caused by *Mycobacterium tuberculosis* complex. *Mycobacterium avium intracellulare* complex, as found in the culture of the patient whose case was reported, is one of the non-tuberculous mycobacteria known as MOTT (“mycobacteria other than tuberculosis”) (1–3). The diagnosis of the case patient should therefore have been: disseminated non-tuberculous mycobacteriosis, caused by *Mycobacterium avium intracellulare* complex—a pathology that is often observed in association with HIV infection and in the course of neoplastic hematological disorders, such as in the case report.

In general, the distinction between TB and MOTT is of far reaching importance. The route of infection, pathogenicity, responsiveness to antimycobacterial medication, and often the clinical presentation are different—factors that influence diagnosis, treatment, and prognosis.

DOI: 10.3238/arztebl.2010.0149a

REFERENCES

1. Konietzko N, Loddenkemper R: Tuberkulose. Stuttgart: Georg Thieme Verlag 1999.

2. Henry MT, Inamdar L, O’Riordan D, Schweiger M, Watson JP: Nontuberculous mycobacteria in non-HIV patients: epidemiology, treatment and response. *Eur Respir J* 2004; 23: 741–6.
DOI: 10.1183/09031936.04.00114004
3. Griffith DE, et al.: An Official ATS/IDSA Statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med* 2007; 175: 367–416.
DOI: 10.1164/rccm.200604-571ST
4. Herzog A: Case report: dangerous errors in the diagnosis and treatment of bony tuberculosis. *Dtsch Arztebl Int* 2009; 106 (36): 573–7.

Dr. med. Robert Kropp
Prof. Dr. med. Volker Schulz
Deutsches Tuberkulose-Archiv
Liegnitzer Str. 5
36100 Petersberg, Germany
dr.robert.kropp@gmx.de

In Reply:

The number of letters underlines the importance of the topic for readers of *Deutsches Ärzteblatt*.

Several correspondents pointed out—entirely correctly—that infection with *Mycobacterium avium intracellulare*, the causative agent of avian tuberculosis, is not a typical form of TB in humans. By contrast to infection with *M. tuberculosis*, no person to person spread exists, nor is the disease notifiable.

For the clinical course, however—and that is the crux of the matter here—no difference exists. Infection of humans with avian TB (*M. avium*) is found almost exclusively in patients with HIV at a late stage of illness and affects mainly the lungs. The clinical presentation in the case report patient was therefore untypical for infection with *M. avium intracellulare*. The patient was not HIV positive, and he did not have neoplastic disease, as Dr Kropp or Professor Dr Schulz assumed.

Because of the lacking clinical differentiation between tuberculosis and infection with *M. avium*, the diagnosis of “tuberculosis of the bone” was chosen—among other reasons, to enable better understanding among non-specialist doctors. The diagnosis suggested by Dr Kropp—“disseminated non-tuberculous mycobacteriosis caused by *Mycobacterium avium intracellulare* complex”—in an atypical case would have created more confusion than understanding. Since the differentiation and treatment decision is always made only after the causative strain has been identified, I do not see the clinical diagnosis of “tuberculosis of the bone” as erroneous or even incorrect in this case, rather in contrast to Dr Schneider. The diagnosis could certainly have been made earlier if the examination had been thorough enough. The specimen taken in the first diagnostic puncture was sent to a pathology laboratory for suspected malignancy, not to a microbiological lab. The process took the wrong direction right from the outset because the wrong clinical diagnosis had been made.

As Dr Heinrich and Dr Birkenmaier correctly state, the diagnosis could have been made earlier by using PCR and cultures—had these but been considered. Dr Hauer’s mention of the interferon-gamma-release assay (IGRA), which is now available, is an important addition.

Although antibiograms are now done routinely in tuberculosis, their reliability for other mycobacteria is not undisputed. Back then, eight years ago, when much of the current knowledge into *M. avium* did not exist, therapy and its duration were decided on the basis of the causative strain.

All the literature citations mentioned in the readers' correspondence were published much later, at a time when more was known about this infection as an opportunistic infection in HIV patients. But even such new insights can protect patients only if the initial diagnosis is correct.

Dr Fischer rightly points out that, on the basis of the histology results alone, without further disease markers or molecular genetic confirmation, a diagnosis of ALL should have not been made under any circumstances, not should the patient have been subjected to chemotherapy, an invasive form of treatment.

My thanks go to Professor Schilling for pointing out the risk of confusion with CRMO (chronic recurrent multifocal osteomyelitis), which is possible because of

lymphocellular infiltration and paravertebral soft tissue inflammation. Because of the clinical course, this initial differential diagnosis became increasingly unlikely, since CRMO does not affect the skull and is not accompanied by psoas abscesses. However, the illness of the patient in this case report bore a merely temporary resemblance to CRMO, which is also characterized by a lengthy process until the correct diagnosis is made.

DOI: 10.3238/arztebl.2010.0149b

REFERENCES

1. Herzog A: Case report: dangerous errors in the diagnosis and treatment of bony tuberculosis. Dtsch Arztebl Int 2009; 106 (36): 573–7.

Dr. med. Alexander Herzog
Klinik für Integrative Onkologie
Kurstr. 16–18
63667 Nidda/Bad Salzhausen, Germany
info@fachklinikdrherzog.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.