Published in final edited form as:

Child Dev. 2010 ; 81(1): 224–236. doi:10.1111/j.1467-8624.2009.01391.x.

Neurodevelopmental Effects of Early Deprivation in Post-Institutionalized Children

Seth D. Pollak,

University of Wisconsin

Charles A. Nelson,

Harvard Medical School / Children's Hospital Boston

Mary F. Schlaak,

University of Wisconsin

Barbara J. Roeber,

University of Wisconsin

Sandi S. Wewerka,

University of Minnesota

Kristen L. Wiik,

University of Minnesota

Kristin A. Frenn,

University of Minnesota

Michelle M. Loman, and

University of Minnesota

Megan R. Gunnar

University of Minnesota

Abstract

The neurodevelopmental sequelae of early deprivation were examined by testing ($N = 132$) 8 and 9 year old children who had endured prolonged versus brief institutionalized rearing or rearing in the natal family. Behavioral tasks included measures that permit inferences about underlying neural circuitry. Children raised in institutionalized settings showed neuropsychological deficits on tests of visual memory and attention, as well as visually mediated learning and inhibitory control. Yet, these children performed at developmentally appropriate levels on similar tests where auditory processing was also involved and on tests assessing executive processes such as rule acquisition and planning. These findings suggest that specific aspects brain-behavioral circuitry may be particularly vulnerable to post-natal experience.

Over the past decade, increased attention has been devoted to the development of children who have spent some or all of their lives in institutional care (Johnson, 2001). The increase in adoption of institutionalized children has heightened concerns about long-term effects of early deprivation. While the deprivation experienced by children in institutional settings is often impossible to accurately quantify, the environments many of these children endure fall below the quality needed to sustain normal physical and behavioral development. As evidence, institutionalized infants/toddlers lose about 1 month of linear growth for every 2–3 months in institutional care (Johnson, 2001) with behavioral development exhibiting similar dramatic delays retardation (Gunnar, 2001). When institutionalized children are placed in families, marked improvements in physical, social, and cognitive functioning is typically observed, yet many of the children maintain persistent behavioral problems (Ames, 1997;

Hodges & Tizard, 1989; Rutter, 1998; Verhulst et al., 1990; Verhulst et al., 1992). The developmental difficulties experienced by many of these children raise questions about the effects of early deprivation including factors such as failure to provide adequate nutrition, medical care, stimulation, and the lack of consistent and supportive caregiving relationships. Although early research emphasized the significance of maternal deprivation, Rutter (1981) rightly noted that many other types of stimulation needed for normal development are also deficient in these environments.

The critical questions that emerge from the plight of these children concern which aspects of inadequate stimulation result in cascading developmental effects, which developmental processes are most affected by inadequate early care, and specification about how the transfer to more normative caregiving can foster growth and recovery following institutionalization. Institutionalized children have experienced highly species atypical deprivation; in many countries the institutional conditions are quite poor: children may be confined to cots, fed gruel through propped up bottles, lack toys or stimulation, and receive very little linguistic stimulation and/or one-to-one interaction with caregivers (Nelson, 2007; Rutter et al., 1998). Even in institutions where basic physical needs were met, lack of individualized care and attention remain prominent. At adoption, children generally move into middle- to upper-middle class families, who are generally highly stable and well educated (Hellerstedt et al., 2007). In short, adoption marks a dramatic termination of deprivation, allowing an examination of the impact of early deprivation/neglect on subsequent development. Adoption into a supportive home can provide a profound natural intervention in the life of a child exposed to significant early adversity (van Ijzendoorn et al., 2005). As a result, internationally-adopted children provide a natural experiment on the impact of different degrees and duration of care on subsequent biobehavioral development. Despite occasional significant adjustment problems of the children, there are very few adoption disruptions for families who adopt internationally (e.g., Brumble, 2007). Furthermore, all studies of children adopted or fostered from institutions have shown that, varying with duration of institutionalization, once out of the institution children begin to show remarkable rebounds in physical and cognitive development (Kreppner et al., 2007; Maclean, 2003). Both the capacity of children to rebound after early institutional care and limitations on recovery imposed by longer periods of institutionalization was recently demonstrated experimentally in a study involving random assignment to high quality foster care for children who began their lives in Romanian institutions (Nelson et al., 2007). Studying post-institutionalized children several years after adoption allows examination of long-term impacts of early experience on children's development.

The present study is specifically motivated by the convergence of behavioral studies of post-institutionalized children and experimental studies of deprivation on the nonhuman primates (see review by Sanchez & Pollak, 2008). In terms of domains where children do not appear to catch-up in the immediate years following adoption, Beckett et al. (2007) reported that children adopted after prolonged period of institutional care from Romania had significantly lower scholastic attainment scores than those adopted early (before 6 months) either within the United Kingdom (UK) or from Romania. A separate research team also concluded that although adopted children showed age-expected development in some domains, adoptions after 12 months of age were associated with problems in school achievement (van Ijzendoorn & Juffer, 2006). Another recent report indicated that children residing within Romanian orphanages had poorly developed language abilities (Windsor et al., 2007). One recent study focused specifically on language and cognitive outcomes at 6 and 11 years of age with a large sample of institutionally-reared Romanian children adopted into UK families. This study noted few negative effects of deprivation if institutionalization ended before the age of 6 months. Even for the children over 18 months, the presence of even very minimal language skills (the child's ability to imitate speech sounds) at the time of arrival

was a strong positive prognostic factor for school-aged cognitive outcomes. Importantly, variations in adoptive parent characteristics were unrelated to differences in the adopted children's cognitive outcomes (Croft, Beckett, et al., 2007). Studies of institutionally-reared children yield consistent evidence that early deprivation can have long-term consequences for cognitive functioning and school readiness. Yet, most extant research has employed gross measures of functioning such as checklists, questionnaires, school records, and global DQ and IQ tests. While such global measures provide some suggestion of which neural systems may have been affected, they are not specific enough to test hypotheses about the development of neural systems. Our goal was to more directly examine brain-behavior processes in systems believed to underlie children's scholastic performance, such as aspects of attention, inhibitory control, working memory, and learning.

These domains are of interest not only because of the persistent difficulties observed in previously neglected children, but because nonhuman primate studies suggest that the prefrontal cortex (PFC) and its associated systems are particularly relevant to understanding cognitive functions and may be especially vulnerable to early experience (e.g. Sanchez et al., 2003). The PFC develops over a protracted period, well into adolescence (Huttenlocker, 1990; Sowell et al., 1999). While the timing of development in different regions has not been fully determined, there is evidence that neuronal organization in the dorsolateral prefrontal cortex (Brodman areas 9 and 10) changes dramatically until 5–7 years of age (Blinkov & Glezer, 1968), and slows thereafter. A similar protracted period has been observed for the neurotransmitter systems mediating prefrontal activity. Bourgeois (2001) has plotted the overproduction and pruning of synapses in the rhesus monkey against that of the human, taking into consideration differences in age compression (i.e., the rate at which each species develops), and has reported remarkable similarity across species. Collectively, these data, coupled with the postmortem data, point to a protracted period of PFC development. Presumably this may allow experience-dependent fine-tuning of attention, learning, emotion, and memory systems (e.g., Black, Jones, Nelson & Greenough, 1998).

Although it is critical to understand whether and how postnatal deprivation/neglect influences brain development in human children, the human neuroscience evidence is sparse. Two studies conducted by Chugani and colleagues reported that post-institutionalized (PI) children from Romania showed significantly decreased metabolism bilaterally in the orbital frontal gyrus, the infralimbic prefrontal cortex, the medial temporal structures (amygdala and head of hippocampus), the lateral temporal cortex, and the brain stem as compared to normal adults and children with chronic epilepsy. Diffusion tensor imaging also suggested that the PI children had reduced white matter tracts between the anterior temporal and frontal lobes (Chugani, Behen, Muzik, Juhász, Nagy, & Chugani 2001; Eluvathingal et al., 2006). These brain imaging findings are consistent with recent work from the Bucharest Early Intervention Project (BEIP; see Zeanah et al., 2003). For example, this group has reported that children reared in institutions show dramatic reductions in brain activity as revealed by the electroencephalogram and the event-related potential (see Marshall et al., 2004; Parker et al., 2005a,b). Importantly, children placed in high-quality foster care before the age of two years show improvement in their EEG (Marshall et al., in press); ERP amplitude also improves with placement in foster care, although it is not time- or sensitive-period dependent (Moulson et al., in press).

The present study sought to determine which domains of cognitive development may be particularly affected by institutional neglect/deprivation using children adopted internationally from orphanages or other institutions. These post-institutionalized children were (a) adopted over age 12 months, (b) spent at least 75% of their lives prior to adoption in institutional care, and (c) resided in their adoptive families for a minimum of 3 years at the time of testing. We compared these children to two different groups. Our first comparison

group consisted of children who were adopted before 8 months predominantly from foster care overseas, having little or no institutional care history. This group helps to control for prenatal factors and heritable factors associated with a child becoming orphaned or abandoned. Our second comparison group consists of non-adopted children who were reared in their families of origin. As noted by Rutter et al. (1997), even when using "normed" neuropsychological evaluations, and especially when using non-standardized tests, it is critical to include non-adopted children who have grown up in families of similar economic and educational histories to the families of adopted children. Furthermore, for domains that yielded differences between the post-institutionalized children and both other groups, we examined whether duration of institutional care correlated with the children's performance. Thus, the strategy employed here is to identify as "institutional deprivation effects" those cognitive functions for which the post-institutionalized group differs from the two comparison groups and, further, where the duration of institutional care is associated with the outcome within the post-institutionalized group. Here, we focus on basic cognitive/learning processes that might underlie school-based learning problems.

Method

Participants

The participants were 132 children ages 8 years, 0 months to 9 years, 11 months. Three groups were examined (see Table 1 for details). A post-institutionalized (PI) group whose selection criteria was adopted 12 months of age or older (range 12 to 78 mos, $M = 23.4$ mos, $Sd = 12.9$) having spent at least 75% of their preadoption lives in institutional care. Over 50% of this group had no experience other than institutional care prior to adoption ($M = 22.1$ mos in institutional care, $Sd = 12.4$). Children in the PI group were adopted from Asia ($n = 19$), Latin America ($n = 1$), Russia/Eastern Europe ($n = 27$) and Africa ($n = 1$). An early adopted predominantly from foster care (EA) group whose selection criteria was adoption at 8 months or earlier (range 2 to 8 mos, $M = 5.2$ mos, $Sd = 1.7$ mos) having spent 2 months or less in institutional care. Nearly 83% of this group had spent all of their preadoption lives in foster care overseas. Those with institutional experience had been adopted before 3 months of age. Children in the EA group were adopted from Asia (67%) and Latin America (33%). The EA group permitted comparison with children who had experienced loss of birth families and adoption into another culture. Finally, the third group was children born and raised in their birth families in the United States (non-adopted, NA). The children were recruited and tested at two sites, the University of Wisconsin ($n = 58$) and the University of Minnesota ($n = 74$). As shown in Table 1, there were no differences between the three groups in terms of the numbers of boys versus girls, child's age, or parent education. As expected, children in the PI group had lower IQ scores than those children in the two (EA and NA) comparison groups, $F(2, 131) = 16.13$, $p < .001$. On average, adoptive parents had higher family incomes than those from control families, $F(2, 131) = 3.99$, $p = .02$.

Recruitment and Screening

The internationally-adopted children were drawn from the Minnesota and the Wisconsin International Adoption Project Registries—registries of families created through international adoption that expressed interest in being contacted about research participation. The non-adopted children were recruited in Wisconsin through fliers and advertisements and in Minnesota from the Institute of Child Development Participant Pool, a registry of children whose parents indicated interest in being contacted about research opportunities in response to a mailing soon after their children were born.

Within 6 months of the present study, the children were all screened to determine if they met the study's group assignment (described above) and exclusion criteria. Criteria for group

assignment were determined through phone interview and parent questionnaire. Children then participated in an extensive developmental profile, parts of which determined exclusion criteria. Children were excluded if their IQs were below the normal range (<78), parents reported congenital abnormalities (e.g., Down/Cerebral Palsy), and failure on the Fetal Alcohol Syndrome (FAS) screener. We screened for IQ using the Vocabulary and Block Design subtests of the Wechsler Intelligence Scale for Children, 3rd ed. (WISC-III; Wechsler, 1998). If children scored below one standard deviation on either WISC-III subtests they were subsequently administered the Leiter International Performance Scale, Revised (Roid & Miller, 1997), a non-verbal IQ assessment, to avoid excluding children with normal IQs whose performance might have been affected by the English language demands of the WISC-III. Children were required to pass either the WISC or the Leiter to be included in this study, if they did not reach criterion on both the WISC and Leiter, then they were excluded. Eight children were assessed with the Leiter and of those, 4 (all PI) were excluded because of IQ's in the study exclusion range.

To screen for possible FAS, we photographed front and side views of children's faces and analyzed these images using FAS Facial Photographic Analysis Software (Astley, 2003). Children scoring in the moderate range or higher on this screener were excluded and images where coders were uncertain were also reviewed by a pediatric dysmorphologist. Reliability among coders was 100%. Parents of children who failed the FAS screener were contacted and referred for further evaluation. Four children (2 PI, 2 EA) were excluded because of possible Fetal Alcohol Syndrome. Note that this exclusion criterion only eliminated children with significant facial dysmorphology due to prenatal alcohol exposure. Children with exposures that did not affect facial morphology would not be identified by this method.

In addition to these assessments, children were also administered the Paragraph Comprehension subtest from the Comprehensive Assessment of Spoken Language battery (CASL). The tasks employed in this study were selected because they required only rudimentary language comprehension. Nonetheless, children scoring two standard deviations below the published norms of this tests were excluded to reduce the possibility that language problems might impair their performance. None of the PI or EA children needed to be excluded because of low CASL scores.

Four PI (8%), 2 EA (3%) and 1 NA (2%) were on stimulant medications for attention problems; these children were tested when parents agreed to withhold medication (e.g. weekends). Across all of the above-mentioned criteria, 1 NA, 4 EA, and 7 PI children were screened but not included in our final sample of 132.

For descriptive purposes, because low birth weight is associated with poorer cognitive functioning, as part of the screening examination parents provided information about the children's birth weight. Parents were asked to check baby books and other (adoption records) information for this information. Parents who were not able to find these records were asked to sign a medical release so that the child's pediatrician could be contacted for this information. Birth weight data were obtained from 97% of the of the NA, 85% of the EA and 41% of the PI children. Notably, in cases in which the PI children were abandoned (typical for Chinese post-institutionalized children, for example) this information was unknown. Analysis of the birth weight data yielded means in grams for the PI, EA and NA children, respectively, of 2,893 ($Sd=1,134$), 2,982 ($Sd=524$), and 3,314 ($Sd=699$), $F(2,94)=5.9$, $p<.01$. Hockbergen post-hoc tests indicated that both PI and EA groups were lower birth weight, on average, than NA children, although PI and EA birth weight did not differ. Because of the large number of EA children from Asia, we also compared EA and PI children only from that area of the world. The results failed to yield any difference between Asian EA and PI children, $t<.01$, df for unequal n 's=5.3, ns .

Procedure

Neuropsychological functioning was assessed through two well-validated test batteries. One of these, the Cambridge Neuropsychological Test Automated Battery (CANTAB; Cambridge Cognition, Cambridge, UK), is a computerized series of neuropsychological tests that cover a wide range of cognitive domains. The second battery is the NEPSY Developmental Neuropsychological Assessment (Korkman, Kirk, & Kemp, 1998), a battery developed specifically for use with children. The CANTAB offers several strengths: it has been used extensively with children (Luciana & Nelson, 1988, 2001; 2002); it has proved sensitive in discriminating various clinical populations from typically developing children; the individual tasks have been studied extensively to confirm their neural correlates (Joyce & Robbins, 1991; Owen, Downes, Sahakian, Polkey, & Robbins, 1990; Owen, Morris, Sahakian, Polkey, & Robbins, 1996; Owen et al., 1997; Owen, Roberts, Polkey, Sahakian, & Robbins, 1991; Owen, Sahakian, Semple, Polkey, & Robbins, 1995; ;Saghal et al., 1991; Sahakian et al., 1988); it is computerized for standardized administration; the stimuli cannot be verbalized and the subtasks require non-verbal responses, thus performance is not confounded with subjects' verbal skills.

Neuropsychological test—Memory functions were evaluated with three different tests. The first, Spatial Working Memory, tests the child's ability to retain spatial information and to manipulate remembered items in working memory. It is a self-ordered task, which also assesses heuristic strategy. The test begins with a number of colored squares displayed on a computer monitor. By touching the boxes and using a process of elimination, the child should find one blue 'token' in each of a number of boxes and use them to fill up an empty column on the right hand side of the screen. The number of boxes is gradually increased, until it is necessary to search a total of eight boxes. The child's score is based upon errors (touching boxes that have already been found to be empty, and revisiting boxes which have already been found to contain a token) as well as response latency, which is age-adjusted. Memory for Faces assesses the ability to recognize faces after a single exposure. Children are presented 16 pictures of children one at a time for 5 seconds each, during which children are instructed to remember the faces and identify the sex of each child. Next the children are presented with pictures of three children, one of which they previously saw. They are asked to identify the picture they saw previously. Thirty minutes later, this procedure is repeated. The score on this test is based on the sum of correct responses in immediate and delay tasks. Next, we administered a Narrative Memory test that assesses the ability to retell a story under both free and cued recall conditions. A story is told to the child. Following the story, children are asked to tell the examiner the same story. After the child has recalled as much as they can, the examiner cues details the child did not mention.

Learning processes were evaluated using the Paired Associates Learning test, which assesses visual memory and new learning. Boxes are displayed on the screen and are opened in a randomized order. One or more of them will contain a pattern. The patterns are then displayed in the middle of the screen, one at a time, and the subject must touch the box where the pattern was originally located. If the subject makes an error, the patterns are re-presented to remind the subject of their locations. The difficulty level increases through the test.

Attentional processes were evaluated using five different tasks. The Auditory Attention and Response Set test assesses children's ability to maintain selective auditory attention, as well as the child's ability to maintain a complex mental set. In the first part of the task, the child learns to manipulate a red item whenever they hear the word red. In the second part of the task, the child then must shift set and respond to red stimuli when hearing the word yellow. Match to Sample Visual Search is a matching test, with a speed/accuracy trade-off. Children

are shown a complex visual pattern in the middle of a computer screen, and then, after a brief delay, a varying number of similar patterns are shown in a circle of boxes around the edge of the screen. Only one of these boxes matches the pattern in the center of the screen, and the subject must indicate which it is by touching it. Reaction time is measured on the basis of the release of the press-pad, which allows for its more accurate measurement. Rapid Visual Information Processing is a test of visual sustained attention. A white box appears in the center of the computer screen, inside which digits appear in a pseudo-random order, at the rate of 100 digits per minute. Children are requested to detect target sequences of digits and to register responses using a press pad. Outcome measures are calculated using Signal Detection Theory. The Visual Attention test assesses the speed and accuracy with which a child can scan an array and locate targets. Finally, the Knock and Tap test measures the child's ability to inhibit immediate impulses evoked by visual stimuli that conflict with a verbal direction. The child learns a pattern of motor responses and then must maintain that cognitive set and inhibit the impulse to imitate the examiner's action.

The working memory and Knock and Tap tasks also involve components of executive functions. We examined two additional aspects of executive function (rule acquisition and manipulation through reversal and planning, here labeled "executive processes". The Intra-Extra Dimensional Shift Set (ID/ED) Test was used to measure rule acquisition and manipulation through reversal. Two artificial dimensions are used in the test: color-filled shapes and white lines. Simple stimuli are made up of just one of these dimensions, whereas compound stimuli are made up of both, namely white lines overlying color-filled shapes. Children begin by seeing two simple color-filled shapes, and must learn which one is correct by touching it. Feedback teaches the child which stimulus is correct, and after six correct responses, the stimuli and/or rules are changed. These shifts are initially intra-dimensional (e.g. color filled shapes remain the only relevant dimension), then later extra-dimensional (white lines become the only relevant dimension). Children progress through the test by satisfying a set criterion of learning at each stage. In addition, Stockings of Cambridge is a version of the Tower of London spatial planning test, which gives a measure of frontal lobe function (Baker et al., 1996; Owen et al., 1990; Robbins, 1996. Children are shown two displays containing three colored balls. The displays are presented in such a way that they can easily be perceived as stacks of colored balls held in stockings or socks suspended from a beam. The child must use the balls in the lower display to copy the pattern shown in the upper display. The balls may be moved one at a time by touching the required ball, then touching the position to which it should be moved. The time taken to complete the pattern and the number of moves required are taken as measures of the child's planning ability.

Results

Our first set of analyses sought to determine which domains of functioning were either minimally affected by early deprivation or may reflect developmental catch-up following family rearing. Although we could not distinguish between these explanations, we could determine tasks in which our three groups of children performed differently. Omnibus tests revealed group differences in memory functioning, $F_{\text{mult}}(6, 252) = 3.31, p = .004$, on tests of attentional functioning, $F_{\text{mult}}(10, 224) = 2.72, p = .004$, and on the test of learning, $F(2, 131) = 12.47, p = .001$. However, all three groups of children performed similarly on tests of executive processes, $F_{\text{mult}}(4, 254) = 1.20, \text{ns}$.

Next, we explored those domains (memory, attention, learning) where multivariate tests suggested group differences to determine which specific tasks discriminated between groups of children. Table 2 shows the descriptive information for these tasks, the pattern of post-hoc comparison between groups and the correlations of scores with duration of institutional care. We included sex in our analyses, but in no instance did we find an interaction of sex

and group. Where main effects of sex were noted, girls performed better than boys; however, as sex differences did not modify the main effects of group, we do not discuss them further. Information on sex differences is available upon request.

Within the memory domain, the groups performed differently on the spatial working memory test, $F(2, 128) = 7.96, p = .001$. Hochberg post-hoc tests revealed that PI children performed more poorly than both EA ($p = .008$) and controls ($p = .001$), but that EA children performed similarly to controls ($p = ns$). The Memory for Faces test resulted in a marginal group difference, $F(2, 128) = 2.95, p = .056$. On this test, PI children scored lower than NA children ($p = .057$), but the EA and NA groups performed similarly. The groups did not differ on auditory memory ability.

In terms of tests of learning, children's performance on the Paired Associates Learning test differed, $F(2, 131) = 12.47, p = .001$. PI children performed scored lower than EA ($p = .003$) and NA ($p < .001$), but EA and NA groups performed similarly ($p = ns$).

Within the attention domain, groups performed similarly on tests of auditory attention $F(2, 116) = 2.30, ns$, Match to Sample $F(2, 116) = 1.75, ns$, and Rapid Visual Processing $F(2, 116) = 1.03, ns$. However, the groups differed on the test of visual attention, $F(2, 116) = 8.96, p = .001$. Here, PI children performed more poorly than EA ($p = .001$) and controls ($p = .006$), but EA children performed similarly to controls ($p = ns$). There was also a marginal difference on the Knock and Tap test, $F(2, 116) = 2.92, p = .06$. Here, PI children showed more impulsivity than controls ($p = .05$), but there were no other differences between groups.

Discussion

The primary goal of this study was to determine which cognitive processes differentiated children who experienced early institutional deprivation and neglect. Post-institutionalized internationally adopted children differed from both groups of comparison children on tests of visual memory and attention, as well as visually-mediated learning and inhibitory control. Yet, these children performed at developmentally appropriate levels on similar tests where auditory processing was also involved. These children also performed well on tests of executive processes involving rule acquisition and manipulation and planning.

This study was not designed to show precisely which neural circuits are affected by early experience, but the tasks that PI children had difficulty with do provide a useful guide for specifying neuroanatomical substrates that can be more directly examined in future studies. Below, we briefly highlight areas of brain circuitry associated with these tasks. Performance on the Memory for Faces task reflects an interaction between prefrontal cortex (PFC) and stimulus-specific visual cortical association areas that mediate visual working memory. The neurophysiological models of visual working memory developed in the nonhuman primate also holds for humans, based upon event-related fMRI (ER-fMRI) studies of facial delay-recognition task similar to the one used in the present study (Druzga & D'Esposito, 2003). Memory tasks such as the one used in the present study involve both visual memory and visual perception. But there is not yet agreement in the field about the neural bases of these processes. Some data suggest that visual memory and visual perception are associated with common neural substrates (Slotnick, 2004), whereas other data suggest that right medial temporal-lobe structures are critically involved in retention, but not in the perception, of new faces (Crane & Milner, 2002).

Spatial Working Memory is a test of the child's ability to retain spatial information and to manipulate remembered items in working memory and also assesses heuristic strategy. The task is considered a sensitive measure of frontal lobe dysfunction based upon studies of

nonhuman primates and patients with frontal lobe damage (Owen et al., 1990; Owen et al., 1995; Owen, Morris, et al., 1996; Passingham, 1985). Owen, Evans, & Petrides (1996) used positron emission tomography with magnetic resonance imaging to demonstrate the existence, within the human brain, of two functionally distinct subdivisions of the lateral frontal cortex, which subserve different aspects of spatial working memory. When the spatial working memory task required the organization and execution of a sequence of spatial moves retained in working memory, significant changes in blood flow were observed in ventrolateral frontal cortex (area 47) bilaterally. By contrast, when the task required active monitoring and manipulation of spatial information within working memory, additional activation foci were observed in mid-dorsolateral frontal cortex (areas 46 and 9). Both of these processing stages were required to successfully complete a spatial working memory task (Owen, Evans, et al., 1996). Another study also linked poor performance on the CANTAB Spatial Working Memory task in children to reduced dorsolateral prefrontal cortex using fMRI (Luna et al., 2002).

It is difficult to link performance on tests of Visual Attention to distinct neural systems. A recent study used ER-fMRI to measure brain activity as subjects oriented visual attention to an upcoming target. This activation was lateralized to the left hemisphere and reflected a widely distributed network that included (a) structures in parietal and temporal cortices and thalamus usually associated with selective attention, (b) ventral-stream object processing structures in occipital, inferior-temporal, and parahippocampal cortex, and (c) structures in medial- and dorsolateral-prefrontal cortex associated with cognitive control (Arrington et al., 2000). In visual attention tasks, subjects will undoubtedly make errors and need to reorient to targets. This same study revealed that brain areas specific to attentional reorientation were right-lateralized and included posterior temporal and inferior parietal regions, as well as prefrontal regions that likely subserve control processes related to inhibition of inappropriate responding (Arrington et al., 2000). In sum, selective visual attention results from distributed brain activity that includes maintenance of the selected object's representation accompanied by suppression of response to ignored objects (Duncan, 1993; Farah, 1990; Phafet et al., 1990), with control of these processes mediated by the medial dorsolateral prefrontal cortex (DL PFC).

Paired Associates Learning is a stringent test for visual episodic memory and associative learning (Sahakian & Owen, 1992). To perform well, children had to learn the locations of a progressively increasing number of abstract stimuli. Although such a complex behavioral task undoubtedly draws upon widely distributed neural systems, ER-fMRI suggests that successful performance on this type of task relies heavily upon medial temporal lobe connections to the frontal lobe (Aizenstein et al., 2000). Similarly, the Knock and Tap test cannot be mapped onto discrete circuitry. The test measures the child's ability to inhibit immediate impulses evoked by visual stimuli that conflict with a verbal direction. The child learns a pattern of motor responses and then must maintain that cognitive set and inhibit the impulse to imitate the examiner's action.

Taken together, the present data suggest delayed maturation of select aspects of frontal circuitry, and perhaps reduced functional connectivity of frontal cortex with other neocortical and subcortical regions, plays a key role in scholastic difficulties among children who experienced early institutionalized deprivation and neglect. Indeed, regions of the PFC have long been associated with cognitive processes similar to the ones assessed in this study (Fletcher & Henson, 2001; Owen et al., 1996). Studies in nonhuman primates, lesion studies, as well as functional neuroimaging studies in humans, have documented that DLPFC is crucial for maintaining and manipulating information in ways assessed by the tasks used here (Carlson et al., 1998; Fuster, 2000; Goldman-Rakic, 1988; McCarthy et al., 1994; Owen, Evans, & Petrides, 1996; Pierrot-Deseilligny et al., 1995; Sweeney et al., 1996).

The distinction between the performance of PI and EA children suggests (but does not prove) that the ontogenesis of the prefrontal cortex also includes an extensive postnatal interval. Such an interpretation is consistent with animal studies which reveal, for example, that the thickness of the prefrontal cortex of the rat brain is not maximal until post natal day 20, which marks the beginning of the postweaning period (e.g. roughly equivalent to adolescence in humans; Vincent et al., 1995). Indeed, PI children are often noted to have problems in attention regulation and emotional control, functions presumably influenced by the development of the PFC and its distributed systems (see review by Gunnar, 2001; Shallice et al., 2002). Consistent with this view, Sanchez et al. (1998) studied rhesus monkeys that were socially deprived between 2 and 12 months of age. These monkeys exhibited cognitive deficits that had also been noted in earlier studies (e.g., Harlow et al., 1971). Sanchez's MRI studies revealed that the animals' performance on executive function tasks was correlated with decreased neuronal development of prefrontal, medial temporal, and amygdala substrates. Studied two years later, these monkeys exhibited increased abnormalities in the prefrontal cortex that was related to the monkeys' performance on learning and working memory tasks (Sanchez et al., 2003). In a separate study, Mathew et al. (2003) reported neuropathological alterations in the prefrontal cortex of adult macaques with early adverse experience.

There are a number of features of the present study that are important to consider in interpreting these data. First, as described earlier, we excluded children with clear developmental delays or indications of fetal alcohol exposure, which would have artificially exaggerated group performance deficits. By avoiding a situation where a subset of PI children may be experiencing global cognitive delays, we were also able to detect specific patterns of processing deficits. In general, executive function tasks are sensitive to perturbations in children's lives, but effects on these tasks are not specific to certain situations or conditions. If we observed broad or diffuse cognitive deficits in the PI sample, it would not be possible to rule out general attentional, IQ, motivational, motor, or emotional problems as driving children's performance. Yet, in the present study we observed that on a majority of tasks, PI children displayed performance equivalent to their peers, which permits stronger interpretation of those specific areas where deficits emerged. Second, this sample of children represents families who volunteered to participate in research; it may be the case that parents who believed their children were doing particularly well or poorly were more likely to participate. Yet, such sampling issues—if they exist—do not readily explain the pattern of results observed in this study. In addition, our unusual sampling procedures (see Hellerstedt et al., 2008) likely resulted in a group of children that are both diverse and representative.

The study of children reared in atypical situations requires opportunistic approaches that are also fraught with interpretive challenges. Our requirement that the EA group be adopted before 8 months of age and that these children had to spend less than 2 months in institutional care meant that none of the EA children were drawn from Eastern Europe, where adoption processes take more time. Most of the EA children spent most of their pre-adoptive lives in foster care overseas, an option that did not exist in Eastern Europe between 1990 and 1998, when the children that we tested were adopted. It is likely that prenatal conditions were less optimal for children who end up being placed in orphanage or institutional care overseas than for children born and raised in their families of origin in the United States. In part, the inclusion of a group of children adopted internationally with little or no institutional experience partly addressed this problem. We obtained birth weight data on the majority of both EA and NA children, and found, as might be expected, that the EA children were lower in birth weight than the NA children. Although we were only able to obtain birth weight data on 41% of the PI children, there was no evidence that birth weight differentiated PI from EA children. While we must be cautious in interpretation of these data

because we lack evidence on the reliability of the information and are missing substantial data for the PI children, the pattern of data suggest that EA and PI children tended to experience prenatal environments that were impoverished. This tends to clarify, but not resolve, the issue of whether pre versus postnatal environmental impacts influenced the outcomes we examined.

In terms of the tasks administered to children, it is critical that measurement involves a sufficient level of task difficulty relative to the ability level of participants. In this regard, the specific tasks used in this study are particularly useful. For example, the test of spatial working memory ability and paired associates learning systematically varies the working memory load, which increases the amount of information that needs to be remembered and the number of trials over which it needs to be maintained. The children's pattern of performance across the tasks that we administered did not reveal clear performance deficits on auditory memory and attentional tasks. One possibility is that the deficits observed here are a function of how much information needs to be kept active in working memory over time, and how precisely that information needs to be encoded. At the same time, firm conclusions about neural activity cannot be made based solely on behavioral data. However, these data do underscore the need to employ specific assessments that have been used in imaging and lesion studies and which may allow more fine-grained analysis of the impact of institutional neglect on neurobehavioral development.

Why did the PI children perform better on tasks that relied primarily on auditory, as compared with visual, information? One possibility is that visual development is more vulnerable to post-natal influences. The auditory system starts functioning during the last trimester of gestation (Birnholtz & Benaceraff, 1983), whereas the visual system does not start functioning until after birth. Thus, both visual and auditory *functional* development reflect experience-dependent processes; the difference is that auditory experience starts before visual experience, when the brain is at a different point in its development (Cf., Saffran, Werker, & Werner, 2006; Kellman & Arterberry, 2006). Sloutsky and colleagues (Robinson & Sloutsky, 2004; Sloutsky & Napolitano, 2003) have demonstrated that young children, unlike adults, exhibit auditory dominance. This may reflect earlier maturation of the auditory system relative to the visual system. It may also be the case that attentional processes more easily engage some stimulus properties. For example, auditory stimuli are serial, transient events that must be perceived quickly whereas visual stimuli are presented episodically for longer periods of time. Thus, attentional systems may allocate more resources to transient relative to stable stimuli. Similarly, Posner, Nissen, and Klein (1976) proposed that attention to visual stimuli must be learned, whereas attention to auditory stimuli is more automatic. Also consistent with the argument that post-natal experience is more likely to influence visual processing, Maurer and colleagues (1999) have demonstrated that the neural circuitry responsible for adults' face expertise is not pre-specified, but requires early visual experience (see also Nelson, 2001; Pascalis, de Haan, & Nelson, 2005). Because infants have poor visual acuity, their cortices are only exposed to low spatial frequency. Thus, early exposure to visual information sets up the neural architecture for more complex visual processing. These studies have demonstrated that when visual input is delayed by as little as two months, permanent visual deficits result (Le Grand et al., 1999; Le Grand, Mondloch, Maurer, and Brent, 2003; Maurer et al., 1999).

Models of the role of experience in neural development, and the mounting information on molecular processes in neural plasticity, indicate that neural activities (i.e., activity-dependent processes) are critical to brain development (see Fox, Levitt, & Nelson, 2008). This implies that in addition to the stimuli available in the environment, active engagement of the environment may be essential in order for some aspects of cognitive development to occur. While these opportunities abound in typical human rearing environments,

institutionalized child rearing may restrict the kinds of dynamic experiences and input necessary for some aspects of neurobehavioral development. It is hoped that the more we understand about the specific aspects of neurodevelopment impacted by early deprivation, the more we can focus on identifying targeted intervention and training experiences that would optimize these children's outcomes.

Acknowledgments

This project was supported by grants from the National Institute of Mental Health to Seth Pollak (R01 MH068858) and Megan Gunnar (R01 MH068857). Support for C.A. Nelson comes from the NIH (NS034458; MH078829) and the Richard David Scott endowment. Infrastructure support was provided by the Waisman Center, University of Wisconsin through the National Institute of Child Health and Human Development (P30-HD03352, M. Seltzer, Director). We acknowledge the research assistance and support provided by Meg Bale. We also greatly appreciate the children and their families whose participation made this research possible. Please direct correspondence to Seth Pollak (spollak@wisc.edu) or Megan Gunnar (gunnar@umn.edu).

References

- Aizenstein HJ, MacDonald AW, Stenger VA, Nebes RD, Larson JK, Ursu S, et al. Complementary category learning systems identified using event-related functional MRI. *Journal of Cognitive Neuroscience*. 2000; 12(6):977–987. [PubMed: 11177418]
- Ames, EW. The Development of Romanian Orphanage Children Adopted to Canada (Final report to the national Welfare Grants Program: Human Resources Development Canada). Burnaby, British Columbia: Simon Fraser University; 1997.
- Arrington CM, Carr TH, Mayer AR, Rao SM. Neural mechanisms of visual attention: Object-based selection of a region in space. *Journal of Cognitive Neuroscience*. 2000; 12(2):106–117. [PubMed: 11506651]
- Astley, S. FAS Facial Photographic Analysis Software. Seattle, WA: FAS Diagnostic & Prevention Network University of Washington; 2003.
- Baker SC, Rogers RD, Owen AM, Frith CD, Dolan RJ, Frackowiak RSJ, Robbins TW. Neural systems engaged by planning: A PET study of the Tower of London task. *Neuropsychologia*. 1996; 34:515–526. [PubMed: 8736565]
- Beckett C, Maughan B, Rutter M, Castle J, Colvert E, Groothues C, et al. Scholastic attainment following severe early institutional deprivation: A study of children adopted from Romania. *Journal of Abnormal Child Psychology*. 2007; 35(6):1063–1073. [PubMed: 17643189]
- Black, JE.; Jones, TA.; Nelson, CA.; Greenough, WT. Neuronal plasticity and the developing brain. In: Alessi, NE.; Coyle, JT.; Harrison, SI.; Eth, S., editors. *Handbook of Child and Adolescent Psychiatric Science and Treatment*. New York: Wiley and Sons; 1998. p. 31–53.
- Blinkov, SM.; Glezer, II. *A Quantitative Handbook*. New York: Plenum Press; 1968. The Human Brain in Figures and Tables.
- Brumble K. Intercountry transracial special needs adoptees: Today's teenagers and young adults. How have they fared? The parent's perspective. *Dissertation Abstracts International Section A: Humanities and Social Sciences*. 2007; 68(4-A):1404.
- Bourgeois JA. Visual hallucinations and psychotic symptoms during treatment with selective serotonin reuptake inhibitors: Is the sigma receptor involved? *Journal of Clinical Psychopharmacology*. 2000; 20:580. [PubMed: 11001245]
- Carlson S, Martinkauppi S, Rama P, Salli E, Korvenoja A, Aronen HJ. Distribution of cortical activation during visuospatial n-back tasks as revealed by functional magnetic resonance imaging. *Cerebral Cortex* (New York, N.Y.: 1991). 1998; 8(8):743–752.
- Chugani HT, Behen ME, Muzik O, Juhasz C, Nagy F, Chugani DC. Local brain functional activity following early deprivation: A study of post-institutionalized Romanian orphans. *Neuroimage*. 2001; 14:1290–1301. [PubMed: 11707085]
- Crane J, Milner B. Do I know you? Face perception and memory in patients with selective amygdalo-hippocampectomy. *Neuropsychologia*. 2002; 40:530–538. [PubMed: 11749983]

- Croft C, Beckett C, Rutter M, Castle J, Colvert E, Groothues C, et al. Early adolescent outcomes of institutionally-deprived and non-deprived adoptees. II: Language as a protective factor and a vulnerable outcome. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*. 2007; 48(1):31–44.
- Downes JJ, Roberts AC, Sahakian BJ, Evenden JL, Morris RG, Robbins TW. *Neuropsychologia*. 1989; 27(11/12):1329–1343. [PubMed: 2615934]
- Druzgal TJ, D'Esposito M. Dissecting contributions of prefrontal cortex and fusiform face area to face working memory. *Journal of Cognitive Neuroscience*. 2003; 15:771–784. [PubMed: 14511531]
- Duncan J. Coordination of what and where in visual attention. *Perception*. 1993; 22(11):1261–1270. [PubMed: 8047413]
- Goldman-Rakic PS. Topography of cognition: Parallel distributed networks in primate association cortex. *Annual Review of Neuroscience*. 1988; 11:137–156.
- Farah, MJ. *Visual Agnosia: Disorders of Object Recognition and What They Tell Us About Normal Vision*. Cambridge, MA: MIT Press; 1990.
- Fletcher PC, Henson RN. Frontal lobes and human memory: Insights from functional neuroimaging. *Brain: A Journal of Neurology*. 2001; 124(Pt 5):849–881. [PubMed: 11335690]
- Fox SE, Levitt P, Nelson CA. How the Timing and Quality of Early Experiences Influence the Development of Brain Architecture. 2008 Manuscript under review.
- Fuster JM. Prefrontal neurons in networks of executive memory. *Brain Research Bulletin*. 2000; 52(5): 331–336. [PubMed: 10922510]
- Gunnar, M. Effects of early deprivation: Findings from orphanage-reared children. In: Nelson, CA.; Luciana, M., editors. *Handbook of Cognitive Neuroscience*. Cambridge: MIT Press; 2001. p. 617–629.
- Harlow HF, Harlow MK, Suomi SJ. From thought to therapy: Lessons from a primate laboratory. *American Scientist*. 1971; 59:538–549. [PubMed: 5004085]
- Hellerstedt WL, Madsen NJ, Gunnar M, Grotevant HD, Lee RM, Johnson DE. The international adoption project: Population-based surveillance of Minnesota parents who adopted children internationally. *Maternal and Child Health Journal*. 2008; 12:162–171. [PubMed: 17562147]
- Hodges J, Tizard B. IQ and behavioural adjustment of ex-institutional adolescents. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*. 1989; 30(1):53–75.
- Huttenlocker PR. Morphometric study of human cerebral cortex development. *Neuropsychologia*. 1990; 28:517–527. [PubMed: 2203993]
- Johnson, DE. The impact of orphanage rearing on growth and development. In: Nelson, CA., editor. *The effects of adversity on neurobehavioral development: Minnesota symposia on child psychology*. Vol. 31. New York: Erlbaum and Associates; 2001.
- Joyce EM, Robbins TW. Frontal lobe function in Korsakoff and non-Korsakoff alcoholics: Planning and spatial working memory. *Neuropsychologia*. 1991; 29:709–723. [PubMed: 1944873]
- Korkman, M.; Kirk, U.; Kemp, S. *NEPSY: A Developmental Neuropsychological Assessment Manual*. San Antonio, TX: Psychological Corporation; 1998.
- Kreppner JM, Rutter M, Beckett C, Castle J, Colvert E, Groothues C, et al. Normality and impairment following profound early institutional deprivation: A longitudinal follow-up into early adolescence. *Developmental Psychology*. 2007; 43(4):931–946. [PubMed: 17605526]
- Le Grand R, Mondloch CJ, Mauer D, Brent HP. Early visual experience and face processing. *Vision Research*. 1999; 39:3480. [PubMed: 10615511]
- Le Grand R, Mondloch CJ, Maurer D, Brent HP. Expert face processing requires visual input to the right hemisphere during infancy. *Nature Neuroscience*. 2003; 6(10):1108–1112.
- Luciana M, Nelson CA. The functional emergence of prefrontally -guided working memory systems in four to eight-year-old children. *Neuropsychologia*. 1988; 36:273–293. [PubMed: 9622192]
- Luciana M, Nelson CA. Assessment of neuropsychological function through use of the Cambridge Neuropsychological Testing Automated Battery: Performance in 4- to 12-year-old children. *Developmental Neuropsychology*. 2002; 22:595–624. [PubMed: 12661972]

- Luciana M, Sullivan J, Nelson CA. Individual differences in phenylalanine levels moderate performance on tests of executive function in adolescents treated early and continuously for PKU. *Child Development*. 2001; 72:1637–1652. [PubMed: 11768137]
- Luna B, Minshew NJ, Garver K, Lazar NA, Thulborn KR, Eddy WF, et al. Neocortical system abnormalities in autism: An fMRI study of spatial working memory. *Neurology*. 2002; 59:834–840. [PubMed: 12297562]
- Maclean K. The impact of institutionalization on child development. *Developmental Psychopathology*. 2003; 15(4):853–884.
- Marshall PJ, Fox NA, the BEIP Core Group. A comparison of the electroencephalogram between institutionalized and community children in Romania. *Journal of Cognitive Neuroscience*. 2004; 16:1327–1338. [PubMed: 15532128]
- Marshall P, Reeb BC, Fox NA, the BEIP Core Group. Effects of early intervention on EEG power and coherence in previously institutionalized children in Romania. *Development and Psychopathology*. in press.
- Mathew SJ, Shungu DC, Mao X, Smith EL, Perera GM, Kegeles LS, et al. A magnetic resonance spectroscopic imaging study of adult nonhuman primates exposed to early-life stressors. *Biological Psychiatry*. 2003; 54(7):727–735. [PubMed: 14512213]
- Maurer D, Lewis TL, Brent HP, Levin AV. Rapid improvement in the acuity of infants after visual input. *Science (New York, N.Y.)*. 1999; 286(5437):108–110.
- McCarthy G, Blamire AM, Puce A, Nobre AC, Bloch G, Hyder F, et al. Functional magnetic resonance imaging of human prefrontal cortex activation during a spatial working memory task. *Proceedings of the National Academy of Sciences of the United States of America*. 1994; 91(18):8690–8694. [PubMed: 8078943]
- Moulson MC, Nelson CA. Early adverse experiences and the neurobiology of facial emotional processing. *Developmental Psychology*. (in press).
- Nelson CA. A neurobiological perspective on early human deprivation. *Child Development Perspectives*. 2007; 1:13–18.
- Nelson CA, Zeanah CH, Fox NA, Marshall PJ, Smyke A, Guthrie D. Cognitive recovery in socially deprived young children: The Bucharest Early Intervention Project. *Science*. 2007; 318:1937–1940. [PubMed: 18096809]
- Owen AM, Downes JJ, Sahakian BJ, Polkey CE, Robbins TW. Planning and spatial working memory deficits following frontal lobe lesions in man. *Neuropsychologia*. 1990; 28:1021–1034. [PubMed: 2267054]
- Owen AM, Evans AC, Petrides M. Evidence for a two-stage model of spatial working memory processing within the lateral frontal cortex: A positron emission tomography study. *Cerebral Cortex*. 1996; 6(1):31–38. [PubMed: 8670636]
- Owen AM, Iddon JL, Summers BA, Robbins TW. Spatial and non-spatial working memory at different stages of Parkinson's disease. *Neuropsychologia*. 1997; 35:519–532. [PubMed: 9106280]
- Owen AM, Morris RG, Sahakian BJ, Polkey CE, Robbins TW. Double dissociations of memory and executive functions in a self-ordered working memory task following frontal lobe excision, temporal lobe excisions or amygdalohippocampectomy in man. *Brain*. 1996; 119:1597–1615. [PubMed: 8931583]
- Owen AM, Roberts AC, Polkey CE, Sahakian BJ, Robbins TW. Extra-dimensional versus intra-dimensional set-shifting performance following frontal lobe excisions, temporal lobe excisions, or amygdalo-hippocampectomy in man. *Neuropsychologia*. 1991; 29:993–1006. [PubMed: 1762678]
- Owen AM, Sahakian BJ, Semple J, Polkey CE, Robbins TW. Visuospatial short-term recognition memory and learning after temporal lobe excision, frontal lobe excision, or amygdalo-hippocampectomy in man. *Neuropsychologia*. 1995; 33:1–24. [PubMed: 7731533]
- Parker SW, Nelson CA, et al. The impact of early institutional rearing on the ability to discriminate facial expressions of emotion: an event-related potential study. *Child Development*. 2005a; 76:54–72. [PubMed: 15693757]
- Parker SW, Nelson CA, the BEIP Core Group. An event-related potential study of the impact of institutional rearing on face recognition. *Development and Psychopathology*. 2005b; 17:621–639. [PubMed: 16262985]

- Pascalis O, Scott LS, Kelly DJ, Dufour RW, Shannon RW, Nicholson E, Petit O, Coleman M, Nelson CA. Plasticity of Face Processing in Infancy. *Proceedings of the National Academy of Sciences*. 2005; 102:5297–5300.
- Phaf RH, Van der Heijden AH, Hudson PT. SLAM: A connectionist model for attention in visual selection tasks. *Cognitive Psychology*. 1990; 22(3):273–341. [PubMed: 2376113]
- Pierrot-Deseilligny C, Rivaud S, Gaymard B, Muri R, Vermersch AI. Cortical control of saccades. *Annals of Neurology*. 1995; 37(5):557–567. [PubMed: 7755349]
- Posner MI, Nissen MJ, Klein RM. Visual dominance: An information-processing account of its origins and significance. *Psychological Review*. 1976; 83(2):157–171. [PubMed: 769017]
- Robbins TW. Dissociating executive functions of the prefrontal cortex. *Philosophical Transactions: Biological Sciences*. 1996; 351:1463–1471. [PubMed: 8941958]
- Roberts AC, De Salvia MA, Wilkonsin LS, Collins P, Muir JL, Everitt BJ, Robbins TW. 6-hydroxydopamine lesions of the prefrontal cortex in monkeys enhance performance on an analog of the Wisconsin card sort test: Possible interactions with subcortical dopamine. *The Journal of Neuroscience*. 1994; 14(5):2531–2544. [PubMed: 8182426]
- Robinson CW, Sloutsky VM. Auditory dominance and its change in the course of development. *Child Development*. 2004; 75(5):1387–1401. [PubMed: 15369521]
- Roid, GH.; Miller, LJ. *Leiter International Performance Scale-Revised*. Wood Dale, IL: Stoelting Co.; 1997.
- Rutter M. Psychological sequelae of brain damage in children. *The American Journal of Psychiatry*. 1981; 138(12):1533–1544. [PubMed: 6795943]
- Rutter M, Dunn J, Plomin R, Simonoff E. Integrating nature and nurture: Implications of person-environment correlations and interactions for developmental psychopathology. *Development and Psychopathology*. 1997; 9:335–364. [PubMed: 9201448]
- Rutter M. Some research considerations on intergenerational continuities and discontinuities: Comment on the special section. *Developmental Psychology*. 1998; 34(6):1269–1273. [PubMed: 9823511]
- Sanchez MM, Hearn EF, Do D, Rilling JK, Herndon JG. Differential rearing affects corpus callosum size and cognitive function of rhesus monkeys. *Brain Research*. 1998; 812:38–49. [PubMed: 9813233]
- Sanchez MM, Ladd CO, Plotsky PM. Early adverse experience as a developmental risk factor for later psychopathology: Evidence from rodent and primate models. *Development and Psychopathology*. 2001; 13:419–450. [PubMed: 11523842]
- Sanchez, MM.; Pollak, SD. Socio-emotional development following early abuse and neglect: Challenges and insights from translational research. In: de Haan, M.; Gunnar, MR., editors. *Handbook of Developmental Social Neuroscience*. Guilford Press; (In press)
- Shallice T, Marzocchi GM, Coser S, Del Savio M, Meuter RF, Rumiati RI. Executive function profile of children with attention deficit hyperactivity disorder. *Developmental Neuropsychology*. 2002; 21:43–71. [PubMed: 12058835]
- Saghal A, Sahakian BJ, Robbins TW, Wray CJ, Lloyd S, Cook JH. Detection of visual memory and learning deficits in Alzheimer's disease using the Cambridge Neuropsychological Test Automated Battery. *Dementia*. 1991; 2:150–158.
- Sahakian BJ, Morris RG, Evendon JL, Heald A, Levy R, Philpot M. A comparative study of visuospatial memory and learning in Alzheimer-type dementia and Parkinson's disease. *Brain*. 1988; 111:695–718. [PubMed: 3382917]
- Sahakian BJ, Owen AM. Computerized assessment in neuropsychiatry using CANTAB: Discussion paper. *Journal of the Royal Society of Medicine*. 1992; 85(7):399–402. [PubMed: 1629849]
- Slotnick SD. Visual memory and visual perception recruit common neural substrates. *Behavioral and Cognitive Neuroscience Reviews*. 2004; 3(4):207–221. [PubMed: 15812107]
- Sloutsky VM, Napolitano AC. Is a picture worth a thousand words? Preference for auditory modality in young children. *Child Development*. 2003; 74(3):822–833. [PubMed: 12795392]
- Sowell ER, Thompson PM, Holmes CJ, Jernigan TL, Toga AW. In vivo evidence for post-adolescent brain maturation in frontal and striatal regions. *Nature Neuroscience*. 1999; 2:859–861.

- Sweeney JA, Mintun MA, Kwee S, Wiseman MB, Brown DL, Rosenberg DR, et al. Positron emission tomography study of voluntary saccadic eye movements and spatial working memory. *Journal of Neurophysiology*. 1996; 75(1):454–468. [PubMed: 8822570]
- VanIjzendoorn MH, Juffer F, Poelhuis CW. Adoption and cognitive development: A meta-analytic comparison of adopted and nonadopted children's IQ and school performance. *Psychological Bulletin*. 2005; 131(2):301–316. [PubMed: 15740423]
- van Ijzendoorn HW, Juffer F. The Emanuel Miller Memorial Lecture 2006: Adoption as intervention. Meta-analytic evidence for massive catch-up and plasticity in physical, socio-emotional, and cognitive development. *Journal of Child Psychology & Psychiatry*. 2006; 27:1228–1245. [PubMed: 17176378]
- Verhulst FC, Althaus M, Versluis-denBieman HJM. Problem behavior in international adoptees: III. diagnosis of child psychiatric disorders. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1990; 29(3):420–428. [PubMed: 2347840]
- Verhulst FC, Althaus M, Versluis-denBieman HJ. Damaging backgrounds: Later adjustment of international adoptees. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1992; 31(3):518–524. [PubMed: 1592786]
- Weschler, D. *Weschler Intelligence Scale for Children*. 3rd Ed.. San Antonio, TX: Harcourt Brace & Co; 1998. (WISC-III)
- Windsor J, Glaze LE, Koga SF, the BEIP Core Group. Language acquisition with limited input: Romanian institution and foster care. *Journal of Speech, Language, and Hearing Research*. 2007; 50:1365–1381.
- Zeanah CH, Nelson CA, Fox NA, Smyke AT, Marshall P, Parker SW, Koga S. Designing research to study the effects of institutionalization on brain and behavioral development: The Bucharest Early Intervention Project. *Development and Psychopathology*. 2003; 15:885–907. [PubMed: 14984131]

Table 1

Sample Characteristics

	Post- Institutionalized (N = 48)	Early- Adopted (N = 40)	Control (N = 44)
Sex (% female)	50	47	55
Age	8 years, 4 months (.5 months)	8 years, 4 months (.5 months)	8 years, 5 months (.5 months)
IQ ^I	105.9 (15.8)	113.7 (16.5)	124.7 (15.7)
Years of parent education	16.3 (2.2)	16.8 (1.6)	16.4 (2.0)
Median Family Income	75–110K	75–110K	50–75K

^I Estimated based on vocabulary and block design subtests of the WISC-III, or the LEITER-R for participants who scored below the normal range on the WISC-III.

Table 2

Descriptive Data on Domains Exhibiting Significant Group Differences

	Post-Institutionalized	Groups Early-Adopted	Control	Post-hoc Comparisons	Duration Institutional Care
Memory Tasks					
Spatial	52.7	40.9	39.0	Pl< Con	r=-.18
Working Memory [^] (% errors)	(18.4)	(19.5)	(15.7)	EA= Con	
Memory for	11.1	12.2	12.6	Pl<Con	r=-.11
Faces ⁺ (scaled score)	(3.1)	(2.8)	(3.3)	EA= Con	
Auditory	9.7	10.8	11.4	No significant	r=-.31 [*]
Narrative Memory ⁺ (scaled score)	(4.1)	(3.3)	(3.2)	differences	
Attention Tasks					
Match to	95.3	96.2	97.4	No significant	r=-.16
Sample [^] (% correct)	(7.3)	(4.6)	(3.4)	difference	
Auditory	9.2	9.6	9.7	No significant	r=-.06
Attention ⁺ (scaled score)	(1.3)	(1.4)	(1.0)	difference	
Rapid	10.5	9.6	8.9	No significant	r=-.04
Visual Processing [^] (% correct)	(6.1)	(3.9)	(4.6)	differences	
Visual	10.1	12.2	11.7	Pl< Con	r=-.33 [*]
Attention ⁺ (scaled score)	(2.3)	(2.7)	(2.1)	EA= Con	
Knock-	28.00	28.69	29.10	Pl< Con	r=-.35 [*]

	Post-Institutionalized	Groups Early-Adopted	Control	Post-hoc Comparisons	Duration Institutional Care
Tap ⁺ (raw score)	(2.93)	(1.81)	(1.05)	EA=Con	
Executive Control Tasks					
ID/ED [^] (total errors)	41.2 (19.3)	38.3 (23.7)	33.7 (17.6)	No significant differences	r=.09
Stockings Of Cambridge [^] (problems solved)	6.8 (1.8)	7.3 (1.7)	7.0 (1.5)	No significant differences	r=-.14
Learning Task					
Paired	14.6	6.9	6.2	PI<Con	r=-.46***
Associates Learning [^] (total errors)	(13.8)	(5.9)	(5.2)	EA=Con	

* p<.05

** p<.01

+ NEPSY subtest

[^] CANTAB subtest.