Published in final edited form as:

Psychol Bull. 2008 November ; 134(6): 912–943. doi:10.1037/a0013740.

Serotonergic Function, Two-Mode Models of Self-Regulation, and Vulnerability to Depression: What Depression Has in Common With Impulsive Aggression

Charles S. Carver, Sheri L. Johnson, and Jutta Joormann

Department of Psychology, University of Miami.

Abstract

Evidence from diverse literatures supports the viewpoint that two modes of self-regulation exist, a lower-order system that responds quickly to associative cues of the moment and a higher-order system that responds more reflectively and planfully; that low serotonergic function is linked to relative dominance of the lower-order system; that how dominance of the lower-order system is manifested depends on additional variables; and that low serotonergic function therefore can promote behavioral patterns as divergent as impulsive aggression and lethargic depression. Literatures reviewed include work on two-mode models; studies of brain function supporting the biological plausibility of the two-mode view and the involvement of serotonergic pathways in functions pertaining to it; and studies relating low serotonergic function to impulsiveness, aggression (including extreme violence), aspects of personality, and depression vulnerability. Substantial differences between depression and other phenomena reviewed are interpreted by proposing that depression reflects both low serotonergic function and low reward sensitivity. The article closes with brief consideration of the idea that low serotonergic function relates to even more diverse phenomena, whose natures depend in part on sensitivities of other systems.

Keywords

serotonin; depression; two-mode models; dual-process models; self-regulation

Depression is a disorder that afflicts countless people worldwide. A substantial literature has examined the role of neurotransmitters in depression; research has also turned to molecular genetics as a tool to gain leverage on neurobiological processes underlying depression. This body of neurobiological work has yielded valuable evidence. An important question remains, however, about how to interpret the evidence in psychological terms. A confluence of ideas from other areas of work suggests an answer to that question. This article reviews some of those literatures (almost exclusively human research) and advances a psychological viewpoint that may help in interpreting the neurobiological evidence.

A central idea examined here is that low serotonergic function is a marker of a deficit in executive control processes, processes that override or inhibit lower-level influences on behavior. A result is that persons with low serotonergic function (and thus diminished executive

Copyright 2008 by the American Psychological Association

Correspondence concerning this article should be addressed to Charles S. Carver or Jutta Joormann, Department of Psychology, University of Miami, Coral Gables, FL 33124-0751; or to Sheri L. Johnson, Department of Psychology, University of California, Berkeley, Berkeley, CA 94720. ccarver@miami.edu, jjoormann@miami.edu, or sljohnson@berkeley.edu.

Sheri L. Johnson is now at the Department of Psychology, University of California, Berkeley.

control) are especially responsive to associative and affective cues of the moment. Lack of executive control ultimately has divergent effects on behavior, however, because what follows depends on other aspects of the person's makeup. In the cases that are most familiar, a deficit in executive control yields impulsive action and is associated with externalizing disorders. The position advanced here is that in depressed persons, the same executive control deficit often (though not always) leads to automatic (and in that sense "impulsive") *inaction*, plus absorption in negative emotions.

It should be clear from this statement that the focus on serotonergic function in this article is not intended to imply that serotonin function is the entire story behind depression (or externalizing disorders either, for that matter). Dopamine function also appears to play a role in depression (Naranjo, Tremblay, & Busto, 2001). Indeed, the very idea that low serotonergic function and deficits in executive control affect different people in quite different ways clearly implies that there is an interaction among causal influences, leading some but not others to experience depression in the context of serotonergic deficits.

The view that is advanced here relies on several quite different sources of information, which are described in separate sections. One source is a set of theories from cognitive, social, personality, motivational, and developmental psychology, which converge on the idea that there exist two modes of processing information and regulating action, which operate simultaneously and in competition with each other. This set of theories forms the psychological perspective from which the other literatures are considered. A second body of evidence, considered next, describes functions of certain brain regions and the interplay among them, which appear consistent with the two-mode models, and thus supports the biological plausibility of the two-mode view.

The review then turns to cognitive and behavioral effects related to differences in serotonergic function, with the goal of mapping these effects onto the two-mode models. First described are studies of cognitive processing of emotional stimuli, impulsiveness in behavior, impulse-related disorders, and self-reports of several aspects of personality, all of which are then shown to reflect principles in the two-mode models. Next is evidence linking low serotonergic function to depression vulnerability. These findings represent a striking contrast to much of what was reviewed just previously, because the behavioral qualities now being related to low serotonergic function are quite different. This divergence between literatures is interpreted by incorporating another variable: sensitivity of approach motivation. The result is an interactive model, with two classes of potentially problematic outcomes (poor impulse control vs. depression).

The review continues by addressing additional neurobiological and behavioral findings concerning depression, which offer further support for the two-mode interpretation. Finally a set of remaining issues and further directions is considered.

Two-Mode Models of Functioning

Several theoretical models have emerged in diverse areas of psychology over the past two decades which posit that that people experience the world through two simultaneous but somewhat distinct modes of processing (Barrett, Tugade, & Engle, 2004; Epstein, 1994; J. St. B. T. Evans, 2008; Kahneman, 2003; Stanovich, 1999; Toates, 2006). These two processing modes may use different aspects of available information (Rudman, Phelan, & Heppen, 2007). There is also evidence that the two modes learn in different ways, and that the two patterns of learning create parallel and competing paths to action, which require continuous arbitration (Daw, Niv, & Dayan, 2005).

In cognitive psychology, this idea helped deal with a clash between two views of cognition. One view treats cognition as sequential symbol processing; another view assumes simultaneous parallel processing (e.g., Bechtel & Abrahamsen, 1991; Dawson, 2005; J. L. McClelland, 1999). Many cognitive psychologists now believe that both views are partly correct, that cognition (broadly conceived) uses two kinds of processes. One process—effortful, top-down, symbolic, and reflective—is used for planning and strategic behavior. The other—automatic, reflexive, bottom-up, and associationist—is used for acts that are heuristic, skilled, or urgent (Norman, 1986; Shastri & Ajjanagadde, 1993; Sloman, 1996; Smolensky, 1988; Smolensky & Legendre, 2006).

Two-mode models (also labeled *dual-process models*) exist in the literature of personality psychology as well. Indeed, Epstein's (1973, 1985, 1990, 1994) cognitive-experiential self theory may have been the first explicit two-mode model in contemporary psychology. It was based on the premise that humans experience reality via two systems. One is a symbolic processor—the rational mind. The other is associative and intuitive, uses short cuts and heuristics, and functions automatically and quickly. Epstein argued that both systems are always at work and that they jointly determine behavior.

Metcalf and Mischel (1999), drawing on several decades' work on delay of gratification, proposed a similar model. Delay of gratification research poses a choice between a smaller, less desired, but immediate reward versus a larger, more desired reward later (Mischel, 1974). Metcalf and Mischel (1999) proposed that two systems determine the ability to restrain in this and many other contexts: a "hot" system (emotional, impulsive, reflexive, and connectionist) and a "cool" system (strategic, flexible, slower, and unemotional). How a person responds to a difficult situation depends on which system presently dominates.

A conceptually similar line of argument about two modes of functioning also exists in social psychology. An early step in this direction was the argument that there are two paths to persuasion, one involving careful and thoughtful processing and the other involving heuristics and associations (Chaiken & Trope, 1999; E. R. Smith & DeCoster, 2000; Wilson, Lindsey, & Schooler, 2000). More recently, an analysis of automaticity in attribution posited two processing modes that are quite similar to those in the cognitive theories (Lieberman, Gaunt, Gilbert, & Trope, 2002). Strack and Deutsch (2004) proposed a farther-reaching model of overt social behavior, in which action is a joint output of two modes of functioning—reflective and impulsive—that occur simultaneously and may be either mutually supportive or in conflict with each other.

The two-mode idea has made a number of other appearances in social psychology as well. There is a rapidly expanding literature on implicit attitudes, implicit self-concepts, and the like (Custers & Aarts, 2005; Greenwald et al., 2002; McConnell, Rydell, Strain, & Mackie, 2008; Rydell & McConnell, 2006). Indeed, the idea that implicit and explicit mental processes play distinct roles in behavior is being applied to a steadily expanding range of topics, including (for example) moral judgments (Haidt, 2001), reactions to cues of stigma (Pryor, Reeder, Yeadon, & Hesson-McInnis, 2004), temporal discounting (McClure, Laibson, Loewenstein, & Cohen, 2004), and addiction (Wiers & Stacy, 2006). Of course, the idea that both implicit and self-attributed self-aspects influence behavior in different ways also traces to classic motivation theory (D. C. McClelland, 1984, 1985; D. C. McClelland, Koestner, & Weinberger, 1989; see also Schultheiss & Pang, 2007).

The two-mode idea also has an important presence in developmental psychology. Rothbart and her colleagues (e.g., Derryberry & Rothbart, 1997; Rothbart, Ahadi, & Evans, 2000; Rothbart, Ahadi, Hershey, & Fisher, 2001; Rothbart & Bates, 1998; Rothbart, Ellis, Rueda, & Posner, 2003; Rothbart & Posner, 1985) have argued for the existence of basic temperament systems

for approach and avoidance and a third temperament generally termed *effortful control* (see also Kochanska & Knaack, 2003; Nigg, 2000, 2003, 2006). Effortful control concerns attentional management (both in terms of sensitivity to new stimuli and in terms of the focusing or persistence of attention during long-lasting tasks) along with inhibitory control (the ability to suppress an approach behavior when doing so is situationally appropriate).

This model postulates that effortful control is superordinate to both approach and avoidance temperaments (e.g., Ahadi & Rothbart, 1994; see also Clark, 2005). The label “effortful” conveys the sense that this is an executive, planful activity, entailing the use of cognitive resources to deter the tendency to react impulsively. Rothbart (e.g., Rothbart & Bates, 1998), Eisenberg (e.g., Eisenberg et al., 2004), and Kochanska (e.g., Kochanska & Knaack, 2003) have seen effortful control as dependent on certain cortical functions (see also Nigg, 2001, 2003, regarding what he calls executive inhibition), and a variety of evidence from neuroimaging studies of both adults and children has supported that argument (e.g., Durston, Thomas, Worden, Yang, & Casey, 2002; Durston, Thomas, Yang, et al., 2002).

Commonalities Among Theories

Two-mode theories emerged in various areas for various reasons. Specific theories stress different issues, but they are similar in many respects (for broad views, see Barrett et al., 2004; J. St. B. T. Evans, 2008; Toates, 2006). Across theories, the two modes could be depicted fairly accurately as follows: A reflexive (or reactive or implicit) system, which is nonverbal and associationist, spontaneously creates action when its schemas or production systems are sufficiently activated. Being association based, it is said by many to be intuitive (e.g., Lieberman, 2000). It is often characterized as “emotional.” This label sometimes seems to imply that emotional experience is subjectively salient in this mode and sometimes seems to imply that this mode is very responsive to emotions triggered by situational cues. This system dominates when speed is needed (as when a situation is emotionally charged) and also when processing resources are diminished. That is, it requires relatively little capacity and thus can function under suboptimal conditions (see also De Neys, 2006). It tends to respond to short-term contingencies without consideration for the future or for broader consequences of the action. Thus, although it can be highly adaptive with respect to short-term contingencies, it may be less adaptive when a longer view is beneficial (cf. Daw et al., 2005; Kurzban & Aktipis, 2007; LeDoux, 2002).

The reflective (or rational or explicit) system, in contrast, operates mostly consciously, uses logical rules, is verbal and deliberative, and thus is comparatively slow. This rational system provides a more cautious, analytic, planful way of proceeding. It anticipates future conditions, makes decisions based on those anticipations, and forms top-down intentions. It is more wide ranging in its search for relevant information. Because the reflective system requires substantial cognitive capacity, it loses efficiency under high mental load or other conditions that limit cognitive capacity.

Although two-mode theorists tend to characterize the modes partly in terms of differences in emotionality, that point requires clarification. The theorists do not say that planful behavior is devoid of emotion. It is clear that planning ahead (as in delay of gratification) implies evaluating the affective consequences of various courses of action. However, an urgent and intense emotion appears likely to compete with, and to potentially short circuit, the tendency to be planful. The two-mode theories typically do not specify in detail the mechanism by which emotional salience or intensity promotes ascendance of the reactive system. It is generally implied, however, that greater subjective intensity means that the emotion cue is strongly activating an associative network within the reflexive system.

Behavioral Implications of Two-Mode Models

As a group, these theories suggest that the two modes, given their different operating characteristics, have differing influences on behavior. Behavior that is managed by what is typically characterized as the associationist, automatic, implicit, or reflexive mode tends to be impulsive. More precisely, this system responds to readily available associative cues of the moment. Behavior that is managed by what is typically characterized as the rational, reflective, sequential, or deliberative mode tends to be planful. It responds to two kinds of parameters—social and temporal—that go beyond immediate cues of incentive or threat.

This difference in operating characteristics tends to translate, for the most part, to a dimension of impulsive responsivity versus planful constraint (Carver, 2005; Nigg, 2001, 2003; Spont, 1992; Strack & Deutsch, 2004). This characterization requires an important qualification, however: Specifically, in principle, each mode of functioning can promote either action or inaction.

The developmental two-mode model is perhaps most explicit on this point. As described earlier, Rothbart and others posit three temperaments. Approach and avoidance temperaments together form the basis of what is termed *reactive control*. Reactive control is relatively reflexive and involuntary. The reactions in the term reactive control are automatic tendencies to approach incentives and avoid threats. These tendencies compete (Figure 1), and the resultant overt action is determined by which reactive tendency is stronger.

Reactive approach and avoidance tendencies can be relatively balanced, or one can dominate over the other. What is called *reactive undercontrol* (Eisenberg et al., 2004) refers to cases in which the automatic system for approach dominates the automatic system for inhibition or withdrawal; what is called *reactive overcontrol* refers to cases in which the latter dominates (Nigg, 2001, 2003, termed this *motivational inhibition*).

Thus, in the absence of executive override (effortful control), a person with a sensitive reactive approach system and an insensitive reactive avoidance system will display impulsive pursuit of incentives. A person with a sensitive reactive avoidance system and an insensitive reactive approach system will display reflexive freezing or avoidance. Reactive undercontrol relates to impulsiveness and externalizing disorders; reactive overcontrol relates to behavioral inhibition and internalizing disorders (Rothbart, Ellis, & Posner, 2004; Valiente et al., 2003).

The development of the capacity for effortful control (which depends on emergent executive functions) changes this situation, because effortful control can counter whatever is the resultant reactive tendency (effortful control is superordinate to reactive control). If enough effortful control capacity is available, the impulsive grabbing of incentives that arises from a sensitive approach system can be restrained (Kochanska & Knaack, 2003; Murray & Kochanska, 2002). This child (or adult) can delay gratification. More generally, impulsive action is dampened.

However, effortful control is, in principle, not entirely a matter of restraining approach (Figure 2). Effortful control sometimes means forcing the production of an action that one does not want to take (overriding a reflexive tendency toward *inaction*). For example, effortful control can lead a sedentary adult to exercise, or can lead a person to stay engaged in a boring task (Sansone, Wiebe, & Morgan, 1999). Effortful control can lead children to look happy when they receive gifts they do not like (Kieras, Tobin, Graziano, & Rothbart, 2005). Effortful control may be required for a spendthrift to inhibit the tendency to buy things, but it may also be required for a tightwad to overcome the tendency *not* to buy things (Rick, Cryder, & Loewenstein, 2008).

Thus, effortful control can have diverging effects, depending on the context. Effortful control can overcome relatively reflexive tendencies triggered by reactive systems. However, exerting effortful control can move a person either toward restraint or toward action, depending on what reflexive response is being overcome.

Two-Mode Models and Brain Functions

Dual-process models in the literatures just discussed are based on behavioral research. However, arguments have also been made on largely neurobiological grounds that effortful and automatic modes of functioning are managed by different brain areas (Bechara, 2005; Rolls, 2005; see also Corbetta & Shulman, 2002; Daw et al., 2005).¹ The automatic or reflexive system is said to include the ventromedial prefrontal cortex (PFC), the lateral temporal cortex, and subcortical regions such as the amygdala and the basal ganglia (Casey, Tottenham, & Fossella, 2002; Posner & DiGirolamo, 2000; Satpute & Lieberman, 2006). Controlled, effortful processing recruits regions that include rostral and dorsal areas of the anterior cingulate cortex (ACC) and the lateral and dorsal PFC.

It should be noted that assignment of brain areas to reflective and reflexive systems is not fully consistent across studies and models, and the picture continues to evolve as more data come in. Thus, the linking of specific brain regions to the two modes is by no means complete. It should also be noted that subcomponents of larger structures likely serve different systems. For example, the orbitofrontal cortex (OFC) has been found to be activated both in (reflexive) responses to stimuli evoking a sad mood and during (effortful) attempts to repair that sad mood (Cooney, Joormann, Atlas, Eugene, & Gotlib, 2007). Thus, some regions cannot be meaningfully assigned to one mode or the other. Nonetheless, there is evidence that does tend to support the two-mode idea. The sections that follow review this evidence; later comes discussion of how this view of the function of these regions helps interpret literature on serotonin and depression.

Automatic, Reflexive Mode

The reflexive mode includes the basal ganglia, which play a key role in automatic aspects of affect, cognition, and behavior. The putamen and caudate nuclei, for example, are involved in implicit learning and motor skill acquisition, responses that are learned slowly but executed quickly. The ventral striatum (a circuit consisting of the nucleus accumbens, parts of the putamen, globus pallidus, and caudate nuclei) is important in reward processing, contributing to the motivational salience of stimuli. Activity in this subcortical area increases in anticipation of reward (Ernst et al., 2004; Knutson, Fong, Adams, Varner, & Hommer, 2001). Craving and compulsive drug seeking are tied to activity in this area (Hyman & Malenka, 2001), and reactivity in this area relates to preference for immediate over delayed rewards (Hariri, Brown, et al., 2006).

The amygdala appears to play a key role in assigning emotional value to stimuli, both aversive and appetitive (Cunningham, Van Bavel, & Johnsen, 2008), although most available evidence concerns aversive stimuli (Davis & Whalen, 2001). For example, viewing faces expressing fear activates the amygdala (Whalen et al., 1998). The amygdala also is activated by novelty: for instance, an unfamiliar face (with no emotion) versus a familiar one (Schwartz, Wright, Shin, Kagan, Whalen, et al., 2003). The amygdala is also reactive to positive stimuli and stimuli that draw interest for reasons other than emotional content (Davis & Whalen, 2001). Reactivity of the amygdala appears to be a fairly stable trait; children characterized as inhibited at 2 years

¹In years past, it would have been common to refer globally to the reflective mode of functioning as depending on cortical or prefrontal functions and the reflexive mode as depending more on subcortical functions (indeed, we have sometimes done so). It has become increasingly apparent, however, that that dichotomy is not tenable.

of age show hyperreactivity of the amygdala to novel stimuli as adults (Schwartz, Wright, Shin, Kagan, & Rauch, 2003).

In light of such evidence, the amygdala is widely viewed as flagging stimuli quickly for emotional relevance (Aggleton, 2000; Anderson & Phelps, 2001; Cunningham et al., 2008; see also Kagan, 2007), thereby rendering the stimulus salient (Spoont, 1992). In this way, the amygdala plays an important role both in conditioning emotional reactions and in early stages of producing emotional reactions (Cahill, Babinsky, Markowitsch, & McGaugh, 1995; Cheng, Knight, Smith, & Helmstetter, 2006; LeDoux, 2000).

Besides emotion processing, subcortical areas are also involved in impulsive action. An example is impulsive aggression (Blair, 2003; Siever et al., 1999). For example, early positron emission tomography (PET) studies (Raine et al., 1994) showed high activity in the amygdala and other subcortical areas in persons who had committed impulsive murders.

Despite the fact that the PFC is linked in many people's minds to higher-order processing and executive control, activation in some PFC regions has been consistently linked to automatic and affective processing (Ochsner & Gross, 2005). For example, some ventral and medial areas of the PFC with strong direct connections to subcortical structures including the amygdala and the basal ganglia are activated in response to emotion-eliciting stimuli. The OFC in particular, with strong reciprocal connections to the amygdala and other subcortical structures, has been linked to learning through stimulus-reinforcement associations and to responses to changing reinforcers (Kringelbach & Rolls, 2004). These PFC regions represent information pertaining to reinforcers and are used to form intuitions about stimulus– outcome probabilities prior to conscious awareness (Bechara, Damasio, & Damasio, 2000).

Deliberative, Reflective Mode

With respect to the effortful mode, regions of the dorsal PFC are activated when people engage in reflective top-down processing, as in reappraising emotional stimuli in order to regulate emotional responses to them (Ochsner, Bunge, Gross, & Gabrieli, 2002). Similarly, effortfully suppressing emotion activates the dorsolateral PFC. As an example, when men viewed erotic pictures, the amygdala and related areas were activated; when they were asked to inhibit their emotional reactions, however, dorsolateral PFC and ACC regions were activated (Beauregard, Lévesque, & Bourgouin, 2001), along with diminished activation of the amygdala. Low activation in the dorsal PFC has also been linked to prolonged activation of subcortical structures in response to an emotional challenge (Siegle, Steinhauer, Thase, Stenger, & Carter, 2002).

It is important to keep in mind that executive control does not always imply reduction of emotion. For example, one study had some participants dampen emotions and others increase them (Ochsner et al., 2004). As would be expected, what happened in the amygdala depended on the task. Efforts to reduce emotion reduced amygdala activity; efforts to increase emotion increased amygdala activity. Efforts to increase also uniquely activated left medial PFC; efforts to reduce uniquely activated regions of right lateral PFC and OFC. However, there were also areas in which both tasks yielded the same effects. Both efforts to increase and efforts to decrease emotion led to increased activity in left lateral PFC and ACC, and dorsal medial PFC.

It was noted earlier that some ventral PFC areas are involved in reflexive processes; there is also evidence, however, that some ventral PFC areas (parts of the OFC and ventral medial PFC) play a role in *regulating* reflexive processes (Drevets, 2000; Pizzagalli et al., 2004)—inhibiting the amygdala (Beauregard et al., 2001; Hariri et al., 2002, 2005) and ventral striatum (Nestler & Carlezon, 2006). For example, activation of some ventral PFC areas has been found during suppression or reappraisal of emotional stimuli (Lévesque et al., 2003; Ochsner et al.,

2004) and when suppressing influences of emotional stimuli on behavior (Beer, Knight, & D'Esposito, 2006). Impulsivity has also been related to low activation in ventral PFC (Brown, Manuck, Flory, & Hariri, 2006). Thus, evidence links ventral PFC and OFC to both postulated systems (Cooney et al., 2007; Lévesque et al., 2003; Ochsner et al., 2002). Perhaps functional differentiation in the OFC links it to both modes (Zald & Kim, 1996); perhaps this region's high degree of interconnectivity with other regions makes it a bridge between systems (Wallis, 2007).

The focus of this section thus far has been on control of emotions and impulses, but dorsal and lateral parts of the PFC actually are involved primarily in control of cognition (Horn, Dolan, Elliott, Deakin, & Woodruff, 2003; Jonides, Smith, Marshuetz, Koeppe, & Reuter-Lorenz, 1998), inhibiting attention to distracting information (Chao & Knight, 1998) and inhibiting response tendencies when a task changes (Jonides et al., 1998). Thus, lateral and dorsal PFC regions are involved in cognitive processes that seem involved in the effortful regulation of emotion. However, these PFC areas are not themselves directly connected to subcortical areas (Clark, Cools, & Robbins, 2004; Lieberman, 2007; Schaefer et al., 2003), though they do connect closely to ventral and medial areas of the PFC, which have strong reciprocal connections to subcortical areas. It is possible that dorsal and lateral areas work through ventral and medial areas to affect amygdala activity (Lieberman, 2007; S. F. Taylor, Phan, Decker, & Liberzon, 2003).

Another cortical area that plays a key role in the regulation of subcortical regions is the ACC. A core function of the ACC is detection of conflict between responses cued by different aspects of available information (Bush, Luu, & Posner, 2000; Carter et al., 1998; Mitchell, 2006), but there is also functional differentiation within the ACC (e.g., Milham & Banich, 2005; Mohanty et al., 2007; van Veen & Carter, 2006). The dorsal ACC seems especially responsive to cognitive aspects of conflict, calling on dorsolateral PFC areas to exert top-down attentional control, to maintain focus on the task at hand by inhibiting processing of task-irrelevant information (Paus, 2001; van Veen & Carter, 2006). The dorsal ACC may be further differentiated, as well, with anterior areas involved in early-stage response evaluation and posterior areas involved in later-stage processes involving the response itself (Milham & Banich, 2005). This leads some to see this area as being involved in cognitive-motor mechanisms (Mohanty et al., 2007). Overall, however, the function of these areas is to minimize distraction and keep the intended behavior on track.

In contrast to these cognitive influences, rostral and ventral areas of the ACC represent its "affective" subdivision. They connect extensively with the OFC and with subcortical systems including amygdala and nucleus accumbens. Pizzagalli et al. (2001) argued that a main function of this subdivision of the ACC is assessing conflict between what the organism is currently doing and incoming information about potentially rewarding and emotional stimuli. Indeed, several studies have shown activation of this area when effortful emotion regulation is required (Bush et al., 2000; Ochsner & Barrett, 2001).

The affective subdivision may integrate salient affective and cognitive information and may change attentional processes in the cognitive subdivision (e.g., Pizzagalli et al., 2001). Indeed, pathways related to attentional processing and amygdala pathways converge within the rostral ACC. It thus has been argued that the rostral ACC may be especially involved in cognitive control in the context of emotional stimuli (Mohanty et al., 2007).

Interplay Between Reflexive and Reflective Systems

Dual-process models hold that both systems are at work and potentially competing with each other. Certainly some of that competition was implied in the preceding section. However, the interplay between the systems has been related more explicitly to a variety of outcomes,

including impulsivity and emotional processing. With regard to impulsivity, impulsivity on tasks such as the go/no-go task has been linked to greater activation in subcortical areas relevant to emotion identification (e.g., amygdala, right insula), coupled with lower activation in ventral PFC areas (Brown et al., 2006; Horn et al., 2003). Thus, when areas of the reflective system are not inhibiting areas of the reflexive system that generate prepotent responses to cues, the person is more impulsive.

There is also evidence linking this impulsivity to emotional processing per se. Brown et al. (2006) found that higher scores on the Barratt Impulsivity Scale predicted more activity in the amygdala and less activity in ventral PFC during viewing of emotional faces (a task that engages amygdala activity). This is consistent with the idea that impulsivity relates more broadly to a general vigilance for immediate emotional relevance, coupled with lower effortful control.

Another study showing the interplay of these systems in regulating emotion and impulsive behavior (Hariri, Brown, et al., 2006) determined behavioral preferences for delayed versus immediate reward in one session, then related preference to activation of brain areas in another session. Initial preference for immediate reward (suggesting relative dominance of the reflexive system) predicted greater ventral striatum responses to both success and failure feedback on the later task (compared with no feedback) and greater activation of medial PFC (after success), indicating relative lack of executive control. In contrast, initial preference for delayed reward (suggesting relative dominance of the deliberative system) predicted greater activation of lateral OFC and dorsolateral PFC after success feedback, indicating relative presence of executive control. Thus, activity in prefrontal areas highly connected with the ventral striatum—the medial PFC and the OFC—related to preference for immediate rewards; activity in dorsolateral PFC (which may be involved in constraining the other areas) related to preference for larger, delayed rewards.

These findings are also consistent with those from a study by McClure et al. (2004) who reported that ventral striatum activity was significantly higher as participants chose smaller, immediate rewards compared with larger, delayed rewards. Activation of medial PFC and OFC also correlated positively with the selection of immediate rewards. In contrast, activation of dorsolateral PFC correlated with selection of larger, delayed rewards.

Consistent with this central role of certain prefrontal areas in delaying reward and controlling impulses, adults with PFC damage have been characterized as resembling children in terms of emotional reactivity and difficulties in regulating emotion (Fox, Henderson, Rubin, Calkins, & Schmidt, 2001). Galvan et al. (2006) recently reported that adolescents show greater activation in ventral striatum and more diffuse recruitment of the OFC in a reward task. They suggested that the fact that subcortical areas develop earlier than dorsal areas of PFC might account for impulsive and risk taking behavior in this age group (see also Steinberg, 2007).

As another example of the interplay between systems, functional connectivity (by within-subject correlations) has been seen between amygdala and dorsolateral, dorsal medial, ACC, and OFC during affect regulation by cognitive reappraisal (Banks, Eddy, Angstadt, Nathan, & Phan, 2007). Indeed, the degree of within-subject coupling between the amygdala and the OFC and dorsal medial cortical areas predicted the degree of affect reduction that occurred during the task.

Section Summary

The overall pattern of effects described in this section appears to be generally consistent with a two-mode picture of self-regulation. It fits the idea that certain brain areas (associated with the reflexive) system quickly identify emotional relevance and initiate emotional reactions,

whereas specific areas of the prefrontal cortex (associated with the deliberative system) sometimes act to inhibit those reactions.

Methods of Human Research on Serotonergic Function

The two-mode viewpoint outlined in the preceding two sections is the perspective from which the rest of the article proceeds. Specifically, the sections that follow briefly review other literatures and interpret them from the two-mode framework. Those literatures also share one more feature: Each of them examines the involvement of serotonergic function in some type of behavior. All those sections raise the possibility that serotonergic function influences the balance between the two processing modes. Before turning to those sections, however, some comment is in order about research methods bearing on serotonergic function in humans (for greater detail, see Manuck, Kaplan, & Lotrich, 2006).

The processes by which serotonin operates in the nervous system are complex and not fully understood (Hensler, 2006; Lesch & Canli, 2006). It is clear, however, that more is involved than levels of available serotonin (e.g., Neumeister et al., 2006), including such factors as sensitivity and density of receptors, efficiency of metabolizing neurotransmitters released into the synaptic cleft, efficiency of re-uptake of neurotransmitter from the cleft, dietary intake of amino acids, and recent history of the cell's firing (thereby depleting available stores).

Thus it can be misleading to think of the serotonergic system in terms of level of serotonin per se, though there are contexts in which that view is appropriate. That is, temporary perturbations can change how much serotonin is available and thus have transient effects on behavior. In any long-term view of the serotonergic system, however (such as a person's chronic experience over an extended period of time, or the tuning of the system by the prolonged use of selective serotonin reuptake inhibitors [SSRIs]), the issue really is the overall functioning of the system, rather than the levels of serotonin per se.

In light of such complexities, researchers have turned to several indirect ways to examine the role of serotonergic function in a given behavior or experience. Some of these methods focus on temporary perturbations of functioning, some on longer-standing patterns.

Acute Tryptophan Depletion or Enhancement

One relatively common and direct procedure temporarily lowers levels of serotonin experimentally, by artificially depleting the person of tryptophan, a precursor to serotonin. Tryptophan can be depleted with a drink (or capsules) containing high levels of 15 amino acids but no tryptophan. A control condition is a drink that has no effect on the balance of tryptophan to other amino acids. The manipulation results in an episode of acute tryptophan depletion with its lowest point several hours later, which in turn temporarily reduces available serotonin. The question of interest in this paradigm is the nature and extent of the response (behavioral, affective, or cognitive) to the lowering of available serotonin. Less frequently, manipulations have been used that work in the opposite direction, enhancing tryptophan levels and thus serotonin availability.

Drug Challenge

Another approach is to assess serotonergic function indirectly, by means of a drug challenge to the serotonergic system (Coccaro & Kavoussi, 1994; Power & Cowen, 1992). A challenge test involves administering precursors, agonists, or releasers of serotonin and assessing subsequent change in a hormone that is sensitive to change in serotonin. In challenge studies of serotonin and aggression, the most commonly used drug in the existing literature is fenfluramine, a serotonin releaser (Lopez Ibor, Saiz Ruiz, & Iglesias, 1988), and the most commonly measured outcome is prolactin response 3–4 hr later (Manuck et al., 2006).

Serotonergic function is inferred from the prolactin response (Cowen, 2002; Flory, Mann, Manuck, & Muldoon, 1998; Hennig, Toll, Schonlau, Rohrmann, & Netter, 2000). Blunted prolactin responses are taken to indicate lower serotonergic functioning. The level of prolactin response then is related to the target behavior or psychological characteristic.²

One further issue is important in this particular literature (Manuck et al., 2006): Women's prolactin response to serotonergic agonists can vary as much as threefold as a function of phase of the menstrual cycle (O'Keane, O'Hanlon, Webb, & Dinan, 1991). Unfortunately, this is often not considered in studies using this method. This is a probable reason for inconsistent results among women when researchers use this methodology.

Variation in Genotype

Another methodological strategy is to relate the behavior of interest to genetic polymorphisms that have independently been associated with serotonergic function (Manuck et al., 2006). Though a large number of genes relate to serotonergic function (Martinowich & Lu, 2007), most of the genetic research discussed here examined the gene that codes the serotonin transporter. The transporter plays a key role in serotonergic transmission by facilitating reuptake of serotonin from the synaptic cleft (Heils et al., 1996). Indeed, under normal circumstances, this is the principal mechanism for actively clearing serotonin from the synaptic cleft.

Transcriptional activity of the transporter gene is believed to be influenced by (or at least associated with) a repetitive sequence in a polymorphic region called 5-HTTLPR, which has a short version and a long version (i.e., which has more repetitions). A variety of indirect evidence links this polymorphism to serotonergic function, though it is important to note that the evidence is not completely uniform (disconfirming findings have been reported, e.g., by Mann et al., 2000, and van Dyck et al., 2004).

Supportive evidence includes the following: The long version is associated in vitro with higher transcription efficiency than that of the short one. In human lymphoblast cells, for example, the long-long combination was linked to serotonin uptake 1.9 to 2.2 times that of the short-short or long-short combination, which had comparable serotonin uptake (Lesch et al., 1996). The alleles have also shown similar ratios in binding levels, in imaging measures of radioligand binding to 5-HTT in vivo (Heinz et al., 2000) and in postmortem calculations of 5-HTT density (Little et al., 1998). Neuroimaging studies in humans and nonhuman primates have found an inverse relation between 5-HTT availability and cerebrospinal fluid concentrations of a 5-HT metabolite (Heinz et al., 2002), suggesting that the 5-HTTLPR is functional and influences serotonergic neurotransmission. One recent study found carriers of the short allele had twice the turnover of brain serotonin as those without the short allele (Barton et al., 2008). Of particular importance, in some of the literatures to be discussed, the 5-HTTLPR polymorphism has been associated with outcomes that resemble those obtained through direct manipulation of the serotonergic system. Thus, although there remain questions about causal mechanisms, the short allele is widely viewed as a marker of low serotonergic function (e.g., Canli & Lesch, 2007).

Two variants of the long allele have been identified relatively recently (Hu, Zhu, Lipsky, & Goldman, 2004). The expression of one variant is comparable with that of the short allele, and

²Fenfluramine is not the only possible drug challenge, and because of its potential toxicity with long-term or repeated usage, newer studies are turning to other challenges. Other drugs used include m-CPP, buspirone, and ipsapirone. Unfortunately, m-CPP and buspirone influence dopamine function as well as serotonin, and ipsapirone and m-CPP both bind to adrenergic receptors (Yatham & Steiner, 1993). For these reasons, even though findings from these challenges are generally comparable with those from fenfluramine (Manuck et al., 2006), consideration of challenge studies here is limited almost entirely to fenfluramine studies.

the two long variants behave differently in response to pharmacological challenge (Hu et al., 2005; Neumeister et al., 2006). Such findings encourage closer attention to variability in this polymorphism (this polymorphic region may have as many as 10 variants in humans; Nakamura, Ueno, Sano, & Tanabe, 2000). Further, polymorphisms at other sites on this gene also appear to have functional effects on transporter expression (Kraft et al., 2007). More generally, it is important to acknowledge that the genetic association studies we describe here evaluate correlations between a phenotype and the single marker that is under study, rather than a causal link.

Nonetheless, most published work on the serotonin transporter is limited to comparisons on 5-HTTLPR. Most published work is further limited to comparisons that involve the long and short allele without further differentiation. Thus, in this article, mention of polymorphisms in the serotonin transporter gene always refers specifically to 5-HTTLPR and primarily to short-versus long-allele carriers. A few studies that do go beyond this are noted later on.

Serotonergic Function and the Brain

This section addresses the idea that serotonergic function plays an important role in differentiating the two modes of functioning introduced in the article's first major section. This idea is advanced by relating serotonergic function to brain areas that were discussed in the second major section as subserving those two modes.

The central serotonergic system projects from the brainstem extensively throughout cortical and subcortical structures. Serotonergic projections originate primarily in the dorsal and median raphe nuclei. Neurons within these nuclei have different but overlapping projection patterns. Serotonergic synapses are densely concentrated in subcortical regions, including the amygdala, ventral striatum, and hypothalamus. Raphe neuron firing is regulated mostly by projections from brainstem nuclei, with some input from thalamus, hypothalamus, and limbic areas, but there is evidence that the ventral OFC has a direct inhibitory effect on raphe activity (Hensler, 2006; Ressler & Nemeroff, 2000).

The focus here is on areas of serotonergic innervation that are most directly relevant to the two-mode idea. Ascending serotonergic projections from the raphe nuclei have a widespread cortical distribution, with particularly dense serotonergic projections to the medial, ventral, and orbital regions of the PFC (Way, Lácan, Fairbanks, & Melega, 2007). Tryptophan depletion has been found to decrease 5-HT₂ receptor binding in the dorsolateral PFC (Yatham et al., 2001), suggesting that receptors in this region are particularly sensitive to serotonergic variation. Tryptophan depletion has also been found to decrease activation in the dorsolateral/medial PFC during a verbal working memory task (P. P. Allen et al., 2006). Tryptophan depletion has been shown to affect ACC activity (K. A. Smith, Morris, Friston, Cowen, & Dolan, 1999), particularly among persons with the short allele of the serotonin transporter gene (Neumeister et al., 2006). Thus, serotonin seems important in the functioning of cortical structures identified in the earlier section on brain function and two-mode models.

A previous section also noted that cortical and subcortical structures are interconnected, forming complex circuits that regulate emotion and behavior. Corticolimbic neural circuits that mediate emotion are densely innervated by serotonergic neurons and exhibit rich expression of serotonin receptors (Parsey et al., 2006). Indeed, serotonin appears critical for the development of this emotional circuitry, and even transient alterations during early development modify neural connections (Hariri & Holmes, 2006).

Within these circuits, serotonergic function is thought to be involved particularly in constraining excitatory influences on amygdala activity (Hariri & Holmes, 2006; Hariri & Weinberger, 2003). It has been argued that increased serotonin levels may thereby decrease

the sensitivity of the amygdala to external stimuli (particularly aversive stimuli; Hariri et al., 2002). Support for this general idea has emerged from studies of the 5-HTTLPR polymorphism (Hariri et al., 2005), drug challenge studies, and neuroimaging studies (Clark et al., 2004).³

In one of the first projects to combine genetic analyses and brain imaging, Hariri et al. (2002, 2005) found that carriers of the short allele of 5-HTTLPR had greater amygdala reactivity to emotional faces than did those with the long allele. Considerable additional evidence that short-allele carriers display amygdala hyperreactivity has accumulated since then (for a meta-analysis, see Munafò, Brown, & Hariri, 2008, who report that 5-HTTLPR may account for up to 10% of phenotypic variance in amygdala activation to a broad range of salient environmental stimuli).⁴ There is also evidence that the short allele of 5-HTTLPR relates to lower resting activity in the ventromedial PFC, an area that constrains amygdala activity (Rao et al., 2007).

Other data relate the 5-HTTLPR polymorphism to both structural and functional variations in this circuit as a whole. Pezawas et al. (2005) found smaller gray matter volume in a specific region of the PFC (the rostral ACC) and in the amygdala in carriers of the short allele compared with those without it. These authors also used functional connectivity analysis, a measure of correlation in activity between regions, derived from fMRI data. Amygdala and ACC activity were correlated overall; the subgenual rostral part of the ACC (BA 32/25/24) was positively correlated with amygdala activity, and the supragenual more caudal part of the ACC (BA 32) was negatively correlated with amygdala activity. These two regions of the ACC also showed strong positive connectivity with each other, suggesting that they may form a feedback loop with the amygdala.

These analyses revealed less coupling of these structures when viewing angry and fearful faces among carriers of the short allele. Carriers of the short allele therefore appear to show less inhibitory regulation of the amygdala than do those with the long allele (see also Heinz et al., 2005). It is noteworthy that the connectivity difference between groups was most noticeable in a PFC area that has the highest density of serotonin transporter terminals in the human cortex and is the target of dense projections from the amygdala.

The observations of amygdala hyperreactivity in carriers of the short allele in other studies may reflect the decoupling of this circuit. If activity of this prefrontal region inhibits activity in the amygdala, less connectivity in the carriers of the short allele would permit greater excitability of the amygdala and deficits in affect regulation (Hariri, Drabant, & Weinberger, 2006).

Pezawas et al. (2005) also asked their participants to complete a self-report measure of harm avoidance, as an indicator of emotionality. Measures of prefrontal activity and amygdala activity per se did not predict individual differences in harm avoidance. However, indices of the connectivity between these areas accounted for almost 30% of the variance in this trait.

³It may be desirable to distinguish here between inhibition and constraint as concepts. Spont (1992) argued that constraint means regulation of information flow so as to prevent being overwhelmed by it, rather than suppression of information flow. Her view of serotonergic function was that it constrains—regulates—communication rather than inhibiting it (see also Depue & Spont, 1986). By limiting the flow of information, constraint serves an important homeostatic function, preventing overshoot of other elements within the overall system (e.g., excessive stress reactivity).

⁴It was initially assumed that low serotonergic function relates to stronger amygdala reactions to emotion-relevant stimuli. Other evidence, however, casts a somewhat different light on the findings. Canli et al. (2005) found that the short allele predicted greater amygdala activity to negative stimuli than to neutral stimuli (replicating previous findings) but also predicted more amygdala activity to a fixation point than to neutral stimuli (replicated by Heinz et al., 2007). Canli et al. (2005, 2006) suggested that the short allele relates to greater tonic activation: that in the absence of cognitive constraints (i.e., at rest) the amygdala of this person is active, and that a nonemotional cognitive event (perceiving a neutral stimulus) reduces the activity (Canli & Lesch, 2007; Shulman et al., 1997), creating a difference between neutral stimuli and fixation. An accumulation of other evidence (Brocke et al., 2006; Canli et al., 2006) suggests that the amygdala in a person with the short allele does not just respond more intensely to emotional stimuli but is more vigilant—more active whenever there is uncertainty or novelty (cf. Davis & Whalen, 2001).

This suggests a link between decoupling of the amygdala–rostral-ACC feedback circuitry and elevation in dispositional emotionality. This circuit as a whole, then, appears to be affected by serotonergic function.⁵

Section Summary

As described earlier, specific brain regions and their interconnections appear to function in ways that resemble the functions of the two-mode behavioral and cognitive models. Evidence reviewed in this section appears to be consistent with the view that serotonergic function plays an important role in these brain areas and their interaction. The data suggest that low serotonin function (as manipulated by acute tryptophan depletion and assessed by serotonin transporter polymorphism) results in less constraint of amygdala activity.

Cognitive and Behavioral Effects of Serotonergic Function

The previous section examined how serotonergic function is involved in brain regions that were identified earlier with the two-mode model. What evidence links serotonergic function to psychological manifestations of the two processing modes? If serotonergic function is related to constraint of areas associated with the reflexive system (e.g., the amygdala), a number of other things should follow. Most simply, high serotonergic function should facilitate cognitive and behavioral inhibition of emotional states and impulses. This does appear to be so.

Serotonergic Function and Emotion-Related Processing

Much of the human research relevant to this idea has examined behavioral effects of serotonergic function experimentally, using acute tryptophan depletion to temporarily reduce serotonin. Tryptophan depletion has been shown to impair performance on cognitive tasks of executive processes such as reversal learning, memory, and attention tasks (Park et al., 1994; Riedel, Klaassen, Deutz, van Someren, & van Praag, 1999). Tryptophan depletion affects performance on behavioral and cognitive tasks that have emotional components. As an example, tryptophan depletion impairs the ability to inhibit responses to previously rewarded cues (Cools, Blackwell, et al., 2005; Park et al., 1994; Rogers et al., 2003). It is noteworthy that the strongest effect of tryptophan depletion found by Cools, Blackwell, et al. (2005) was among people who reported being highly impulsive. This pattern is consistent with the two-mode model.

A particular focus has been on how cognitive tasks that involve emotional stimuli are affected by tryptophan depletion. For example, in one study acute tryptophan depletion led female participants to show slower processing of happy words (but not sad ones) in an affective go/no-go task (F. C. Murphy, Smith, Cowen, Robbins, & Sahakian, 2002). This study also found impairment on the first round of a reversal task involving reward and punishment, but no impairment on a task without affective stimuli.

Tryptophan depletion generally does not affect mood per se among persons with no personal or family history of depression (Benkelfat, Ellenbogen, Dean, & Palmour, 1994; Park et al., 1994). However, the impairment in suppressing emotion-relevant stimuli suggests that tryptophan depletion might exaggerate mood change in response to a stressor. Consistent with that possibility, tryptophan depletion has led persons to report more sadness during exposure to uncontrollable stress (aversive noise), whereas the effect was only minor when the noise was controllable (Richell, Deakin, & Anderson, 2005).

⁵This discussion of serotonergic function was framed in terms of circuitry that seems relevant to two-mode models. A question that can be raised but not answered at this point is whether the effect of serotonergic function in this circuitry is better characterized as activating the deliberative system, which then constrains the reflexive system, or as deactivating the reflexive system, which then permits the deliberative system to play a larger role (cf. Gotlib & Hamilton, 2008). For present purposes, the end result is the same in either case.

Evidence for the role of serotonin in inhibition of responses to emotion-relevant stimuli has also emerged in one study of tryptophan supplements (which enhance serotonin availability). Women given tryptophan supplements for 14 days showed decreased recognition of disgusted faces and a diminished startle response to negative pictures, as well as an increase in the ability to recognize subtle facial expressions of happiness, compared with a control group (S. E. Murphy, Longhitano, Ayres, Cowen, & Harmer, 2006). The effect did not emerge for men.

Behavioral Effects of SSRIs in Nonclinical Samples

Researchers have also manipulated the serotonergic system by administration of SSRIs. The vast majority of such studies are of patients under treatment. However, in clinical samples an important question arises as to whether any effect observed reflects the change in serotonergic function or relates instead to other changes that follow from improved psychiatric status. Given such issues, effects of SSRIs in nonclinical samples are of particular interest. Several such studies have been published.

Knutson et al. (1998) administered 4 weeks of SSRI to a group of 26 nonclinical participants and compared them with a control group of 25. SSRI led to less negative affect (from the Positive and Negative Affect Schedule, PANAS; Watson, Clark, & Tellegen, 1988). There was no change in positive affect, and no tests were reported for specific affects. Those with SSRI also displayed more cooperative behavior in a group task than did control participants.

In another small study of nonclinical participants, Tse and Bond (2002) had roommates make ratings of the participants and measured several qualities of behavior during a mixed-motive game with another person. Those receiving SSRI ($n = 10$) were rated by roommates as being less submissive than control participants ($n = 10$), and they showed more dominant eye-contact behavior during the mixed-motive game than did control participants. However, they also displayed more cooperative, affiliative behavior during the game.

Impulsiveness

The idea that serotonin function plays a role in constraining emotion also has implications for behaviors involving impulsive responses to emotions. Evidence of behavioral impulsiveness comes from a number of sources (e.g., Crockett, Clark, Tabibnia, Lieberman, & Robbins, 2008; Walderhaug et al., 2002), but of particular importance is work bearing on impulsive responses to anger. That literature includes both studies of low-level aggression in the laboratory and studies of violence in naturalistic settings. Some of the former research is noted here; the latter studies are taken up in a later section.

An important conceptual point is made in a study of aggressiveness by Cleare and Bond (1995). Participants were pre-assessed as being either high or low in aggressive tendencies. Those high in aggressive tendencies became more aggressive, hostile, and quarrelsome after tryptophan depletion, and less so after tryptophan enhancement, but there was no effect for those low in aggressive tendencies. Similar results were reported by Finn, Young, Pihl, and Ervin (1998). This pattern suggests that effects of low serotonergic function on aggression are less about aggression per se and more about the release of existing habitual tendencies to be aggressive (see also Manuck et al., 2006; Spoont, 1992). This is consistent with the idea that serotonergic function promotes effortful control over underlying vulnerabilities.

A later study (Bjork, Dougherty, Moeller, & Swann, 2000) reinforced this point: Tryptophan depletion in this case led to greater aggressive response to provocation among men high in aggressiveness but had an opposite effect among those low in aggressiveness. Thus, the effect of serotonergic function on aggression appears to depend at least partly on dominant preexisting tendencies.

Section Summary

The studies described in this section appear to be relatively consistent in suggesting that experimentally increasing serotonergic function reduces responsiveness to negative emotional stimuli, decreases aggression, and increases cooperativeness; there also were increases in what might be thought of as social potency (see also Spont, 1992; Young & Leyton, 2002). Reducing serotonergic function impaired performances on tasks with emotional elements; it also increased expression of aggression and deterioration of cooperativeness, though these effects often occurred only among persons who were relatively high in aggressive tendencies by disposition.

Serotonergic Function, Impulse-Related Clinical Disorders, and Violence

Evidence linking serotonergic function to lack of executive control, impulsiveness, and aggression in nonclinical samples suggests that low serotonergic function should also relate to disorders that are characterized by these qualities. There are in fact intriguing parallels between phenomena described in the preceding section and certain phenomena that are more clinically relevant. This section considers some of those parallels, though it is not an attempt to review externalizing disorders in general (for broader treatment see, e.g., Krueger et al., 2002). Specifically, evidence is briefly considered here regarding serotonergic involvement in attention-deficit/hyperactivity disorder (ADHD) and externalizing behaviors such as conduct disorder, disruptive disorders, and antisocial personality disorder.⁶

ADHD

There is a substantial basis for believing that ADHD is actually a disorder of inability to control behavioral impulses (Barkley, 1997; Nigg, 2001, 2003; Quay, 1988). Symptoms include distractibility, difficulty following instructions, disorganization, and talking out of turn. Evidence of actual deficits in attention has generally been lacking (Barkley, 1997). Barkley (1997) argued that this disorder rests on deficits in executive functions and consequent deficits in motor control (for evidence that problems labeled *attention problems* reflect primarily difficulties with response inhibition, even in unselected children, see Friedman et al., 2007).

A more recent review led Nigg (2001, 2003) to a similar conclusion. Indeed, Nigg explicitly utilized a two-mode approach in analyzing ADHD. Inhibition by anxiety, uncertainty, or fear he termed *motivational inhibition*. What he termed *executive inhibition* resembles the concept of effortful control (though he applied it only to inhibitory tendencies). His review led him to conclude that ADHD reflects executive deficits. Consistent with this, an association has been found between the 5-HTTLPR polymorphism and ADHD (Beitchman et al., 2003).

Conduct Disorder, Antisocial Personality Disorder, Violence

A lack of executive control has also been linked to conduct disorder and other displays of antisocial tendencies. Dennis and Brotman (2003) found that a measure of effortful control predicted lower levels of (mother-reported) aggression in a high risk sample (younger siblings of youths adjudicated through the family court system). Other evidence ties weak effortful control to externalizing disorders as well (Rothbart et al., 2004; Valiente et al., 2003).

There is also evidence linking this sort of behavioral problem to serotonergic function. Within an ADHD group, one study related aggression to low serotonergic function assessed by fenfluramine challenge (Halperin et al., 1994). A more recent study of children with ADHD

⁶Apart from space considerations, studies of serotonergic function in substance-related disorders are not addressed because of the clear evidence that exposure to alcohol and drugs itself influences the function of neurotransmitter systems and the complexity of evaluating pre- and postdiagnostic data within humans (Koob & Le Moal, 2008).

found a prospective link from low serotonergic function (again by fenfluramine challenge) to antisocial personality disorder 9 years later (Flory, Newcorn, Miller, Harty, & Halperin, 2007). Another developmental study tested boys with disruptive behavior disorders (ages 7–11 years) with fenfluramine challenge and reevaluated the boys' clinical status nearly 7 years later (Halperin et al., 2006). Adjusting for baseline aggression, low serotonin function prospectively predicted higher adolescent aggression and antisocial behavior. Regardless of level of aggression in childhood, no child who had displayed high serotonin function in childhood was particularly aggressive in adolescence, suggesting that high serotonin function may have a protective role.

A good deal of research has also examined serotonergic function assessed by drug challenge in adults with clinical conditions reflecting impulsive aggression (for a more extensive review, see Manuck et al., 2006).⁷ O'Keane et al. (1992) found that convicted murderers with antisocial personality disorder had a blunted response to fenfluramine, compared with that of control participants. Lower serotonergic function has long been linked to history of fighting and assault (Coccaro, Kavoussi, Cooper, & Hauger, 1997), domestic violence (George et al., 2001), and impulsive aggression more generally, particularly among men (Cleare & Bond, 1997; Coccaro, Kavoussi, Hauger, Cooper, & Ferris, 1998). Consistent with the idea that impulsivity is a key element in these disorders, a serotonin releaser was found in one study to cause lower impulsivity on a delayed reward task among persons with conduct disorders (Cherek & Lane, 2000).

Genetic evidence also connects serotonergic function to violent and antisocial behavior. One recent study of young adults, some with personality disorders and others without, related the short allele of the 5-HTTLPR polymorphism to traits pertaining to borderline personality disorder and antisocial personality disorder (Lyons-Ruth et al., 2007), though another study (Sakai et al., 2007) found no link from this polymorphism to adolescent conduct problems. A sample of men in China convicted of extremely violent crimes was more likely to include short-allele carriers of the 5-HTTLPR polymorphism than was a control group (Liao, Hong, Shih, & Tsai, 2004), though the polymorphism did not relate to antisocial personality disorder within the criminal group. Another study (men only) found that the combination of the 5-HTTLPR polymorphism short allele and adverse childhood environment predicted violent adult behavior (Reif et al., 2007).

Other studies also make more differentiated points regarding serotonin and impulsive aggression. M. C. Dolan, Anderson, and Deakin (2001) linked low serotonergic function to higher impulsivity and higher aggression in male aggressive offenders. Interestingly, both impulsivity and aggression also related to higher anxiety in this sample, arguing against a path in which impulsive aggression is a product of low fear. This pattern thus fits Nigg's (2001) description of lack of executive inhibition rather than lack of motivational inhibition.

M. C. Dolan, Deakin, Roberts, and Anderson (2002) examined men with aggressive personality disorders (from a maximum security psychiatric hospital) and a control sample (from hospital staff) on some of the same outcomes and some different ones. Men diagnosed with psychopathy (the majority of the aggressive group) had poorer executive function and greater self-report impulsivity than did the others. The psychopathic subset of aggressors had lower serotonergic response than did the nonpsychopathic subset, but it did not differ from that of the control group (the authors noted, however, that the control group was likely not representative of the broader community). Impulsivity, but not aggression, correlated with serotonergic function.

⁷Similar conclusions come from studies of borderline personality disorder, where serotonergic function has been related to impulsiveness and emotional reactivity (see Carver & Miller, 2006; Loeper & Paris, 2000).

M. C. Dolan and Anderson (2003) examined criminal offenders with diagnoses of personality disorders, relating response to fenfluramine challenge with scores on the Psychopathy Checklist. Lower serotonergic function did not predict overall psychopathy scores but did relate to scores on the impulsive–antisocial component, which combines ratings of irresponsibility, adolescent antisocial behavior, adult antisocial behavior, impulsiveness, and poor behavioral control.

There is also research linking impulsive aggression to brain activity that occurs in response to serotonin. New et al. (2002) tested control participants and persons with a history of impulsive aggression by PET scan under a serotonin releaser and under placebo. Scans revealed differences between groups in drug response. Patients showed less activation (vs. placebo) than did control participants in left medial posterior OFC, leading New et al. (2002) to conclude that impulsive aggression by the patients reflects lack of activity in areas involved in planning and long-term regulation of behavior. Consistent with these findings, the OFC has also been implicated in the control of aggressive impulses in other research (Dutton, 2002).

Raine and Yang (2006) reviewed a variety of imaging research relating to antisocial behavior. They concluded that the evidence indicates impairments of a number of brain areas in antisocial populations, including both dorsolateral and orbitofrontal regions of the PFC, amygdala, and anterior cingulate. This pattern suggests problems in the circuitry constraining emotion, consistent with a two-mode view.

Suicidality

Suicidal behavior has long been linked to serotonin, via serotonin metabolites in cerebrospinal fluid (e.g., Ågren, 1980). Over 1,000 studies have been conducted on personality traits and suicidality (Brezo, Paris, & Turecki, 2006). Although negative affective traits such as neuroticism and hopelessness are involved in development of suicidal ideation, a large body of research suggests that suicidal acts and completed suicides are reliably tied to impulsivity, particularly impulsive aggression (for a detailed review, see Brezo et al., 2006). Mann, Waternaux, Haas, and Malone (1999) suggested that impulsiveness in particular is a risk factor for suicide, rendering a person who has suicidal feelings more likely to act on them.

Cross-sectional studies suggest that lower cerebral-spinal fluid 5-HIAA (a serotonin metabolite) is related to greater lethality of suicide attempts (Placidi et al., 2001). Although prospective research is limited, several studies of high risk samples have now found that lower cerebrospinal levels of serotonin among persons hospitalized for an initial suicide attempt predict prospectively a three- to fourfold increase in risk of subsequent death by suicide (for a review, see Mann & Currier, 2007).

Section Summary

A good deal of evidence links low serotonergic function, assessed in various ways, to disorders and other kinds of problem behaviors that are characterized by deficient impulse control. Those problems often result in disruptive behavior, sometimes extending to impulsive violence, and sometimes extending to suicide. This pattern is consistent with the two-mode model, and with the idea that effortful control is necessary to restrain behaviors of this sort.

Personality Traits and Serotonergic Function

Much research has also examined relationships of serotonergic function to personality as measured by self-report, using several different procedures to assess serotonergic function. Some of this work has focused on qualities of personality pertaining to aggression and impulsiveness; other studies have examined a broader spectrum of personality qualities.

Trait Hostility and Impulsiveness

As noted in the preceding section, an extensive literature links serotonergic function to impulsive aggression. Hostility as a trait has also been related to low serotonergic function in nonclinical samples (Cleare & Bond, 1997; Depue, 1995; Netter, Hennig, & Rohrmann, 1999). Depue (1995) reported a more extensive array of associations, relating low serotonergic function (assessed by fenfluramine challenge) as well to the Control–Impulsivity facet scale from the Constraint factor of the Multidimensional Personality Questionnaire (Tellegen, 1985), the Aggression facet of the Multidimensional Personality Questionnaire's Negative Emotionality factor (but not other facets), two sensation seeking subscales, and several indices of impulsiveness. Depue (1995) reported that low serotonergic function related most strongly to hostility subscales that he characterized as reflecting impulsive, action-oriented aggression (irritability and assaultiveness), as opposed to more passive, cognitive forms (resentment and suspiciousness).

A more recent study (Hennig, Reuter, Netter, Burk, & Landt, 2005) yielded results that resemble these conceptually, by using both a drug challenge and a genetic method. The drug challenge was a single dose of SSRI, with the outcome being change in cortisol as an indirect indicator of serotonin function. High responders to the SSRI scored high on what Hennig et al. (2005) called aggressive hostility (marked by assaultiveness and negativism). What they called neurotic hostility (marked by irritability and resentment, but also guilt) did not relate to cortisol response. Hennig et al. (2005) also assessed the tryptophan hydroxylase gene, which expresses an enzyme that helps synthesize serotonin. Variations in that gene related to aggressive but not neurotic hostility. Hennig et al. (2005) noted that aggressive hostility was not related to a separate measure of neuroticism except for the neuroticism facet concerning aggression. They concluded that aggressive hostility resembles psychoticism (defined by impulsivity and lack of social connection), not neuroticism.⁸

Personality and Serotonergic Function Assessed by Drug Challenge

Studies have also related serotonergic function as reflected in responses to drug challenge to personality in community samples, assessing personality more broadly. Manuck et al. (1998) assessed prolactin responses to fenfluramine and related them to the Neuroticism–Extroversion–Openness Personality Inventory (NEO–PI), plus additional measures. Among men, blunted prolactin response (suggesting low serotonergic function) related to higher neuroticism (assessed by NEO–PI) and the Neuroticism facet Angry Hostility (other facets were not tested); blunted prolactin response also related to low conscientiousness (assessed by NEO–PI) and to separate measures of impulsiveness and life history of aggression. No effects emerged in the full sample of women (perhaps because of variation in menstrual cycle), but analyses limited to postmenopausal women yielded effects for conscientiousness and impulsiveness. The association with neuroticism was replicated in a larger sample by the same research group (Flory, Manuck, Matthews, & Muldoon, 2004), but in this case no tests were reported for conscientiousness or any facet of neuroticism.⁹

⁸A good deal of the evidence links low serotonergic function to traits related to negative emotions. There is far less evidence concerning serotonergic function and positive affects, and the evidence that does exist is mixed. Zald and Depue (2001) found an inverse relation between prolactin response to fenfluramine and ambient positive (and negative) feelings; Flory et al. (2004) found a positive relation between prolactin response to fenfluramine and positive feelings.

⁹Just as neuroticism has facets, so does conscientiousness, and there is reason to believe that some facets relate to serotonergic function but not others. This would compromise research on conscientiousness that did not test facets. Roberts, Chernyshenko, Stark, and Goldberg (2005) found six factors in multiple measures of conscientiousness, only moderately inter-correlated, which differed substantially in their relations to criterion measures. Indeed, in other research (Moon, 2001) conscientiousness was unrelated to a study outcome, but two facets had substantial and opposite relationships with the outcome. Just as there is merit in examining how facets of neuroticism relate to markers of serotonergic function, it also seems important to test facets of conscientiousness separately.

Several studies have related serotonergic function assessed by drug challenge to the Tridimensional Personality Questionnaire (TPQ; Cloninger, 1987), with mixed results. Two studies found links between low levels of harm avoidance and lower serotonergic function (Gerra et al., 2000; Hansenne & Ansseau, 1999). One found the opposite (Hennig et al., 2000). J. Evans, Platts, Lightman, and Nutt (2000) found no link from serotonergic function to any TPQ scale. Apart from this inconsistency in results, there are measurement issues concerning the TPQ that urge caution in interpreting findings relating to it (Carver & Miller, 2006; B. P. O'Connor, 2002).

SSRIs and Personality

A previous section described behavioral results from a slim literature on SSRI effects in nonclinical samples. Two studies from that literature also examined effects on personality self-reports. Knutson et al. (1998) found that 4 weeks of SSRI led to lower self-report scores on the Assaultiveness and Irritability scales of the Buss–Durkee Hostility Inventory (BDHI; Buss & Durkee, 1957). It is of interest that these are the same two subscales as Depue (1995) had earlier found to be related to low serotonergic function as assessed by drug challenge.

Another small study of nonclinical participants (Tse & Bond, 2001) examined effects of SSRI on an abbreviated version of the Temperament and Character Inventory (Cloninger, Svrakic, & Pryzbeck, 1993). SSRI led to an increase on a measure of self-direction (reflecting such qualities as responsibility, purposefulness, and resourcefulness) but produced no other difference.

Personality and the Serotonin Transporter Polymorphism

There is also a substantial literature on the 5-HTTLPR polymorphism and personality as assessed by broad-ranging self-report inventories, which permitted investigation of diverse possible associations. This work began with several large-scale studies in which the data were examined thoroughly. Lesch et al. (1996) studied a sample of 505 normal participants, mostly men. They found that the short allele (identified with low serotonergic function) related positively to neuroticism and inversely to agreeableness (both assessed by NEO–PI–Revised [NEO–PI–R]). Not all facets of neuroticism related to the polymorphism, however. In analysis of Neuroticism facet scales, the polymorphism related most strongly to Angry Hostility, Depression, and Impulsiveness scales; a link with the Anxiety scale was barely significant and there was no link to the Vulnerability scale.

The Impulsiveness facet in neuroticism requires further clarification. Its inclusion in that trait domain scale reflects the fact that negative feelings, themselves the hallmark of neuroticism, sometimes lead to impulsive actions such as overindulgence (which is well represented in the items). There is some question as to whether impulsivity as assessed by this scale is a good representative of impulsivity as a construct.

A second major study examined 397 participants, mostly women (Greenberg et al., 2000). Again, the 5-HTTLPR polymorphism related to both neuroticism and agreeableness, with an additional weak but significant association for conscientiousness. Analysis of Neuroticism facets revealed the strongest relations again for Angry Hostility and Depression scales. Associations with Anxiety, Impulsiveness, and Vulnerability scales were all nonsignificant.

A third large-scale study was reported by Sen, Villafuerte, et al. (2004). This sample of 419 participants yielded an association of the polymorphism with NEO–PI neuroticism and a weaker but significant association with agreeableness. As in the other studies, the facet of neuroticism most strongly related to the polymorphism was the Depression scale. The short

allele related significantly to Anxiety and Self-Consciousness scores in this sample, to the Angry Hostility scores marginally ($p = .052$), and not to Impulsiveness or Vulnerability scores.

Many smaller-scale studies have been conducted, with varying outcomes. There have by now been several meta-analyses of the relationship of variation in 5-HTTLPR to neuroticism. In one metaanalysis (Sen, Burmeister, & Ghosh, 2004), with all studies combined there was no association. When the studies were separated by personality measure, 5-HTTLPR related to NEO-PI-R Neuroticism, but not Temperament and Character Inventory (TCI) or TPQ Harm Avoidance. Schinka, Busch, and Robichaux-Kenne (2004) reached a similar conclusion from their meta-analysis. Munafò, Clark, and Flint (2005a) found the opposite: a significant relation with TPQ/TCI Harm Avoidance, but not with NEO-PI Neuroticism. However, reanalysis of their data using the procedures used in the other meta-analyses reversed the pattern, with an association now emerging for NEO-PI Neuroticism (Munafò, Clark, & Flint, 2005b).

Unfortunately, these meta-analyses have two serious problems. First, none examined neuroticism facets. Because facet analyses were very informative in the earlier large-scale studies, this appears a serious omission. Second, traits other than neuroticism were disregarded. Thus, there was no test of a link from genetic difference to agreeableness, which had been found in all three large-scale studies (Greenberg et al., 2000; Lesch et al., 1996; Sen, Villafuerte, et al., 2004). The pattern presented by these three large-scale studies as a group is a picture relating low serotonergic function to disagreeable, dysphoric surliness.

Schmitz, Hennig, Kuepper, and Reuter (2007) took a more fine-grained approach to examining neuroticism in genetic research. They collected data on three measures of neuroticism and closely related traits (though again other traits were not considered), and tested the measures item by item against the polymorphism. Twelve items significantly related to genotype; item content focused on depression and stress reactivity. Consistent with that finding, the 5-HTTLPR polymorphism has been linked to differences in cortisol response to a standard laboratory stressor (Gotlib, Joormann, Minor, & Hallmayer, 2008).

Section Summary

As a group, the bulk of the findings reviewed in this section appear consistent with the idea that serotonergic function relates to a group of traits with agreeableness and constraint at one end and impulsivity, along with aggressive and depressive tendencies, at the other end (Spoont, 1992). The pattern appears consistent with the view that serotonergic pathways are involved in impulse control (Depue, 1995; Depue & Collins, 1999; Depue & Spoont, 1986; Manuck, Flory, Muldoon, & Ferrell, 2003; Soubrié, 1986; Spoont, 1992; Zuckerman, 2005), particularly impulses reflecting strong negative emotions. High serotonergic function appears to relate to consideration of the future consequences of one's behavior (conscientiousness) and to positive social connection (agreeableness, cooperation, affiliation). This includes social potency (even dominance), but a positive social engagement rather than dominance by force (cf. Manuck et al., 2006). The literature reviewed here is quite different from that reviewed by Spoont (1992), but the conclusions from the two reviews are in many ways quite similar.¹⁰

Interpretation in Terms of Two-Mode Models

The preceding sections related serotonergic function to cognitive processing of emotional stimuli, overt behaviors such as impulsiveness and aggression, disorders involving poor

¹⁰An early view of serotonergic function related it specifically to the experience of anxiety (Handley, 1995). Indeed, it remains common to refer to "anxiety-related traits" in discussing this literature (e.g., George et al., 2001; Hennig et al., 2000; Lesch & Canli, 2006; Munafò et al., 2005a; Schinka, 2005; Schinka et al., 2004). The tendency to invoke anxiety doubtlessly reflects observed associations of serotonergic function with global measures of neuroticism. However, it seems clear from this review that anxiety is not the only emotion that must be considered here (see also Panksepp & Cox, 1986; Soubrié, 1986).

impulse control, and a range of personality traits. This evidence appears amenable to interpretation in terms of two-mode models, as follows.

Relations between low serotonergic function and hostile and depressive trait tendencies can be seen as reflecting frequent intrusion of strong emotions; as noted earlier, this has been argued to characterize reflexive-mode processing. Frequent strong emotions would also be consistent with brain mechanisms reviewed earlier that yield a relative lack of executive control over emotional reactions. Relations between low serotonergic function and impulsive aggression (or other problematic impulsive actions) appear to reflect a lack of effortful control by the deliberative system over spontaneous action (see also Wilkowski & Robinson, 2008, regarding the role of effortful control in aggression).

Relations between serotonergic function and agreeableness (found frequently) and conscientious and related qualities (found less frequently) both concern qualities of personality that are addressed by two-mode models. Both sets of associations imply that high serotonergic functioning permits a relatively broader perspective on behavioral decisions—broader than the impulsive grabbing of incentives or fleeing from threats. Conscientiousness entails a broadened temporal perspective: taking future contingencies into account in one's actions. Agreeableness entails a broadened social perspective: taking the needs of others into account in one's actions. Indeed, consistent with the view that these traits reflect operation of the reflective mode, some argue that both of these traits have their temperamental origins in effortful control (Ahadi & Rothbart, 1994; Caspi & Shiner, 2006; Jensen-Campbell et al., 2002).

In further support of the idea that taking future contingencies into account entails effortful control, trait self-control (a closely related concept) has been linked to higher grades (Tangney, Baumeister, & Boone, 2004; Wolfe & Johnson, 1995), less procrastination (Steel, 2007), and less criminality (Krueger, 2002). In further support of the idea that taking into account the needs of others entails effortful control, trait self-control has been linked to accommodative behaviors in intimate relationships, which helps to sustain those relationships (Finkel & Campbell, 2001), an effect that held even when controlling for level of commitment to the relationship. Similarly, trait self-control has been linked to relationship satisfaction and relationship longevity among both romantic partners and best friends (Vohs, Lasaleta, & Fennis, in press).

As noted earlier, Hennig et al. (2005) suggested that the low serotonergic pattern resembles psychoticism, a trait characterized by impulsive social disconnection. Impulsive social disconnection seems an apt characterization of the combination of low agreeableness and low conscientiousness, both of which have been linked to psychopathy (Derefinko & Lynam, 2006).

Serotonin and Depression Vulnerability

Previous sections of the article relate low serotonin function to behavioral qualities such as reactivity to emotional information, poor impulse control, and uncooperativeness. A section on personality related serotonin function to traits that are conceptually similar to those behaviors, traits such as aggressiveness, impulsiveness, and disagreeableness. Correlates of low serotonin function from these various sources appear to share the property of poor control over (usually emotion-driven) responding. In contrast, this section turns away from impulsive behavior to what appears to be a quite different quality: depression.

Some information on depression has already been reviewed: The section on traits included evidence linking low serotonergic function to depressive tendencies. That evidence came from correlations with broad personality inventories, however, and is more suggestive than definitive. This section reviews research that makes a case for a link between low serotonergic

function and the more explicit assessment of elevated depressive symptoms and diagnosis of depression.

Depression and Response to Drug Challenge and Manipulated Serotonin Availability

Several studies have found blunted prolactin response to fenfluramine among depressed persons (Flory et al., 1998; Lichtenberg et al., 1992; O'Keane & Dinan, 1991; Siever, Murphy, Slater, de la Vega, & Lipper, 1984). There is also evidence of a blunted response to serotonin challenge (including citalopram) among persons with a history of depression who are not currently depressed and are not medicated (Bhagwagar, Whale, & Cowen, 2002; Flory et al., 1998). This evidence is not entirely consistent, however. Occasional studies have found elevated rather than blunted responses among depressed persons, though generally in small samples (e.g., Strickland et al., 2002).

Evidence of the involvement of serotonin in depression also comes from studies of acute tryptophan depletion (temporarily reducing available serotonin) among persons who were formerly depressed (see Moore et al., 2000, for a review). This procedure causes a temporary increase in depressive symptoms among some formerly depressed persons (Booij & Van der Does, 2007; Booij et al., 2002; Booij, Van der Does, et al., 2006; Delgado, Charney, Price, & Aghajanian, 1990; Delgado et al., 1999; Lam et al., 1996; Moreno et al., 1999; Munafò, Hayward, & Harmer, 2006; Neumeister et al., 2006; Neumeister et al., 2004; K. A. Smith, Clifford, Hockney, Clark, & Cowen, 1997; but see Leyton et al., 1997, for a nonreplication). In one case, tryptophan depletion induced cognitive deficits on the Mini-Mental State Examination (Porter et al., 2005). In other studies, tryptophan depletion produced memory impairments (Sambeth et al., 2007). In another study, tryptophan depletion increased cerebral glucose metabolism in the OFC cortex, anterior and posterior cingulate cortices, and ventral striatum in remitted participants but not in control participants (Neumeister et al., 2004). The return of depression symptoms has been found among remitted persons who are medication free (Neumeister et al., 2006) as well as among those taking SSRIs (Delgado et al., 1999).

Persons whose first-degree relatives have a history of major depression also appear to be at elevated risk of negative affect after tryptophan depletion, compared with those with no family history of major depression (Benkelfat et al., 1994; Klaassen et al., 1999; Neumeister et al., 2002). Depression family history has been shown to increase risk of vegetative symptoms of depression after tryptophan depletion (Klaassen et al., 1999).

Serotonin Transporter Polymorphism and Depression

Early studies of the serotonin transporter gene looked for a direct (i.e., unmoderated) link between variations in 5-HTTLPR and clinical depression. Several studies found such relations (Collier et al., 1996; Hoefgen et al., 2005), but others did not. A meta-analysis of earlier studies (Anguelova, Benkelfat, & Turecki, 2003) reached a negative conclusion. Hoefgen et al. (2005) noted that that meta-analysis involved samples of great diversity, in contrast to their homogenous sample. They suggested that obtaining small effects (such as they found) may require substantial samples and also less variability attributable to sources other than the polymorphism.

A number of other studies have investigated the possibility of an interaction between this polymorphism and adverse experience (a diathesis–stress approach). For example, Caspi et al. (2003) reported that this polymorphism interacted with stressful life events to predict depressive symptoms and depression diagnosis: Negative life events had an adverse effect on those carrying at least one short allele, but not among those with two long alleles. This suggests that the short allele marks a genetic vulnerability: A factor present before onset, which interacts with triggers to increase the risk of onset (Monroe & Simmons, 1991). Without adversity, this

genetic makeup did not yield depression, but this genetic makeup rendered people reactive to adversity.

A number of other studies have yielded similar results. These studies have examined children (Kaufman et al., 2004, 2006), adolescents (Eley et al., 2004), schoolteachers (Wilhelm et al., 2006), young adults (Kendler, Kuhn, Prescott, & Riley, 2005), and adults exposed to hurricanes (Kilpatrick et al., 2007). In all of those cases, the short allele of the serotonin transporter polymorphism was associated with elevated risk of depression, but only in combination with recent stressful life events or histories of maltreatment (and low social support, in the case of Kilpatrick et al., 2007).

Another recent study of Korean elders (Kim et al., 2007) found an interaction between serotonin transporter gene and stressful life events, such that carriers of the short allele who had had higher levels of adversity over the preceding year were more likely to be depressed. At 2.5 year follow-up of those not depressed at baseline, those with two short alleles who had experienced more negative events in the preceding year were more likely to be depressed.

Other studies have yielded partial replications of this pattern, but with qualifications. One study examined a large general population sample (Grabe et al., 2005), finding effects among women, but not men, and in interaction with other variables. In this case, the short allele was related to reports of both mental and physical distress among women who were unemployed, and (independently) among those who had from one to three chronic diseases. Another study of adolescents found that the short allele predicted depressive symptoms in interaction with environmental stress among female participants but had the opposite effect among male participants (Sjöberg et al., 2006).

S. E. Taylor et al. (2006) found an interaction on self-reported depressive symptoms: Among persons reporting adverse early family environment, those homozygous for the short allele (but not heterozygous) reported elevated depressive symptoms. Among persons reporting particularly benign early environment, those homozygous for the short allele reported the lowest depressive symptoms. S. E. Taylor et al. (2006) concluded that the short-short genotype is not so much a risk factor for depression as it is a magnifier of the impact of the environment on the person's well-being. If the person is exposed to great adversity, the result may be enhanced symptoms of depression. If the person is instead exposed to a very supportive environment, the result may be a positive emotional life.

Jacobs et al. (2006) examined a large sample of women over a 15-month span, using self-reports of depressive symptoms as the outcome. They found an interaction of genotype with neuroticism, such that the presence of the short allele appeared to enhance the relation of neuroticism with depressive symptoms. This effect was independent of adverse life events, though there also was an interaction of neuroticism with adverse life events. The authors concluded that the polymorphism moderates the way in which persons respond to their daily life experiences. That is, neuroticism is associated with more emotional responses to adversity, but the impact of neuroticism on that emotional response is greater if the person carries the short allele of the serotonin transporter gene. Importantly, this suggests that low serotonergic function enhances stress reactivity even within a given level of neuroticism.

Zalsman et al. (2006) examined 191 persons with mood disorder and 125 healthy control participants, in a triallelic design (grouping the less expressive long allele with the short allele). They found that genotype did not distinguish patients from control participants, but among patients it did distinguish those with more severe depressive symptoms. The latter finding emerged both as a main effect of genotype and an interaction of genotype with life stresses.

Not all attempts to replicate the diathesis–stress effect have been successful. For example, in a sample of 1,206 twins, Gillespie, Whitfield, Williams, Heath, and Martin (2005) found that stress predicted major depressive disorder, but genotype played no role. Similar results were obtained by Surtees et al. (2005). Gillespie et al. pointed out that earlier demonstrations of the interaction had generally used young adults and that only 20% of their sample was below age 30. They speculated that genetic differences may contribute to vulnerability to different degrees at different ages. However, that interpretation does not fit the findings of Grabe et al. (2005) or Kim et al. (2007) described earlier.

Although most genetic research pertaining to serotonergic function and depression has examined the 5-HTTLPR polymorphism, there have also been a more limited number of studies of other polymorphisms. An example is a study of the tryptophan hydroxylase 1 gene (Jokela, Rääkkönen, Lehtimäki, Rontu, & Keltikangas-Järvinen, 2007). As noted earlier, tryptophan is involved in the synthesis of serotonin, and genes influencing tryptophan would ultimately influence serotonin. This study found that depressive symptoms were more common, both cross-sectionally and prospectively, in persons with a combination of low social support and the A allele of the gene. Another example is a study of the serotonin receptor 2A gene (Jokela, Keltikangas-Järvinen, et al., 2007). In this study, the lowest symptoms of depression were among people with the combination of the T genotype of the T102C polymorphism and high maternal nurturance.

A methodologically sophisticated study that combined diagnostic status, the serotonin transporter polymorphism, and tryptophan depletion was conducted by Neumeister et al. (2006). They gathered PET data from persons with remitted recurrent major depressive disorder who had not taken antidepressants in at least 3 months (plus never-depressed control participants). A significant interaction emerged among depression status, genotype, and tryptophan depletion predicting neural responses. Persons with no depression history generally did not have strong responses to tryptophan depletion. Among persons with remitted depression, however, those with the short allele displayed greater activation of the hippocampus, the amygdala, and ACC in response to tryptophan depletion.

Summary of Association Research

The results outlined in the previous sections can be summarized as follows. Despite some nonreplications and qualifications, research with diverse methodologies links low serotonergic function to elevated symptoms of depression and to clinical depression. Depressed persons have displayed a blunted response to drug challenge (compared with other persons), and manipulations that reduce available serotonin have induced symptoms among previously depressed persons. Similarly, the short allele of the 5-HTTLPR has been linked to elevated risk of depression, though that link has often depended on the additional presence of adverse life experiences. Thus, there seems to be solid evidence that connects low serotonergic function to depression.

Gender

An issue that should be mentioned in this context concerns the gender difference in depression. Women are roughly twice as likely as men to experience clinical depression in their lifetimes (Kessler et al., 2003). Heritability estimates for depression are also higher among women than among men (Kendler, Gatz, Gardner, & Pedersen, 2006). This suggests that at least some of the gender difference in depression risk may reflect biological vulnerabilities.

In light of this difference, it is of considerable interest that gender differences also exist pertaining to serotonergic function. For example, there is considerable evidence that women are more susceptible than are men to negative mood changes after acute tryptophan depletion

(Booij & Van der Does, 2007; Booij et al., 2002; Ellenbogen, Young, Dean, Palmour, & Benkelfat, 1996; Moreno, McGahuey, Freeman, & Delgado, 2006; K. A. Smith et al., 1997; Walderhaug et al., 2007). Also, as noted earlier, two studies relating the short allele of the 5-HTTLPR polymorphism in interaction with stress to depressive symptoms found this association among women but not men (Grabe et al., 2005; Sjöberg et al., 2006). A recent analysis pooling nine studies found that tryptophan depletion creates greater memory impairments among women than among men (Sambeth et al., 2007).

This is not to say that the short allele has behavioral consequences only among women. However, the pattern of consequences may differ by gender. For example, one study has found that the short allele predicts greater aggression in response to stress among men but not women (Verona, Joiner, Johnson, & Bender, 2006). Another found that the short allele of the serotonin transporter gene interacted with adverse childhood environment among men to produce violent adult behavior (Reif et al., 2007). Perhaps the effects of low serotonin function depend partly on interactions of serotonin with sex hormones (L. H. O'Connor & Fischette, 1986).

Serotonin and Symptom Profiles

Another important question is whether serotonergic function relates selectively to some particular aspect of depressive symptomatology or whether the relationship is more general. Several studies have provided evidence on this question. In research on persons who have recovered from depression, acute tryptophan depletion has been found to have an impact on both interview and self-report measures of anxiety and depressive symptoms as well as on self-reported somatic symptoms of depression (Booij, Swenne, et al., 2006; Spillmann et al., 2001). Other studies have found that acute tryptophan depletion produces effects of comparable size for mood and vegetative symptoms (Riedel, Klaassen, & Schmitt, 2002) and for REM sleep deficits that are typically associated with depression (Moore et al., 1998).

Evidence linking suicidality to low serotonergic function was described earlier. Some of the studies that related depression to the serotonin transporter polymorphism also examined suicidality. Caspi et al. (2003) found that the short allele was linked to higher risk of suicidality in interaction with life stressors. Zalsman et al. (2006) found that genotype did not relate to suicidality within the depressed group, but they did not test for an interaction with life events, nor did they compare the depressed group with the control group.

One might also ask whether there are differences between major depressive disorder and depression associated with bipolar disorder. Recent research (Sher et al., 2003; Sobczak, Honig, van Duinen, & Riedel, 2002) suggests that hormonal responses to fenfluramine and to tryptophan are blunted to a similar degree among people with unipolar and bipolar depression. Unaffected family members of persons with bipolar disorder have also shown atypical cognitive (Sobczak, Riedel, Booij, Het Rot, & Deutz, 2002) and mood responses (Quintin et al., 2001) to serotonin challenge; this suggests that deficits in serotonergic function in bipolar disorder do not reflect medication effects or scars of previous episodes.

In sum, there is substantial evidence of the involvement of serotonergic function as a contributor to risk of depression. The possibility that serotonergic function relates selectively to some symptoms but not others among depression-vulnerable persons is intriguing. However, support for that possibility is generally lacking.

Differentiating Vulnerabilities: Impulse Problems Versus Depression

An earlier section linked low serotonergic function to behavioral problems in which a salient feature was poor impulse control. Given various evidence reviewed prior to that section, it is not surprising that low serotonergic function relates to disorders involving impulsiveness. That

is, in diverse contexts, markers of low serotonergic function have been linked to impulse expression, particularly impulsive reactions to emotion cues.

It is less obvious at first glance why low serotonergic function would be a vulnerability factor for depression. Although depression has sometimes been linked to impulsive qualities (Peluso et al., 2007), it is more frequently associated with lethargy, an absence of behavioral engagement. How can we account for this very substantial difference in presentation?

Earlier in the article, the balance of reflexive to deliberative control over behavior was characterized in terms of relative responsiveness to salient cues (as argued by Spont, 1992). Deficits in deliberative control enhance responsiveness to cues of the moment, particularly emotion cues. What kind of behavior follows from that responsiveness, however, depends on what the salient cues are. That, in turn, depends partly on the person's emotional and motivational state (or sensitivity). Thus, low serotonergic function and the resulting deficits in effortful control may have divergent effects in people who vary systematically in other ways.

Interaction Between Systems

What other systematic variation might interact with serotonin function to yield such a divergence? A prime candidate is variation in the degree of sensitivity, or engagement, of a general incentive approach system. When poor executive oversight is combined with moderately high incentive sensitivity (a fairly reactive approach system), the result is impulsiveness. When poor executive oversight is combined with low incentive sensitivity (a nonreactive approach system), the result is inaction: lack of effort toward potential rewards. In both cases, the effects of variation in level of incentive sensitivity (high and low, respectively) are amplified by the absence of effortful override (Figure 2, earlier).

In the case of depression vulnerability, the intersection of low serotonergic function and nonreactive incentive system could create difficulties in two ways. First, a lack of incentive sensitivity means that the person is not strongly motivated to approach potentially rewarding contexts. A relative lack of effortful control will amplify this problem, such that the depression-vulnerable person will have greater difficulty overriding this lack of motivation so as to engage in actions. This combination thus should yield apathy, passivity, and fatigue, which characterize many cases of depression. Second, if a deficit in executive processes leaves the amygdala relatively uncontrolled and thus hyperactive, the salience of emotional associations should also be enhanced, which presumably are relatively negative in the person vulnerable to depression. Thus, low serotonergic function would have two adverse effects. It would hamper the ability to keep negative emotions at a distance. It would also hamper the ability to engage in effortful pursuit of incentives (Salamone, Correa, Farrar, & Mingote, 2007).

Blunted Incentive System in Depression

Evidence that depression is associated with a blunted approach system comes from several sources. One of them is an earlier literature using EEG to examine anterior cortical asymmetry and motivational processes. This technique has far poorer spatial resolution than do newer imaging techniques, and thus it is far less specific about what areas are activated. However, it has been used to make a case concerning partial localization of approach and avoidance systems in left and right hemispheric areas, respectively (e.g., Davidson, 1998).

Previously depressed (Henriques & Davidson, 1990) and clinically depressed persons (Henriques & Davidson, 1991) have been found to have lower activation in left anterior cortical areas than do nondepressed persons, with no difference in right anterior activation. The lower activation was in areas that previous research had related to approach sensitivities (e.g., Coan & Allen, 2003; Harmon-Jones & Allen, 1997; Sutton & Davidson, 1997). This general pattern

regarding depression has been replicated a number of times (J. J. Allen, Iacono, Depue, & Arbisi, 1993; Gotlib, Ranganath, & Rosenfeld, 1998), though not all attempts have been successful (Reid, Duke, & Allen, 1998). These findings as a group suggest that depression relates to low engagement of areas pertaining to approach. One recent study has even linked early-onset depression to a deficit in left prefrontal responsiveness to a task that had been designed specifically to engage approach motivation (Shankman, Klein, Tenke, & Bruder, 2007).

Behavioral research also suggests that depression relates to blunted incentive sensitivity. For example, depressed persons were found to be less responsive to reward (though not to punishment) than were nondepressed persons (Henriques & Davidson, 2000; Henriques, Glowacki, & Davidson, 1994; see also Depue & Iacono, 1989). There is also evidence that dysphoric persons engage more effort at easy task levels than do nondysphoric persons, as though they overestimated the effort that is required by the task, but that dysphoric persons are more likely to disengage effort when the task actually is demanding (Brinkmann & Gendolla, 2008).

Other evidence relates self-reports of low incentive sensitivity to depression (Campbell-Sills, Liverant, & Brown, 2004; Pinto-Meza et al., 2006). Indeed, three separate studies have found that self-reports of low incentive sensitivity predicted worse course of depression over time (Campbell-Sills et al., 2004; Kasch, Rottenberg, Arnow, & Gotlib, 2002; McFarland, Shankman, Tenke, Bruder, & Klein, 2006).

Blunted approach motivation may also be reflected in low dopaminergic function. Dopaminergic pathways are believed to be critical in the engagement of goal-directed effort (Farrar et al., 2007; Salamone et al., 2007; Salamone, Correa, Mingote, & Weber, 2005; Salamone, Correa, Mingote, Weber, & Farrar, 2006). A weakly functioning dopaminergic incentive system yields less “wanting” for appetitive outcomes (Berridge, 2007) and less engagement of effort in pursuit of them (Salamone et al., 2005, 2006, 2007). A recent comprehensive review reported a range of evidence for deficits in the function of dopamine among depressed persons, drawing from pharmacological studies, genetic studies, and dopamine challenge studies (Dunlop & Nemeroff, 2007).

Animal research has long implicated subcortical reward pathways in behaviors that mimic depressive symptoms, such as diminished motivation, apathy, and poor appetite (Nestler & Carlezon, 2006). Some brain studies in humans also link depression to deficits in subcortical regions involved in reward sensitivity, such as the caudate. For example, in a recent fMRI study, children diagnosed with depression displayed less activation of the caudate throughout a task involving reward, compared with children without depression (Forbes, Fox, Cohn, Galles, & Kovacs, 2006). Adults with depression have been found to show hyperreactivity of the caudate and the putamen to an amphetamine challenge that was designed to intensify dopaminergic activity (Tremblay et al., 2005); this pattern is consistent with there being a baseline deficit in dopaminergic activity.

Incentive System in Impulsive Disorders

What about impulse-related problems? A very substantial literature has addressed the role of dopamine and approach motivation (in interaction with serotonin) in externalizing disorders. Review of that literature, which includes studies using genetic and psychophysiological indices of various kinds, is well beyond the scope of this article. It is worth noting, however, that there is evidence linking externalizing disorders to elevated approach tendencies.

The concepts of reactive undercontrol (Eisenberg et al., 2004) and motivational inhibition (Nigg, 2001, 2003) are also relevant here. As noted earlier, evidence links reactive undercontrol

to externalizing disorders in childhood (Rothbart, Ellis, & Posner, 2004; Valiente et al., 2003). Similarly, Nigg has held that the core issue in conduct disorder (and by implication psychopathy as well) is lack of a sufficiently reactive threat system. In principle, however, “sufficiently reactive” means only that the threat system is more reactive than is the incentive system, reactive enough to hold it in check. This could mean an insensitive threat system in combination with no special sensitivity of approach; it could also mean a relatively engaged incentive system with no special deficit of avoidance.

There is at least some evidence of both a weak threat system and a strong incentive system in psychopathy. In one study, various measures of psychopathy correlated with lower Anxiety facet scores on the NEO-PI-R and with higher scores on Assertiveness and excitement seeking facets of Extraversion, both reflecting incentive pursuit (Derefinko & Lynam, 2006). Another study relating self-reports of incentive and threat sensitivities to diverse externalizing symptoms found that self-reports of high reward sensitivity predicted primary and secondary psychopathy and impulsive ADHD symptoms (Hundt, Kimbrel, Mitchell, & Nelson-Gray, 2008). Low threat sensitivity predicted secondary psychopathy and inattentive ADHD symptoms.

Section Summary

The question of how to interpret the very different manifestations of low serotonergic function in impulsive disorders and depression prompts the search for an interaction. There is support for such an interaction: between the serotonergic system (and associated executive deficits) and a general system of incentive sensitivity, in the form of evidence of a blunted approach system in depression.

Two-Mode View of Depression: Additional Evidence Concerning Brain Function

The link between two-mode models and depression has been made here in terms of the involvement of serotonergic function in brain functions that are consistent with the two-mode models, and the involvement of serotonergic function in risk of depression. Evidence from other sources further supports that link. This section reviews one source: neurobiological research on depression. Specifically, evidence suggests involvement in depression of brain areas that are strongly influenced by serotonin function.

Elevated Amygdala Activity in Depression

Findings from both structural and functional imaging studies have indicated that depression involves hyperactivity in brain areas that have been related to the reflexive mode (Mayberg, 1997, 2002). More specifically, in various imaging studies, depression has been related to elevations in blood flow, blood oxygenation, and glucose metabolism in the amygdala (Drevets et al., 1992, 2002). Indeed, amygdala activation has been found to correlate positively with depression severity (Abercrombie et al., 1998; Drevets et al., 1992). Amygdala activation normalizes following antidepressant treatment (Sheline et al., 2001).

Depression has also been related to heightened amygdala responses to fearful or negative faces (e.g., Breiter et al., 1996), a pattern that seems to diminish with antidepressants (Drevets, 2001; Sheline et al., 2001; for an extended discussion, see Davidson, Pizzagalli, Nitschke, & Putnam, 2002). Similarly, processing of negative words has been related to greater amygdala activation among depressed persons than among control participants (Siegle et al., 2002). Finally, persons who report being ruminators, a quality that renders them vulnerable to depression (Nolen-Hoeksema, 2000; Nolen-Hoeksema, Morrow, & Fredrickson, 1993), show

greater and longer amygdala activation when asked to temporarily increase their negative affect, compared with other people (Ray et al., 2005; Siegle et al., 2002).

Hypoactivation in Areas of PFC in Depression

Amygdala hyperactivity is one potential source of problems relevant to depression, by rendering existing negative emotions more salient (Spoont, 1992). A complementary source of problems, suggested by literatures reviewed earlier, is a link between depression and low activity in parts of the PFC believed to regulate affect by inhibiting activation in subcortical areas or activation in ventral parts of the PFC. Evidence fits this picture, as well (Levin, Heller, Mohanty, Herrington, & Miller, 2007; Mayberg, 2004).

A number of studies have found that dorsolateral PFC activation in depressed persons is low, both at baseline and during tasks (e.g., Baxter et al., 1989; Bench, Friston, Brown, Frackowiak, & Dolan, 1993; Davidson, 2000; Drevets, 1998; Siegle et al., 2002), while activation in ventral PFC and OFC are high (R. J. Dolan, Bench, Brown, Scott, & Frackowiak, 1994; R. J. Dolan et al., 1993). Antidepressant medication normalizes both of these differences (Kennedy et al., 2001; Mayberg, 1997). Other studies have found abnormally low cerebral blood flow and metabolism among depressed persons in areas of the lateral and dorsolateral PFC and the ACC (for reviews, see Drevets, 1999, 2000; Drevets & Raichle, 1998). These functional differences may be due to structural abnormalities associated with depression, such as lower neuron and glial count and size (e.g., Cotter et al., 2002; Rajkowska, 2000).

The key point here is the potential impact of this pattern on the affect-regulatory corticolimbic circuitry as a whole. Low activation in regions of the PFC that either project to subcortical structures, or project to other parts of the PFC that play an important role in inhibiting activity in subcortical regions, may reduce the constraining influence on those structures. The result can be a prolonged activation of the subcortical areas in response to challenges (cf. Siegle et al., 2002). This may be expressed as sustained negative affect. Indeed, Mayberg (1997, 2003) has proposed a model of depression involving disruption of corticolimbic pathways, based on evidence of abnormalities in these pathways among depressed patients and changes in them after treatment.

Section Summary

Neurobiological studies suggest that several aspects of brain function which are influenced by serotonergic function may also relate to depression. The effects include amygdala hyperreactivity and diminished functionality of specific prefrontal areas. This evidence suggests multiple processes: Depression likely involves subcortical functions yielding high negative emotion. Depression may also involve deficits in executive control over these systems, minimizing the likelihood that they will be overridden.

Two-Mode View of Depression: Additional Behavioral Evidence

The perspective of two-mode models also provides a different angle on some of the behavioral phenomena known to be related to depression. From this perspective, people who are vulnerable to depression should be deficient in executive control of their actions or emotions (or aspects thereof). Additional evidence concerning psychological and behavioral manifestations of depression supports the plausibility of that view. Indeed, the idea that depression can be usefully viewed from a two-mode framework has already appeared in several places (Barrett et al., 2004; Beevers, 2005; Dalgleish et al., 2007; Haefel et al., 2007).

Implicit System and Vulnerability

Beevers (2005) proposed a two-mode model of depression vulnerability that assumes many of the same conceptual underpinnings as we suggest here but that has a rather different focus. Beevers focused primarily on how the implicit system might tend to induce and maintain depression. He argued that persons with many negative associations in the implicit self therefore have a “pattern-completion” mechanism (a reflexive, connectionist mechanism) that is negatively biased when processing information about the self. When experiences are assimilated to this schema, they evoke the negative feelings that are linked to these negative associations. For that reason, these people are likely to experience negative feelings more frequently than are other people.

Several findings from other sources are consistent with that position. For example, negative responses on implicit measures of self-constructs did better than responses on explicit measures at predicting negative emotional reactions to laboratory stressors (Haefel et al., 2007). Similarly, Conner and Barrett (2005) have confirmed that people who have negative implicit self-concepts experience more negative affect in daily life than do people whose implicit self-concepts are less negative.

Indeed, negative associations in the implicit self can multiply over time, if the person develops an automatic tendency to think negatively about the self (Verplanken, Friborg, Wang, Trafimow, & Woolf, 2007). Such a tendency leads to further encoding of negative associations. Reports of habitual negative thinking about the self have been shown to relate to lower implicit self-esteem and to predict variance in depressive symptoms over and above that predicted by explicit negative content of thoughts about the self (Verplanken et al., 2007).

Beevers (2005) argued that, because of the automatic pattern-completion tendency, people who are depressed must engage in an effortful (reflective) corrective process to counter the negative associations. If the effortful process fails to occur, the implicit self maintains control over experience. As experiences continue to be assimilated to the negative implicit self, negative affect continues to be evoked and depression becomes more likely. Maintenance of an existing depression also becomes more likely.

Deficits in Effortful Control

Effortful processing requires both a reason to do so and the capacity to do so (Barrett et al., 2004). With respect to the former, Beevers (2005) held that a negative association triggers an effortful correction only if the association is unexpected, and thus calls for attentional resources to sort things out. However, a negative association would be unexpected only if the person’s explicit self-view was positive. Inasmuch as depressed people typically describe themselves negatively on explicit self-view indices (Abramson et al., 2002), a negative implicit association would generally not violate their expectancies and would not trigger effortful override. Thus, a prompt for effortful processing may often be lacking.¹¹

Other views of depression also emphasize the idea of cognitive control deficits. Hertel (2000), for example, viewed negative thoughts, biased processing of emotional information, and ruminative thinking in depression as automatic patterns activated by negative mood states.

¹¹ Authors differ in where conceptually to assign the risk (and why). Haefel et al. (2007) argued that explicit interpretations are the ultimate risk factor, because they represent the final interpretation of the event. Thus, they suggested that people with positive implicit self-construal can be vulnerable if they have negative explicit self-construal. Beevers (2005) viewed the negative implicit self-construal as the real culprit, but its impact is great only if not overridden. On the other hand, he saw the override as unlikely if the explicit self-construal is negative. Thus, as a practical matter, those conceptions differ only with respect to the case of a positive implicit self and a negative explicit self. Our position is that persons who are vulnerable to depression are impaired in the ability to readily evoke the reflective viewpoint that the explicit self-construal depends on. Thus the positive self-construal is unlikely to be brought to bear. Our view therefore would seem somewhat closer to that of Beevers (2005) than to that of Haefel et al. (2007).

Those patterns dominate unless overridden by effortful cognitive control (e.g., Hertel & Rude, 1991).

Deficits in effortful cognitive control can yield diverse outcomes. In some cases, it can keep the person from moving on. For example, depression has been related to impaired shift in behavior after error feedback (Compton et al., 2008; Holmes & Pizzagalli, 2007). Similarly, there is evidence that depressive rumination interferes with disengaging from previously relevant but no longer relevant information, as in task switching (Whitmer & Banich, 2007).

In other cases, deficits in effortful cognitive control can keep the person from remaining task focused. Thus, there is also broad evidence of mental restlessness and distractibility among depressed persons. Smallwood and his colleagues have repeatedly found that depression relates to mind wandering across diverse tasks (for a review, see Smallwood, O'Connor, Sudberry, & Obonsawin, 2007; see also Smallwood & Schooler, 2006). Greater mind wandering has also been linked to lower levels of working memory capacity (Kane et al., 2007). The distraction of mind wandering among depressed persons interferes with completion of planful activities.

Other phenomena related to depression also fit this view of executive control deficits (see also Levin et al., 2007). For example, depression is well known to be associated with autobiographical recall of memories that are very general in character rather than specific (Williams et al., 2007). A recent set of studies has yielded convincing evidence that this effect stems from executive control deficits (Dalgleish et al., 2007).

There is also evidence that depressed persons have trouble ignoring negative information (Frings, Wentura, & Holtz, 2007) and disengaging from negative information once attended to (Caseras, Garner, Bradley, & Mogg, 2007; Joormann & Gotlib, 2008; Joormann, Yoon, & Zetsche, 2007). Several measures have been developed to assess the capacity to effortfully suppress attention to negative material. Examples are the emotion Stroop task, measures of negative priming (Joormann, 2004; Wentura, 1999), and attention tasks with a long stimulus duration (Derryberry & Reed, 2002; Vasey, 2003, cited in Lonigan, Vasey, Phillips, & Hazen, 2004). Depressed persons show negative biases on the emotion Stroop, negative priming tasks, and attention tasks with long stimulus durations (Goeleven, De Raedt, Baert, & Koster, 2006; Gotlib et al., 2004; Gotlib & Krasnoperova, 1998; Joormann, 2004). All of this represents further evidence of deficits among depressed people in effortful cognitive control, in this case, specifically effortful control over thoughts.

Cognitive Effects of Tryptophan Depletion in Depression

Do these cognitive control deficits stem from low serotonergic function? Information on that question comes from studies of acute tryptophan depletion. As noted earlier, among healthy persons, acute tryptophan depletion affects cognitive processes involved in inhibiting responses to emotion-relevant stimuli (e.g., S. E. Murphy et al., 2006). Munafò et al. (2006) found that recovered but medicated depressed persons had particularly strong increases in Stroop interference for threat cues after acute tryptophan depletion; the effect of tryptophan depletion was not seen among healthy control participants nor among persons with remitted depression who were unmedicated.

Other studies have shown that such effects among formerly depressed persons are more pronounced for tasks with negatively valenced stimuli than for tasks without emotion-relevant stimuli. For example, Booij, Van der Does, et al. (2006) studied the effects of acute tryptophan depletion on processing of words in a Stroop task. Tryptophan depletion speeded processing of neutral and sad words, but it slowed processing of happiness-related words.

Hayward, Goodwin, Cowen, and Harmer (2005) also examined the effects of low-dose tryptophan depletion on persons with and without a history of depression, for emotion-relevant and other tasks. Recovered depressed people with acute tryptophan depletion displayed more elevation of emotion-potentiated startle and slower recognition of happy facial expressions, compared with other people. Tryptophan depletion and diagnostic group, however, did not interact in predicting non-emotion-relevant tasks such as a counting Stroop. Again, tryptophan depletion (and thus serotonin reduction) rendered formerly depressed people more responsive to negative emotional cues.

Restoring Effortful Control and Impediments to Doing So

If effortful cognitive control among depressed persons is poor, how can it be enhanced? Beevers (2005) pointed out that much of cognitive therapy involves training the person to use effortful and rule-based processing to identify and correct underlying negative biases. Although it is common to think of cognitive therapy as being a simple self-corrective process, it has always been clear that more is involved than merely providing positive self-statements. The two-mode view provides a good context for understanding effects of that intervention (cf. Daw et al., 2005). The intervention is, in effect, an attempt to restore effortful management of what is currently automatic behavior (see also Wood & Neal, 2007). This view helps account for the fact that cognitive therapy is a substantially more arduous process than one might initially assume (cf. Strunk, DeRubeis, Chiu, & Alvarez, 2007). There is evidence that it is possible to increase working memory capacity (Friedman et al., 2008; Olesen, Westerberg, & Klingberg, 2004), and it may be that such changes themselves are an outcome of cognitive therapy techniques.

Even if a person with depression tries to use such corrective processes, there are reasons why their implementation may fail. For one, the person must have the attentional capacity to do so (Barrett et al., 2004). Evidence suggests that persons with depression do not easily deploy effortful control. If the person's cognitive resources are stretched thin and thus unavailable, the correction is even less likely to take place.

Ironically, one thing that can deplete cognitive resources is negative affect. Thus, the emotional distress experienced in depression can help create the very conditions that make it difficult to counter further negative implicit processing. Beevers (2005) noted that even when an effortful correction is attempted, it may be insufficient. He further reasoned that a downward spiral may be induced, as cognitive resources are stretched by the need to counter negative thoughts. Indeed, there is evidence that the attempt to suppress or inhibit information attaches negative valence to that information (Fenske & Raymond, 2006).

Consistent with that reasoning, Wenzlaff and Bates (1998) have shown that diminished working memory capacity (induced by a cognitive load) led depressed persons to display more negative bias on a scrambled sentence task. Intriguingly, negativity under cognitive load was a better predictor of depressive symptoms over time than was negativity without cognitive load. Hence, the cognitive resources available to correct implicit biases may be particularly important for the course of depressive symptoms.

Section Summary

A good deal of evidence suggests that persons with depression have difficulties exerting effortful cognitive control. Their attention is drawn to negative emotional cues (perhaps because of their many negative associations in memory), which they then have trouble disengaging from. They are prone to rumination about negativity but also prone to mind wandering rather than task engagement. Those recovered from depression experience recurrence of sensitivity to negative cues when their serotonin levels are artificially lowered.

These various findings are consistent with a two-mode view in which depression relates to limited executive function.

Future Directions

An earlier section of the article suggests an interactive view of vulnerability to depression versus vulnerability to disorders involving impulsivity. It was suggested there that both tend to reflect low serotonergic function, and thus low effortful control, but that they differ in level of engagement of the system responsible for eager, effortful approach. Psychopathy and impulsive aggression were characterized as reflecting low serotonergic function and relatively high approach motivation (perhaps true of other impulse-related problems as well; cf. Depue & Lenzenweger, 2005). Depression was characterized as reflecting low serotonergic function and low approach motivation.

Further Interactions?

A broad question for the future is whether other interactions should also be explored more fully (Depue & Lenzenweger, 2005; Nigg, 2006). For example, perhaps overt expression of a vulnerability to anxiety disorders may also depend on poor executive control (Lonigan et al., 2004). Consistent with this idea, serotonin has been implicated in the development of anxiety disorders (see Leonardo & Hen, 2006, for review). There is also evidence that the 5-HTTLPR polymorphism (in a triallelic design) interacts with daily stress to influence reports of anxious mood (Gunthert et al., 2007). Cools, Calder, and colleagues (2005) found that experimental reduction of serotonin (via tryptophan depletion) led to increased amygdala activation in response to fearful faces but did so only among persons who had previously reported being high in threat sensitivity.

These findings all suggest an interactive combination of a highly sensitive threat system and low serotonergic functioning.¹² How the two-mode model might help in the understanding of anxiety-based problems, and in distinguishing the etiology of anxiety disorders from that of depression, would seem to be an important topic for further investigation.

Low Serotonin Function as Amplifier

Many observers have noted that the attempt to link any given neurotransmitter to the operation of a single behavioral system is likely to be a great oversimplification; many have also noted that effects of one neurotransmitter are likely to depend on the person's level of another neurotransmitter (Depue, 1995; Depue & Lenzenweger, 2005; Zuckerman, 2005). Indeed, what was known as permissive theory (Prange, Wilson, Lynn, Alltop, & Stikeleather, 1974) long ago argued specifically that low serotonin function enhances risk of depression in interaction with other neurotransmitters (see Howland & Thase, 1999). Characterizing one neurotransmitter as having an overall effect is hazardous.

Nonetheless, it does not seem too far an extrapolation from the evidence to suggest that low serotonergic function promotes a stronger manifestation of whatever tendencies the person has at the reflexive or implicit level of functioning (for similar conclusions, see Depue, 1995; Nigg, 2006; Spont, 1992). In an incentive-sensitive person, low serotonergic function amplifies the pursuit of incentives. In an incentive-insensitive person, low serotonergic function exaggerates the lack of effortful engagement. In a threat-sensitive person, low serotonergic function may enhance vigilance to threat.

¹²It is perhaps noteworthy in this context that SSRIs are effective for anxiety disorders as well as depression.

Low serotonergic function may also promote a stronger manifestation of the pattern of conditioned associations the person has acquired over time (S. E. Taylor et al., 2006). In a person who experienced childhood maltreatment (and who thus has many negative associations with the self in memory), low serotonergic function may enhance the negative aftereffects of that maltreatment (Caspi et al., 2003). In a person who had a very supportive early family environment (and who thus has many positive elements in associative memory), low serotonergic function may enhance the positive reverberations of that environment (S. E. Taylor et al., 2006).

The specific cases of depression, impulsive disorders, and anxiety disorders are only three possibilities, reflecting interactions of a serotonergic system with two other systems. A more complete understanding of the role of serotonin in behavior will require a more elaborated understanding of how serotonergic function interacts with effects of other neurotransmitters. The idea that diverse disorders follow from diverse combinations of system sensitivities (Depue & Lenzenweger, 2005; Lenzenweger & Willett, 2007) is very intriguing and seems worthy of much more examination.

Effortful Control and Overcoming Inaction

Another sort of interaction also deserves more attention than it has yet received. As noted earlier (Figure 2), the two-mode reasoning that guided this review suggests that high serotonergic function, and the associated high degree of effortful control, does more than restrain impulses. In theory (and in limited evidence), effortful control can also cause a person who is unmotivated to act to take action. In principle, then, effortful control should have divergent effects, depending on the behavioral context. Effortful control (and related constructs such as executive control, working memory capacity, and serotonergic function) should be involved in countermanding tendencies triggered by inhibitory reactive systems as well as impulsive reactive systems. This idea also deserves closer attention.

Conclusion

Recent years have seen an explosion of interest in the role of serotonergic function in depression and elsewhere. The developing literatures use a wide array of research paradigms. The functions of serotonin have been interpreted in terms of diverse constructs, ranging from neuroticism and anxiety-related traits, to cortical control over amygdala activity, to effortful cognitive processes. Review of these literatures suggests that it is useful to conceptualize these various functions in terms of two-mode models of self-regulation. Viewed through this lens, the evidence suggests that serotonergic function can be linked to impulsivity versus constraint in the personality literature, effortful control processes in the cognitive and developmental literatures, and executive control over the amygdala and other subcortical areas in the neurobiological literature. This two-mode picture helps organize what is known about the experience of depression, and it may also be useful in suggesting new areas of investigation.

This article was prompted by evidence that low serotonergic function is a risk factor in depression. Though that evidence is essentially neurobiological in nature, it has implications that are cognitive, behavioral, and phenomenological. Ideas and evidence from several literatures that are psychological in nature appear to foster a deeper understanding of the meaning of this evidence regarding biological vulnerability. This confluence suggests another message as well. Increasingly it is held that biological concepts and methods are key resources to inform psychological theory (e.g., Cacioppo et al., 2007; Winkelman, Knutson, Paulus, & Trujillo, 2007). We would hold, however (see also Posner & Rothbart, 2007), that interpretation of neurobiological evidence also benefits from considering the findings through the lens of psychological principles.

Acknowledgments

Work on the article was supported by the National Science Foundation (BCS0544617), the National Cancer Institute (CA64710), the National Institute of Mental Health (MH076021), and a NARSAD Young Investigator Award.

References

- Abercrombie HC, Schaefer SM, Larson CL, Oakes TR, Lindgren KA, Holden JE. Metabolic rate in the right amygdala predicts negative affect in depressed patients. *NeuroReport* 1998;9:3301–3307. [PubMed: 9831467]
- Abramson, LY.; Alloy, LB.; Hankin, BL.; Haefel, GJ.; MacCoon, DJ.; Gibb, BE. Cognitive vulnerability—stress models of depression in a self-regulatory and psychobiological context. In: Gotlib, IH., editor. *Handbook of depression*. New York: Guilford Press; 2002. p. 268–294.
- Aggleton, JP. *The amygdala: A functional analysis*. 2nd ed. Oxford, United Kingdom: Oxford University Press; 2000.
- Ågren H. Symptom patterns in unipolar and bipolar depression correlating with monoamine metabolites in the cerebrospinal fluid: II. Suicide. *Psychiatry Research* 1980;3:225–236. [PubMed: 6171841]
- Ahadi, SA.; Rothbart, MK. Temperament, development and the big five. In: Halverson, CF., Jr.; Kohnstamm, GA.; Martin, RP., editors. *The developing structure of temperament and personality from infancy to adulthood*. Hillsdale, NJ: Erlbaum; 1994. p. 189–207.
- Allen JJ, Iacono WG, Depue RA, Arbisi P. Regional electroencephalographic asymmetries in bipolar seasonal affective disorder before and after exposure to bright light. *Biological Psychiatry* 1993;33:642–646. [PubMed: 8329494]
- Allen PP, Cleare AJ, Lee F, Fusar-Poli P, Tunstall N, Fu CH, et al. Effect of acute tryptophan depletion on pre-frontal engagement. *Psychopharmacology* 2006;187:486–497. [PubMed: 16823591]
- Anderson AK, Phelps EA. Lesions of the human amygdala impair enhanced perception of emotionally salient events. *Nature* 2001 May 17;411:305–309. [PubMed: 11357132]
- Anguelova M, Benkelfat C, Turecki G. A systematic review of association studies investigating genes coding for serotonin receptors and the serotonin transporter: I. Affective disorders. *Molecular Psychiatry* 2003;8:574–591. [PubMed: 12851635]
- Banks SJ, Eddy KT, Angstadt M, Nathan PJ, Phan KL. Amygdala-frontal connectivity during emotion regulation. *Social Cognitive and Affective Neuroscience* 2007;2:303–312. [PubMed: 18985136]
- Barkley RA. Behavioral inhibition, sustained attention, and executive functions: Constructing a unifying theory of ADHD. *Psychological Bulletin* 1997;121:65–94. [PubMed: 9000892]
- Barrett LF, Tugade MM, Engle RW. Individual differences in working memory capacity and dual-process theories of the mind. *Psychological Bulletin* 2004;130:553–573. [PubMed: 15250813]
- Barton DA, Esler MD, Dawood T, Lambert EA, Haikerwal D, Brenchley C, et al. Elevated brain serotonin turnover in patients with depression: Effect of genotype and therapy. *Archives of General Psychiatry* 2008;65:38–46. [PubMed: 18180427]
- Baxter LR, Schwartz JM, Phelps ME, Mazziotta JC, Guze BH, Selin CE, et al. Reduction of prefrontal cortex glucose metabolism common to three types of depression. *Archives of General Psychiatry* 1989;46:243–250. [PubMed: 2784046]
- Beauregard M, Lévesque J, Bourgouin P. Neural correlates of conscious self-regulation of emotion. *Journal of Neuroscience* 2001;21:6993–7000. [PubMed: 11549709]
- Bechara A. Decision making, impulse control, and loss of willpower to resist drugs: A neurocognitive perspective. *Nature Neuroscience* 2005;8:1458–1463.
- Bechara A, Damasio H, Damasio AR. Emotion, decision making and the orbitofrontal cortex. *Cerebral Cortex* 2000;10:295–307. [PubMed: 10731224]
- Bechtel, W.; Abrahamsen, A. *Connectionism and the mind: An introduction to parallel processing in networks*. Cambridge, United Kingdom: Basil Blackwell; 1991.
- Beer JS, Knight RT, D'Esposito M. Controlling the integration of emotion and cognition: The role of frontal cortex in distinguishing helpful from hurtful emotional information. *Psychological Science* 2006;17:448–453. [PubMed: 16683934]

- Beevers CG. Cognitive vulnerability to depression: A dual process model. *Clinical Psychology Review* 2005;25:975–1002. [PubMed: 15905008]
- Beitchman JH, Davidge KM, Kennedy JL, Atkinson L, Lee V, Sharp S, et al. The serotonin transporter gene in aggressive children with and without ADHD and nonaggressive matched controls. *Annals of the New York Academy of Sciences* 2003;1008:248–251. [PubMed: 14998889]
- Bench CJ, Friston KJ, Brown RG, Frackowiak RS, Dolan RJ. Regional cerebral blood flow in depression measured by positron emission tomography: The relationship with clinical dimensions. *Psychological Medicine* 1993;23:579–590. [PubMed: 7901863]
- Benkelfat C, Ellenbogen MA, Dean P, Palmour RM. Mood-lowering effect of tryptophan depletion: Enhanced susceptibility in young men at genetic risk for major affective disorders. *Archives of General Psychiatry* 1994;51:687–697. [PubMed: 8080345]
- Berridge KC. The debate over dopamine's role in reward: The case for incentive salience. *Psychopharmacology* 2007;191:391–431. [PubMed: 17072591]
- Bhagwagar Z, Whale R, Cowen PJ. State and trait abnormalities in serotonin function in major depression. *British Journal of Psychiatry* 2002;180:24–28. [PubMed: 11772847]
- Bjork JM, Dougherty DM, Moeller FG, Swann AC. Differential behavioral effects of plasma tryptophan depletion and loading in aggressive and nonaggressive men. *Neuropsychopharmacology* 2000;22:357–369. [PubMed: 10700655]
- Blair RJR. Neurobiological basis of psychopathy. *British Journal of Psychiatry* 2003;182:5–7. [PubMed: 12509310]
- Booij L, Swenne CA, Brosschot JF, Haffmans PMJ, Thayer JF, Van der Does AJW. Tryptophan depletion affects heart rate variability and impulsivity in remitted depressed patients with a history of suicidal ideation. *Biological Psychiatry* 2006;60:507–514. [PubMed: 16603135]
- Booij L, Van der Does AJW. Cognitive and serotonergic vulnerability to depression: Convergent findings. *Journal of Abnormal Psychology* 2007;116:86–94. [PubMed: 17324019]
- Booij L, Van der Does W, Benkelfat C, Bremner JD, Cowen PJ, Fava M, et al. Predictors of mood response to acute tryptophan depletion: A reanalysis. *Neuropsychopharmacology* 2002;27:852–861. [PubMed: 12431859]
- Booij L, Van der Does AJW, Haffmans PMJ, Riedel WJ, Fekkes D, Blom MJB. The effects of high-dose and low-dose tryptophan depletion on mood and cognitive functions of remitted depressed patients. *Journal of Psychopharmacology* 2006;19:267–275. [PubMed: 15888512]
- Breiter HC, Etcoff NL, Whalen PJ, Kennedy WA, Rauch SL, Buckner RL, et al. Response and habituation of the human amygdala during visual processing of facial expression. *Neuron* 1996;17:875–887. [PubMed: 8938120]
- Brezo J, Paris J, Turecki G. Personality traits as correlates of suicidal ideation, suicide attempts, and suicide completions: A systematic review. *Acta Psychiatrica Scandinavica* 2006;113:180–206. [PubMed: 16466403]
- Brinkmann K, Gendolla GHE. Does depression interfere with effort mobilization? Effects of dysphoria and task difficulty on cardiovascular response. *Journal of Personality and Social Psychology* 2008;94:146–157. [PubMed: 18179324]
- Brocke B, Armbruster D, Müller J, Hensch T, Jacob CP, Lesch K-P, et al. Serotonin transporter gene variation impacts innate fear processing: Acoustic startle response and emotional startle. *Molecular Psychiatry* 2006;11:1106–1112. [PubMed: 17033630]
- Brown SM, Manuck SB, Flory JD, Hariri AR. Neural basis of individual differences in impulsivity: Contributions of corticolimbic circuits for behavioral arousal and control. *Emotion* 2006;6:239–245. [PubMed: 16768556]
- Bush G, Luu P, Posner MI. Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Sciences* 2000;4:215–222. [PubMed: 10827444]
- Buss AH, Durkee A. An inventory for assessing different kinds of hostility. *Journal of Consulting and Clinical Psychology* 1957;21:353–356.
- Cacioppo JT, Amaral DG, Blanchard JJ, Cameron JL, Carter CS, Crews D, et al. Social neuroscience: Progress and implications for mental health. *Perspectives on Psychological Science* 2007;2:99–123.
- Cahill L, Babinsky R, Markowitsch HJ, McGaugh JL. The amygdala and emotional memory. *Nature* 1995 September 28;377:295–296. [PubMed: 7566084]

- Campbell-Sills L, Liverant GI, Brown TA. Psychometric evaluation of the Behavioral Inhibition/Behavioral Activation Scales in a large sample of outpatients with anxiety and mood disorders. *Psychological Assessment* 2004;16:244–254. [PubMed: 15456380]
- Canli T, Lesch K. Long story short: The serotonin transporter in emotion regulation and social cognition. *Nature Neuroscience* 2007;10:1103–1109.
- Canli T, Omura K, Haas BW, Fallgatter A, Constable RT, Lesch KP. Beyond affect: A role for genetic variation of the serotonin transporter in neural activation during a cognitive attention task. *Proceedings of the National Academy of Sciences of the United States of America* 2005;102:12224–12229. [PubMed: 16093315]
- Canli T, Qiu M, Omura K, Congdon E, Haas BW, Amin Z, et al. Neural correlates of epigenesis. *Proceedings of the National Academy of Sciences of the United States of America* 2006;103:16033–16038. [PubMed: 17032778]
- Carter CS, Braver TS, Barch DM, Botvinick MM, Noll D, Cohen JD. Anterior cingulate cortex, error detection, and the online monitoring of performance. *Science* 1998 May 1;280:747–749. [PubMed: 9563953]
- Carver CS. Impulse and constraint: Perspectives from personality psychology, convergence with theory in other areas, and potential for integration. *Personality and Social Psychology Review* 2005;9:312–333. [PubMed: 16223354]
- Carver CS, Miller CJ. Relations of serotonin function to personality: Current views and a key methodological issue. *Psychiatry Research* 2006;144:1–15. [PubMed: 16914207]
- Caseras X, Garner M, Bradley BP, Mogg K. Biases in visual orienting to negative and positive scenes in dysphoria: An eye movement study. *Journal of Abnormal Psychology* 2007;116:491–497. [PubMed: 17696705]
- Casey BJ, Tottenham N, Fossella J. Clinical, imaging, lesion and genetic approaches toward a model of cognitive control. *Developmental Psychobiology* 2002;40:237–254. [PubMed: 11891636]
- Caspi, A.; Shiner, RL. Personality development. In: Damon, W.; Lerner, R.; Eisenberg, N., editors. *Handbook of child psychology: Vol. 3. Social, emotional, and personality development*. 6th ed. New York: Wiley; 2006. p. 300–365.
- Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, et al. Influence of life stress on depression: Moderation by a polymorphism in the 5-HTT gene. *Science* 2003 July 18;301:386–389. [PubMed: 12869766]
- Chaiken, SL.; Trope, Y., editors. *Dual-process theories in social psychology*. New York: Guilford Press; 1999.
- Chao LL, Knight RT. Contribution of human prefrontal cortex to delay performance. *Journal of Cognitive Neuroscience* 1998;10:167–177. [PubMed: 9555105]
- Cheng DT, Knight DC, Smith CN, Helmstetter FJ. Human amygdala activity during the expression of fear responses. *Behavioral Neuroscience* 2006;120:1187–1195. [PubMed: 17201461]
- Cherek DR, Lane SD. Fenfluramine effects on impulsivity in a sample of adults with and without history of conduct disorder. *Psychopharmacology* 2000;152:149–156. [PubMed: 11057518]
- Clark LA. Temperament as a unifying basis for personality and psychopathology. *Journal of Abnormal Psychology* 2005;114:505–521. [PubMed: 16351374]
- Clark L, Cools R, Robbins TW. The neuropsychology of ventral prefrontal cortex: Decision-making and reversal learning. *Brain and Cognition* 2004;55:41–53. [PubMed: 15134842]
- Cleare AJ, Bond AJ. The effect of tryptophan depletion and enhancement on subjective and behavioural aggression in normal male subjects. *Psychopharmacology* 1995;118:72–81. [PubMed: 7597125]
- Cleare AJ, Bond AJ. Does central serotonergic function correlate inversely with aggression? A study using D-fenfluramine in healthy subjects. *Psychiatry Research* 1997;69:89–95. [PubMed: 9109176]
- Cloninger CR. A systematic method for clinical description and classification of personality variants: A proposal. *Archives of General Psychiatry* 1987;44:573–588. [PubMed: 3579504]
- Cloninger CR, Svrakic DM, Przybeck TR. A psychobiological model of temperament and character. *Archives of General Psychiatry* 1993;50:975–990. [PubMed: 8250684]
- Coan JA, Allen JJB. Frontal EEG asymmetry and the behavioral activation and inhibition systems. *Psychophysiology* 2003;40:106–114. [PubMed: 12751808]

- Coccaro EF, Kavoussi RJ. Neuropsychopharmacologic challenge in biological psychiatry. *Clinical Chemistry* 1994;40:319–327. [PubMed: 7906209]
- Coccaro EF, Kavoussi RJ, Cooper TB, Hauger RL. Central serotonin activity and aggression: Inverse relationship with prolactin response to d-fenfluramine, but not CSF 5-HIAA concentration, in human subjects. *American Journal of Psychiatry* 1997;154:1430–1435. [PubMed: 9326827]
- Coccaro EF, Kavoussi RJ, Hauger RL, Cooper TB, Ferris CF. Cerebrospinal fluid vasopressin levels: Correlates with aggression and serotonin function in personality-disordered subjects. *Archives of General Psychiatry* 1998;55:708–714. [PubMed: 9707381]
- Collier DA, Stöber G, Li T, Heils A, Catalano M, di Bella D, et al. A novel functional polymorphism within the promoter of the serotonin transporter gene: Possible role in susceptibility to affective disorders. *Molecular Psychiatry* 1996;1:453–460. [PubMed: 9154246]
- Compton RJ, Lin M, Vargas G, Carp J, Fineman SL, Quandt LC. Error detection and posterror behavior in depressed undergraduates. *Emotion* 2008;8:58–67. [PubMed: 18266516]
- Conner T, Barrett LF. Implicit self-attitudes predict spontaneous affect in daily life. *Emotion* 2005;5:476–488. [PubMed: 16366751]
- Cools R, Blackwell A, Clark L, Menzies L, Cox S, Robbins TW. Tryptophan depletion disrupts the motivational guidance of goal-directed behavior as a function of trait impulsivity. *Neuropsychopharmacology* 2005;30:1362–1373. [PubMed: 15770237]
- Cools R, Calder AJ, Lawrence AD, Clark L, Bullmore E, Robbins TW. Individual differences in threat sensitivity predict serotonergic modulation of amygdala response to fearful faces. *Psychopharmacology* 2005;180:670–679. [PubMed: 15772862]
- Cooney RE, Joormann J, Atlas LY, Eugene F, Gotlib IH. Neural correlates of affect regulation through mood-incongruent recall. *NeuroReport* 2007;18:1771–1774. [PubMed: 18090309]
- Corbetta M, Shulman GL. Control of goal-directed and stimulus-driven attention in the brain. *Nature Reviews Neuroscience* 2002;3:201–215.
- Cotter D, Mackay D, Chana G, Beasley C, Landau S, Everall IP. Reduced neuronal size and glial cell density in area 9 of the dorsolateral prefrontal cortex in subjects with major depressive disorder. *Cerebral Cortex* 2002;12:386–394. [PubMed: 11884354]
- Cowen PJ. Cortisol, serotonin, and depression: All stressed out. *British Journal of Psychiatry* 2002;180:99–100. [PubMed: 11823315]
- Crockett MJ, Clark L, Tabibnia G, Lieberman MD, Robbins TW. Serotonin modulates behavioral reactions to unfairness. *Science* June 27;2008 320:1739. [PubMed: 18535210]
- Cunningham WA, Van Bavel JJ, Johnsen IR. Affective flexibility: Evaluation processing goals shape amygdala activity. *Psychological Science* 2008;19:152–160. [PubMed: 18271863]
- Custers R, Aarts H. Positive affect as implicit motivator: On the nonconscious operation of behavioral goals. *Journal of Personality and Social Psychology* 2005;89:129–142. [PubMed: 16162049]
- Dalgleish T, Golden AJ, Barrett LF, Yeung CA, Murphy V, Tchanturia K, et al. Reduced specificity of autobiographical memory and depression: The role of executive control. *Journal of Experimental Psychology: General* 2007;136:23–42. [PubMed: 17324083]
- Davidson RJ. Anterior electrophysiological asymmetries, emotion, and depression: Conceptual and methodological conundrums. *Psychophysiology* 1998;35:607–614. [PubMed: 9715104]
- Davidson RJ. Affective style, psychopathology, and resilience: Brain mechanisms and plasticity. *American Psychologist* 2000;55:1196–1214. [PubMed: 11280935]
- Davidson RJ, Pizzagalli D, Nitschke JB, Putnam K. Depression: Perspectives from affective neuroscience. *Annual Review of Psychology* 2002;53:545–574.
- Davis M, Whalen PJ. The amygdala: Vigilance and emotion. *Molecular Psychiatry* 2001;6:13–34. [PubMed: 11244481]
- Daw ND, Niv Y, Dayan P. Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nature Neuroscience* 2005;8:1704–1711.
- Dawson, MRW. Connectionism: A hands-on approach. Malden, MA: Blackwell; 2005.
- Delgado PL, Charney DS, Price LH, Aghajanian GK. Serotonin function and the mechanism of antidepressant action: Reversal of antidepressant-induced remission by rapid depletion of plasma tryptophan. *Archives of General Psychiatry* 1990;47:411–418. [PubMed: 2184795]

- Delgado PL, Miller HL, Salomon RM, Licinio J, Krystal JH, Moreno FA, et al. Tryptophan-depletion challenge in depressed patients treated with desipramine or fluoxetine: Implications for the role of serotonin in the mechanism of antidepressant action. *Biological Psychiatry* 1999;46:212–220. [PubMed: 10418696]
- De Neys W. Dual processing in reasoning: Two systems but one reasoner. *Psychological Science* 2006;17:428–433. [PubMed: 16683931]
- Dennis TA, Brotman LM. Effortful control, attention, and aggressive behavior in preschoolers at risk for conduct problems. *Annals of the New York Academy of Sciences* 2003;1008:252–255. [PubMed: 14998890]
- Depue RA. Neurobiological factors in personality and depression. *European Journal of Personality* 1995;9:413–439.
- Depue RA, Collins PF. Neurobiology of the structure of personality: Dopamine, facilitation of incentive motivation, and extraversion. *Behavioral and Brain Sciences* 1999;22:491–517. [PubMed: 11301519]
- Depue RA, Iacono WG. Neurobehavioral aspects of affective disorders. *Annual Review of Psychology* 1989;40:457–492.
- Depue, RA.; Lenzenweger, MF. A neurobiological dimensional model of personality disturbance. In: Lenzenweger, MF.; Clarkin, JF., editors. *Major theories of personality disorder*. 2nd ed. New York: Guilford Press; 2005. p. 391–453.
- Depue RA, Spont MR. Conceptualizing a serotonin trait: A behavioral dimension of constraint. *Annals of the New York Academy of Sciences* 1986;487:47–62. [PubMed: 2436540]
- Derefinko KJ, Lynam DR. Convergence and divergence among self-report psychopathy measures: A personality-based approach. *Journal of Personality Disorders* 2006;20:261–280. [PubMed: 16776555]
- Derryberry D, Reed MA. Anxiety-related attentional biases and their regulations by attentional control. *Journal of Abnormal Psychology* 2002;111:225–236. [PubMed: 12003445]
- Derryberry D, Rothbart MK. Reactive and effortful processes in the organization of temperament. *Development and Psychopathology* 1997;9:633–652. [PubMed: 9448999]
- Dolan MC, Anderson IM. The relationship between serotonergic function and the Psychopathy Checklist: Screening Version. *Journal of Psychopharmacology* 2003;17:216–222. [PubMed: 12870570]
- Dolan MC, Anderson IM, Deakin JFW. Relationship between 5-HT function and impulsivity and aggression in male offenders with personality disorders. *British Journal of Psychiatry* 2001;178:352–359. [PubMed: 11282815]
- Dolan MC, Deakin JFW, Roberts N, Anderson I. Serotonergic and cognitive impairment in impulsive aggressive personality disordered offenders: Are there implications for treatment? *Psychological Medicine* 2002;32:105–117. [PubMed: 11885567]
- Dolan RJ, Bench CJ, Brown RG, Scott LC, Frackowiak RS. Neuropsychological dysfunction in depression: The relationship to regional cerebral blood flow. *Psychological Medicine* 1994;24:849–857. [PubMed: 7892353]
- Dolan RJ, Bench CJ, Liddle PF, Friston KJ, Frith CD, Grasby PM, et al. Dorsolateral prefrontal cortex dysfunction in the major psychoses: Symptom or disease specificity? *Journal of Neurology, Neurosurgery, and Psychiatry* 1993;56:1290–1294.
- Drevets WC. Functional neuroimaging studies of depression: The anatomy of melancholia. *Annual Review of Medicine* 1998;49:341–361.
- Drevets WC. Prefrontal cortical-amygdalar metabolism in major depression. *Annals of the New York Academy of Sciences* 1999;29:614–637. [PubMed: 10415674]
- Drevets WC. Neuroimaging studies of mood disorders. *Biological Psychiatry* 2000;48:813–829. [PubMed: 11063977]
- Drevets, WC. Neuroimaging studies of major depression. In: Morihisa, JM., editor. *Advances in brain imaging*. Washington, DC: American Psychiatric Publishing; 2001. p. 123–170.
- Drevets WC, Price JL, Bardgett ME, Reich T, Todd RD, Raichle ME. Glucose metabolism in the amygdala in depression: Relationship to diagnostic subtype and plasma cortisol levels. *Pharmacology, Biochemistry, and Behavior* 2002;71:431–447.

- Drevets WC, Raichle ME. Reciprocal suppression of regional cerebral blood flow during emotional versus higher cognitive processes: Implications for interactions between emotion and cognition. *Cognition and Emotion* 1998;12:353–385.
- Drevets WC, Videen TO, Price JL, Preskorn SH, Carmichael ST, Raichle ME. A functional anatomical study of unipolar depression. *Journal of Neuroscience* 1992;12:3628–3641. [PubMed: 1527602]
- Dunlop BW, Nemeroff CB. The role of dopamine in the pathophysiology of depression. *Archives of General Psychiatry* 2007;64:327–337. [PubMed: 17339521]
- Durston S, Thomas KM, Worden MS, Yang Y, Casey BJ. The effect of preceding context on inhibition: An event-related fMRI study. *NeuroImage* 2002;16:449–453. [PubMed: 12030830]
- Durston S, Thomas KM, Yang Y, Ulug AM, Zimmerman RD, Casey BJ. A neural basis for the development of inhibitory control. *Developmental Science* 2002;5:F9–F16.
- Dutton DG. The neurobiology of abandonment homicide. *Aggression and Violent Behavior* 2002;7:407–421.
- Eisenberg N, Cumberland A, Spinrad TL, Fabes RA, Shepard SA, Reiser M, et al. The relations of regulation and emotionality to children's externalizing and internalizing problem behavior. *Child Development* 2001;72:1112–1134. [PubMed: 11480937]
- Eisenberg N, Spinrad TL, Fabes RA, Reiser M, Cumberland A, Shepard SA, et al. The relations of effortful control and impulsivity to children's resiliency and adjustment. *Child Development* 2004;75:25–46. [PubMed: 15015673]
- Eley TC, Sugden K, Corsico A, Gregory AM, Sham P, McGuffin P, et al. Gene-environment interaction analysis of serotonin system markers with adolescent depression. *Molecular Psychiatry* 2004;9:908–915. [PubMed: 15241435]
- Ellenbogen MA, Young SN, Dean P, Palmour RM, Benkelfat C. Mood response to acute tryptophan depletion in healthy volunteers: Sex differences and temporal stability. *Neuropsychopharmacology* 1996;15:465–474. [PubMed: 8914119]
- Epstein S. The self-concept revisited: Or a theory of a theory. *American Psychologist* 1973;28:404–416. [PubMed: 4703058]
- Epstein S. The implications of cognitive-experiential self theory for research in social psychology and personality. *Journal for the Theory of Social Behavior* 1985;15:283–310.
- Epstein, S. Cognitive-experiential self-theory. In: Pervin, L., editor. *Handbook of personality: Theory and research*. New York: Guilford Press; 1990. p. 165-192.
- Epstein S. Integration of the cognitive and the psychodynamic unconscious. *American Psychologist* 1994;49:709–724. [PubMed: 8092614]
- Ernst M, Nelson EE, McClure EB, Monk CS, Munson S, Eshel N, et al. Choice selection and reward anticipation: An fMRI study. *Neuropsychologia* 2004;42:1585–1597. [PubMed: 15327927]
- Evans J, Platts H, Lightman S, Nutt D. Impulsiveness and the prolactin response to d-fenfluramine. *Psychopharmacology* 2000;149:147–152. [PubMed: 10805609]
- Evans, J. St. B. T Dual-processing accounts of reasoning, judgment, and social cognition. *Annual Review of Psychology* 2008;59:255–278.
- Farrar AM, Pereira M, Velasco F, Hockemeyer J, Müller CE, Salamone JD. Adenosine A_{2A} receptor antagonism reverses the effects of dopamine receptor antagonism on instrumental output and effort-related choice in the rat: Implications for studies of psychomotor slowing. *Psychopharmacology* 2007;191:579–586. [PubMed: 17072593]
- Fenske MJ, Raymond JE. Affective influences of selective attention. *Current Directions in Psychological Science* 2006;15:312–316.
- Finkel EJ, Campbell WK. Self-control and accommodation in close relationships: An interdependence analysis. *Journal of Personality and Social Psychology* 2001;81:263–277. [PubMed: 11519931]
- Finn PR, Young SN, Pihl RO, Ervin FR. The effects of acute plasma tryptophan manipulation on hostile mood: The influence of trait hostility. *Aggressive Behavior* 1998;24:173–185.
- Flory JD, Mann JJ, Manuck SB, Muldoon MF. Recovery from major depression is not associated with normalization of serotonergic function. *Biological Psychiatry* 1998;43:320–326. [PubMed: 9513746]

- Flory JD, Manuck SB, Matthews KA, Muldoon MF. Serotonergic function in the central nervous system is associated with daily ratings of positive mood. *Psychiatry Research* 2004;129:11–19. [PubMed: 15572180]
- Flory JD, Newcorn JH, Miller C, Harty S, Halperin JM. Serotonergic function in children with attention-deficit hyperactivity disorder: Relationship to later antisocial personality disorder. *The British Journal of Psychiatry* 2007;190:410–414. [PubMed: 17470955]
- Forbes E, Fox NA, Cohn JF, Galles SF, Kovacs M. Children's affect regulation during a disappointment: Psychophysiological responses and relation to parent history of depression. *Biological Psychology* 2006;71:264–277. [PubMed: 16115722]
- Fox NA, Henderson HA, Rubin KH, Calkins SD, Schmidt LA. Continuity and discontinuity of behavioral inhibition and exuberance: Psychophysiological and behavioral influences across the first four years of life. *Child Development* 2001;72:1–21. [PubMed: 11280472]
- Friedman NP, Haberstick BC, Willcutt EG, Miyake A, Young SE, Corley RP, et al. Greater attention problems during childhood predict poorer executive functioning in late adolescence. *Psychological Science* 2007;18:893–900. [PubMed: 17894607]
- Friedman NP, Miyake A, Young SE, DeFries JC, Corley RP, Hewitt JK. Individual differences in executive functions are almost entirely genetic in origin. *Journal of Experimental Psychology: General* 2008;137:201–225. [PubMed: 18473654]
- Frings C, Wentura D, Holtz M. Dysphorics cannot ignore unpleasant information. *Cognition and Emotion* 2007;21:1525–1534.
- Galvan A, Hare TA, Parra CE, Penn J, Voss K, Glover G, Casey BJ. Earlier development of the accumbens relative to orbitofrontal cortex might underlie risk-taking behavior in adolescents. *Journal of Neuroscience* 2006;26:6885–6892. [PubMed: 16793895]
- George DT, Umhau JC, Phillips MJ, Emmela D, Ragan PW, Shoaf SE, et al. Serotonin, testosterone, and alcohol in the etiology of domestic violence. *Psychiatry Research* 2001;104:27–37. [PubMed: 11600187]
- Gerra G, Zaimovic A, Timpano M, Zambelli U, Delsignore R, Brambilla F. Neuroendocrine correlates of temperamental traits in humans. *Psychoneuroendocrinology* 2000;25:479–496. [PubMed: 10818282]
- Gillespie NA, Whitfield JB, Williams B, Heath AC, Martin NG. The relationship between stressful life events, the serotonin transporter (5-HTTLPR) genotype and major depression. *Psychological Medicine* 2005;35:101–111. [PubMed: 15842033]
- Goeleven E, De Raedt R, Baert S, Koster EHW. Deficient inhibition of emotional information in depression. *Journal of Affective Disorders* 2006;93:149–157. [PubMed: 16647141]
- Gotlib IH, Hamilton JP. Neuroimaging and depression: Current status and unresolved issues. *Current Directions in Psychological Science* 2008;17:159–163.
- Gotlib IH, Joormann J, Minor KL, Hallmayer J. Biological stress reactivity mediates the relation between genotype and risk for depression. *Biological Psychiatry* 2008;63:847–851. [PubMed: 18005940]
- Gotlib IH, Kasch KL, Traill S, Joormann J, Arnow BA, Johnson SL. Coherence and specificity of information-processing biases in depression and social phobia. *Journal of Abnormal Psychology* 2004;113:386–398. [PubMed: 15311984]
- Gotlib IH, Krasnoperova E. Biased information processing as a vulnerability factor for depression. *Behavior Therapy* 1998;29:603–617.
- Gotlib IH, Ranganath C, Rosenfeld JP. Frontal EEG alpha asymmetry, depression, and cognitive functioning. *Cognition and Emotion* 1998;12:449–478.
- Grabe HJ, Lange M, Wolff B, Völzke H, Lucht M, Freyberger HJ, et al. Mental and physical distress is modulated by a polymorphism in the 5-HT transporter gene interacting with social stressors and chronic disease burden. *Molecular Psychiatry* 2005;10:220–224. [PubMed: 15263905]
- Greenberg BD, Li Q, Lucas FR, Hu S, Sirota LA, Benjamin J, et al. Association between the serotonin transporter promoter polymorphism and personality traits in a primarily female population sample. *American Journal of Medical Genetics (Neuropsychiatric Genetics)* 2000;96:202–216. [PubMed: 10893498]

- Greenwald AG, Banaji MR, Rudman LA, Farnham SD, Nosek BA, Mellott DS. A unified theory of implicit attitudes, stereotypes, self-esteem, and self-concept. *Psychological Review* 2002;109:3–25. [PubMed: 11863040]
- Gunther KC, Conner TS, Armeli S, Tennen H, Covault J, Kranzler HR. Serotonin transporter gene polymorphism (5-HTTLPR) and anxiety reactivity in daily life: A daily process approach to gene-environment interaction. *Psychosomatic Medicine* 2007;69:762–768. [PubMed: 17942837]
- Haefel GJ, Abramson LY, Brazy PC, Shah JY, Teachman BA, Nosek BA. Explicit and implicit cognition: A preliminary test of a dual-process theory of cognitive vulnerability to depression. *Behaviour Research and Therapy* 2007;45:1155–1167. [PubMed: 17055450]
- Haidt J. The emotional dog and its rational tail: A social intuitionist approach to moral judgment. *Psychological Review* 2001;108:814–834. [PubMed: 11699120]
- Halperin JM, Kalmar JH, Schulz KP, Marks DJ, Sharma V, Newcorn JH. Elevated childhood serotonergic function protects against adolescent aggression in disruptive boys. *Journal of the American Academy of Child & Adolescent Psychiatry* 2006;45:833–840. [PubMed: 16832320]
- Halperin JM, Sharma V, Siever LJ, Schwartz ST, Matier K, Wornell G, et al. Serotonergic function in aggressive and nonaggressive boys with attention deficit hyperactivity disorder. *American Journal of Psychiatry* 1994;151:243–248. [PubMed: 8296897]
- Handley SL. 5-hydroxytryptamine pathways in anxiety and its treatment. *Pharmacology & Therapeutics* 1995;66:103–148. [PubMed: 7630926]
- Hansenne M, Ansseau M. Harm avoidance and serotonin. *Biological Psychology* 1999;51:77–81. [PubMed: 10579422]
- Hariri AR, Brown SM, Williamson DE, Flory JD, de Wit H, Manuck SB. Preference for immediate over delayed rewards is associated with magnitude of ventral striatal activity. *The Journal of Neuroscience* 2006;26:13213–13217. [PubMed: 17182771]
- Hariri AR, Drabant EM, Munoz KE, Kolachana BS, Mattay VS, Egan MF, et al. A susceptibility gene for affective disorders and the response of the human amygdala. *Archives of General Psychiatry* 2005;62:146–152. [PubMed: 15699291]
- Hariri AR, Drabant EM, Weinberger DR. Imaging genetics: Perspectives from studies of genetically driven variation in serotonin function and corticolimbic affective processing. *Biological Psychiatry* 2006;59:888–897. [PubMed: 16442081]
- Hariri AR, Holmes A. Genetics of emotional regulation: The role of the serotonin transporter in neural function. *Trends in Cognitive Science* 2006;10:182–191.
- Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, et al. Serotonin transporter genetic variation and the response of the human amygdala. *Science* 2002 July 19;297:400–403. [PubMed: 12130784]
- Hariri AR, Weinberger DR. Functional neuroimaging of genetic variation in serotonergic neurotransmission. *Genes, Brain, and Behavior* 2003;2:341–349.
- Harmon-Jones E, Allen JJB. Behavioral activation sensitivity and resting frontal EEG asymmetry: Covariation of putative indicators related to risk for mood disorders. *Journal of Abnormal Psychology* 1997;106:159–163. [PubMed: 9103728]
- Hayward G, Goodwin GM, Cowen PJ, Harmer CJ. Low-dose tryptophan depletion in recovered depressed patients induces changes in cognitive processing without depressive symptoms. *Biological Psychiatry* 2005;57:517–524. [PubMed: 15737667]
- Heils A, Teufel A, Petri S, Stober G, Riederer P, Bengel D, et al. Allelic variation of human serotonin transporter gene expression. *Journal of Neurochemistry* 1996;66:2621–2624. [PubMed: 8632190]
- Heinz A, Braus DF, Smolka MN, Wrase J, Puls I, Hermann D, et al. Amygdala-prefrontal coupling depends on a genetic variation of the serotonin transporter. *Nature Neuroscience* 2005;8:20–21.
- Heinz A, Jones DW, Bissette G, Hommer D, Ragan P, Knable M, et al. Relationship between cortisol and serotonin metabolites and transporters in alcoholism. *Pharmacopsychiatry* 2002;35:127–134. [PubMed: 12163982]
- Heinz A, Jones DW, Mazzanti C, Goldman D, Ragan P, Hommer D, et al. A relationship between serotonin transporter genotype and in vivo protein expression and alcohol neurotoxicity. *Biological Psychiatry* 2000;47:643–649. [PubMed: 10745057]

- Heinz A, Smolka MN, Braus DF, Wrase J, Beck A, Flor H, et al. Serotonin transporter genotype (5-HTTLPR): Effects of neutral and undefined conditions on amygdala activation. *Biological Psychiatry* 2007;61:1011–1014. [PubMed: 17157270]
- Hennig J, Reuter M, Netter P, Burk C, Landt O. Two types of aggression are differentially related to serotonergic activity and the A779C TPH polymorphism. *Behavioral Neuroscience* 2005;119:16–25. [PubMed: 15727508]
- Hennig J, Toll C, Schonlau P, Rohrmann S, Netter P. Endocrine responses after d-fenfluramine and ipsapirone challenge: Further support for Cloninger's tridimensional model of personality. *Neuropsychobiology* 2000;41:38–47. [PubMed: 10629434]
- Henriques JB, Davidson RJ. Regional brain electrical asymmetries discriminate between previously depressed and healthy control subjects. *Journal of Abnormal Psychology* 1990;99:22–31. [PubMed: 2307762]
- Henriques JB, Davidson RJ. Left frontal hypoactivation in depression. *Journal of Abnormal Psychology* 1991;100:535–545. [PubMed: 1757667]
- Henriques JB, Davidson RJ. Decreased responsiveness to reward in depression. *Cognition and Emotion* 2000;14:711–724.
- Henriques JB, Glowacki JM, Davidson RJ. Reward fails to alter response bias in depression. *Journal of Abnormal Psychology* 1994;103:460–466. [PubMed: 7930045]
- Hensler JG. Serotonergic modulation of the limbic system. *Neuroscience Biobehavioral Review* 2006;30:203–214.
- Hertel, PT. The cognitive-initiative account of depression-related impairments in memory. In: Medin, D., editor. *The psychology of learning and motivation*. Vol. Vol. 39. New York: Academic Press; 2000. p. 47-71.
- Hertel PT, Rude SS. Depressive deficits in memory. Focusing attention improves subsequent recall. *Journal of Experimental Psychology: General* 1991;120:301–309. [PubMed: 1836493]
- Hoefgen B, Schulze TG, Ohlraun S, von Widdern O, Höfels S, Gross M, et al. The power of sample size and homogenous sampling: Association between the 5-HTTLPR serotonin transporter polymorphism and major depressive disorder. *Biological Psychiatry* 2005;57:247–251. [PubMed: 15691525]
- Holmes AJ, Pizzagalli DA. Task feedback effects on conflict monitoring and executive control: Relationship to subclinical measures of depression. *Emotion* 2007;7:68–76. [PubMed: 17352564]
- Horn NR, Dolan M, Elliott R, Deakin JFW, Woodruff PWR. Response inhibition and impulsivity: An fMRI study. *Neuropsychologia* 2003;41:1959–1966. [PubMed: 14572528]
- Howland, RH.; Thase, ME. Affective disorders: Biological aspects. In: Millon, T.; Blaney, PH.; Davis, RD., editors. *Oxford textbook of psychopathology*. New York: Oxford University Press; 1999. p. 166-202.
- Hu X, Oroszi G, Chun J, Smith TL, Goldman D, Schuckit MA. An expanded evaluation of the relationship of four alleles to the level of response to alcohol and the alcoholism risk. *Alcoholism, Clinical and Experimental Research* 2005;29:8–16.
- Hu X, Zhu G, Lipsky R, Goldman D. HTTLPR allele expression is codominant, correlating with gene effects on fMRI and SPECT imaging, intermediate phenotypes, and behavior. *Biological Psychiatry* 2004;55:191S.
- Hundt NE, Kimbrel NA, Mitchell JT, Nelson-Gray RO. High BAS, but not low BIS, predicts externalizing symptoms in adults. *Personality and Individual Differences* 2008;44:565–575.
- Hyman SE, Malenka RC. Addiction and the brain: The neurobiology of compulsion and its persistence. *Nature Reviews Neuroscience* 2001;2:695–703.
- Jacobs N, Kenis G, Peeters F, Derom C, Villetinck R, van Os J. Stress-related negative affectivity and genetically altered serotonin transporter function. *Archives of General Psychiatry* 2006;63:989–996. [PubMed: 16953001]
- Jensen-Campbell LA, Rosseli M, Workman KA, Santisi M, Rios JD, Bojan D. Agreeableness, conscientiousness and effortful control processes. *Journal of Research in Personality* 2002;36:476–489.

- Jokela M, Keltikangas-Järvinen L, Kivimäki M, Puttonen S, Elovainio M, Rontu R, et al. Serotonin receptor 2A gene and the influence of childhood maternal nurturance on adulthood depressive symptoms. *Archives of General Psychiatry* 2007;64:356–360. [PubMed: 17339524]
- Jokela M, Räikkönen K, Lehtimäki T, Rontu R, Keltikangas-Järvinen L. Tryptophan hydroxylase 1 gene (TPH1) moderates the influence of social support on depressive symptoms in adults. *Journal of Affective Disorders* 2007;100:191–197. [PubMed: 17134762]
- Jonides J, Smith EE, Marshuetz C, Koeppel RA, Reuter-Lorenz PA. Inhibition in verbal working memory revealed by brain activation. *Proceedings of the National Academy of Sciences of the United States of America* 1998;95:8410–8413. [PubMed: 9653200]
- Joormann J. Attentional bias in dysphoria: The role of inhibitory processes. *Cognition and Emotion* 2004;18:125–147.
- Joormann J, Gotlib IH. Updating the contents of working memory in depression: Interference from irrelevant negative material. *Journal of Abnormal Psychology* 2008;117:182–192. [PubMed: 18266496]
- Joormann J, Yoon KL, Zetsche U. Cognitive inhibition in depression. *Applied and Preventive Psychology* 2007;12:128–139.
- Kagan J. A trio of concerns. *Perspectives on Psychological Science* 2007;2:361–376.
- Kahneman D. A perspective on judgment and choice: Mapping bounded rationality. *American Psychologist* 2003;58:697–720. [PubMed: 14584987]
- Kane MJ, Brow LH, McVay JC, Sylvia PJ, Myin-Germeys I, Kwapil TR. For whom the mind wanders, and when. *Psychological Science* 2007;18:614–621. [PubMed: 17614870]
- Kasch KL, Rottenberg J, Arnow BA, Gotlib IH. Behavioral activation and inhibition systems and the severity and course of depression. *Journal of Abnormal Psychology* 2002;111:589–597. [PubMed: 12428772]
- Kaufman J, Yang B-Z, Douglas-Palumberi H, Grasso D, Lipschitz D, Houshyar S, et al. Brain-derived neurotrophic factor-5-HTTLPR gene interactions and environmental modifiers of depression in children. *Biological Psychiatry* 2006;59:673–680. [PubMed: 16458264]
- Kaufman J, Yang B-Z, Douglas-Palumberi H, Houshyar S, Lipschitz D, Krystal JH, et al. Social supports and serotonin transporter gene moderate depression in maltreated children. *Proceedings of the National Academy of Sciences of the United States of America* 2004;101:17316–17321. [PubMed: 15563601]
- Kendler KS, Gatz M, Gardner CO, Pedersen NL. A Swedish national twin study of lifetime major depression. *American Journal of Psychiatry* 2006;163(1):109–114. [PubMed: 16390897]
- Kendler KS, Kuhn JW, Prescott CA, Riley B. The interaction of stressful life events and a serotonin transporter polymorphism in the prediction of episodes of major depression: A replication. *Archives of General Psychiatry* 2005;62:529–535. [PubMed: 15867106]
- Kennedy SH, Evans KR, Krüger S, Mayberg HS, Meyer JH, McCann S, et al. Changes in regional brain glucose metabolism measured with positron emission tomography after paroxetine treatment of major depression. *American Journal of Psychiatry* 2001;158:899–905. [PubMed: 11384897]
- Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al. The epidemiology of major depressive disorder: Results from the National Comorbidity Survey Replication (NCS-R). *Journal of the American Medical Association* 2003;289:3095–3105. [PubMed: 12813115]
- Kieras JE, Tobin RM, Graziano WG, Rothbart MK. You can't always get what you want: Effortful control and children's responses to undesirable gifts. *Psychological Science* 2005;16:391–396. [PubMed: 15869699]
- Kilpatrick DG, Koenen KC, Ruggiero KJ, Acierno R, Galea S, Resnick HS, et al. The serotonin transporter genotype and social support and moderation of posttraumatic stress disorder and depression in hurricane-exposed adults. *American Journal of Psychiatry* 2007;164:1693–1699. [PubMed: 17974934]
- Kim J-M, Stewart R, Kim S-W, Yang S-J, Shin I-S, Kim Y-H, et al. Interactions between life stressors and susceptibility genes (5-HTTLPR and BDNF) on depression in Korean elders. *Biological Psychiatry* 2007;62:423–428. [PubMed: 17482146]

- Klaassen T, Riedel WJ, Honig A, Van Someren A, Deutz NEP, Van Praag HM. Mood effects of 24-hour tryptophan depletion in healthy first degree relatives of patients with affective disorders. *Biological Psychiatry* 1999;46:489–497. [PubMed: 10459398]
- Knutson B, Fong GW, Adams CM, Varner JL, Hommer D. Dissociation of reward anticipation and outcome with event-related fMRI. *NeuroReport* 2001;12:3683–3687. [PubMed: 11726774]
- Knutson B, Wolkowitz OM, Cole SW, Chan T, Moore EA, Johnson RC, et al. Selective alteration of personality and social behavior by serotonergic intervention. *American Journal of Psychiatry* 1998;155:373–379. [PubMed: 9501748]
- Kochanska G, Knaack A. Effortful control as a personality characteristic of young children: Antecedents, correlates, and consequences. *Journal of Personality* 2003;71:1087–1112. [PubMed: 14633059]
- Koob GF, Le Moal M. Addiction and the brain antireward system. *Annual Review of Psychology* 2008;59:29–53.
- Kraft JB, Peters EJ, Slager SL, Jenkins GD, Reinalda MS, McGrath PJ, et al. Analysis of association between the serotonin transporter and antidepressant response in a large clinical sample. *Biological Psychiatry* 2007;61:734–742. [PubMed: 17123473]
- Kringelbach ML, Rolls ET. The functional neuroanatomy of the human orbitofrontal cortex: Evidence from neuroimaging and neuropsychology. *Progress in Neurobiology* 2004;72:341–372. [PubMed: 15157726]
- Krueger RF. Personality from a realist's perspective: Personality traits, criminal behaviors, and the externalizing spectrum. *Journal of Research in Personality* 2002;36:564–572.
- Krueger RF, Hicks BM, Patrick CJ, Carlson SR, Iacono WG, McGue M. Etiologic connections among substance dependence, antisocial behavior, and personality: Modeling the externalizing spectrum. *Journal of Abnormal Psychology* 2002;111:411–424. [PubMed: 12150417]
- Kurzban R, Aktipis CA. Modularity and the social mind: Are psychologists too self-ish? *Personality and Social Psychology Review* 2007;11:131–149. [PubMed: 18453459]
- Lam RW, Zis AP, Grewal A, Delgado PL, Charney DS, Krystal JH. Effects of rapid tryptophan depletion in patients with seasonal affective disorder in remission after light therapy. *Archives of General Psychiatry* 1996;53:41–44. [PubMed: 8540776]
- LeDoux JE. Emotion circuits in the brain. *Annual Review of Neuroscience* 2000;23:155–184.
- LeDoux, JE. *Synaptic self: How our brains become who we are*. New York: Viking; 2002.
- Lenzenweger MF, Willett JB. Predicting individual change in personality disorder features by simultaneous individual change in personality dimensions linked to neurobehavioral systems: The longitudinal study of personality disorders. *Journal of Abnormal Psychology* 2007;116:684–700. [PubMed: 18020716]
- Leonardo ED, Hen R. Genetics of affective and anxiety disorders. *Annual Review of Psychology* 2006;57:117–137.
- Lesch K-P, Bengel D, Heils A, Sabol SZ, Greenberg BD, Petri S, et al. Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science* 1996 November 29;274:1527–1531. [PubMed: 8929413]
- Lesch, K-P.; Canli, T. 5-HT_{1A} receptor and anxiety-related traits. In: Canli, T., editor. *Biology of personality and individual differences*. New York: Guilford Press; 2006. p. 273-294.
- Lévesque J, Fanny E, Joannette Y, Paquette V, Mensour B, Beaudoin G, et al. Neural circuitry underlying voluntary suppression of sadness. *Biological Psychiatry* 2003;53:502–510. [PubMed: 12644355]
- Levin RL, Heller W, Mohanty A, Herrington JD, Miller GA. Cognitive deficits in depression and functional specificity of regional brain activity. *Cognitive Therapy and Research* 2007;31:211–233.
- Leyton M, Young SN, Blier P, Ellenbogen MA, Palmour RM, Ghadirian A-M, et al. The effect of tryptophan depletion on mood in medication-free, former patients with major affective disorder. *Neuropsychopharmacology* 1997;16:294–297. [PubMed: 9094147]
- Liao D-L, Hong C-J, Shih H-L, Tsai S-J. Possible association between serotonin transporter promoter region polymorphism and extremely violent crime in Chinese males. *Neuropsychobiology* 2004;50:284–287. [PubMed: 15539857]
- Lichtenberg P, Shapira B, Gillon D, Kindler S, Cooper TB, Newman ME, et al. Hormone responses to fenfluramine and placebo challenge in endogenous depression. *Psychiatry Research* 1992;43:137–146. [PubMed: 1410069]

- Lieberman MD. Intuition: A social cognitive neuroscience approach. *Psychological Bulletin* 2000;126:109–137. [PubMed: 10668352]
- Lieberman MD. Social cognitive neuroscience: A review of core processes. *Annual Review of Psychology* 2007;58:259–289.
- Lieberman, MD.; Gaunt, R.; Gilbert, DT.; Trope, Y. Reflection and reflexion: A social cognitive neuroscience approach to attributional inference. In: Zanna, M., editor. *Advances in experimental social psychology*. Vol. Vol. 34. San Diego, CA: Academic Press; 2002. p. 199–249.
- Little KY, McLaughlin DP, Zhang L, Livermore CS, Dalack GW, McFinton PR, et al. Cocaine, ethanol, and genotype effects on human midbrain serotonin transporter binding sites and mRNA levels. *American Journal of Psychiatry* 1998;155:207–213. [PubMed: 9464199]
- Lonigan CJ, Vasey MW, Phillips BM, Hazen RA. Temperament, anxiety, and the processing of threat-relevant stimuli. *Journal of Clinical Child and Adolescent Psychology* 2004;33:8–20. [PubMed: 15028537]
- Looper KJ, Paris J. What dimensions underlie Cluster B personality disorders? *Comprehensive Psychiatry* 2000;41:432–437. [PubMed: 11086148]
- Lopez Ibor JJ Jr, Saiz Ruiz J, Iglesias LM. The fenfluramine challenge test in the affective spectrum: A possible marker of endogeneity and severity. *Pharmacopsychiatry* 1988;21:9–14. [PubMed: 3283785]
- Lyons-Ruth K, Holmes BM, Sasvari-Szekely M, Ronai Z, Nemoda Z, Pauls D. Serotonin transporter polymorphism and borderline or antisocial traits among low-income young adults. *Psychiatric Genetics* 2007;17:339–343. [PubMed: 18075474]
- Mann JJ, Currier D. A review of prospective studies of biologic predictors of suicidal behavior in mood disorders. *Archives of Suicide Research* 2007;11:3–16. [PubMed: 17178639]
- Mann JJ, Huang YY, Underwood MD, Kassir SA, Oppenheim S, Kelly TM, et al. A serotonin transporter gene promoter polymorphism (5-HTTLPR) and prefrontal cortical binding in major depression and suicide. *Archives of General Psychiatry* 2000;57:729–738. [PubMed: 10920459]
- Mann JJ, Waternaux C, Haas GL, Malone KM. Toward a clinical model of suicidal behavior in psychiatric patients. *American Journal of Psychiatry* 1999;156:181–189. [PubMed: 9989552]
- Manuck SB, Flory JD, McCaffery JM, Matthews KA, Mann JJ, Muldoon MF. Aggression, impulsivity, and central nervous system serotonergic responsivity in a nonpatient sample. *Neuropsychopharmacology* 1998;19:287–299. [PubMed: 9718592]
- Manuck, SB.; Flory, JD.; Muldoon, MF.; Ferrell, RE. A neurobiology of intertemporal choice. In: Loewenstein, G.; Read, D.; Baumeister, RF., editors. *Time and decision: Economic and psychological perspectives on intertemporal choice*. New York: Russell Sage Foundation; 2003. p. 139–172.
- Manuck, SB.; Kaplan, JR.; Lotrich, FE. Brain serotonin and aggressive disposition in humans and nonhuman primates. In: Nelson, RJ., editor. *Biology of aggression*. New York: Oxford University Press; 2006. p. 65–102.
- Martinowich K, Lu B. Interaction between BDNF and serotonin: Role in mood disorders. *Neuropsychopharmacology Reviews* 2007;33:73–83. [PubMed: 17882234]
- Mayberg HS. Limbic-cortical dysregulation: A proposed model of depression. *Journal of Neuropsychiatry and Clinical Neurosciences* 1997;9:471–481. [PubMed: 9276848]
- Mayberg, HS. Mapping mood: An evolving emphasis on frontallimbic interactions. In: Stuss, DT.; Knight, RT., editors. *Principles of frontal lobe function*. New York: Oxford University Press; 2002. p. 376–391.
- Mayberg HS. Positron emission tomography imaging in depression: A neural systems perspective. *Neuroimaging Clinics of North America* 2003;13:805–815. [PubMed: 15024963]
- Mayberg, HS. Depression: A neuropsychiatric perspective. In: Panksepp, J., editor. *Textbook of biological psychiatry*. New York: Wiley; 2004. p. 197–229.
- McClelland, DC. Human motivation. Glenview, IL: Scott, Foresman; 1984.
- McClelland DC. How motives, skills, and values determine what people do. *American Psychologist* 1985;40:812–825.
- McClelland DC, Koestner R, Weinberger J. How do self-attributed and implicit motives differ? *Psychological Review* 1989;96:690–702.

- McClelland, J.L. Cognitive modeling, connectionist. In: Wilson, R.W.; Keil, F.C., editors. *The MIT encyclopedia of the cognitive sciences*. Cambridge, MA: MIT Press; 1999. p. 137-139.
- McClure SM, Laibson DI, Loewenstein G, Cohen JD. Separate neural systems value immediate and delayed monetary rewards. *Science* 2004 October 15;306:503-507. [PubMed: 15486304]
- McConnell AR, Rydell RJ, Strain LM, Mackie DM. Forming implicit and explicit attitudes toward individuals: Social group association cues. *Journal of Personality and Social Psychology* 2008;94:792-807. [PubMed: 18444739]
- McFarland BR, Shankman SA, Tenke CE, Bruder GE, Klein DN. Behavioral activation system deficits predict the six-month course of depression. *Journal of Affective Disorders* 2006;91:229-234. [PubMed: 16487598]
- Metcalf J, Mischel W. A hot/cool-system analysis of delay of gratification: Dynamics of willpower. *Psychological Review* 1999;106:3-19. [PubMed: 10197361]
- Milham MP, Banich MT. Anterior cingulate cortex: An fMRI analysis of conflict specificity and functional differentiation. *Human Brain Mapping* 2005;25:328-335. [PubMed: 15834861]
- Mischel, W. Processes in delay of gratification. In: Berkowitz, L., editor. *Advances in experimental social psychology*. Vol. 7. New York: Academic Press; 1974. p. 249-292.
- Mitchell RLC. Anterior cingulate activity and level of cognitive conflict: Explicit comparisons. *Behavioral Neuroscience* 2006;120:1395-1401. [PubMed: 17201486]
- Mohanty A, Engels AS, Herrington JD, Heller W, Ho MR, Banich MT, et al. Differential engagement of anterior cingulate cortex subdivisions for cognitive and emotional function. *Psychophysiology* 2007;44:343-351. [PubMed: 17433093]
- Monroe SM, Simmons AD. Diathesis-stress theories in the context of life stress research: Implications for the depressive disorders. *Psychological Bulletin* 1991;110:406-425. [PubMed: 1758917]
- Moon H. The two faces of conscientiousness: Duty and achievement striving in escalation of commitment dilemmas. *Journal of Applied Psychology* 2001;86:535-540.
- Moore P, Gillin JC, Bhatti T, DeModena A, Seifritz E, Clark C, et al. Rapid tryptophan depletion, sleep electroencephalogram, and mood in men with remitted depression on serotonin reuptake inhibitors. *Archives of General Psychiatry* 1998;55:534-539. [PubMed: 9633672]
- Moore P, Landolt H-P, Seifritz E, Clark C, Bhatti T, Kelsoe J, et al. Clinical and physiological consequences of rapid tryptophan depletion. *Neuropsychopharmacology* 2000;23:601-622. [PubMed: 11063917]
- Moreno FA, Gelenberg AJ, Heninger GR, Potter RL, McKnight KM, Allen J, et al. Tryptophan depletion and depressive vulnerability. *Biological Psychiatry* 1999;46:498-505. [PubMed: 10459399]
- Moreno FA, McGahuey CA, Freeman MP, Delgado PL. Sex differences in depressive response during monoamine depletions in remitted depressive subjects. *Journal of Clinical Psychiatry* 2006;67:1618-1623. [PubMed: 17107256]
- Munafò MR, Brown SM, Hariri AR. Serotonin transporter (5-HTTLPR) genotype and amygdala activation: A meta-analysis. *Biological Psychiatry* 2008;63:852-857. [PubMed: 17949693]
- Munafò MR, Clark T, Flint J. Does measurement instrument moderate the association between the serotonin transporter gene and anxiety-related personality traits? A meta-analysis. *Molecular Psychiatry* 2005a;10:415-419.
- Munafò MR, Clark T, Flint J. Promise and pitfalls in the meta-analysis of genetic association studies: A response to Sen and Shinka. *Molecular Psychiatry* 2005b;10:895-897.
- Munafò MR, Hayward G, Harmer C. Selective processing of social threat cues following acute tryptophan depletion. *Journal of Psychopharmacology* 2006;20:33-39.
- Murphy FC, Smith KA, Cowen PJ, Robbins TW, Sahakian BJ. The effects of tryptophan depletion on cognitive and affective processing in healthy volunteers. *Psychopharmacology* 2002;163:42-53. [PubMed: 12185399]
- Murphy SE, Longhitano C, Ayres RE, Cowen PJ, Harmer CJ. Tryptophan supplementation induces a positive bias in the processing of emotional material in healthy female volunteers. *Psychopharmacology* 2006;187:121-130. [PubMed: 16767422]
- Murray KT, Kochanska G. Effortful control: Factor structure and relation to externalizing and internalizing behaviors. *Journal of Abnormal Child Psychology* 2002;30:503-514. [PubMed: 12403153]

- Nakamura M, Ueno S, Sano A, Tanabe H. The human serotonin transporter gene linked polymorphism (5-HTTLPR) shows ten novel allelic variants. *Molecular Psychiatry* 2000;5:32–38. [PubMed: 10673766]
- Naranjo CA, Tremblay LK, Busto UE. The role of the brain reward system in depression. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 2001;25:781–823. [PubMed: 11383978]
- Nestler EJ, Carlezon WJ. The mesolimbic dopamine reward circuit in depression. *Biological Psychiatry* 2006;59:1151–1159. [PubMed: 16566899]
- Netter P, Hennig J, Rohrmann S. Psychobiological differences between the aggression and psychoticism dimension. *Pharmacopsychiatry* 1999;32:5–12. [PubMed: 10071177]
- Neumeister A, Hu X, Luckenbaugh DA, Schwarz M, Nugent AC, Bonne O, et al. Differential effects of 5-HTTLPR genotypes on the behavioral and neural responses to tryptophan depletion in patients with major depression and controls. *Archives of General Psychiatry* 2006;63:978–986. [PubMed: 16953000]
- Neumeister A, Konstantinidis A, Stastny J, Schwarz MJ, Vitouch O, Willeit M, et al. Association between serotonin transporter gene promoter polymorphism (5-HTTLPR) and behavioral responses to tryptophan depletion in healthy women with and without family history of depression. *Archives of General Psychiatry* 2002;59:613–620. [PubMed: 12090814]
- Neumeister A, Nugent AC, Waldeck T, Geraci M, Schwarz M, Bonne O, et al. Neural and behavioral responses to tryptophan depletion in unmedicated patients with remitted major depressive disorder and controls. *Archives of General Psychiatry* 2004;61:765–773. [PubMed: 15289275]
- New AS, Hazlett EA, Buchsbaum MS, Goodman M, Reynolds D, Metropoulou V, et al. Blunted prefrontal cortical ¹⁸fluorodeoxyglucose positron emission tomography response to meta-chlorophenylpiperazine in impulsive aggression. *Archives of General Psychiatry* 2002;59:621–629. [PubMed: 12090815]
- Nigg JT. On inhibition/disinhibition in developmental psychopathology: Views from cognitive and personality psychology as a working inhibition taxonomy. *Psychological Bulletin* 2000;126:220–246. [PubMed: 10748641]
- Nigg JT. Is ADHD a disinhibitory disorder? *Psychological Bulletin* 2001;127:571–598. [PubMed: 11548968]
- Nigg JT. Response inhibition and disruptive behaviors: Toward a multiprocess conception of etiological heterogeneity for ADHD combined type and conduct disorder early-onset type. *Annals of the New York Academy of Sciences* 2003;1008:170–182. [PubMed: 14998883]
- Nigg JT. Temperament and developmental psychopathology. *Journal of Child Psychology and Psychiatry* 2006;47:395–422. [PubMed: 16492265]
- Nolen-Hoeksema S. The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *Journal of Abnormal Psychology* 2000;109:504–511. [PubMed: 11016119]
- Nolen-Hoeksema S, Morrow J, Fredrickson BL. Response styles and the duration of episodes of depressed mood. *Journal of Abnormal Psychology* 1993;102:20–28. [PubMed: 8436695]
- Norman, DA. Reflections on cognition and parallel distributed processing. In: McClelland, JL.; Rumelhart, DE.; the PDP Research Group. , editors. *Parallel distributed processing: Explorations in the microstructure of cognition: Vol. 2. Psychological and biological models*. Cambridge, MA: MIT Press; 1986. p. 531-546.
- Ochsner, KN.; Barrett, LF. A multiprocess perspective on the neuroscience of emotion. New York: Guilford Press; 2001.
- Ochsner KN, Bunge SA, Gross JJ, Gabrieli JD. Rethinking feelings: An fMRI study of the cognitive regulation of emotion. *Journal of Cognitive Neuroscience* 2002;14:1215–1229. [PubMed: 12495527]
- Ochsner KN, Gross JJ. The cognitive control of emotion. *Trends in Cognitive Sciences* 2005;9:242–249. [PubMed: 15866151]
- Ochsner KN, Ray RD, Cooper JC, Robertson ER, Chopra S, Gabrieli JDE, Gross JJ. For better or for worse: Neural systems supporting the cognitive down- and up-regulation of negative emotion. *NeuroImage* 2004;23:483–499. [PubMed: 15488398]

- O'Connor BP. The search for dimensional structure differences between normality and abnormality: A statistical review of published data on personality and psychopathology. *Journal of Personality and Social Psychology* 2002;83:962–982. [PubMed: 12374447]
- O'Connor LH, Fischette CT. Hormone effects on serotonin-dependent behaviors. *Annals of the New York Academy of Sciences* 1986;474:437–444. [PubMed: 3555242]
- O'Keane V, Dinan TG. Prolactin and cortisol responses to *d*-fenfluramine in major depression: Evidence for diminished responsivity of central serotonergic function. *American Journal of Psychiatry* 1991;148:1009–1015. [PubMed: 1853948]
- O'Keane V, Moloney E, O'Neill H, O'Connor A, Smith C, Dinan TG. Blunted prolactin responses to *d*-fenfluramine in sociopathy: Evidence for subsensitivity of central serotonergic function. *British Journal of Psychiatry* 1992;160:643–646. [PubMed: 1591573]
- O'Keane V, O'Hanlon M, Webb M, Dinan T. *d*-fenfluramine/prolactin response throughout the menstrual cycle: Evidence for an oestrogen-induced alteration. *Clinical Endocrinology* 1991;34:289–292. [PubMed: 1879060]
- Olesen PJ, Westerberg H, Klingberg T. Increased prefrontal and parietal activity after training of working memory. *Nature Neuroscience* 2004;7:75–79.
- Panksepp J, Cox J. An overdue burial for the serotonin theory of anxiety. *Behavioral and Brain Sciences* 1986;9:340–341.
- Park SB, Coull JT, McShane RH, Young AH, Sahakian BJ, Robbins TW, et al. Tryptophan depletion in normal volunteers produces selective impairments in learning and memory. *Neuropharmacology* 1994;33:575–588. [PubMed: 7984295]
- Parsey RV, Hastings RS, Oquendo MA, Huang YY, Simpson N, Arcement J, et al. Lower serotonin transporter binding potential in the human brain during major depressive episodes. *American Journal of Psychiatry* 2006;163:52–58. [PubMed: 16390889]
- Paus T. Primate anterior cingulate cortex, where motor control, drive and cognition interface. *Nature Reviews Neuroscience* 2001;2:417–424.
- Peluso MAM, Hatch JP, Glahn DC, Monkul ES, Sanches M, Najt P, et al. Trait impulsivity in patients with mood disorders. *Journal of Affective Disorders* 2007;100:227–231. [PubMed: 17097740]
- Pezawas L, Meyer-Lindenberg A, Drabant EM, Verchinski BA, Munoz KE, Kolachana BS, et al. 5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: A genetic susceptibility mechanism for depression. *Nature Neuroscience* 2005;8:828–834.
- Pinto-Meza A, Caseras X, Soler J, Puigdemont D, Perez V, Torrubia R. Behavioural inhibition and behavioural activation systems in current and recovered major depression participants. *Personality and Individual Differences* 2006;40:215–226.
- Pizzagalli DA, Oakes TR, Fox AS, Chung MK, Larson CL, Abercrombie HC, et al. Functional but not structural subgenual prefrontal cortex abnormalities in melancholia. *Molecular Psychiatry* 2004;9:325. [PubMed: 14699431]
- Pizzagalli D, Pascual-Marqui RD, Nitschke JB, Oakes TR, Larson CL, Abercrombie HC, et al. Anterior cingulate activity as a predictor of degree of treatment response in major depression: Evidence from brain electrical tomography analysis. *American Journal of Psychiatry* 2001;158:405–415. [PubMed: 11229981]
- Placidi GPA, Oquendo MA, Malone KM, Huang Y-Y, Ellis SP, Mann JJ. Aggressivity, suicide attempts, and depression: Relationship to cerebrospinal fluid monoamine metabolite levels. *Biological Psychiatry* 2001;50:783–791. [PubMed: 11720697]
- Porter RJ, Phipps AJ, Gallagher P, Scott A, Stevenson PS, O'Brien JT. Effects of acute tryptophan depletion on mood and cognitive functioning in older recovered depressed subjects. *American Journal of Geriatric Psychiatry* 2005;13:607–615. [PubMed: 16009737]
- Posner MI, DiGirolamo GJ. Cognitive neuroscience: Origins and promise. *Psychological Bulletin* 2000;126:873–889. [PubMed: 11107880]
- Posner MI, Rothbart MK. Research on attention networks as a model for the integration of psychological science. *Annual Review of Psychology* 2007;58:1–23.
- Power AC, Cowen PJ. Neuroendocrine challenge tests: Assessment of 5-HT function in anxiety and depression. *Molecular Aspects of Medicine* 1992;13:205–220. [PubMed: 1435104]

- Prange AJ Jr, Wilson IC, Lynn CW, Alltop LB, Stikeleather RA. L-tryptophan in mania. Contribution to a permissive hypothesis of affective disorders. *Archives of General Psychiatry* 1974;30:56–62. [PubMed: 4587067]
- Pryor JB, Reeder GD, Yeadon C, Hesson-McInnis M. A dual-process model of reactions to perceived stigma. *Journal of Personality and Social Psychology* 2004;87:436–452. [PubMed: 15491270]
- Quay HC. Theories of ADHD. *Journal of the American Academy of Child & Adolescent Psychiatry* 1988;27:262–263. [PubMed: 3360739]
- Quintin P, Benkelfat C, Launay JM, Arnulf I, Pointereau-Bellenger A, Barbault S, et al. Clinical and neurochemical effect of acute tryptophan depletion in unaffected relatives of patients with bipolar affective disorder. *Biological Psychiatry* 2001;50:184–190. [PubMed: 11513817]
- Raine A, Buchsbaum MS, Stanley J, Lottenberg S, Abel L, Stoddard J. Selective reductions in prefrontal glucose metabolism in murderers. *Biological Psychiatry* 1994;36:365–373. [PubMed: 7803597]
- Raine A, Yang Y. Neural foundations to moral reasoning and antisocial behavior. *Social Cognitive and Affective Neuroscience* 2006;1:203–213. [PubMed: 18985107]
- Rajkowska G. Postmortem studies in mood disorders indicate altered numbers of neurons and glial cells. *Biological Psychiatry* 2000;48:766–777. [PubMed: 11063973]
- Rao H, Gillihan SJ, Wang J, Korczykowski M, Sankoorikal GMV, Kaercher KA, et al. Genetic variation in serotonin transporter alters resting brain function in healthy individuals. *Biological Psychiatry* 2007;62:600–606. [PubMed: 17481593]
- Ray RD, Oschner KN, Cooper JC, Robertson ER, Gabrieli JDE, Gross JJ. Individual differences in trait rumination and the neural systems supporting cognitive reappraisal. *Cognitive, Affective, & Behavioral Neuroscience* 2005;5:156–168.
- Reid SA, Duke LM, Allen JJB. Resting frontal electroencephalographic asymmetry in depression: Inconsistencies suggest the need to identify mediating factors. *Psychophysiology* 1998;35:389–404. [PubMed: 9643053]
- Reif A, Rösler M, Freitag CM, Schneider M, Eujen A, Kissling C, et al. Nature and nurture predispose to violent behavior: Serotonergic genes and adverse childhood environment. *Neuropsychopharmacology* 2007;32:2375–2383. [PubMed: 17342170]
- Ressler KJ, Nemeroff CB. Role of serotonergic and noradrenergic systems in the pathophysiology of depression and anxiety disorders. *Depression and Anxiety* 2000;12:2–19. [PubMed: 11098410]
- Richell RA, Deakin JFW, Anderson IM. Effect of acute tryptophan depletion on the response to controllable and uncontrollable noise stress. *Biological Psychiatry* 2005;57:295–300. [PubMed: 15691531]
- Rick SI, Cryder CE, Loewenstein G. Tightwads and spendthrifts. *Journal of Consumer Research* 2008;34:767–782.
- Riedel WJ, Klaassen T, Deutz NEP, van Someren A, van Praag HM. Tryptophan depletion in normal volunteers produces selective impairment in memory consolidation. *Psychopharmacology* 1999;141:362–369. [PubMed: 10090643]
- Riedel WJ, Klaassen T, Schmitt JAJ. Tryptophan, mood, and cognitive function. *Brain, Behavior, and Immunity* 2002;16:581–589.
- Roberts BW, Chernyshenko OS, Stark S, Goldberg LR. The structure of conscientiousness: An empirical investigation based on seven major personality questionnaires. *Personnel Psychology* 2005;58:103–139.
- Rogers RD, Tunbridge EM, Bhagwagar Z, Drevets WC, Sahakian BJ, Carter CS. Tryptophan depletion alters the decision-making of healthy volunteers through altered processing of reward cues. *Neuropsychopharmacology* 2003;28:153–162. [PubMed: 12496952]
- Rolls, ET. *Emotion explained*. Oxford, United Kingdom: Oxford University Press; 2005.
- Rothbart MK, Ahadi SA, Evans DE. Temperament and personality: Origins and outcomes. *Journal of Personality and Social Psychology* 2000;78:122–135. [PubMed: 10653510]
- Rothbart MK, Ahadi SA, Hershey K, Fisher P. Investigations of temperament at three to seven years: The Children's Behavior Questionnaire. *Child Development* 2001;72:1394–1408. [PubMed: 11699677]

- Rothbart, MK.; Bates, JE. Temperament. In: Damon, W., editor; Eisenberg, N., editor. Handbook of child psychology: Vol 3. Social, emotional and personality development. 5th ed. New York: Wiley; 1998. p. 105-176.(Series Ed.) & (Vol. Ed.)
- Rothbart, MK.; Ellis, LK.; Posner, MI. Temperament and self-regulation. In: Baumeister, RF.; Vohs, KD., editors. Handbook of self-regulation: Research, theory, and applications. New York: Guilford Press; 2004. p. 357-370.
- Rothbart MK, Ellis LK, Rueda MR, Posner MI. Developing mechanisms of temperamental effortful control. *Journal of Personality* 2003;71:1113–1143. [PubMed: 14633060]
- Rothbart, MK.; Posner, M. Temperament and the development of self-regulation. In: Hartlage, LC.; Telzow, CF., editors. The neuropsychology of individual differences: A developmental perspective. New York: Plenum; 1985. p. 93-123.
- Rudman LA, Phelan JE, Heppen JB. Developmental sources of implicit attitudes. *Personality and Social Psychology Bulletin* 2007;33:1700–1713. [PubMed: 18000104]
- Rydell RJ, McConnell AR. Understanding implicit and explicit attitude change: A systems of reasoning analysis. *Journal of Personality and Social Psychology* 2006;91:995–1008. [PubMed: 17144760]
- Sakai JT, Lessem JM, Haberstick BC, Hopfer CJ, Smolen A, Ehringer MA, et al. Case-control and within-family tests for association between 5HTTLPR and conduct problems in a longitudinal adolescent sample. *Psychiatric Genetics* 2007;17:207–214. [PubMed: 17621163]
- Salamone JD, Correa M, Farrar A, Mingote SM. Effort-related functions of nucleus accumbens dopamine and associated forebrain circuits. *Psychopharmacology* 2007;191:461–482. [PubMed: 17225164]
- Salamone JD, Correa M, Mingote SM, Weber SM. Beyond the reward hypothesis: Alternative functions of nucleus accumbens dopamine. *Current Opinion in Pharmacology* 2005;5:34–41. [PubMed: 15661623]
- Salamone JD, Correa M, Mingote SM, Weber SM, Farrar AM. Nucleus accumbens dopamine and the forebrain circuitry involved in behavioral activation and effort-related decision making: Implications for understanding anergia and psychomotor slowing in depression. *Current Psychiatry Reviews* 2006;2:267–280.
- Sambeth A, Blokland A, Harmer CJ, Kilkens TOC, Nathan PJ, Porter RJ, et al. Sex differences in the effect of acute tryptophan depletion on declarative episodic memory: A pooled analysis of nine studies. *Neuroscience and Biobehavioral Reviews* 2007;31:516–529. [PubMed: 17229463]
- Sansone C, Wiebe DJ, Morgan C. Self-regulating interest: The moderating role of hardiness and conscientiousness. *Journal of Personality* 1999;67:701–733. [PubMed: 10444855]
- Satpute AB, Lieberman MD. Integrating automatic and controlled processes into neurocognitive models of social cognition. *Brain Research* 2006;1079:86–97. [PubMed: 16490183]
- Schaefer A, Collete F, Philippot P, Van der Linden M, Laureys S, Delfiore G, et al. Neural correlates of “hot” and “cold” emotional processing: A multilevel approach to the functional anatomy of emotion. *NeuroImage* 2003;18:938–949. [PubMed: 12725769]
- Schinka JA. Measurement scale does moderate the association between the serotonin transporter gene and trait anxiety: Comments on Munafò et al. *Molecular Psychiatry* 2005;10:892–893. [PubMed: 15983622]
- Schinka JA, Busch RM, Robichaux-Keene N. A meta-analysis of the association between the serotonin transporter gene polymorphism (5-HTTLPR) and trait anxiety. *Molecular Psychiatry* 2004;9:197–202. [PubMed: 14966478]
- Schmitz A, Hennig J, Kuepper Y, Reuter M. The association between neuroticism and the serotonin transporter polymorphism depends on structural differences between personality measures. *Personality and Individual Differences* 2007;42:789–799.
- Schultheiss, OC.; Pang, JS. Measuring implicit motives. In: Robins, RW.; Fraley, RC.; Krueger, R., editors. Handbook of research methods in personality psychology. New York: Guilford Press; 2007. p. 322-344.
- Schwartz CE, Wright CI, Shin LM, Kagan J, Rauch SL. Inhibited and uninhibited infants “grown up”: Adult amygdalar response to novelty. *Science* 2003 June 20;300:1952–1953. [PubMed: 12817151]
- Schwartz CE, Wright CI, Shin LM, Kagan J, Whalen PJ, McMullin KG, et al. Differential amygdalar response to novel versus newly familiar neutral faces: A functional MRI probe developed for studying inhibited temperament. *Biological Psychiatry* 2003;53:854–862. [PubMed: 12742672]

- Sen S, Burmeister M, Ghosh D. Meta-analysis of the association between a serotonin transporter promoter polymorphism (5-HTTLPR) and anxiety-related personality traits. *American Journal of Medical Genetics Part B (Neuropsychiatric Genetics)* 2004a;127B:85–89.
- Sen S, Villafuerte S, Nesse R, Stoltenberg SF, Hopcian J, Gleiberman L, et al. Serotonin transporter and GABA(A) alpha 6 receptor variants are associated with neuroticism. *Biological Psychiatry* 2004b; 55:244–249. [PubMed: 14744464]
- Shankman SA, Klein DN, Tenke CE, Bruder GE. Reward sensitivity in depression: A biobehavioral study. *Journal of Abnormal Psychology* 2007;116:95–104. [PubMed: 17324020]
- Shastri L, Ajjanagadde V. From simple associations to systematic reasoning: A connectionist representation of rules, variables, and dynamic bindings using temporal synchrony. *Behavioral and Brain Sciences* 1993;16:417–494.
- Sheline YI, Barch DM, Donnelly JM, Ollinger JM, Snyder AZ, Mintun MA. Increased amygdala response to masked emotional faces in depressed subjects resolves with antidepressant treatment: An fMRI study. *Biological Psychiatry* 2001;50:651–658. [PubMed: 11704071]
- Sher L, Oquendo MA, Li S, Ellis S, Brodsky BS, Malone KM, et al. Prolactin response to fenfluramine administration in patients with unipolar and bipolar depression and healthy controls. *Psychoneuroendocrinology* 2003;28:559–573. [PubMed: 12689612]
- Shulman GL, Fiez JA, Corbetta M, Buckner RL, Miezin FM, Raichle ME, et al. Decreases in cerebral cortex: Common blood flow changes across visual tasks. *Journal of Cognitive Neuroscience* 1997;9:648–663.
- Siegle GJ, Steinhauer SR, Thase ME, Stenger VA, Carter CS. Can't shake that feeling: Event-related fMRI assessment of sustained amygdala activity in response to emotional information in depressed individuals. *Biological Psychiatry* 2002;51:693–707. [PubMed: 11983183]
- Siever LJ, Buchsbaum MS, New AS, Spiegel-Cohen J, Wei T, Hazlett EA, et al. d,l-fenfluramine response in impulsive personality disorder assessed with [18F]fluorodeoxyglucose positron emission tomography. *Neuropsychopharmacology* 1999;20:413–423. [PubMed: 10192822]
- Siever L, Murphy DL, Slater S, de la Vega E, Lipper S. Plasma prolactin changes following fenfluramine in depressed patients compared with controls: An evaluation of central serotonergic responsivity in depression. *Life Sciences* 1984;34:1029–1039. [PubMed: 6700367]
- Sjoberg RL, Nilsson KW, Nordquist N, Ohrvik J, Leppert J, Lindstrom L, et al. Development of depression: Sex and the interaction between environment and a promoter polymorphism of the serotonin transporter gene. *International Journal of Neuropsychopharmacology* 2006;9:443–449. [PubMed: 16212676]
- Sloman SA. The empirical case for two forms of reasoning. *Psychological Bulletin* 1996;119:3–22.
- Smallwood J, O'Connor RC, Sudberry MV, Obonsawin MC. Mind-wandering and dysphoria. *Cognition and Emotion* 2007;21:816–842.
- Smallwood J, Schooler JW. The restless mind. *Psychological Bulletin* 2006;132:946–958. [PubMed: 17073528]
- Smith ER, DeCoster J. Dual-process models in social and cognitive psychology: Conceptual integration and links to underlying memory systems. *Personality and Social Psychology Review* 2000;4:108–131.
- Smith KA, Clifford EM, Hockney RA, Clark DM, Cowen PJ. Effect of tryptophan depletion on mood in male and female volunteers: A pilot study. *Human Psychopharmacology* 1997;12:111–117.
- Smith KA, Morris JS, Friston KJ, Cowen PJ, Dolan RJ. Brain mechanisms associated with depressive relapse and associated cognitive impairment following acute tryptophan depletion. *British Journal of Psychiatry* 1999;164:525–529. [PubMed: 10616631]
- Smolensky P. On the proper treatment of connectionism. *Behavioral and Brain Sciences* 1988;11:1–23.
- Smolensky, P.; Legendre, G. *The harmonic mind: From neural computation to optimality-theoretic grammar*. Vol. Vols. 1 & 2. Cambridge, MA: MIT Press; 2006.
- Sobczak S, Honig A, van Duinen MA, Riedel WJ. Serotonergic dysregulation in bipolar disorders: A literature review of serotonergic challenge studies. *Bipolar Disorders* 2002;4:347–356. [PubMed: 12519094]

- Sobczak S, Riedel WJ, Booij I, Het Rot AM, Deutz NEP. Cognition following acute tryptophan depletion: Difference between first-degree relatives of bipolar disorder patients and matched healthy control volunteers. *Psychological Medicine* 2002;32:503–512. [PubMed: 11989995]
- Soubrié P. Reconciling the role of central serotonin neurons in human and animal behavior. *Behavioral and Brain Sciences* 1986;9:319–364.
- Spillmann MK, Van der Does AJW, Rankin MA, Vuolo RD, Alpert JE, Nierenberg AA, et al. Tryptophan depletion in SSRI-recovered depressed outpatients. *Psychopharmacology* 2001;155:123–127. [PubMed: 11400999]
- Spoont MR. Modulatory role of serotonin in neural information processing: Implications for human psychopathology. *Psychological Bulletin* 1992;112:330–350. [PubMed: 1454898]
- Stanovich, KE. Who is rational? Studies of individual differences in reasoning. Mahwah, NJ: Erlbaum; 1999.
- Steel P. The nature of procrastination: A meta-analytic and theoretical review of quintessential self-regulatory failure. *Psychological Bulletin* 2007;133:65–94. [PubMed: 17201571]
- Steinberg L. Risk taking in adolescence: New perspectives from brain and behavioral science. *Current Directions in Psychological Science* 2007;16:55–59.
- Strack F, Deutsch R. Reflective and impulsive determinants of social behavior. *Personality and Social Psychology Review* 2004;8:220–247. [PubMed: 15454347]
- Strickland PL, Deakin JFW, Percival C, Dixon J, Gater RA, Goldberg DP. Bio-social origins of depression in the community: Interactions between social adversity, cortisol and serotonin neurotransmission. *British Journal of Psychiatry* 2002;180:168–173. [PubMed: 11823330]
- Strunk DR, DeRubeis RJ, Chiu AW, Alvarez J. Patients' competence in and performance of cognitive therapy skills: Relation to the reduction of relapse risk following treatment for depression. *Journal of Consulting and Clinical Psychology* 2007;75:523–530. [PubMed: 17663607]
- Surtees PG, Wainwright NWJ, Willis-Owen SAG, Luben R, Day NE, Flint J. Social adversity, the serotonin transporter (5-HTTLPR) polymorphism, and major depressive disorder. *Biological Psychiatry* 2005;59:224–229. [PubMed: 16154545]
- Sutton SK, Davidson RJ. Prefrontal brain asymmetry: A biological substrate of the behavioral approach and inhibition systems. *Psychological Science* 1997;8:204–210.
- Tangney JP, Baumeister RF, Boone AL. High self-control predicts good adjustment, less pathology, better grades, and interpersonal success. *Journal of Personality* 2004;72:271–324. [PubMed: 15016066]
- Taylor SE, Way BM, Welch WT, Hilmert CJ, Lehman BJ, Eisenberger NI. Early family environment, current adversity, the serotonin transporter promoter polymorphism, and depressive symptomatology. *Biological Psychiatry* 2006;60:671–676. [PubMed: 16934775]
- Taylor SF, Phan KL, Decker LR, Liberzon I. Subjective rating of emotionally salient stimuli modulates neural activity. *NeuroImage* 2003;18:650–659. [PubMed: 12667842]
- Tellegen, A. Structure of mood and personality and their relevance to assessing anxiety, with an emphasis on self-report. In: Tuma, AH.; Maser, JD., editors. *Anxiety and the anxiety disorders*. Hillsdale, NJ: Erlbaum; 1985. p. 681–706.
- Toates F. A model of the hierarchy of behaviour, cognition, and consciousness. *Consciousness and Cognition: An International Journal* 2006;15:75–118.
- Tremblay LK, Naranjo CA, Graham SJ, Herrmann N, Mayberg HS, Hevenor S, et al. Functional neuroanatomical substrates of altered reward processing in major depressive disorder revealed by a dopaminergic probe. *Archives of General Psychiatry* 2005;62:1228–1236. [PubMed: 16275810]
- Tse WS, Bond AJ. Serotonergic involvement in the psychosocial dimension of personality. *Journal of Psychopharmacology* 2001;15:195–198. [PubMed: 11565628]
- Tse WS, Bond AJ. Serotonergic intervention affects both social dominance and affiliative behavior. *Psychopharmacology* 2002;161:324–330. [PubMed: 12021836]
- Valiente C, Eisenberg N, Smith CL, Reiser M, Fabes RA, Losoya S, et al. The relations of effortful control and reactive control to children's externalizing problems: A longitudinal assessment. *Journal of Personality* 2003;71:1171–1196. [PubMed: 14633062]

- van Dyck CH, Malison RT, Staley JK, Jacobsen LK, Seibyl JP, Laruelle M, et al. Central serotonin transporter availability measured with [¹²³I]beta-CIT SPECT in relation to serotonin transporter genotype. *American Journal of Psychiatry* 2004;161:525–531. [PubMed: 14992979]
- van Veen V, Carter CS. Conflict and cognitive control in the brain. *Current Directions in Psychological Science* 2006;15:237–246.
- Vasey, MW. Effortful control as a moderator of the relation between anxiety and attentional bias for threat in youth. Ohio State University; 2003. Unpublished raw data
- Verona E, Joiner TE, Johnson F, Bender TW. Gender specific gene-environment interactions on laboratory-assessed aggression. *Biological Psychology* 2006;71:33–41. [PubMed: 16360879]
- Verplanken B, Friborg O, Wang CE, Trafimow D, Woolf K. Mental habits: Metacognitive reflection on negative self-thinking. *Journal of Personality and Social Psychology* 2007;92:526–541. [PubMed: 17352607]
- Vohs, KD.; Lasaleta, JD.; Fennis, B. Self-regulation in the interpersonal sphere. In: Forgas, JP.; Baumeister, RF.; Tice, D., editors. *The psychology of self-regulation*. New York: Psychology Press; (in press).
- Walderhaug E, Lunde H, Nordvik JE, Landrø NI, Refsum H, Magnusson A. Lowering of serotonin by rapid tryptophan depletion increases impulsiveness in normal individuals. *Psychopharmacology* 2002;164:385–391. [PubMed: 12457268]
- Walderhaug E, Magnusson A, Neumeister A, Lappalainen J, Lunde H, Refsum H, et al. Interactive effects of sex and 5-HTTLPR on mood and impulsivity during tryptophan depletion in healthy people. *Biological Psychiatry* 2007;62:593–599. [PubMed: 17544379]
- Wallis JD. Orbitofrontal cortex and its contribution to decision-making. *Annual Review of Neuroscience* 2007;30:31–56.
- Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology* 1988;54:1063–1070. [PubMed: 3397865]
- Way BM, Lácán G, Fairbanks LA, Melega WP. Architectonic distribution of the serotonin transporter within the orbitofrontal cortex of the vervet monkey. *Neuroscience* 2007;148:937–948. [PubMed: 17766046]
- Wentura D. Activation and inhibition of affective information: Evidence for negative priming in the evaluation task. *Cognition and Emotion* 1999;13:65–91.
- Wenzlaff RM, Bates DE. Unmasking a cognitive vulnerability to depression: How lapses in mental control reveal depressive thinking. *Journal of Personality and Social Psychology* 1998;75:1559–1571. [PubMed: 9914666]
- Whalen PJ, Rauch SL, Etcoff NL, McInerney SC, Lee MB, Jenike MA. Masked presentations of emotional facial expressions modulate amygdala activity without explicit knowledge. *Journal of Neuroscience* 1998;18:411–418. [PubMed: 9412517]
- Whitmer AJ, Banich MT. Inhibition versus switching deficits in different forms of rumination. *Psychological Science* 2007;18:546–553. [PubMed: 17576269]
- Wiers, RW.; Stacy, AW. *Handbook of implicit cognition and addiction*. Thousand Oaks, CA: Sage Publishers; 2006.
- Wilhelm K, Mitchell PB, Niven H, Finch A, Wedgwood L, Scimone A, et al. Life events, first depression onset, and the serotonin transporter gene. *British Journal of Psychiatry* 2006;188:210–215. [PubMed: 16507960]
- Wilkowski BM, Robinson MD. The cognitive basis of trait anger and reactive aggression. *Personality and Social Psychology Review* 2008;12:3–21. [PubMed: 18453470]
- Williams JMG, Barnhofer T, Crane C, Watkins E, Hermans D, Raes F, et al. Autobiographical memory specificity and emotional disorder. *Psychological Bulletin* 2007;133:122–148. [PubMed: 17201573]
- Wilson TD, Lindsey S, Schooler TY. A model of dual attitudes. *Psychological Review* 2000;107:101–126. [PubMed: 10687404]
- Winkielman P, Knutson B, Paulus M, Trujillo JL. Affective influence on judgments and decision making: Moving towards core mechanisms. *Review of General Psychology* 2007;11:179–192.

- Wolfe RN, Johnson SD. Personality as a predictor of college performance. *Educational and Psychological Measurement* 1995;55:177–185.
- Wood W, Neal DT. A new look at habits and the habit-goal interface. *Psychological Review* 2007;114:843–863. [PubMed: 17907866]
- Yatham LN, Liddle PF, Shiah IS, Lam RW, Adam MJ, Zis AP, et al. Effects of rapid tryptophan depletion on brain 5-HT(2) receptors: A PET study. *British Journal of Psychiatry* 2001;178:448–453. [PubMed: 11331561]
- Yatham LN, Steiner M. Neuroendocrine probes of serotonergic function: A critical review. *Life Sciences* 1993;53:447–463. [PubMed: 8341131]
- Young SN, Leyton M. The role of serotonin in human mood and social interaction insight from altered tryptophan levels. *Pharmacology, Biochemistry and Behavior* 2002;71:857–865.
- Zald DH, Depue RA. Serotonergic functioning correlates with positive and negative affect in psychiatrically healthy males. *Personality and Individual Differences* 2001;30:71–86.
- Zald DH, Kim SW. Anatomy and function of the orbital frontal cortex I: Anatomy, neurocircuitry, and obsessive-compulsive disorder. *Journal of Neuropsychiatry and Clinical Neurosciences* 1996;8:125–138. [PubMed: 9081547]
- Zalsman G, Huang Y, Oquendo MA, Burke AK, Hu X, Brent DA, et al. Association of a triallelic serotonin transporter gene promoter region (5-HTTLPR) polymorphism with stressful life events and severity of depression. *American Journal of Psychiatry* 2006;163:1588–1593. [PubMed: 16946185]
- Zuckerman, M. *Psychobiology of personality*. 2nd ed. New York: Cambridge University Press; 2005.

Figure 1.

Three temperamental influences on behavior. A: A reactive system for approaching rewards and a reactive system for avoiding threats or punishment compete for ascendancy; in the absence of effortful control, the resultant of that competition is expressed in behavior. B: The engagement of an effortful control system permits the resultant arising from the competition of the reactive systems to be overridden, thus dampening the role of the reactive systems in determining behavior. Adapted from various statements by Rothbart, Eisenberg, and others.

	Action	Restraint
Effortful control dominant	Planful, effortful action	Planful, effortful restraint
Reflexive system dominant	Impulsive approach	Reflexive inhibition or withdrawal

Figure 2.

Either a dominant reflexive system or a dominant effortful control system can ultimately result in either restraint or action.