Published in final edited form as:

Antiviral Res. 2008 November ; 80(2): 150–157. doi:10.1016/j.antiviral.2008.05.012.

INTRAMUSCULARLY ADMINISTERED NEURAMINIDASE INHIBITOR PERAMIVIR IS EFFECTIVE AGAINST LETHAL H5N1 INFLUENZA VIRUS IN MICE

David A. Boltz^a, Natalia A. Ilyushina^a, C. Shane Arnold^b, Y. Sudhakar Babu^b, Robert G. Webster^{a,c}, and Elena A. Govorkova^{a,*}

^a Department of Infectious Diseases, St. Jude Children's Research Hospital, 332 N., Lauderdale St., Memphis, TN 38105–2794, USA

^b BioCryst Pharmaceuticals, Inc., 2190 Parkway Lake Dr, Birmingham, AL 35244, USA

^c Department of Pathology, University of Tennessee Health Science Center, Memphis, Tennessee 38105, USA

Abstract

The replication efficiency and multi-organ dissemination of some influenza A (H5N1) viruses requires a rapid (re)evaluation of the available antiviral strategies. We assessed five regimens of the neuraminidase (NA) inhibitor peramivir in mice inoculated with H5N1 virus. The regimens differed by: (1) frequency of administration on first day (once vs twice); (2) duration of administration (1 day vs 8 days); (3) route of administration (intramuscular [IM] injection alone or followed by oral administration). In all regimens, BALB/c mice were administered 30 mg/kg peramivir IM one hour after lethal challenge with 5 MLD₅₀ of A/Vietnam/1203/04 (H5N1) influenza virus. When given only on the day of inoculation, a single IM injection produced a 33% survival rate, which increased to 55% with two injections. Eight-day regimens significantly increased survival; two IM injections followed by 7 daily IM injections was the most effective regimen (100% survival; inhibition of replication in lungs and brain). When this 8-day regimen began at 24 hours after inoculation, 78% of mice survived; 56% survived when treatment began at 48 after hours. Anti-HA antibody titer differed with the peramivir regimen and corresponded to the severity of disease. Overall, our results demonstrate that IM administration of peramivir is effective in promoting the survival of mice infected with systemically replicating H5N1 virus.

Keywords

Peramivir; Neuraminidase inhibitor; H5N1 influenza virus; antibody; parenteral administration

Introduction

Since 1997, highly pathogenic avian influenza A (H5N1) viruses have posed a threat to human health, causing a fatality rate >60% (WHO, 2008). Although mild and even asymptomatic

*Correspondence should be addressed to: Elena A. Govorkova, Department of Infectious Diseases, St. Jude Children's Research Hospital, 332 N Lauderdale, Memphis, TN 38105–2794; elena.govorkova@stjude.org; Telephone: 901–495–2243; Fax: 901–523–2622.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

infections have been observed, human H5N1 virus infection is commonly associated with severe clinical disease, including pneumonia with progressive respiratory failure, gastrointestinal symptoms, hepatic and renal dysfunction, and encephalitis, (Yuen et al., 1998; de Jong et al., 2005; Ungchusak et al., 2005). Systemic spread and multi-organ failure contribute to the high mortality rates observed. H5N1 virus has been isolated from the cerebrospinal fluid, feces, serum, and plasma of patients (Chutinimitkul et al., 2006; de Jong et al., 2006; Buchy et al., 2007) and from post-mortem tissues. Moreover, detection of viral RNA in intestinal tissues suggests that viral replication can occur in the gastrointestinal tract (Gu et al., 2007; Uiprasertkul et al., 2005).

Neuraminidase (NA) inhibitors which target the conserved residues of the NA active site of both influenza A and B viruses (Colman, 1994), are recommended for the treatment of H5N1 influenza virus infection (Schunemann et al., 2007). NA inhibitors approved for clinical use are orally administered oseltamivir and inhaled zanamivir, both of which are effective against potentially pandemic avian influenza strains under experimental conditions (Leneva et al., 2000; Govorkova et al., 2001; Yen et al., 2005). Information about drug efficacy in humans is limited. Treatment often starts late in the course of H5N1 virus infection, with suboptimal dosages and duration of treatment (Chotpitayasunondh et al., 2004; Beigel et al., 2005). Oseltamivir-resistant H5N1 influenza viruses with mutations at positions H274Y and N294S of the NA have been isolated from patients during (de Jong et al., 2005; Le et al., 2005) and before (Saad et al., 2007) antiviral treatment. Thus far, such isolates remain sensitive to zanamivir (Le et al., 2005; McKimm-Breschkin, 2005; Mishin et al., 2005); however, new H5N1 isolates with mutations not previously reported to be associated with resistance show decreased sensitivity to oseltamivir and zanamivir in vitro (Hurt et al., 2007; McKimm-Breschkin et al., 2007). New approaches for the control of infection with highly pathogenic influenza viruses must be explored; these may include novel antiviral drugs, combinations of antivirals, or optimization of the existing antiviral regimens (dosage, duration and route of administration).

Peramivir (BCX-1812, RWJ-270201), a unique cyclopentane NA inhibitor, is currently under development. Peramivir, generated by using structure-based drug design, has three chemical groups that interact with the active-site residues of the influenza virus NA protein, resulting in tight binding and a slow rate of dissociation (Babu et al., 2000; Bantia et al., 2001). Peramivir potently inhibits the NA activity of various influenza A and B viruses, in vivo and in vitro (Babu et al., 2000; Bantia et al., 2001; Drusano et al., 2001; Sidwell et al., 2001a; Smee et al., 2001). Peramivir has also shown efficacy in cell culture and in mice against influenza viruses with pandemic potential, including H5N1 and H9N2 virus subtypes (Govorkova et al., 2001). Although orally administered peramivir was effective in mice, it did not protect humans against contemporary human influenza A virus in clinical trials; likely resulting from the relatively low blood concentrations observed (Barroso et al., 2005).

Because of the low absorption of oral peramivir, injection of peramivir is being investigated. No injectable anti-influenza drug is currently approved for clinical use. Although intravenously administered zanamivir is well tolerated in humans and effectively prevents experimental influenza A infection (Calfee et al., 1999; Cass et al., 1999; Fritz et al., 1999), only the inhaled formulation of zanamivir is commercially available. Parenterally administered peramivir, now in clinical trials against seasonal influenza A infection has produced high blood levels of the drug in human volunteers and was well tolerated (Kilpatrick et al., 2007). Pre-clinical studies in mice showed that intramuscularly injected peramivir is effective against infection with contemporary influenza A viruses of H1N1 and H3N2 subtypes (Bantia et al., 2006) and with sub-lethal doses of H5N1 subtype (Yun et al., 2008); however, 100% survival was not achieved against H5N1 infection. It is unknown what regimens of parenterally administered peramivir will completely protect against H5N1 virus infection, although relatively rapid production of

a high serum drug concentration could possibly offer an advantage against systemically replicating viruses.

In this study, we evaluated the efficacy of five different regimens of peramivir against infection with the highly pathogenic A/Vietnam/1203/04 (H5N1) influenza virus in a mouse model. Peramivir administered IM for 8 days prevented the deaths of mice inoculated with lethal dose of H5N1 virus and completely eliminated viral spread to internal organs.

2. Materials and Methods

2.1 Neuraminidase inhibitors

Peramivir {[1S,2S,3R,4R,1'S]-3-[1'-acetylamino-2'-ethyl]butyl-4-[(aminoimino)-methyl]amino-2-hydroxycyclopentane-1-carboxylic acid; BCX-1812} and oseltamivir carboxylate, the active metabolite of oseltamivir {[3R,4R,5S]-4-acetamido-5-amino-3-[1-ethylpropoxy]-1-cyclohexane-1-carboxylic acid} were synthesized by BioCryst Pharmaceuticals (Birmingham, AL). Zanamivir (4-guanidino-Neu5Ac2en) was provided by the R. W. Johnson Pharmaceutical Research Institute. Compounds were provided as lyophilized powder and were maintained at 4°C. The compounds were dissolved in distilled water and aliquots were stored at -20°C until used.

2.2 Viruses and cells

The H5N1 influenza viruses A/Vietnam/1203/04 and A/HongKong/213/03 were obtained from the World Health Organization collaborating laboratories. Stock viruses were grown in the allantoic cavities of 10-day-old embryonated chicken eggs for 32 h at 37°C, and aliquots were stored at -70°C until used. Virus titer was determined by calculating the 50% egg infectious dose (EID₅₀) per ml of virus stock (Reed and Muench, 1938). Experiments with highly pathogenic H5N1 influenza viruses were conducted in an animal biosafety level 3+ containment facility approved by the U.S. Department of Agriculture. Madin-Darby canine kidney (MDCK) cells were obtained from the American Type Culture Collection (Manassas, VA) and were grown in minimum essential medium (MEM) supplemented with 5% fetal bovine serum, 5 mM L-glutamine, 0.2% sodium bicarbonate, 100 U/ml penicillin, 100 µg/ml streptomycin sulfate, and 100 µg/ml kanamycin sulfate in a humidified atmosphere of 5% CO₂ at 37°C.

2.3 NA enzyme inhibition assay

We used a modified fluorescence-based NA enzyme inhibition assay (Potier et al., 1979). Briefly, NA inhibitors at concentrations ranging from 0.00005 µM to 10 µM were incubated with H5N1 influenza virus preparations representing a standard amount of NA activity before the addition of the substrate 2'-(4-methylumbelliferyl)-d-N-acetylneuraminic acid (MUNANA, Sigma, St. Louis, MO). The reaction was stopped after 1 h of incubation at 37°C and the fluorescence of the released 4-methylumbelliferone was measured in a Fluoroskan II (Labsystems, Helsinki, Finland) spectrophotometer using excitation and emission wavelengths of 355 and 460 nm, respectively. The IC₅₀ was defined as the concentration of NA inhibitor necessary to reduce NA activity by 50% relative to that in a reaction mixture containing virus but no inhibitor.

2.4 Drug susceptibility in cell culture

The drug susceptibility of A/Vietnam/1203/04 (H5N1) virus was determined by a plaque reduction assay (Hayden et al., 1980). MDCK cells were inoculated with virus diluted in MEM to yield 50 to 80 plaques per well and then were overlaid with infection medium [MEM with 0.3% BSA] containing NA inhibitors at concentrations of 0.0001 µM to 10 µM. Plaque size

was measured after 3 days of incubation at 37°C. At least two independent experiments were performed to determine the concentration of compound required to reduce plaque size by 50% relative to plaque size in untreated wells (EC₅₀).

2.5 Drug efficacy in vivo

Female 6- to 8-week-old BALB/c mice (weight, 18 to 20 g; Jackson Laboratories, Bar Harbor, ME) were lightly anesthetized by inhalation of isoflurane and inoculated intranasally with 5 MLD₅₀ of A/Vietnam/1203/04 (H5N1) influenza virus in 50 µl of PBS. Peramivir was administered starting one hour post-inoculation (p.i.) to groups of 10–15 BALB/c mice according to five different regimens: (1) a single intramuscular (IM) injection of 30 mg/kg [Single IM]; (2) two IM injections, each 30 mg/kg [2x IM], administered during 1 day; (3) a single IM injection of 30 mg/kg on the first day and on each of the next 7 days [Single IM + 7d IM]; (4) a single IM injection of 30 mg/kg on the first day followed by oral administration of 30 mg/kg once daily for 7 days [Single IM + 7d oral]; and (5) two IM injections of 30 mg/kg of peramivir during the first day followed by one IM injection of 30 mg/kg daily for 7 days [2x IM + 7d IM]. The design of experiments is shown in Figure 1. All studies were conducted under applicable laws and guidelines and after approval from the St. Jude Children's Research Hospital Animal Care and Use Committee.

2.6 Efficacy of delayed treatment

To determine the effects of delayed treatment, we initiated the 2x IM + 7d IM regimen 24, 48, or 72 hours after mice were inoculated with 5 MLD₅₀ of A/Vietnam/1203/04 (H5N1) influenza virus. Control (inoculated, untreated) animals received sterile PBS on the same schedule. The groups of 13 BALB/c mice were observed daily for 21 days for clinical signs of infection and for survival. Three mice in each group were killed on day 4 p.i. to determine virus titers in the lungs as described below.

2.7 Virus titers in lungs and brain

On days 3, 6, and 9 after inoculation with A/Vietnam/1203/04 (H5N1) influenza virus, three mice in each experimental and control group were humanely killed. The brains and lungs were removed, washed thoroughly in a large volume of cold sterile PBS, homogenized, and suspended in 1 ml of PBS. The cellular debris was cleared by centrifugation at 2000 × g for 10 min. Supernatant was serially diluted and inoculated into 10-day-old embryonated chicken eggs. The lower limit of virus detection was 0.75 log₁₀ EID₅₀/ml. For calculation of the mean, samples with a virus titer <0.75 log₁₀EID₅₀/ml were assigned a value of 0. Virus titers in each organ were calculated by the method of Reed and Muench (1938) and were expressed as mean log₁₀EID₅₀/ml ± SD.

2.8 Emergence of drug-resistant variants

The RNeasy Kit (Qiagen, Chatsworth, CA) was used to extract viral RNA from the lungs and brains of mice on days 6 and 9 p.i., and the One Step RT-PCR kit (Qiagen, Chatsworth, CA) was used according to the protocol provided. Universal primers were used for amplification of the NA and HA (HA1 region) genes (Hoffmann et al., 2001). The sequences were determined by the Hartwell Center for Bioinformatics and Biotechnology at St. Jude Children's Research Hospital by using BigDye Terminator (v. 3) chemistry and synthetic oligonucleotides. Samples were analyzed on Applied Biosystems 3700 DNA analyzers.

2.9 Anti-HA antibody response

Serum samples were collected from mice 21 days p.i., treated with receptor-destroying enzyme, heat-inactivated at 56°C for 30 min, and tested by hemagglutination inhibition (HI) assay with 0.5% packed chicken red blood cells (CRBC).

2.10 Statistical analysis

Mean virus titers in mouse organs were compared by unpaired two-tailed t-test. The Kaplan-Meier method was used to estimate the probability of survival and the log-rank test to compare survival estimates of the placebo and treatment groups (Venables and Ripley, 1997). The proportional hazards model was used to determine the death hazard ratio of the treatment and placebo groups (Cox, 1972).

3. Results

3.1 Susceptibility of H5N1 virus to NA inhibitors in vitro

To compare the susceptibility of A/Vietnam/1203/04 (H5N1) influenza virus to three different NA inhibitors in vitro, we performed NA inhibition and plaque reduction assays in MDCK cells. Overall, the mean IC_{50} and EC_{50} values obtained with peramivir (0.6 ± 0.2 nM and 0.3 ± 0.1 nM, respectively) were comparable to those for zanamivir (0.9 ± 0.2 nM and 0.7 ± 0.1 nM) and oseltamivir carboxylate (0.3 ± 0.1 nM and 0.5 ± 0.1 nM), demonstrating the high susceptibility of this H5N1 influenza virus to all three NA inhibitors in vitro (data not shown).

3.2. Effect of peramivir on survival and disease signs after challenge with lethal H5N1 virus

We evaluated the effect of five different regimens of peramivir on the lethality and clinical signs of A/Vietnam/1203/04 (H5N1) virus infection in mice (Figure 1). Untreated inoculated control mice exhibited progressive weight loss with a mean day of death of 9.2. The survival rate of treated mice varied with the regimens. A single IM injection prevented death in 33% of animals, and two IM injections (2x IM) prevented death in 55% (Table 1). Minimal weight loss was observed on day 6 p.i. in mice receiving peramivir for one day; however, weight loss was maximal on day 9 p.i. Prolonging peramivir therapy from a 1-day to an 8-day regimen significantly lowered the risk of death: the single IM + 7d oral and single IM + 7d IM regimens prevented death in 66% and 88% of animals, respectively ($P < 0.001$). The 2x IM + 7d IM regimen had the greatest efficacy: no weight loss and 100% survival (Table 1).

Despite differences in survival among the peramivir regimens (Figure 2), drug administration significantly delayed death in all treatment groups ($P < 0.01$). The single IM and the 2x IM treatment regimens resulted in a mean day of death of 12.7 and 13.1 respectively; however, in the single IM group mortality occurred one day earlier than in the 2x IM group, with the highest incidence of death on day 14 p.i. (Figure 2A). Administration of peramivir for eight days increased not only the likelihood of survival but also the duration of survival (Figure 2B).

Fifty percent of control animals developed severe neurological signs (hind limb paresis and ataxia) before death. A single IM injection of peramivir did not prevent neurological complications in 5 of 9 mice (Table 1). However, the frequency of neurological signs was decreased by the 2x IM, single IM + 7d oral, and single IM + 7d IM regimens and neurological signs were inhibited by the 2x IM + 7d IM regimen.

3.3 Reduction of lung and brain virus titers by peramivir treatment

We compared the efficacy with which different peramivir regimens inhibited the replication of neurotropic A/Vietnam/1203/04 (H5N1) virus in mice. Neither single-day peramivir regimen inhibited pulmonary virus replication in inoculated mice; viral titers were comparable to those in the control group on days 3, 6, and 9 p.i. (Figure 3A). All of the 8-day regimens significantly reduced ($P < 0.05$) virus replication in mouse lungs on days 3 and 6 p.i. (Figure 3A). Day-9 pulmonary virus titers did not differ significantly between controls and the single IM + 7d oral treatment group but were significantly reduced ($P < 0.05$) in the single IM + 7d IM and 2x IM + 7d IM treatment groups. Notably, the 2x IM + 7d IM regimen was the only

drug regimen that completely inhibited virus replication in the lungs on all days studied (limit of virus detection, $0.75 \log_{10} \text{EID}_{50}/\text{ml}$).

In untreated inoculated mice, H5N1 virus was detectable in the brain as early as day 3 p.i.; the brain virus titer was slightly increased on day 6 p.i. and was dramatically increased (as high as $5.5 \log_{10} \text{EID}_{50}/\text{ml}$) on day 9 p.i. Virus was undetectable in the brains of all treated mice on days 3 and 6 p.i. Both of the 1-day regimens and the single IM + 7d oral regimen significantly reduced ($P < 0.05$) brain virus titers on days 3, 6 and 9 p.i. (Figure 3B). Importantly, both of the 8-day IM regimens completely inhibited virus replication in the brains of mice (virus was not detectable on day 3, 6, or 9 p.i.). Virus replication in both the lung and brain was inhibited only in the 2x IM + 7d IM treatment group, indicating that this regimen is best for controlling systemic A/Vietnam/1203/04 (H5N1) virus spread.

3.4 Emergence of peramivir-resistant variants during treatment

To monitor the emergence of resistant variants during peramivir treatment, we isolated RNA from virus detected in the lungs and brains of mice days 6 and 9 p.i. Sequence analysis of the dominant virus population identified no differences between the amino acids encoded by the NA and HA (HA1 region) genes of the wild-type A/Vietnam/1203/04 (H5N1) virus (challenge virus) and virus isolated from the organs of treated mice (data not shown).

3.5 Antibody production after peramivir treatment

To serologically confirm H5N1 virus infection and to compare the effect of the drug regimens on production of anti-HA antibodies, we collected serum 21 days p.i. for HI assay. Because A/Vietnam/1203/04 (H5N1) virus tends to induce low anti-HA titers, we included A/Hong Kong/213/03 (H5N1) virus, whose HA antigen is better suited to serologic antibody detection, in the assay. Anti-HA antibody titers to both H5N1 viruses were observed in surviving mice given the single IM, 2x IM, and single IM + 7d oral regimens (Figure 4). In the single IM + 7d IM treatment group, only anti-HA antibodies against A/Hong Kong/213/03 (H5N1) were detected, confirming previous observations that this heterologous virus better detects anti-HA antibodies (Hoffmann et al., 2005). Mice that underwent the 2x IM + 7d IM treatment had decreased or undetectable anti-HA serum antibodies. Overall, our data showed that the single IM and the single IM + 7d oral peramivir regimens did not protect animals from H5N1 infection, although they reduced the severity of disease. Multiple IM injections of peramivir were required to completely inhibit the spread of H5N1 viral infection.

3.6. Effect of delayed treatment with peramivir

To assess the therapeutic potential of peramivir we examined the efficacy of the drug when given during the course of established infection. Mice were inoculated with 5 MLD_{50} of A/Vietnam/1203/04 (H5N1) virus and treatment with the 2x IM + 7d IM regimen was initiated 24, 48, or 72 hours p.i. All untreated inoculated mice died between days 8 and 10 p.i. Treatment started either 24 or 48 hours p.i. significantly increased survival (78% and 56% of animals survived, respectively) and decreased weight loss ($P < 0.01$) (Table 2). However, when treatment began 72 hours p.i., mice lost weight progressively and died between days 8 and 12 p.i. Notably, initiation of treatment 24 and 48 hours p.i. also reduced neurological signs. Although initiation of peramivir treatment 24 hours p.i. significantly inhibited virus replication in the lungs ($P < 0.05$), lung titers did not differ significantly from those in controls when treatment began 48 or 72 hours p.i. (Table 2).

4. Discussion

There is limited information about the efficacy of parenterally administered anti-influenza drugs against influenza A (H5N1) viruses. This study evaluated different regimens of

intramuscularly administered peramivir in mice lethally challenged with the A/Vietnam/1203/04 (H5N1) influenza virus. We found that a single IM injection of peramivir protected 33% of animals from death, and IM administration for 8 days was required to achieve 100% survival and inhibit viral replication in the lungs and brains. Delayed IM administration also promoted survival if peramivir was given within 48 hours of virus inoculation.

In the current study, A/Vietnam/1203/04 (H5N1) influenza virus was highly susceptible to peramivir in both enzymatic and cell-culture–based assays. Because virus susceptibility in vitro does not always predict results in vivo (Sidwell et al., 1998, 2001a), studies in an animal model are warranted. In the mouse animal model, we observed that the frequency and duration of peramivir administration are important factors in the adequate control of infection caused by the highly pathogenic H5N1 viruses. We found that a single IM dose of peramivir was inadequate to completely protect mice from mortality caused by the highly pathogenic A/Vietnam/1203/04 (H5N1) virus. A survival rate of 33% was achieved with a single IM dose which increased to 55% with twice daily IM injections. Against seasonal influenza strains, a single IM injection of peramivir significantly reduced weight loss and mortality in mice inoculated with mouse-adapted A/Victoria/3/75 (H3N2) virus and completely protected against lethal challenge with A/NWS/33 (H1N1) virus (Bantia et al., 2006). The difference in effectiveness of 1-day IM treatment with peramivir between seasonal influenza and the H5N1 influenza virus may be explained by the greater virulence of H5N1 viruses than of H1N1 and H3N2 viruses which affects the efficacy of NA inhibitors in experimental animal models (Sidwell and Smeeth, 2000; Yen et al., 2005; Ilyushina et al., 2007).

The survival rate of mice improved as the duration of peramivir administration increased from 1 day to 8 days. We observed that the IM route of peramivir administration offered greater benefit than the oral route, as the single IM + 7d oral regimen was less effective than the single IM + 7d IM and 2x IM + 7d IM regimens (lower survival rates, greater weight loss, and higher lung and brain virus titers). Recent studies also showed that IM injections of peramivir promoted survival in mice and ferrets infected with A/Vietnam/1203/04 (H5N1) virus (Yun et al., 2008), although low virus doses were used (9%–36% survival of control untreated animals) for the evaluation of antiviral efficacy. The comparison of bioavailability and tissue distribution of oral and IM peramivir has not been reported, but low bioavailability of orally administered peramivir may account for its lesser efficacy against a virus that spreads systemically. Although we did not directly compare the efficacy of the different NA inhibitors against lethal H5N1 virus infection in mice, we found that the single IM + 7d oral peramivir regimen was as effective as oseltamivir in preventing death from A/Vietnam/1203/04 (H5N1) virus infection. An 8-day regimen of orally administered oseltamivir 10 mg/kg/day was needed to provide an 80% survival rate in mice inoculated with 5 MLD₅₀ of A/Vietnam/1203/04 (H5N1) virus (Yen et al., 2005).

The prevention of death by intramuscularly injected peramivir corresponded to the inhibition of viral replication in the lungs and inhibition of virus spread to the brain. In animal models of lethal influenza virus infection, H5N1 virus has been isolated from multiple organs, including lungs, brain, liver, spleen, and kidney (Maines et al., 2005; Yen et al., 2005; Govorkova et al., 2007). In our study, mice receiving 1-day regimens survived while viral replication in the lungs was inhibited; however, the subsequent increase of lung titers to equal those in control untreated mice was associated with death. The survival rate was 100% only when viral replication in the lungs was completely inhibited by the 2x IM + 7d IM peramivir regimen. We suggest that the spread of A/Vietnam/1203/04 (H5N1) virus to the brain contributed to the deaths of infected treated animals late in the course of infection. Neurological signs observed in 50% of untreated inoculated mice were associated with isolation of virus from the brain early during infection. In all peramivir regimens except for the single IM regimen, the frequency of neurological

symptoms was significantly reduced, the survival rate was increased, and virus spread to the brain was decreased or prevented.

The emergence of resistant mutants during antiviral treatment is a major concern. NA inhibitor-resistant H5N1 viruses carrying the H274Y or N294S mutation have been identified in infected patients after oseltamivir treatment (de Jong et al., 2005; Le et al., 2005). The development of resistance to peramivir in vivo has not been described, although influenza A and B variants resistant to peramivir have emerged in vitro under drug selection pressure (Smee et al., 2001; Baz et al., 2007; Baum et al., 2003). We found no mutations in the NA or HA genes of viruses isolated from mouse lungs and brains that would reduce susceptibility to NA inhibitors. However, these studies were designed primarily to evaluate the efficacy of parentally administered peramivir in mice inoculated with H5N1 virus; issues of receptor specificity in mouse lungs and analysis of the dominant virus population may limit any conclusions we might draw about the emergence of resistance.

In previous studies, orally administered peramivir or oseltamivir did not affect cellular immune response or prevent the development of humoral immunity in mice during H1N1 influenza virus infection (Burger et al., 2000; Sidwell et al., 2001b). IM administration of peramivir did not inhibit the development of serum anti-HA antibodies to H5N1 virus in mice. However, the levels of anti-HA antibodies differed with the peramivir regimens and corresponded to the severity of disease. High titers of anti-HA antibodies were associated with peramivir regimens that did not prevent severe infection as indicated by weight loss and high pulmonary viral titers, whereas antibody titers were undetectable and pulmonary virus replication was inhibited in mice receiving peramivir IM for eight days. Although no peramivir regimen tested completely prevented H5N1 virus infection, all of the regimens controlled virus replication and virus spread from the primary site of infection. Sufficient viral replication in the lungs and a high viral load is required to trigger an adequate immune response.

We also studied the efficacy of delayed peramivir treatment in a mouse model of lethal H5N1 virus infection. We observed 78% and 56% survival rates when peramivir was administered IM 24 or 48 hours, respectively, after inoculation. Govorkova et al (2001) reported similar survival rates when peramivir was orally administered during H5N1 infection. A higher dose of peramivir (30 mg/kg vs 10 mg/kg) was required to achieve similar efficacy against A/Vietnam/1203/04 (H5N1) virus than against A/Hong Kong/156/97 virus (H5N1) when treatment was delayed (Govorkova et al., 2001). Viral titers in the lung were significantly lower and the survival rate was higher when peramivir treatment was delayed for 24 hours vs. 48 or 72 hours. Pulmonary virus titers in mice treated 48 and 72 hours after inoculation did not differ from those in controls, although treatment continued for only 48 and 24 hours, respectively, before organ collection. We can propose that earlier initiation of treatment will be the most beneficial regimen for IM peramivir, although pathogenicity of the virus can also affect the outcome of treatment and better results can be achieved with less pathogenic H5N1 influenza virus.

In summary, these studies show that IM administered peramivir is effective against systemically replicating H5N1 virus, promoting survival and inhibiting viral replication in lungs and brain. The efficacy of IM peramivir against highly pathogenic H5N1 virus infection is dependent on the timing of treatment initiation, the duration of treatment and the route of drug administration. The IM administration of peramivir enhanced its antiviral effect, suggesting that this route might be successful for the treatment of human H5N1 virus infections.

Acknowledgments

We thank the WHO Global Influenza Surveillance Network for providing the H5N1 viruses; Jennifer L. McClaren, David Carey and Cedric Proctor for excellent technical assistance; and Sharon Naron for editorial assistance. This

study was supported by the National Institute of Allergy and Infectious Diseases (Contract No. HHSN266200700005C), by BioCryst Pharmaceuticals, Inc., USA, and by the American Lebanese Syrian Associated Charities (ALSAC).

References

- Babu YS, Chand P, Bantia S, Kotian PL, Dehghani A, El-Kattan Y, Lin TH, Hutchison TL, Elliot AJ, Parker CD, Ananth SL, Horn LL, Laver GW, Montgomery JA. BCX-1812 (RWJ-270201): discovery of a novel, highly potent, orally active and selective influenza neuraminidase inhibitor through structure-based drug design. *J Med Chem* 2000;43:3482–3486. [PubMed: 11000002]
- Bantia S, Parker CD, Ananth SL, Horn LL, Andries K, Chand P, Kotian PL, Dehghani A, El-Kattan Y, Lin T, Hutchison TL, Montgomery JA, Kellog DL, Babu YS. Comparison of the anti-influenza virus activity of RWJ-270201 with those of oseltamivir and zanamivir. *Antimicrob Agents Chemother* 2001;45:1162–1167. [PubMed: 11257030]
- Bantia S, Arnold CS, Parker CD, Upshaw R, Chand P. Anti-influenza virus activity of peramivir in mice with single intramuscular injection. *Antiviral Res* 2006;69:39–45. [PubMed: 16325932]
- Barroso L, Treanor J, Gubareva L, Hayden FG. Efficacy and tolerability of the oral neuraminidase inhibitor peramivir in experimental human influenza: randomized, controlled trials for prophylaxis and treatment. *Antivir Ther* 2005;10:901–910. [PubMed: 16430195]
- Baum EZ, Wagaman PC, Ly I, Turchi I, Le J, Bucher D, Bush K. A point mutation in influenza B neuraminidase confers resistance to peramivir and loss of slow binding. *Antiviral Res* 2003;59:13–22. [PubMed: 12834856]
- Baz M, Abed Y, Boivin G. Characterization of drug-resistant recombinant influenza A/H1N1 viruses selected in vitro with peramivir and zanamivir. *Antiviral Res* 2007;74:159–62. [PubMed: 17137644]
- Beigel JH, Farrar J, Han AM, Hayden FG, Hyer R, de Jong MD, Lochindarat S, Nguyen TK, Nguyen TH, Tran TH, Nicoll A, Touch S, Yuen KY. Writing Committee of the World Health Organization (WHO) Consultation on Human Influenza A/H5. Avian influenza A (H5N1) infection in humans. *N Engl J Med* 2005;353:1374–1385. [PubMed: 16192482]
- Buchy P, Mardy S, Vong S, Toyoda T, Aubin JT, Miller M, Touch S, Sovann L, Dufourcq JB, Richner B, Tu PV, Tien NT, Lim W, Peiris JS, Van der Werf S. Influenza A/H5N1 virus infection in humans in Cambodia. *J Clin Virol* 2007;39:164–168. [PubMed: 17526430]
- Burger RA, Billingsley JL, Huffman JH, Bailey KW, Kim CU, Sidwell RW. Immunological effects of the orally administered neuraminidase inhibitor oseltamivir in influenza virus-infected and uninfected mice. *Immunopharm* 2000;47:45–52.
- Calfee DP, Peng AW, Hussey EK, Lobo M, Hayden FG. Safety and efficacy of once daily intranasal zanamivir in preventing experimental human influenza A infection. *Antivir Ther* 1999;4:143–149. [PubMed: 12731753]
- Cass LM, Efthymiopoulos C, Bye A. Pharmacokinetics of zanamivir after intravenous, oral, inhaled or intranasal administration to healthy volunteers. *Clin Pharmacokinet Suppl* 1999;36:1–11.
- Chotpitayasunondh T, Ungchusak K, Hanshaoworakul W, Chunsuthiwat S, Sawanpanyalert P, Kijphati R, Lochindarat S, Srisan P, Suwan P, Osothanakorn Y, Anantasetagoon T, Kanjanawasri S, Tanupattarachai S, Weerakul J, Chaiwirattana R, Maneerattanaporn M, Poolsavathitkool R, Choekphaibulkit K, Apisarnthanarak A, Dowell SF. Human disease from influenza A (H5N1), Thailand, 2004. *Emerg Infect Dis* 2005;11:201–209. [PubMed: 15752436]
- Chutinimitkul S, Bhattarakosol P, Srisuratanon S, Eiamudomkan A, Kongsomboon K, Damrongwatanapokin S, Chaisingh A, Suwannakarn K, Chieochansin T, Theamboonlers A, Poovorawan Y. H5N1 influenza A virus and infected human plasma. *Emerg Infect Dis* 2006;12:1041–1043. [PubMed: 16752481]
- Colman PM. Influenza virus neuraminidase: structure, antibodies, and inhibitors. *Protein Sci* 1994;3:1687–1696. [PubMed: 7849585]
- Cox DR. Regression models and life-tables. *J R Stat Soc B* 1972;34:187–220.
- de Jong MD, Bach VC, Phan TQ, Vo MH, Tran TT, Nguyen BH, Beld M, Le TP, Truong HK, Nguyen VV, Tran TH, Do QH, Farrar J. Fatal avian influenza A (H5N1) in a child presenting with diarrhea followed by coma. *N Engl J Med* 2005;352:686–691. [PubMed: 15716562]

- de Jong MD, Simmons CP, Thanh TT, Hien VM, Smith GJ, Chau TN, Hoang DM, Chau NV, Khanh TH, Dong VC, Qui PT, Cam BV, Ha do Q, Guan Y, Peiris JS, Chinh NT, Hien TT, Farrar J. Fatal outcome of human influenza A (H5N1) is associated with high viral load and hypercytokinemia. *Nat Med* 2006;12:1203–1207. [PubMed: 16964257]
- Drusano GL, Preston SL, Smee D, Bush K, Bailey K, Sidwell RW. Pharmacodynamic evaluation of RWJ-270201, a novel neuraminidase inhibitor, in a lethal murine model of influenza predicts efficacy for once-daily dosing. *Antimicrob Agents Chemother* 2001;45:2115–2118. [PubMed: 11408232]
- Fritz RS, Hayden FG, Calfee DP, Cass LM, Peng AW, Alvord WG, Strober W, Straus SE. Nasal cytokine and chemokine responses in experimental influenza A virus infection: results of a placebo-controlled trial of intravenous zanamivir treatment. *J Infect Dis* 1999;180:586–593. [PubMed: 10438343]
- Govorkova EA, Leneva IA, Goloubeva OG, Bush K, Webster RG. Comparison of efficacies of RWJ-270201, zanamivir, and oseltamivir against H5N1, H9N2, and other avian influenza viruses. *Antimicrob Agents Chemother* 2001;45:2723–2732. [PubMed: 11557461]
- Govorkova EA, Ilyushina NA, Boltz DA, Douglas A, Yilmaz N, Webster RG. Efficacy of oseltamivir therapy in ferrets inoculated with different clades of H5N1 influenza virus. *Antimicrob Agents Chemother* 2007;51:1414–1424. [PubMed: 17296744]
- Gu J, Xie Z, Gao Z, Liu J, Korteweg C, Ye J, Lau LT, Lu J, Gao Z, Zhang B, McNutt MA, Lu M, Anderson VM, Gong E, Yu AC, Lipkin WI. H5N1 infection of the respiratory tract and beyond: a molecular pathology study. *Lancet* 2007;370:1137–1145. [PubMed: 17905166]
- Hayden FG, Cote KM, Douglas RG. Plaque inhibition assay for drug susceptibility testing of influenza viruses. *Antimicrob Agents Chemother* 1980;17:865–870. [PubMed: 7396473]
- Hoffmann E, Stech J, Guan Y, Webster RG, Perez DR. Universal primer set for the full-length amplification of all influenza A viruses. *Arch Virol* 2001;146:2275–2289. [PubMed: 11811679]
- Hoffmann E, Lipatov AS, Webby RJ, Govorkova EA, Webster RG. Role of specific hemagglutinin amino acids in the immunogenicity and protection of H5N1 influenza virus vaccines. *Proc Natl Acad Sci USA* 2005;102:12915–12920. [PubMed: 16118277]
- Hurt AC, Selleck P, Komadina N, Shaw R, Brown L, Barr IG. Susceptibility of highly pathogenic A (H5N1) avian influenza viruses to the neuraminidase inhibitors and adamantanes. *Antiviral Res* 2007;73:228–231. [PubMed: 17112602]
- Ilyushina NA, Hoffmann E, Salomon R, Webster RG, Govorkova EA. Amantadine-oseltamivir combination therapy for H5N1 influenza virus infection in mice. *Antivir Ther* 2007;12:363–370. [PubMed: 17591026]
- Kilpatrick, JM.; Harman, LA.; Collis, PJ.; Aitee, G.; Mead, E.; Alexander, WJ. Pharmacokinetics and safety of peramivir by intramuscular administration. *Options for the Control of Influenza VI*; Toronto, Canada: 2007. p. 231
- Le QM, Kiso M, Someya K, Sakai YT, Nguyen TH, Nguyen KH, Pham ND, Ngyen HH, Yamada S, Muramoto Y, Horimoto T, Takada A, Goto H, Suzuki T, Suzuki Y, Kawaoka Y. Avian flu: isolation of drug-resistant H5N1 virus. *Nature* 2005;437:1108. [PubMed: 16228009]
- Leneva IA, Roberts N, Govorkova EA, Goloubeva OG, Webster RG. The neuraminidase inhibitor GS4104 (oseltamivir phosphate) is efficacious against A/Hong Kong/156/97 (H5N1) and A/Hong Kong/1074/99 (H9N2) influenza viruses. *Antiviral Res* 2000;48:101–115. [PubMed: 11114412]
- Maines TR, Lu XH, Erb SM, Edwards L, Guarner J, Greer PW, Nguyen DC, Szretter KJ, Chen LM, Thawatsupha P, Chittaganpitch M, Waicharoen S, Nguyen DT, Nguyen T, Nguyen HH, Kim JH, Hoang LT, Kang C, Phuong LS, Lim W, Zaki S, Donis RO, Cox NJ, Katz JM, Tumpey TM. Avian influenza (H5N1) viruses isolated from humans in Asia in 2004 exhibit increased virulence in mammals. *J Virol* 2005;79:11788–11800. [PubMed: 16140756]
- McKimm-Breschkin JL. Management of influenza virus infections with neuraminidase inhibitors: detection, incidence, and implications of drug resistance. *Treat Respir Med* 2005;4:107–116. [PubMed: 15813662]
- McKimm-Breschkin JL, Selleck PW, Usman TB, Johnson MA. Reduced sensitivity of influenza A (H5N1) to oseltamivir. *Emerg Infect Dis* 2007;13:1354–1357. [PubMed: 18252107]
- Mishin VP, Hayden FG, Gubareva LV. Susceptibilities of antiviral-resistant influenza viruses to novel neuraminidase inhibitors. *Antimicrob Agents Chemother* 2005;49:4515–4520. [PubMed: 16251290]

- Potier M, Mameli L, Blisle M, Dallaire L, Melanon SB. Fluorometric assay of neuraminidase with a sodium (4-methylumbelliferyl--D-N-acetylneuraminate) substrate. *Anal Biochem* 1979;94:287–296. [PubMed: 464297]
- Reed LJ, Muench H. A simple method for estimating fifty percent endpoints. *Am J Hyg* 1938;27:493–497.
- Saad, MD.; Boynton, BR.; Earhart, KC.; Mansour, MM.; Niman, HL.; Elsayed, NM.; Nayel, AL.; Abdelghani, AS.; Essmat, HM.; Labib, EM.; Ayoub, EA.; Monteville, MR. Options for the Control of Influenza VI. Toronto, Canada: 2007. Detection of oseltamivir resistance mutation N294S in humans with influenza A H5N1; p. 228
- Schunemann HJ, Hill SR, Kakad M, Bellamy R, Uyeki TM, Hayden FG, Yazdanpanah Y, Beigel J, Chotpitayasunondh T, Del Mar C, Farrar J, Tran TH, Ozbay B, Sugaya N, Fukuda K, Shindo N, Stockman L, Vist GE, Croisier A, Nagjdaliyev A, Roth C, Thomson G, Zucker H, Oxman AD. WHO Rapid Advice Guideline Panel on Avian Influenza. WHO Rapid Advice Guidelines for pharmacological management of sporadic human infection with avian influenza A (H5N1) virus. *Lancet Infect Dis* 2007;7:21–31. [PubMed: 17182341]
- Sidwell RW, Huffman JH, Barnard DL, Bailey KW, Wong MH, Morrison A, Syndergaard T, Kim CU. Inhibition of influenza virus infections in mice by GS4104, an orally effective influenza virus neuraminidase inhibitor. *Antiviral Res* 1998;37:107–120. [PubMed: 9588843]
- Sidwell RW, Smee DF. In vitro and in vivo assay systems for study of influenza virus inhibitors. *Antiviral Res* 2000;48:1–16. [PubMed: 11080536]
- Sidwell RW, Smee DF, Huffman JH, Barnard DL, Bailey KW, Morrey JD, Babu YS. In vivo influenza virus-inhibitory effects of the cyclopentane neuraminidase inhibitor RWJ-270201. *Antimicrob Agents Chemother* 2001a;45:749–757. [PubMed: 11181355]
- Sidwell RW, Smee DF, Bailey KW, Burger RA. Primary immune system effects of the orally administered cyclopentane neuraminidase inhibitor RWJ-270201 in influenza virus-infected mice. *Int Immunopharmacol* 2001b;1:1211–1218. [PubMed: 11407315]
- Smee DF, Huffman JH, Morrison AC, Barnard DL, Sidwell RW. Cyclopentane neuraminidase inhibitors with potent in vitro anti-influenza virus activities. *Antimicrob Agents Chemother* 2001;45:743–748. [PubMed: 11181354]
- Uiprasertkul M, Puthavathana P, Sangsiriwut K, Pooruk P, Srisook K, Peiris M, Nicholls JM, Choekphaibulkit K, Vanprapar N, Auewarakul P. Influenza A H5N1 replication sites in humans. *Emerg Infect Dis* 2005;11:1036–1041. [PubMed: 16022777]
- Ungchusak K, Auewarakul P, Dowell SF, Kitphati R, Auwanit W, Puthavathana P, Uiprasertkul M, Boonnak K, Pittayawonganon C, Cox NJ, Zaki SR, Thawatsupha P, Chittaganpitch M, Khontong R, Simmerman JM, Chunsutthiwat S. Probable person-to-person transmission of avian influenza A (H5N1). *N Engl J Med* 2005;352:333–340. [PubMed: 15668219]
- Venables, WN.; Ripley, BD. Modern applied statistics. Springer; New York, NY: 1997. p. 223–242.p. 297–321.p. 345–350.
- World Health Organization. H5N1 avian influenza: timeline of major events. World Health Organization. ; 2008.
http://www.who.int/csr/disease/avian_influenza/country/cases_table_2008_02_15/en/index.html
- Yen HL, Monto AS, Webster RG, Govorkova EA. Virulence may determine the necessary duration and dosage of oseltamivir treatment for highly pathogenic A/Vietnam/1203/04 influenza virus in mice. *J Infect Dis* 2005;192:665–672. [PubMed: 16028136]
- Yuen KY, Chan PK, Peiris M, Tsang DN, Que TL, Shortridge KF, Cheung PT, To WK, Ho ET, Sung R, Cheng AF. Clinical features and rapid viral diagnosis of human disease associated with avian influenza A H5N1 virus. *Lancet* 1998;351:467–471. [PubMed: 9482437]
- Yun NE, Linde NS, Zacks MA, Barr IG, Hurt AC, Smith JN, Dziuba N, Holbrook MR, Zhang L, Kilpatrick JM, Arnold CS, Paessler S. Injectable peramivir mitigates disease and promotes survival in ferrets and mice infected with the highly virulent influenza virus, A/Vietnam/1203/04 (H5N1). *Virology* 2008;374:198–209. [PubMed: 18234269]

Figure 1.
Experimental design. BALB/c mice were administered peramivir by intramuscular injection (arrows) or by oral gavage (circles).

Figure 2.

Effect of 1- and 8-day peramivir regimens on mouse survival. BALB/c mice were administered peramivir for 1 day (A) or for 8 days (B). The Kaplan-Meier method was used to estimate the probability of survival, which expressed as survival distribution function. 1- probability of 100% survival.

Figure 3.

Effect of peramivir regimens on virus titers in internal organs. Virus titers were determined at 3, 6, and 9 days p.i. (dpi) in the lungs (A) and brain (B). Each data point represents the mean virus titer $\pm$ SD ($\log_{10}$ EID₅₀/ml) from the lungs or brains of 3 mice. Dotted line indicates minimum level of detection ($0.75 \log_{10}$ EID₅₀/ml). * indicate $P < 0.05$.

Figure 4.

Serum antibody responses in mice inoculated with A/Vietnam/1203/04 (H5N1) virus and treated with peramivir. Hemagglutination inhibition (HI) titers were tested by using A/Vietnam/1203/04 (H5N1) and A/Hong Kong/213/03 (H5N1) viruses. Drug regimens are shown as mg/kg (see Figures 1 and 2). Sera were obtained 21 days p.i. Values are mean HI titers (log₂) ± SE.

Table 1
Effect of peramivir regimens in mice inoculated with A/Vietnam/1203/04 (H5N1) influenza virus

Peramivir regimen ^a	Number of survivors/Total (%)	Days to death, mean ± SE ^c	Hazard ratio(P value) ^d	Mean weight change (% ± SE) ^e			Neurological signs ^f
				6 days p.i.	9 days p.i.	15 days p.i.	
1-day administration:							
single IM	3/9 (33) **	12.7 ± 0.7 *	0.08 **	-3.1 ± 1.5	-14.5 ± 2.9	-4.1 ± 1.8	5/9 (13)
2x IM ^b	5/9 (55) **	13.1 ± 0.6 *	0.05 **	-0.7 ± 1.1	-3.6 ± 2.0 **	-2.3 ± 2.8	2/9 (13.5)
8-day administration:							
single IM + 7d oral	6/9 (66) **	11.7 ± 0.4 *	0.04 **	1.8 ± 0.7 *	-1.0 ± 3.0 **	-3.0 ± 6.8	2/9 (11)
single IM + 7d IM	8/9 (88) **	17.0 ± 0.0 *	0.01 **	3.8 ± 0.5 *	3.8 ± 0.6 **	1.2 ± 3.2	2/9 (15)
2x IM + 7d IM ^b	10/10 (100) **	> 21 *	0.00 **	1.6 ± 1.2 *	2.7 ± 1.5 **	5.8 ± 3.8	0/10 (NA)
Control	0/10 (0)	9.2 ± 0.3	1	-6.2 ± 2.8	-19.1 ± 1.0	NA	5/10 (9)

^aPeramivir (30 mg/kg) or PBS was administered by IM injection or oral gavage, starting 1 h after inoculation of 6-week-old BALB/c mice with 5 MLD₅₀ of A/Vietnam/1203/04 (H5N1) virus. Survival was observed for 21 days.

^bTwo IM injections given 12 hours apart, starting 1 h after inoculation.

^cEstimated by the log-rank test (Venables et al., 1997).

^dDeath hazard ratio (vs. placebo group) was estimated by the proportional hazards model (Cox, 1972)

^eLoss or gain of weight was calculated for each mouse as a percentage of weight on day 0 before virus inoculation. NA – not applicable: all mice died, and therefore groups cannot be statistically compared for weight change.

^fAxaxia or hind limb paresis. Shown are the number of mice with neurological signs/total number. The median day p.i. when the signs were observed is shown in parentheses.

* $P < 0.01$.

** $P < 0.001$ compared to placebo-treated control group.

Table 2
Effect of delayed treatment with peramivir in mice inoculated with A/Vietnam/1203/04 (H5N1) influenza virus

Initiation of treatment (hours p.i.) ^a	Number of survivors/total (%)	Days to death, mean $\pm$ SE ^b	Hazard ratio (P value) ^c	Mean weight change (% $\pm$ SE) ^d			Neurological signs ^e	Viral titer in lungs ^f
				6 days p.i.	9 days p.i.	15 days p.i.		
24	7/9 (78)**	11.9 $\pm$ 0.2*	0.05*	0.4 $\pm$ 0.7*	-2.3 $\pm$ 2.1**	1.0 $\pm$ 1.1	0/9 (NA)	2.5 $\pm$ 0.2*
48	5/9 (56)**	11.2 $\pm$ 0.4*	0.12*	-0.5 $\pm$ 0.6*	-3.4 $\pm$ 2.4**	-1.4 $\pm$ 3.3	0/9 (NA)	5.4 $\pm$ 0.3
72	0/8 (0)	9.2 $\pm$ 0.6	1	-1.3 $\pm$ 1.4	-6.3 $\pm$ 4.5**	NA	3/8 (9.3)	5.8 $\pm$ 0.4
Control	0/10 (0)	8.8 $\pm$ 0.3	1	-6.5 $\pm$ 2.0	-15.1 $\pm$ 1.7	NA	5/10 (9.0)	5.8 $\pm$ 0.8

^aTwo peramivir injections (each 30 mg/kg) were administered on the day of inoculation followed by one injection daily for 7 days (2x IM + 7d IM).

^bEstimated by the log rank test (Venables et al., 1997).

^cDeath hazard ratio (vs. placebo group) was estimated by the proportional hazards model (Cox, 1972).

^dThe loss or gain of weight was calculated for each mouse as a percentage of weight on day 0 before virus inoculation. NA – not applicable; all mice died and therefore groups cannot be compared statistically.

^eAtaxia and hind limb paresis. Shown are the number of mice with neurological signs/total number. The median day p.i. when the signs were observed is shown in parentheses.

^fDetermined on day 4 p.i. Values are mean log₁₀ EID₅₀/ml $\pm$ SE from three individual mice.

* $P < 0.01$,

** $P < 0.001$ compared to placebo-treated control group.