CORRESPONDENCE

Vaccination Against Classical Influenza in Health-Care Workers: Self-Protection and Patient Protection

by Dr. med. Sabine Wicker, Prof. Dr. rer. med. Holger F. Rabenau, Prof. Dr. med. Volkhard A. Kempf, Dr. med. Christian Brandt in volume 36/2009

Restricting Fundamental Rights

The authors propagate compulsory vaccination of medical personnel against influenza. They also demand that staff who are not willing to comply—and only these—should wear face masks, for the patients' benefit; they report high vaccination rates as a result of this stigmatization. One can just imagine how a sick person would react to individual healthcare workers wearing face masks, their fear and alienation. As they do not know the background, patients will immediately associate an uncalculable threat.

The act of imposing restrictions on people's fundamental right of respect for human dignity and physical integrity is legitimized by means of the selected literature. Tom Jefferson, from the independent Cochrane Collaboration, is cited as the influenza expert. Jefferson's conclusion, however, is devastating. In an interview with *Der Spiegel*, he responded to the question of whether one good reason exists to have the flu vaccine: "I do not see any reason." His systematic review of studies showed that 93% of flu-like infections are not due to influenza virus and that therefore the number of 10 000 annual flu deaths, as reported by the Robert Koch-Institute, is also incorrect (1).

The tolerability of the vaccine is always emphasized—in spite of gaps in the data. A previously healthy, 45 year man came to see me with vitiligo on both hands and lower arms. The medical history offered nothing but a flu jab 4 weeks previously. In my experience, only few doctors would assume a possible association in this scenario, much less report such an association. This applies even more to autoimmune diseases with a long latency period. The existing, passive notification system means that such cases would never be captured by the Drug Committee of German Physicians.

DOI: 10.3238/arztebl.2010.0193a

REFERENCES

1. Demicheli V, Di Pietrantonj C, Jefferson T: Vaccines for preventing influenza in healthy adults. *Cochrane Database of Systematic Reviews*. 2007.
2. Wicker S, Rabenau HF, Kempf V, Brandt C: Vaccination against classical influenza in health-care workers: self-protection and patient protection. *Dtsch Arztebl Int* 2009; 106 (36): 567–72.

Dr. med. Eckart Lattmann
 Rotenberg 9
 21386 Betzendorf, Germany
 eckart.lattmann@googlemail.com

A Step Further Towards Former GDR-Style Rules

Having read the article about influenza vaccination of medical personnel I can't help but perceive the German healthcare sector as one step further towards a situation resembling that in the former German Democratic Republic. I left that country for political reasons, and I would rather not see a repeat performance in this day and age. The symbolism of having to wear compulsory face masks for critical medical personnel for higher (socioethical) reasons, by offering vaccination when this is quite obviously clashing with the staff's personal convictions, can probably be understood only by someone who has first-hand experience of similar situations within a system that is closed to the outside.

However, the question that is obvious to me is what limitations are there in terms of what are "tolerable" therapeutic interventions for medical staff? Whatever the limits are, they should be respected and adhered to in the future, thus balancing personal rights with (often only seemingly) general interest. If, as the authors have found, people were not convinced on the basis of arguments in favour of widespread use of influenza vaccination, then it is particularly in the interest of achieving hospital hygiene that democratic principles—such as discussions—are not easily sacrificed—or else the medically beneficial face mask may too easily turn into a symbolic muzzle.

DOI: 10.3238/arztebl.2010.0193b

REFERENCES

1. Wicker S, Rabenau HF, Kempf V, Brandt C: Vaccination against classical influenza in health-care workers: self-protection and patient protection. *Dtsch Arztebl Int* 2009; 106 (36): 567–72.

Dr. med. Stefan Kothe
 Feldmühlenweg 43
 59494 Soest, Germany
 weissflogkothe@t-online.de

Not a Safe Protective Measure Against Infection

Masks used to cover mouth and nose are particularly suitable for sick patients—if acceptable—in order to reduce the aerosol transmission route of infection, especially during coughing fits.

For medical personnel, such face masks—as described, the shape of surgical masks—should be subject to general discussion only if used as a means of prevention, especially with regard to patients and third parties.

For medical personnel looking after flu patients, however, such face masks are not a safe protective measure against infection and, according to the rules of the responsible insurers, would therefore have to be replaced with standardized FFP masks—meeting the recommended particle protection classes 1, 2, or 3—depending in the exposure risk (see also comments of the GUV (German statutory accident insurances) dated 30

April 2009—press information from the BGW (German Social Accident Insurance Institution for the health and welfare services)—and decision of the committee for biological working materials—ABAS—609 of December 2009).

DOI: 10.3238/arztebl.2010.0193c

REFERENCES

1. Wicker S, Rabenau HF, Kempf V, Brandt C: Vaccination against classical influenza in health-care workers: self-protection and patient protection. *Dtsch Arztebl Int* 2009; 106 (36): 567–72.

Götz J. Friedenberger
Cheruskerstr. 32
10829 Berlin, Germany

In Reply:

We thank our correspondents for their letters and wish to take this opportunity to clear up possible misunderstandings. We wish to respond to individual letters as follows.

Dr Lattmann has slightly missed the point. It was important to point out again that the authors do not propose “compulsory vaccination,” but that “mandatory vaccination should be considered for medical personnel who look after immunosuppressed patients.” We would very much welcome clear regulations for dealing with these particularly vulnerable patients. In personnel with chronic viral infections (HIV or hepatitis B/C infection), relevant regulations have been in place for many years. We had already suspected that the discussion surrounding other, acute infections—such as influenza A—is often emotionally led. With regard to the case report of vitiligo after flu vaccination, we wish to point out that a merely temporal association of the initial manifestation of a disease after vaccination does not necessarily imply a causal association.

We are surprised about the comment made by Dr Kothe, that measures instigated by occupational physicians and hospitals in order to avoid nosocomial infections remind him of the regime in the former GDR. Just for information, in countries such as the United States (which certainly do not have a socialist regime), such measures have been established in the healthcare system for many years. Infection protection measures within hospitals (for example, by means of vaccination campaigns or by wearing face masks) are not a step backwards: they prevent nosocomial infections and protect staff as well as patients. Phrases such as

“compulsory vaccinations” and “muzzle” seem to be redolent of an emotional approach to the vaccination debate. None the less, the authors thank Dr Lattmann and Dr Kothe for their comments, which clarify that further information campaigns and interdisciplinary discussion are urgently required. The letter by Friedberg provides a welcome opportunity to focus in greater detail in the problems associated with choosing the right type of face mask. The question of which mask is appropriate is currently the subject of controversial discussion (1–4). Recent data have, however, shown a comparable protective effect of surgical masks and FFP2 masks (2, 3). Compared with surgical masks, FFP2 masks have been less well accepted by medical personnel, because of the increased resistance to breathing and possible skin irritations (1). The necessary protective measures for medical personnel who look after patients with infections transmitted as aerosols will have to be further evaluated, in order to achieve the best possible protection of, as well as good acceptance among, medical staff.

DOI: 10.3238/arztebl.2010.0194

REFERENCES

1. Jefferson T, Del Mar C, Dooley L, et al.: Physical interventions to interrupt or reduce the spread of respiratory viruses: systematic review. *BMJ* 2009; 339: 3675.
2. Johnson DF, Druce JD, Birch C, Grayson ML: A quantitative assessment of the efficacy of surgical and N95 masks to filter influenza virus in patients with acute influenza infection. *CID* 2009; 49: 2757.
3. Loeb M, Dafoe N, Mahony J, et al.: Surgical mask vs N95 respirator for preventing influenza among health care workers: A randomized trial. *JAMA* 2009; 302:1865–71.
4. Shine KI, Rogers B, Goldfrank LR: Novel H1N1 influenza and respiratory protection for health care workers. *N Engl J Med* 2009; 361: 1823–5.
5. Wicker S, Rabenau HF, Kempf V, Brandt C: Vaccination against classical influenza in health-care workers: self-protection and patient protection. *Dtsch Arztebl Int* 2009; 106 (36): 567–72.

Prof. Dr. rer. med. Holger F. Rabenau
Prof. Dr. med. Volkhard A.J. Kempf
Dr. med. Christian Brandt

Dr. med. Dr. med. habil. Sabine Wicker
Betriebsärztlicher Dienst
Klinikum der Johann Wolfgang Goethe-Universität Frankfurt
Theodor-Stern-Kai 7
60590 Frankfurt am Main, Germany
Sabine.Wicker@kgu.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.