Published in final edited form as:

Ann Hum Genet. 2010 March ; 74(2): 97–109. doi:10.1111/j.1469-1809.2009.00560.x.

Genome-wide association study confirms SNPs in *SNCA* and the *MAPT* region as common risk factors for Parkinson disease

Todd L. Edwards¹, William K. Scott¹, Cherylyn Almonte¹, Amber Burt¹, Eric H. Powell¹, Gary W. Beecham¹, Liyong Wang¹, Stephan Züchner¹, Ioanna Konidari¹, Gaofeng Wang¹, Carlos Singer³, Fatta Nahab³, Burton Scott⁴, Jeffrey M. Stajich⁴, Margaret Pericak-Vance¹, Jonathan Haines², Jeffery M. Vance¹, and Eden R. Martin¹

¹John P. Hussman Institute for Human Genomics and the Dr. John T. Macdonald Foundation Department of Human Genetics, Miller School of Medicine, University of Miami

²Center for Human Genetics Research, Vanderbilt University Medical Center, Vanderbilt University

³Department of Neurology, Miller School of Medicine, University of Miami

⁴Department of Medicine, Duke University Medical Center, Duke University

SUMMARY

Parkinson disease (PD) is a chronic neurodegenerative disorder with a cumulative prevalence of greater than one per thousand. To date three independent genome-wide association studies (GWAS) have investigated the genetic susceptibility to PD. These studies have also implicated several genes as PD risk loci with strong, but not genome-wide significant, associations.

In this study, we combined data from two previously published GWAS of Caucasian subjects with our GWAS of 604 cases and 619 controls for a joint analysis with a combined sample size of 1752 cases and 1745 controls. SNPs in *SNCA* (rs2736990, p-value = 6.7×10^{-8} ; genome-wide adjusted p = 0.0109, odds ratio (OR) = 1.29 [95% CI: 1.17–1.42] G vs. A allele, population attributable risk percent (PAR%) = 12%) and the *MAPT* region (rs11012, p-value = 5.6×10^{-8} ; genome-wide adjusted p = 0.0079, OR = 0.70 [95% CI: 0.62–0.79] T vs. C allele, PAR% = 8%) were genome-wide significant. No other SNPs were genome-wide significant in this analysis. This study confirms that *SNCA* and the *MAPT* region are major genes whose common variants are influencing risk of PD.

Keywords

Parkinson disease; Association study; Alpha-synuclein; Microtubule associated protein tau

INTRODUCTION

Parkinson disease [PD (OMIM 168600)] is a chronic neurodegenerative disorder with a cumulative prevalence of greater than one per thousand (Kuopio et al. 1999) with at least 1.5 million cases in the United States and 6 million worldwide (Thomas & Beal 2007). Some genetic contributions to PD are well recognized. Mutations with high penetrance were initially identified in PD due to their easily detectable effects in relatively rare, early-onset,

Corresponding Author: Eden R. Martin, John P. Hussman Institute for Human Genomics, 1501 NW 10th Avenue. BRB 305 (M-860), Miami FL 33136, Telephone: 305-243-2283, Fax: 305-243-2396, Emartin1@med.miami.edu.

Conflicts of Interest: None

Mendelian forms of PD (Bonifati et al. 2003; Kitada et al. 1998; Paisan-Ruiz et al. 2004; Polymeropoulos et al. 1997; Zimprich et al. 2004). These known mutations explain less than 10% of PD cases (Lesage & Brice 2009). Over the last several years significant effort has been focused on investigating the contributions of common variants to PD risk and age-at-onset. Candidate gene approaches to identify genetic associations with PD have been used to follow up family-based genome-wide linkage studies to identify genomic regions containing risk loci (reviewed by Lesage et al (Lesage & Brice 2009)). This focused approach restricts the number of association tests performed, but is limited to identifying loci detectable by linkage analysis (Risch & Merikangas 1996). More recently three genome-wide association studies (GWAS) have been conducted in PD, albeit with results that did not reach genome-wide significance (Fung et al. 2006a; Maraganore et al. 2005; Pankratz et al. 2009). While genes that achieve genome-wide significance in a complex disease GWAS are important, the presence of genetic heterogeneity, unobserved environmental interactions and subphenotypes with distinct genetic etiologies can all reduce the apparent contribution of important genes to below the threshold for genome-wide significance, requiring that large samples be assembled for adequate power.

Associations with PD have been replicated in the candidate gene and GWAS contexts, including those described early in PD association studies, such as alpha-synuclein (*SNCA*, Entrez Gene ID (EGI):6622) (Farrer et al. 2001; Kruger et al. 1999; Maraganore et al. 2006; McCulloch et al. 2008; Mueller et al. 2005; Myhre et al. 2008; Sutherland et al. 2009) and the microtubule-associated protein tau (*MAPT*, EGI:4137) inversion region on chromosome 17 (Fidani et al. 2006; Fung et al. 2006b; Goris et al. 2007; Healy et al. 2004; Kwok et al. 2004; Levecque et al. 2004; Mamah et al. 2005; Martin et al. 2001; Scott et al. 2001; Skipper et al. 2004; Vandrovcova et al. 2007; Winkler et al. 2007; Zabetian et al. 2007; Zappia et al. 2003), as well as ubiquitin-specific protease 24 (*USP24*, EGI:23358) (Haugarvoll et al. 2009; Li et al. 2006; Oliveira et al. 2005), ELAV-like 4 (*ELAVL4*, EGI:1996) (DeStefano et al. 2008; Haugarvoll et al. 2007; Nouredine et al. 2005), monoamine oxidase B (*MAOB*, EGI:4129) (Kurth et al. 1993), Apolipoprotein E (*APOE*, EGI:348) (Rubinshtein et al. 1994), and the mitochondrial haplogroups (Autere et al. 2004; Gaweda-Walerych et al. 2008; Ghezzi et al. 2005; Huerta et al. 2005; Kosel et al. 1998; Pyle et al. 2005; Ross et al. 2003; van der Walt et al. 2003). The consistency of results, particularly for *SNCA* and *MAPT*, suggest that the failure to reach genome-wide significance in previous studies is due to the relatively small GWAS datasets.

We have conducted a GWAS of PD at the Miami Institute for Human Genomics (MIHG) in a sample of 604 unrelated cases and 619 unrelated controls using 491,376 autosomal and sex chromosome SNPs. To increase the power of the analysis we included data from two of the three previous PD GWAS: The National Institute of Neurological Disorders and Stroke (NINDS) (Fung et al. 2006a), and the joint dataset from the Progeni/GenePD studies that was genotyped at the Center for Inherited Disease Research (CIDR) (Pankratz et al. 2009), and excluding the Mayo clinic GWAS which used discordant sibling pairs as subjects for analysis (Maraganore et al. 2005). This provided a joint analysis dataset with a combined sample size of 1,752 cases and 1,745 controls with genotypes at 422,322 SNPs after imputation and sample and SNP quality control procedures.

We demonstrate in our Caucasian-based population that the *SNCA* and *MAPT* regions are the strongest genetic contributors to PD risk, reaching genome-wide significance and establishing these factors without controversy. In addition, several genes replicated in all three datasets, but with less stringent significance. Although they did not achieve genome-wide significance in the joint analysis, the consistency of their effects makes them strong candidates and may provide additional insight into the pathological mechanisms of PD.

MATERIALS AND METHODS

Samples

Samples in the MIHG GWAS include individuals with PD collected by one of 13 ascertainment centers in the PD Genetics Collaboration (Scott et al. 2001) or by the Morris K. Udall Parkinson Disease Center of Excellence (J.M. Vance, PI) ascertainment core. These participants were recruited by participating movement disorder and neurology clinics, referrals, and advertisements. Unaffected spouse and friend controls were recruited when available and willing to participate. All participants provided written informed consent, in accord with protocols established by institutional review boards at each center.

All individuals with PD were examined by a board-certified neurologist. A neurological exam and standard clinical evaluation was performed on all participants with PD. Affected individuals exhibited at least two cardinal symptoms of PD, e.g. bradykinesia, resting tremor, and rigidity and no other causes of Parkinsonism or atypical clinical features. Unaffected individuals had no symptoms of PD upon physical examination and self-reported symptom questionnaire (Rocca et al. 1998). Individuals were excluded if there was a history of encephalitis, neuroleptic therapy within one year before diagnosis, evidence of normal pressure hydrocephalus, or a clinical course with unusual features suggesting atypical or secondary Parkinsonism. Additionally, a blood sample, family history, medical history, and standard cognitive test (Blessed Orientation Memory Concentration (BOMC) (Katzman et al. 1983) test or Modified Mini Mental Status exam (3MS) (Folstein et al. 1975)) were obtained for each individual. To ensure diagnostic consistency across sites, clinical data for all participants were reviewed by a panel consisting of a board-certified neurologist with fellowship training in movement disorders, a board certified neurologist and medical geneticist, and a certified physician assistant.

Genotyping

Genotypes for 635 PD cases and 255 PD controls were generated using the Illumina Infinium 610-quad BeadChip (Illumina, San Diego, CA, USA) and the Illumina Infinium II assay protocol (Gunderson et al. 2005). Additionally, we included 223 cognitively-normal controls with no PD symptoms by self-reported symptom questionnaire (Rocca et al. 1998) from a previous GWAS (Beecham et al. 2009) of late-onset Alzheimer disease (LOAD) genotyped using the Illumina HumanHap 550 BeadChip, and another 164 cognitively-normal controls from a second LOAD study with no self-reported PD symptoms by questionnaire (Rocca et al. 1998) genotyped using the 1M-Duo Infinium HD BeadChip. Genotypes were determined using Illumina BeadStudio Genotyping Module version 3.2.33, samples with 99% genotyping efficiency were used to redefine genotype clusters, per the manufacturer's recommendation. Concordance of genotype calls for two CEPH samples with six replicates each was 99.98%.

Quality Control

Samples with genotyping efficiency of greater than 98% were included in subsequent QC and statistical analysis steps. One case and seven control samples were removed for low efficiency. Population stratification was assessed using Structure (Falush et al. 2003; Pritchard et al. 2000) and Eigenstrat (Price et al. 2006). For Structure analysis, 5000 independent autosomal SNPs with minor allele frequency (MAF) > 0.25 were chosen using PLINK (Purcell et al. 2007) with an r^2 threshold of 0.2 using 10,000 iterations of burn-in and 15,000 iterations of estimation. These analyses indicated that no stratification was present in our sample (Supplemental Figure 1a). For Eigenstrat analysis, 30,000 independent autosomal SNPs with MAF>0.25 were used to generate plots of principal component loadings for samples and to remove outliers (6 cases, 7 controls) using the top ten principal

components over five iterations with a threshold of six standard deviations (Supplemental Figure 1b).

Further quality control steps for the MIHG samples included checks for duplicated and related samples using mean identity by state (22 cases removed as duplicate samples). Two cases and nine controls were identified for excess or insufficient heterozygosity at sex chromosomes and were removed. Finally SNPs with $MAF < 0.01$ were removed in the combined case-control dataset (21,976 SNPs), significant deviation from Hardy-Weinberg equilibrium (HWE) in controls (78 SNPs with $p < 1.00 \times 10^{-7}$), and differential missing rate between cases and controls (29 SNPs with $p < 1 \times 10^{-5}$) were removed.

Since controls from the AD GWAS dataset were genotyped on different Illumina chips, we examined all controls for homogeneity of genotype frequencies. This filter is used to detect where genotyping error rates differ across studies, and is a critical QC step when multiple genotyping experiments are combined into a single data set for analysis. A 4 degree of freedom (df) Fisher's exact chi-square test with 10,000 permutations was used at all SNPs to test for frequency differences for the three genotypes and the three genotyping chips in the study. SNPs were removed if the p-value for this test was less than 0.001 (546 SNPs). After all QC procedures, 604 cases and 619 controls with 491,376 SNPs were available for association analysis in the MIHG GWAS dataset. The sex and age distributions for these samples are described in Table 1.

Data were downloaded from dbGAP (<http://www.ncbi.nlm.nih.gov/sites/entrez?db=gap>) for the CIDR (Pankratz et al. 2009) and NINDS (Fung et al. 2006a) PD GWAS. Imputation of SNP genotypes from these and the MIHG data was performed independently using the software package Impute (Marchini et al. 2007). Samples were removed from these GWAS for genotyping efficiency < 0.98 (CIDR: 9 cases, 4 controls removed; NINDS 11 cases, 9 controls removed). SNP data for these GWAS were removed for genotyping efficiency < 0.98 (CIDR: 7,886 SNPs removed; NINDS: 16,886 SNPs removed), $MAF < 0.01$ (CIDR: 7,676 SNPs removed; NINDS: 4,596 SNPs removed), HWE $p < 10^{-7}$ (CIDR: 790 SNPs removed; NINDS: 67 SNPs removed), and missing at random by disease status $p < 10^{-5}$ (CIDR: 29 SNPs removed). Eigenstrat and Structure were used to ensure that the merged dataset did not contain stratified samples (Supplemental Figures 2a, b). The Fisher's exact test for homogeneity described above was run for all 5 genotyping platforms using an 8 df test, removing SNPs for $p < 0.001$. Imputation was performed on each dataset independently after QC filters had been applied. The remaining high-quality samples and genotypes from each study were used to impute SNPs based on the HapMap reference panel. Individual genotypes with probability of 90% or greater were included. Imputed SNPs with greater than 5% missing genotypes were excluded from analysis. Upon merging the imputed GWAS files in PLINK, Hardy-Weinberg, minor allele frequency, SNP genotyping efficiency, and tests for differential missing rate by phenotype filters were applied as described above. The final dataset for analysis contained 1,752 cases and 1,745 controls with genotypes at 498,571 genotyped and imputed SNPs.

Association analysis of the genotype data was conducted with PLINK (Purcell et al. 2007). Cochran-Armitage (Armitage 1955) trend tests were calculated at each SNP to assess allelic association. Additional analyses evaluating dominant, recessive and genotypic exposures were performed using logistic regression in PLINK. To avoid over-correcting for multiple comparisons by using the conservative Bonferroni correction, we used PLINK to generate empirically adjusted p-values based on 10,000 permutations to adjust for multiple tests (Purcell et al. 2007). Additional logistic regression analysis fitting covariates for age of onset/age at exam, sex or history of smoking was conducted to assess confounding by these variables at all SNPs. If there was an indication that an individual had reported any history

of smoking, they were scored as a smoker, a non-smoker if they reported not smoking, and missing otherwise.

The population proportional attributable risk (PAR%) presented in the Abstract was calculated using the formula for retrospective studies, $[(\text{SNP allele frequency}) \times (\text{OR} - 1)] / [1 + (\text{SNP allele frequency}) \times (\text{OR} - 1)]$, using the SNP allele frequency in controls as the exposure frequency, and the odds ratio (OR) in place of the relative risk (Woodward 2005).

Assessment of associated SNPs and the presence of chromosome 17q21.31 alleles in the H1–H2 haplotype clades in *MAPT* was accomplished using rs1981997 as a haplotype tag SNP, because the major (G) and minor (A) alleles of this SNP are fixed in the H1 and H2 clades respectively (Stefansson et al. 2005). A 2-locus haplotype association analysis was performed using PLINK with SNPs in the *MAPT* region to determine which alleles were on the H1 haplotype, which has previously been associated with PD (Farrer et al. 2002; Golbe et al. 2001; Maraganore et al. 2001; Martin et al. 2001; Refenes et al. 2009; Skipper et al. 2004; Zabetian et al. 2007).

In addition to the statistical analysis of the joint sample, as a final check against cryptic bias arising due to genotyping error rate heterogeneity across studies, we evaluated association with meta-analysis techniques using METAL (Abecasis & Willer 2007).

RESULTS

In the MIHG sample no SNPs were statistically significantly associated with PD at the genome-wide level using permutation tests with a multiple testing corrected threshold of $p < 0.05$ (Table 2). *SNCA* showed the strongest association in the MIHG study (*SNCA* intron, rs356220 $p = 2.7 \times 10^{-6}$, OR = 1.48; 95% CI [1.25–1.74]). Other top associations included SNPs in the chromosome 1p22 gene chloride channel accessory 4 (*CLCA4*, EGI:22802, intron, rs1543467 $p = 3.0 \times 10^{-6}$, OR = 1.51; 95% CI [1.27–1.80]), the chromosome 3q26 gene neuroligin 1 (*NLGN1*, EGI:22871, intron, rs976683 $p = 1 \times 10^{-5}$, OR = 1.49; 95% CI [1.25–1.78]) and the chromosome 20p11 gene solute carrier family 24, member 3 (*SLC24A3*, EGI:57419, intronic SNPs, rs1406968 $p = 1.2 \times 10^{-5}$, OR = 1.47; 95% CI [1.24–1.75]; rs4813368 $p = 1.2 \times 10^{-5}$, OR = 1.48; 95% CI [1.24–1.76]; r^2 for rs1406968:rs4813368 = 0.94). Neither sex nor age-at-onset (age-at-exam in controls) were confounders for the top associations in the MIHG sample.

Two SNPs were statistically significant after empirical adjustment for multiple comparisons (permutation $p < 0.05$) in the joint analysis of the MIHG, CIDR and NINDS samples (Table 3, Supplemental Table 1); notably these SNPs were genotyped, not imputed, in all three datasets. These SNPs are in the genes pleckstrin homology domain-containing protein, family M, member 1 (*PLEKHM1*, EGI:9842, promoter, rs11012 $p = 5.6 \times 10^{-8}$, OR = 0.71; 95% CI [0.62–0.79]), and alpha-synuclein (*SNCA* intron, rs2736990 $p = 6.7 \times 10^{-8}$, OR = 1.30; 95% CI [1.18–1.43]). In addition to rs2736990 in *SNCA*, there were two SNPs with $9 \times 10^{-5} > p\text{-value} > 1 \times 10^{-5}$ (rs11931074, rs356220), two SNPs with $9 \times 10^{-4} > p\text{-value} > 1 \times 10^{-4}$ (rs365188, rs1866995), two with $9 \times 10^{-3} > p\text{-value} > 1 \times 10^{-3}$ (rs2583985, rs3775439) as well as three more SNPs with $p < 0.05$ (rs3857059, rs12502363, rs3822095), out of 15 total *SNCA* SNPs. The SNP rs356220, which was the most associated SNP from the MIHG study, had a p -value of 9.7×10^{-5} in the joint sample. The LD in *SNCA* is weaker than that at chromosome 17q21.31, due to the large inversion there which inhibits recombination. The 3' terminus of *PLEKHM1* is 400 kilobases from the 3' terminus of *MAPT* on chromosome 17, and rs11012 is in linkage disequilibrium (LD; mean $r^2 = 0.75$; Supplemental Figure 3) with SNPs in *MAPT*. Strong association ($p < 1 \times 10^{-5}$) with PD spans the entire region around *MAPT*, including SNPs in intramembrane protease 5 (*IMP5* exonic SNPs, rs12373139 $p =$

2.9×10^{-6} , OR = 0.75; 95% CI [0.67–0.85]; rs12185268 $p = 3.6 \times 10^{-6}$ OR = 0.76; 95% CI [0.67–0.85]), *MAPT* intron (rs8070723 $p = 4.5 \times 10^{-6}$, OR = 0.76; 95% CI [0.67–0.85]) and chromosome 17 open reading frame 69 (C17orf69 exon:Y132C, rs393152 $p = 3.5 \times 10^{-6}$, OR = 0.76; 95% CI [0.67–0.85]). The smallest-to-largest ranking of the p-values for association with these SNPs and PD varied greatly within the three datasets (rs11012 ranked 290 in MIHG, 11 in CIDR and 78,027 in NINDS; rs2736990 ranked 38 in MIHG, 440 in CIDR and 41,827 in NINDS) illustrating that real associations might be deep within the ranked associations in underpowered GWAS studies, and increased sample size will help true association rise to the top. Neither sex nor age-at-onset (age-at-exam in controls) were confounders for the top associations in the joint analysis. Additionally, no SNP was significant after correction for multiple tests in the dominant, recessive or genotypic analyses.

Haplotype analysis with associated chromosome 17q21.31 SNPs and the H1–H2 haplotype clade tag SNP rs1981997 revealed high D' and r^2 values for all SNPs in the region, and alleles at those SNPs with positive effect sizes were in strong LD with the G allele of rs1981997, indicating their presence in the H1 haplotype clade (Supplemental Table 3, Supplemental Figure 3).

Additional strong association signals in the joint data were observed in SNPs in the chromosome 9q33 gene deleted in bladder cancer 1 (*DBCI*, EGI:1620, intron, rs4837628 $p = 1.07 \times 10^{-6}$, OR = 0.79; 95% CI [0.72–0.87]), the chromosome 14q13 gene neuronal PAS domain protein 3 (*NPAS3*, EGI:64067, intron, rs974002 $p = 9.9 \times 10^{-6}$, OR = 1.32; 95% CI [1.17–1.49]), and in imputed SNPs in the chromosome 5q35 gene glucose fructose-6-phosphate transaminase 2 (*GFPT2*, EGI:9945, intronic SNPs, rs17080196 $p = 7.7 \times 10^{-6}$, OR = 0.77; 95% CI 0.69–0.87; rs7703402 (promoter) $p = 8.4 \times 10^{-6}$, OR = 0.76; 95% CI [0.66–0.85]; rs6864729 $p = 8.9 \times 10^{-6}$, OR = 0.77; 95% CI [0.69–0.87]; rs2303012 9.5×10^{-6} , OR = 0.77; 95% CI [0.69–0.87]; Table 3, Supplemental Table 1). However, none of these associations survive multiple testing adjustments.

Weaker associations in the joint data that replicated at an uncorrected $p \leq 0.05$ with consistent effect directions in each of the three GWAS were observed at several loci (Table 4, Supplemental Table 2). These associations were observed at rs12063142 in the chromosome 1p36.13 gene taste receptor 1, member 2 (*TAS1R2*, intron, rs12063142 $p = 4.83 \times 10^{-7}$, OR = 0.76; 95% CI [0.69–0.85]), rs974002 in *NPAS3*, the chromosome 8q21 gene matrix metalloproteinase 16 (*MMP16*, EGI:4325, intron, rs3851539 $p = 3.8 \times 10^{-5}$, OR = 1.21; 95% CI [1.11–1.34]), chromosome 12q24 gene kinase suppressor of ras 2 (*KSR2*, EGI:283455, intron, rs7960736 $p = 6.3 \times 10^{-5}$, OR = 1.22; 95% CI [1.11–1.34]), chromosome 14q24 SNP rs11159221 ($p = 1 \times 10^{-4}$, OR = 0.79; 95% CI [0.69–0.89]), chromosome 2q31 gene WAS/WASL-interacting protein family, member1 (*WASPIP*, EGI:7456, intron, rs1991601 $p = 3 \times 10^{-4}$, OR = 1.23; 95% CI [1.09–1.38]), chromosome 11q32 gene family with sequence similarity 55, family A (*FAM55A*, EGI:120400, intron, rs1080074 $p = 4 \times 10^{-4}$, OR = 1.29; 95% CI [1.12–1.48]), and the chromosome 15q22 gene RAR-related orphan receptor A (*RORA*, EGI:6095, intron, rs1863270 $p = 5 \times 10^{-4}$, OR = 1.19; 95% CI [1.08–1.33]).

Results from each GWAS and the joint analysis for SNPs in the top 10 genes from the PDgene database (www.pdgene.org) are summarized in Table 5. The PDgene database is a free online meta-analysis summary of PD genetic association studies. No associations with $p < 2.00 \times 10^{-3}$ were observed in genes other than *SNCA* and *MAPT*. The strongest association among all SNPs in these genes was in monoamine oxidase B (*MAOB* intron, rs209766 $p = 0.0026$, OR = 0.86; 95% CI [0.76–0.96]). Weaker associations were also detected in leucine-rich repeat kinase 2 (*LRRK2*, EGI:120892, intron, rs1907632 $p = 0.0321$, OR = 1.15; 95%

CI [1.01–1.30]), USP24 (intron, rs12065953 $p = 0.0078$, OR = 1.46; 95% CI [1.10–1.94]), and ELAVL4 (intron, rs17105974 $p = 0.0418$, OR = 0.82; 95% CI [0.65–0.99]).

Results from the meta-analysis of the results from the three GWAS were consistent with those observed in the joint analysis (Pearson's correlation coefficient for p -values from the two analyses = 0.95). The SNPs rs11012 (meta-analysis p -value = 7.33×10^{-8}) in the chromosome 17q21.31 region and rs2736990 (meta-analysis p -value = 7.88×10^{-8}) in *SNCA* were the most strongly associated SNPs in the meta-analysis.

DISCUSSION

This study unambiguously identifies *SNCA* and *MAPT* as risk factors for idiopathic PD. The power to declare association at genome-wide levels of significance in any of these GWAS studies is low for the effects observed in this study; however, in aggregate these datasets provide conclusive evidence for major risk genes and highlight other genes of interest. Our study illustrates the utility of freely available online data resources and large collaborative studies that are better powered to detect associations with modest effects in GWAS data. It is only with joint analysis of three independent GWAS datasets that SNPs in the *SNCA* and the *MAPT* region reached genome-wide significance. The exact mechanisms by which variation in *SNCA* and *MAPT* influence risk for PD are unknown (Devine & Lewis 2008). Joint analysis with trend tests of allelic association also suggests other candidates that show strong, but not genome-wide evidence of association: *DBCI*, *NPAS3*, and *GFPT2*.

The *SNCA* gene product α -synuclein (AS) is an abundant brain protein which is localized in axon terminals where it may mediate synaptic processes (Cabin et al. 2002). PD is considered a synucleinopathy in which aggregations of precipitated filamentous AS (which form Lewy bodies) are a common finding in PD brains at autopsy (Kosaka et al. 1976; Lewy 1912; Okazaki et al. 1961; Tretiakoff 1919; Spillantini et al. 1997). Mutations in *SNCA* have been shown to cosegregate with PD with an autosomal dominant mode of inheritance (Chartier-Harlin et al. 2004; Ibanez et al. 2004; Kruger et al. 1998; Polymeropoulos et al. 1997; Singleton et al. 2003; Zarranz et al. 2004). Other variation in *SNCA* has been reported to increase the risk of PD (Kruger et al. 1999).

The *MAPT* protein (tau) is critical for the assembly and stabilization of the microtubule network, which is essential for axonal transport in neurons (Garcia & Cleveland 2001). Neurological disorders where aggregations of tau are observed in brain tissue are known as tauopathies. The most common tauopathy is AD, where hyperphosphorylated tau accumulates in intraneuronal neurofibrillary tangles (NFTs) (Goedert et al. 1989). Additionally, progressive supranuclear palsy (PSP) and corticobasal degeneration (CBD) are neurodegenerative diseases characterized by tau deposition in neurons and glia (Dickson et al. 2002; Litvan et al. 1996) and parkinsonism. A further set of disorders referred to as frontotemporal dementia and parkinsonism linked to chromosome 17 (FTDP-17) is also characterized by the abnormal deposition of tau (Ingram & Spillantini 2002). Haplotypes in the chromosome 17q21.31 region have been associated with progressive supranuclear palsy (PSP), corticobasal degeneration (CBD) (Pittman et al. 2005), and PD (Farrer et al. 2002; Golbe et al. 2001; Healy et al. 2004; Maraganore et al. 2001; Martin et al. 2001; Refenes et al. 2009; Skipper et al. 2004; Zabetian et al. 2007). In the current study we report haplotypic associations of alleles on the H1 haplotype clade which is consistent with previous investigations of PD, PSP, and CBD. The H1 clade has also been shown to have higher expression levels of tau than H2 (Kwok et al. 2004), which may provide some insight into PD pathology as previous studies have shown that tau mediates and promotes the polymerization of AS (Giasson et al. 2003).

The gene Diacylglycerol kinase, theta (*DGKQ*) was previously observed to be associated with PD in the Pankratz et al. study (Pankratz et al. 2008). This gene is thought to be active in the phosphatidylinositol signaling pathway and is expressed in the brain.

Interestingly, associations between SNPs in *NPAS3* and smoking cessation have been observed in two of three GWAS datasets investigating genetic differences between the ability of individuals to quit smoking (Uhl et al. 2008). Given the known relationship between smoking and PD (Allam et al. 2004; DORN 1959), this gene may merit further investigation in PD studies. We saw no evidence of confounding by or interaction with history of smoking at *NPAS3* SNPs (data not shown).

The MIHG data and the CIDR data were analyzed for interactions with and confounding by history of smoking. Although no SNPs were significant after correction for multiple tests (data not shown), these and other exposures may explain or modify genetic susceptibilities to PD. Previous research in PD families has observed interactions with smoking and the genes nitric oxide synthase 2A (*NOS2A*) (Hancock et al. 2006), glutathione S-transferase omega 1 (*GSTO1*) (Wahner et al. 2007) and *SNCA* (McCulloch et al. 2008). This study only included two *NOS2A* SNPs, neither of which were included in the previous study, and which do not provide sufficient coverage of the gene to evaluate the previous findings of interaction. Similarly, the report of interaction with smoking with the Rep1 variant in *SNCA* was not replicable with the SNPs in this study. The interaction with history of smoking and *GSTO1* at rs4925 was not replicated.

DBC1 is a gene that has been observed to be deleted in some bladder cancer cell lines (Habuchi et al. 1998). The *DBC1* gene product is detectable in several tissues, including brain, spinal cord, cortex and cerebellum (<http://www.genecards.org/>). The *DBC1* protein inhibits cell proliferation by negative regulation of the G1/S transition (Nishiyama et al. 2001). Additionally, this protein mediates non-apoptotic cell death (Wright et al. 2004), and is a regulator of components of the plasminogen pathway (Louhelainen et al. 2006). The relationship between PD and *DBC1* is not obvious based on the known biology of this gene; however, the cell-death phenotypes for this gene in sensitive brain regions might play a role in PD pathophysiology.

It should also be noted that the *GFPT2* results were obtained by imputation in the three GWAS samples used here. These genotypes are likely to be assigned with higher error rates than assayed SNPs. *GFPT2* is the rate-limiting enzyme for the entry of glucose into the hexosamine biosynthesis pathway and is thus a biologically important gene (Oki et al. 1999). Energy production in the central nervous system is a mixture of glycolysis and electron transport in mitochondria. As mitochondria are clearly affected in PD (Beal 2007; Schapira 2008), enzymes such as *GFPT2* may become increasingly important in energy production and thereby cell survival. Additionally, variants in this gene have been associated with type II diabetes mellitus (T2DM) and diabetic nephropathy in Caucasian and African American samples (Zhang et al. 2004); T2DM has been suggested as a risk factor for PD (Driver et al. 2008; Hu et al. 2007).

Despite the intriguing results of our study, there are some limitations. The error rate of imputed genotypes is likely to be higher than that of assayed genotypes, although this effect should be mild for imputed genotypes with high posterior probabilities (Marchini et al. 2007). An additional possible underlying source of heterogeneity is subtle differences in the PD trait definition used by the three studies. It is possible that PD is composed of several subphenotypes, each with a distinct genetic etiology. This analysis would have optimal power to detect only associations that underlie the common features of the trait definitions for all three GWAS. The power of the joint analysis is also not optimal for the magnitude of

effects seen in this study ($\geq 80\%$ power for the combined sample at $\alpha=1\times 10^{-7}$ and MAF 0.25 for odds ratios <0.59 or >1.59 ; $\geq 80\%$ power for the MIHG sample at $\alpha=1\times 10^{-7}$ and MAF 0.25 for odds ratios <0.39 or >2.14); however, with the addition of larger PD studies more genes might be conclusively associated. These results support the meta-analysis of Pankratz et al. of the CIDR GWAS results and the NINDS GWAS results to the extent that the *MAPT* region is prominently represented in the top associations (Pankratz et al. 2009); however, the overall similarity among top hits is sparse. A final limitation is the lack of detailed covariate data across datasets for exposures such as smoking, pesticide exposure and coffee drinking for model adjustment and gene $\times$ environment interaction analysis. This is an important point, as the value of these datasets would increase greatly with the inclusion of well-documented environmental and subphenotype data, such as diabetes status, depression and dementia. Future studies of PD should include important environmental factors in the data collection protocol.

In conclusion, *SNCA* and *MAPT* are significantly associated at a genome-wide level with idiopathic PD. Several other biologically plausible genes are associated with PD but do not meet the genome-wide significance threshold. PD is a complex phenotype with substantial variation in age of onset and clinical course. It is possible that more genes are associated with subsets of PD cases or are associated with more modest effects than those detected here. These results suggest that observing additional PD-gene associations will require pooling datasets to achieve very large samples of thousands of cases and controls to obtain statistically significant p-values for loci of more modest effects after correction for multiple tests. Collaborative projects with coordinated ascertainment are necessary to advance PD genetic epidemiology.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the patients and control subjects who participated in this study. We thank the members of the PD Genetics Collaboration: Martha A. Nance, Ray L. Watts, Jean P. Hubble, William C. Koller, Kelly Lyons, Rajesh Pahwa, Matthew B. Stern, Amy Colcher, Bradley C. Hiner, Joseph Jankovic, William G Ondo, Fred H. Allen, Jr., Christopher G. Goetz, Gary W. Small, Donna Masterman, Frank Mastaglia, and Jonathan L. Haines who contributed subjects to this study. Some of the samples used in this study were collected while the Udall PDRCE was at Duke University. This work was supported by National Institutes of Health grant NS39764. We also thank the investigators from the Pankratz et al (accession number: phs000126.v1.p1), and the Fung et al (accession number: phs000089.v1.p1) studies for making their data available on dbGAP.

References

1. Abecasis G, Willer C. METAL, Meta Analysis Helper. 2007 Ref Type: Unpublished Work.
2. Allam MF, Campbell MJ, Hofman A, Del Castillo AS, Fernandez-Crehuet NR. Smoking and Parkinson's disease: systematic review of prospective studies. *Mov Disord.* 2004; 19(6):614–621. [PubMed: 15197698]
3. Armitage P. Tests for Linear Trends in Proportions and Frequencies. *Biometrics.* 1955; 11(3):375–386.
4. Autere J, Moilanen JS, Finnila S, Soininen H, Mannermaa A, Hartikainen P, Hallikainen M, Majamaa K. Mitochondrial DNA polymorphisms as risk factors for Parkinson's disease and Parkinson's disease dementia. *Hum.Genet.* 2004; 115(1):29–35. [PubMed: 15108120]
5. Beal MF. Mitochondria and neurodegeneration. *Novartis.Found.Symp.* 2007; 287:183–192. [PubMed: 18074639]

6. Beecham GW, Martin ER, Li YJ, Slifer MA, Gilbert JR, Haines JL, Pericak-Vance MA. Genome-wide association study implicates a chromosome 12 risk locus for late-onset Alzheimer disease. *Am.J Hum.Genet.* 2009; 84(1):35–43. [PubMed: 19118814]
7. Bonifati V, Rizzu P, van Baren MJ, Schaap O, Breedveld GJ, Krieger E, Dekker MC, Squitieri F, Ibanez P, Joosse M, van Dongen JW, Vanacore N, van Swieten JC, Brice A, Meo G, van Duijn CM, Oostra BA, Heutink P. Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science.* 2003; 299(5604):256–259. [PubMed: 12446870]
8. Cabin DE, Shimazu K, Murphy D, Cole NB, Gottschalk W, McIlwain KL, Orrison B, Chen A, Ellis CE, Paylor R, Lu B, Nussbaum RL. Synaptic vesicle depletion correlates with attenuated synaptic responses to prolonged repetitive stimulation in mice lacking alpha-synuclein. *J Neurosci.* 2002; 22(20):8797–8807. [PubMed: 12388586]
9. Chartier-Harlin MC, Kachergus J, Roumier C, Mouroux V, Douay X, Lincoln S, Levecque C, Larvor L, Andrieux J, Hulihan M, Waucquier N, Defebvre L, Amouyel P, Farrer M, Destee A. Alpha-synuclein locus duplication as a cause of familial Parkinson's disease. *Lancet.* 2004; 364(9440):1167–1169. [PubMed: 15451224]
10. DeStefano AL, Latourelle J, Lew MF, Suchowersky O, Klein C, Golbe LI, Mark MH, Growdon JH, Wooten GF, Watts R, Guttman M, Racette BA, Perlmutter JS, Marlor L, Shill HA, Singer C, Goldwurm S, Pezzoli G, Saint-Hilaire MH, Hendricks AE, Gower A, Williamson S, Nagle MW, Wilk JB, Massood T, Huskey KW, Baker KB, Itin I, Litvan I, Nicholson G, Corbett A, Nance M, Drasby E, Isaacson S, Burn DJ, Chinnery PF, Pramstaller PP, Al-Hinti J, Moller AT, Ostergaard K, Sherman SJ, Roxburgh R, Snow B, Slevin JT, Cambi F, Gusella JF, Myers RH. Replication of association between ELAVL4 and Parkinson disease: the GenePD study. *Hum.Genet.* 2008; 124(1):95–99. [PubMed: 18587682]
11. Devine MJ, Lewis PA. Emerging pathways in genetic Parkinson's disease: tangles, Lewy bodies and LRRK2. *FEBS J.* 2008; 275(23):5748–5757. [PubMed: 19021752]
12. Dickson DW, Bergeron C, Chin SS, Duyckaerts C, Horoupian D, Ikeda K, Jellinger K, Lantos PL, Lippa CF, Mirra SS, Tabaton M, Vonsattel JP, Wakabayashi K, Litvan I. Office of Rare Diseases neuropathologic criteria for corticobasal degeneration. *J Neuropathol.Exp.Neurol.* 2002; 61(11): 935–946. [PubMed: 12430710]
13. Dorn HF. Tobacco consumption and mortality from cancer and other diseases. *Public Health Rep.* 1959; 74(7):581–593. [PubMed: 13668008]
14. Driver JA, Smith A, Buring JE, Gaziano JM, Kurth T, Logroscino G. Prospective cohort study of type 2 diabetes and the risk of Parkinson's disease. *Diabetes Care.* 2008; 31(10):2003–2005. [PubMed: 18599528]
15. Falush D, Stephens M, Pritchard JK. Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics.* 2003; 164(4):1567–1587. [PubMed: 12930761]
16. Farrer M, Maraganore DM, Lockhart P, Singleton A, Lesnick TG, de AM, West A, de SR, Hardy J, Hernandez D. alpha-Synuclein gene haplotypes are associated with Parkinson's disease. *Hum.Mol.Genet.* 2001; 10(17):1847–1851. [PubMed: 11532993]
17. Farrer M, Skipper L, Berg M, Bisceglia G, Hanson M, Hardy J, Adam A, Gwinn-Hardy K, Aasly J. The tau H1 haplotype is associated with Parkinson's disease in the Norwegian population. *Neurosci.Lett.* 2002; 322(2):83–86. [PubMed: 11958849]
18. Fidani L, Kalinderi K, Bostantjopoulou S, Clarimon J, Goulas A, Katsarou Z, Hardy J, Kotsis A. Association of the Tau haplotype with Parkinson's disease in the Greek population. *Mov Disord.* 2006; 21(7):1036–1039. [PubMed: 16552760]
19. Folstein MF, Folstein SE, McHugh PR. Mini-Mental State - Practical Method for Grading Cognitive State of Patients for Clinician. *Journal of Psychiatric Research.* 1975; 12(3):189–198. [PubMed: 1202204]
20. Fung HC, Scholz S, Matarin M, Simon-Sanchez J, Hernandez D, Britton A, Gibbs JR, Langefeld C, Stiebert ML, Schymick J, Okun MS, Mandel RJ, Fernandez HH, Foote KD, Rodriguez RL, Peckham E, de Vrieze FW, Gwinn-Hardy K, Hardy JA, Singleton A. Genome-wide genotyping in Parkinson's disease and neurologically normal controls: first stage analysis and public release of data. *Lancet Neurol.* 2006a; 5(11):911–916. [PubMed: 17052657]

21. Fung HC, Xiomerisiou G, Gibbs JR, Wu YR, Eerola J, Gurbali V, Hellstrom O, Chen CM, Duckworth J, Papadimitriou A, Tienari PJ, Hadjigeorgiou GM, Hardy J, Singleton AB. Association of tau haplotype-tagging polymorphisms with Parkinson's disease in diverse ethnic Parkinson's disease cohorts. *Neurodegener. Dis.* 2006b; 3(6):327–333. [PubMed: 17192721]
22. Garcia ML, Cleveland DW. Going new places using an old MAP: tau, microtubules and human neurodegenerative disease. *Curr. Opin. Cell Biol.* 2001; 13(1):41–48. [PubMed: 11163132]
23. Gaweda-Walerych K, Maruszak A, Safranow K, Bialecka M, Klodowska-Duda G, Czyzewski K, Slawek J, Rudzinska M, Styczynska M, Opala G, Drozdik M, Canter JA, Barcikowska M, Zekanowski C. Mitochondrial DNA haplogroups and subhaplogroups are associated with Parkinson's disease risk in a Polish PD cohort. *J Neural Transm.* 2008; 115(11):1521–1526. [PubMed: 18810306]
24. Ghezzi D, Marelli C, Achilli A, Goldwurm S, Pezzoli G, Barone P, Pellecchia MT, Stanzione P, Brusa L, Bentivoglio AR, Bonuccelli U, Petrozzi L, Abbruzzese G, Marchese R, Cortelli P, Grimaldi D, Martinelli P, Ferrarese C, Garavaglia B, Sangiorgi S, Carelli V, Torroni A, Albanese A, Zeviani M. Mitochondrial DNA haplogroup K is associated with a lower risk of Parkinson's disease in Italians. *Eur. J Hum. Genet.* 2005; 13(6):748–752. [PubMed: 15827561]
25. Giasson BI, Forman MS, Higuchi M, Golbe LI, Graves CL, Kotzbauer PT, Trojanowski JQ, Lee VM. Initiation and synergistic fibrillization of tau and alpha-synuclein. *Science.* 2003; 300(5619):636–640. [PubMed: 12714745]
26. Goedert M, Spillantini MG, Potier MC, Ulrich J, Crowther RA. Cloning and sequencing of the cDNA encoding an isoform of microtubule-associated protein tau containing four tandem repeats: differential expression of tau protein mRNAs in human brain. *EMBO J.* 1989; 8(2):393–399. [PubMed: 2498079]
27. Golbe LI, Lazzarini AM, Spychala JR, Johnson WG, Stenroos ES, Mark MH, Sage JJ. The tau A0 allele in Parkinson's disease. *Mov Disord.* 2001; 16(3):442–447. [PubMed: 11391737]
28. Goris A, Williams-Gray CH, Clark GR, Foltynie T, Lewis SJ, Brown J, Ban M, Spillantini MG, Compston A, Burn DJ, Chinnery PF, Barker RA, Sawcer SJ. Tau and alpha-synuclein in susceptibility to, and dementia in, Parkinson's disease. *Ann. Neurol.* 2007; 62(2):145–153. [PubMed: 17683088]
29. Gunderson KL, Steemers FJ, Lee G, Mendoza LG, Chee MS. A genome-wide scalable SNP genotyping assay using microarray technology. *Nat. Genet.* 2005; 37(5):549–554. [PubMed: 15838508]
30. Habuchi T, Luscombe M, Elder PA, Knowles MA. Structure and methylation-based silencing of a gene (DBCCR1) within a candidate bladder cancer tumor suppressor region at 9q32–q33. *Genomics.* 1998; 48(3):277–288. [PubMed: 9545632]
31. Hancock DB, Martin ER, Fujiwara K, Stacy MA, Scott BL, Stajich JM, Jewett R, Li YJ, Hauser MA, Vance JM, Scott WK. NOS2A and the modulating effect of cigarette smoking in Parkinson's disease. *Ann. Neurol.* 2006; 60(3):366–373. [PubMed: 16823855]
32. Haugarvoll K, Toft M, Ross OA, Stone JT, Heckman MG, White LR, Lynch T, Gibson JM, Wszolek ZK, Uitti RJ, Aasly JO, Farrer MJ. ELAVL4, PARK10, and the Celts. *Mov Disord.* 2007; 22(4):585–587. [PubMed: 17230446]
33. Haugarvoll K, Toft M, Skipper L, Heckman MG, Crook JE, Soto A, Ross OA, Hulihan MM, Kachergus JM, Sando SB, White LR, Lynch T, Gibson JM, Uitti RJ, Wszolek ZK, Aasly JO, Farrer MJ. Fine-mapping and candidate gene investigation within the PARK10 locus. *Eur. J Hum. Genet.* 2009; 17(3):336–343. [PubMed: 18854859]
34. Healy DG, bou-Sleiman PM, Lees AJ, Casas JP, Quinn N, Bhatia K, Hingorani AD, Wood NW. Tau gene and Parkinson's disease: a case-control study and meta-analysis. *J Neurol. Neurosurg. Psychiatry.* 2004; 75(7):962–965. [PubMed: 15201350]
35. Hu G, Jousilahti P, Bidel S, Antikainen R, Tuomilehto J. Type 2 diabetes and the risk of Parkinson's disease. *Diabetes Care.* 2007; 30(4):842–847. [PubMed: 17251276]
36. Huerta C, Castro MG, Coto E, Blazquez M, Ribacoba R, Guisasola LM, Salvador C, Martinez C, Lahoz CH, Alvarez V. Mitochondrial DNA polymorphisms and risk of Parkinson's disease in Spanish population. *J Neurol. Sci.* 2005; 236(1–2):49–54. [PubMed: 15975594]

37. Ibanez P, Bonnet AM, Debarges B, Lohmann E, Tison F, Pollak P, Agid Y, Durr A, Brice A. Causal relation between alpha-synuclein gene duplication and familial Parkinson's disease. *Lancet*. 2004; 364(9440):1169–1171. [PubMed: 15451225]
38. Ingram EM, Spillantini MG. Tau gene mutations: dissecting the pathogenesis of FTDP-17. *Trends Mol.Med*. 2002; 8(12):555–562. [PubMed: 12470988]
39. Katzman R, Brown T, Fuld P, Peck A, Schechter R, Schimmel H. Validation of A Short Orientation-Memory-Concentration Test of Cognitive Impairment. *American Journal of Psychiatry*. 1983; 140(6):734–739. [PubMed: 6846631]
40. Kitada T, Asakawa S, Hattori N, Matsumine H, Yamamura Y, Minoshima S, Yokochi M, Mizuno Y, Shimizu N. Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature*. 1998; 392(6676):605–608. [PubMed: 9560156]
41. Kosaka K, Oyanagi S, Matsushita M, Hori A. Presenile dementia with Alzheimer-, Pick- and Lewy-body changes. *Acta Neuropathol*. 1976; 36(3):221–233. [PubMed: 188300]
42. Kosel S, Grasbon-Frodl EM, Mautsch U, Egensperger R, von EU, Frishman D, Hofmann S, Gerbitz KD, Mehraein P, Graeber MB. Novel mutations of mitochondrial complex I in pathologically proven Parkinson disease. *Neurogenetics*. 1998; 1(3):197–204. [PubMed: 10737123]
43. Kruger R, Kuhn W, Muller T, Woitalla D, Graeber M, Kosel S, Przuntek H, Epplen JT, Schols L, Riess O. Ala30Pro mutation in the gene encoding alpha-synuclein in Parkinson's disease. *Nat.Genet*. 1998; 18(2):106–108. [PubMed: 9462735]
44. Kruger R, Vieira-Saecker AM, Kuhn W, Berg D, Muller T, Kuhn N, Fuchs GA, Storch A, Hungs M, Woitalla D, Przuntek H, Epplen JT, Schols L, Riess O. Increased susceptibility to sporadic Parkinson's disease by a certain combined alpha-synuclein/apolipoprotein E genotype. *Ann.Neurol*. 1999; 45(5):611–617. [PubMed: 10319883]
45. Kuopio AM, Marttila RJ, Helenius H, Rinne UK. Changing epidemiology of Parkinson's disease in southwestern Finland. *Neurology*. 1999; 52(2):302–308. [PubMed: 9932948]
46. Kurth JH, Kurth MC, Poduslo SE, Schwankhaus JD. Association of a monoamine oxidase B allele with Parkinson's disease. *Ann.Neurol*. 1993; 33(4):368–372. [PubMed: 8489207]
47. Kwok JB, Teber ET, Loy C, Hallupp M, Nicholson G, Mellick GD, Buchanan DD, Silburn PA, Schofield PR. Tau haplotypes regulate transcription and are associated with Parkinson's disease. *Ann.Neurol*. 2004; 55(3):329–334. [PubMed: 14991810]
48. Lesage S, Brice A. Parkinson's disease: from monogenic forms to genetic susceptibility factors. *Hum.Mol.Genet*. 2009; 18(R1):R48–R59. [PubMed: 19297401]
49. Levecque C, Elbaz A, Clavel J, Vidal JS, Amouyel P, Alperovitch A, Tzourio C, Chartier-Harlin MC. Association of polymorphisms in the Tau and Saitohin genes with Parkinson's disease. *J Neurol.Neurosurg.Psychiatry*. 2004; 75(3):478–480. [PubMed: 14966169]
50. Lewy, FH. *Handbuch der Neurologie Band III*. Berlin: Springer; 1912.
51. Li Y, Schrodi S, Rowland C, Tacey K, Catanese J, Grupe A. Genetic evidence for ubiquitin-specific proteases USP24 and USP40 as candidate genes for late-onset Parkinson disease. *Hum.Mutat*. 2006; 27(10):1017–1023. [PubMed: 16917932]
52. Litvan I, Hauw JJ, Bartko JJ, Lantos PL, Daniel SE, Horoupian DS, McKee A, Dickson D, Baner C, Tabaton M, Jellinger K, Anderson DW. Validity and reliability of the preliminary NINDS neuropathologic criteria for progressive supranuclear palsy and related disorders. *J Neuropathol. Exp. Neurol*. 1996; 55(1):97–105. [PubMed: 8558176]
53. Louhelainen JP, Hurst CD, Pitt E, Nishiyama H, Pickett HA, Knowles MA. DBC1 re-expression alters the expression of multiple components of the plasminogen pathway. *Oncogene*. 2006; 25(16):2409–2419. [PubMed: 16369496]
54. Mamah CE, Lesnick TG, Lincoln SJ, Strain KJ, de AM, Bower JH, Ahlskog JE, Rocca WA, Farrer MJ, Maraganore DM. Interaction of alpha-synuclein and tau genotypes in Parkinson's disease. *Ann.Neurol*. 2005; 57(3):439–443. [PubMed: 15732111]
55. Maraganore DM, de AM, Elbaz A, Farrer MJ, Ioannidis JP, Kruger R, Rocca WA, Schneider NK, Lesnick TG, Lincoln SJ, Hulihan MM, Aasly JO, Ashizawa T, Chartier-Harlin MC, Checkoway H, Ferrarese C, Hadjigeorgiou G, Hattori N, Kawakami H, Lambert JC, Lynch T, Mellick GD, Papapetropoulos S, Parsian A, Quattrone A, Riess O, Tan EK, Van BC. Collaborative

- analysis of alpha-synuclein gene promoter variability and Parkinson disease. *JAMA*. 2006; 296(6): 661–670. [PubMed: 16896109]
56. Maraganore DM, de AM, Lesnick TG, Strain KJ, Farrer MJ, Rocca WA, Pant PV, Frazer KA, Cox DR, Ballinger DG. High-resolution whole-genome association study of Parkinson disease. *Am.J Hum.Genet.* 2005; 77(5):685–693. [PubMed: 16252231]
 57. Maraganore DM, Hernandez DG, Singleton AB, Farrer MJ, McDonnell SK, Hutton ML, Hardy JA, Rocca WA. Case-Control study of the extended tau gene haplotype in Parkinson's disease. *Ann.Neurol.* 2001; 50(5):658–661. [PubMed: 11706972]
 58. Marchini J, Howie B, Myers S, McVean G, Donnelly P. A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat.Genet.* 2007; 39(7):906–913. [PubMed: 17572673]
 59. Martin ER, Scott WK, Nance MA, Watts RL, Hubble JP, Koller WC, Lyons K, Pahwa R, Stern MB, Colcher A, Hiner BC, Jankovic J, Ondo WG, Allen FH Jr, Goetz CG, Small GW, Masterman D, Mastaglia F, Laing NG, Stajich JM, Ribble RC, Booze MW, Rogala A, Hauser MA, Zhang F, Gibson RA, Middleton LT, Roses AD, Haines JL, Scott BL, Pericak-Vance MA, Vance JM. Association of single-nucleotide polymorphisms of the tau gene with late-onset Parkinson disease. *JAMA*. 2001; 286(18):2245–2250. [PubMed: 11710889]
 60. McCulloch CC, Kay DM, Factor SA, Samii A, Nutt JG, Higgins DS, Griffith A, Roberts JW, Leis BC, Montimurro JS, Zabetian CP, Payami H. Exploring gene-environment interactions in Parkinson's disease. *Hum.Genet.* 2008; 123(3):257–265. [PubMed: 18210157]
 61. Mueller JC, Fuchs J, Hofer A, Zimprich A, Lichtner P, Illig T, Berg D, Wullner U, Meitinger T, Gasser T. Multiple regions of alpha-synuclein are associated with Parkinson's disease. *Ann.Neurol.* 2005; 57(4):535–541. [PubMed: 15786467]
 62. Myhre R, Toft M, Kachergus J, Hulihan MM, Aasly JO, Klungland H, Farrer MJ. Multiple alpha-synuclein gene polymorphisms are associated with Parkinson's disease in a Norwegian population. *Acta Neurol.Scand.* 2008; 118(5):320–327. [PubMed: 18485051]
 63. Nishiyama H, Gill JH, Pitt E, Kennedy W, Knowles MA. Negative regulation of G(1)/S transition by the candidate bladder tumour suppressor gene DBCCR1. *Oncogene*. 2001; 20(23):2956–2964. [PubMed: 11420708]
 64. Nouredine MA, Qin XJ, Oliveira SA, Skelly TJ, van der WJ, Hauser MA, Pericak-Vance MA, Vance JM, Li YJ. Association between the neuron-specific RNA-binding protein ELAVL4 and Parkinson disease. *Hum.Genet.* 2005; 117(1):27–33. [PubMed: 15827745]
 65. Okazaki H, Lipkin LE, Aronson SM. Diffuse intracytoplasmic ganglionic inclusions (Lewy type) associated with progressive dementia and quadriplegia in flexion. *J Neuropathol.Exp.Neurol.* 1961; 20:237–244. [PubMed: 13730588]
 66. Oki T, Yamazaki K, Kuromitsu J, Okada M, Tanaka I. cDNA cloning and mapping of a novel subtype of glutamine:fructose-6-phosphate amidotransferase (GFAT2) in human and mouse. *Genomics*. 1999; 57(2):227–234. [PubMed: 10198162]
 67. Oliveira SA, Li YJ, Nouredine MA, Zuchner S, Qin X, Pericak-Vance MA, Vance JM. Identification of risk and age-at-onset genes on chromosome 1p in Parkinson disease. *Am.J Hum.Genet.* 2005; 77(2):252–264. [PubMed: 15986317]
 68. Paisan-Ruiz C, Jain S, Evans EW, Gilks WP, Simon J, van der BM, Lopez de MA, Aparicio S, Gil AM, Khan N, Johnson J, Martinez JR, Nicholl D, Carrera IM, Pena AS, de SR, Lees A, Marti-Masso JF, Perez-Tur J, Wood NW, Singleton AB. Cloning of the gene containing mutations that cause PARK8-linked Parkinson's disease. *Neuron*. 2004; 44(4):595–600. [PubMed: 15541308]
 69. Pankratz N, Wilk JB, Latourelle JC, DeStefano AL, Halter C, Pugh EW, Doheny KF, Gusella JF, Nichols WC, Foroud T, Myers RH. Genomewide association study for susceptibility genes contributing to familial Parkinson disease. *Hum.Genet.* 2009; 124(6):593–605. [PubMed: 18985386]
 70. Pittman AM, Myers AJ, bou-Sleiman P, Fung HC, Kaleem M, Marlowe L, Duckworth J, Leung D, Williams D, Kilford L, Thomas N, Morris CM, Dickson D, Wood NW, Hardy J, Lees AJ, de SR. Linkage disequilibrium fine mapping and haplotype association analysis of the tau gene in progressive supranuclear palsy and corticobasal degeneration. *J Med.Genet.* 2005; 42(11):837–846. [PubMed: 15792962]

71. Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, Pike B, Root H, Rubenstein J, Boyer R, Stenroos ES, Chandrasekharappa S, Athanassiadou A, Papapetropoulos T, Johnson WG, Lazzarini AM, Duvoisin RC, Di IG, Golbe LI, Nussbaum RL. Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science*. 1997; 276(5321): 2045–2047. [PubMed: 9197268]
72. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. *Nat.Genet*. 2006; 38(8): 904–909. [PubMed: 16862161]
73. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus genotype data. *Genetics*. 2000; 155(2):945–959. [PubMed: 10835412]
74. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am.J Hum.Genet*. 2007; 81(3):559–575. [PubMed: 17701901]
75. Pyle A, Foltynie T, Tiangyou W, Lambert C, Keers SM, Allcock LM, Davison J, Lewis SJ, Perry RH, Barker R, Burn DJ, Chinnery PF. Mitochondrial DNA haplogroup cluster UKJT reduces the risk of PD. *Ann.Neurol*. 2005; 57(4):564–567. [PubMed: 15786469]
76. Refenes N, Bolbrinker J, Tagaris G, Orlacchio A, Drakoulis N, Kreutz R. Role of the H1 haplotype of microtubule-associated protein tau (MAPT) gene in Greek patients with Parkinson's disease. *BMC Neurol*. 2009; 9:26. [PubMed: 19558713]
77. Risch N, Merikangas K. The future of genetic studies of complex human diseases. *Science*. 1996; 273(5281):1516–1517. [PubMed: 8801636]
78. Rocca WA, Maraganore DM, McDonnell SK, Schaid DJ. Validation of a telephone questionnaire for Parkinson's disease. *Journal of Clinical Epidemiology*. 1998; 51(6):517–523. [PubMed: 9636001]
79. Ross OA, McCormack R, Maxwell LD, Duguid RA, Quinn DJ, Barnett YA, Rea IM, El-Agnaf OM, Gibson JM, Wallace A, Middleton D, Curran MD. mt4216C variant in linkage with the mtDNA TJ cluster may confer a susceptibility to mitochondrial dysfunction resulting in an increased risk of Parkinson's disease in the Irish. *Exp.Gerontol*. 2003; 38(4):397–405. [PubMed: 12670626]
80. Rubinsztein DC, Hanlon CS, Irving RM, Goodburn S, Evans DG, Kellar-Wood H, Xuereb JH, Bandmann O, Harding AE. Apo E genotypes in multiple sclerosis, Parkinson's disease, schwannomas and late-onset Alzheimer's disease. *Mol.Cell Probes*. 1994; 8(6):519–525. [PubMed: 7700274]
81. Schapira AH. Mitochondria in the aetiology and pathogenesis of Parkinson's disease. *Lancet Neurol*. 2008; 7(1):97–109. [PubMed: 18093566]
82. Scott WK, Nance MA, Watts RL, Hubble JP, Koller WC, Lyons K, Pahwa R, Stern MB, Colcher A, Hiner BC, Jankovic J, Ondo WG, Allen FH Jr, Goetz CG, Small GW, Masterman D, Mastaglia F, Laing NG, Stajich JM, Slotterbeck B, Booze MW, Ribble RC, Rampersaud E, West SG, Gibson RA, Middleton LT, Roses AD, Haines JL, Scott BL, Vance JM, Pericak-Vance MA. Complete genomic screen in Parkinson disease: evidence for multiple genes. *JAMA*. 2001; 286(18):2239–2244. [PubMed: 11710888]
83. Singleton AB, Farrer M, Johnson J, Singleton A, Hague S, Kachergus J, Hulihan M, Peuralinna T, Dutra A, Nussbaum R, Lincoln S, Crawley A, Hanson M, Maraganore D, Adler C, Cookson MR, Muenter M, Baptista M, Miller D, Blancato J, Hardy J, Gwinn-Hardy K. alpha-Synuclein locus triplication causes Parkinson's disease. *Science*. 2003; 302(5646):841. [PubMed: 14593171]
84. Skipper L, Wilkes K, Toft M, Baker M, Lincoln S, Hulihan M, Ross OA, Hutton M, Aasly J, Farrer M. Linkage disequilibrium and association of MAPT H1 in Parkinson disease. *Am.J Hum.Genet*. 2004; 75(4):669–677. [PubMed: 15297935]
85. Spillantini MG, Schmidt ML, Lee VM, Trojanowski JQ, Jakes R, Goedert M. Alpha-synuclein in Lewy bodies. *Nature*. 1997; 388(6645):839–840. [PubMed: 9278044]
86. Stefansson H, Helgason A, Thorleifsson G, Steinthorsdottir V, Masson G, Barnard J, Baker A, Jonasdottir A, Ingason A, Gudnadottir VG, Desnica N, Hicks A, Gylfason A, Gudbjartsson DF, Jonsdottir GM, Sainz J, Agnarsson K, Birgisdottir B, Ghosh S, Olafsdottir A, Cazier JB, Kristjansson K, Frigge ML, Thorgeirsson TE, Gulcher JR, Kong A, Stefansson K. A common inversion under selection in Europeans. *Nat.Genet*. 2005; 37(2):129–137. [PubMed: 15654335]

87. Sutherland GT, Halliday GM, Silburn PA, Mastaglia FL, Rowe DB, Boyle RS, O'Sullivan JD, Ly T, Wilton SD, Mellick GD. Do polymorphisms in the familial Parkinsonism genes contribute to risk for sporadic Parkinson's disease? *Mov Disord.* 2009; 24(6):833–838. [PubMed: 19224617]
88. Thomas B, Beal MF. Parkinson's disease. *Hum.Mol.Genet.* 2007; 16(Spec No. 2):R183–R194. [PubMed: 17911161]
89. Tretiakoff, MC. Thesis. University of Paris; 1919.
90. Uhl GR, Liu QR, Drgon T, Johnson C, Walther D, Rose JE, David SP, Niaura R, Lerman C. Molecular genetics of successful smoking cessation: convergent genome-wide association study results. *Arch.Gen.Psychiatry.* 2008; 65(6):683–693. [PubMed: 18519826]
91. van der Walt JM, Nicodemus KK, Martin ER, Scott WK, Nance MA, Watts RL, Hubble JP, Haines JL, Koller WC, Lyons K, Pahwa R, Stern MB, Colcher A, Hiner BC, Jankovic J, Ondo WG, Allen FH Jr, Goetz CG, Small GW, Mastaglia F, Stajich JM, McLaurin AC, Middleton LT, Scott BL, Schmechel DE, Pericak-Vance MA, Vance JM. Mitochondrial polymorphisms significantly reduce the risk of Parkinson disease. *Am.J Hum.Genet.* 2003; 72(4):804–811. [PubMed: 12618962]
92. Vandrovcsa J, Pittman AM, Malzer E, bou-Sleiman PM, Lees AJ, Wood NW, de SR. Association of MAPT haplotype-tagging SNPs with sporadic Parkinson's disease. *Neurobiol.Aging.* 2007
93. Wahner AD, Glatt CE, Bronstein JM, Ritz B. Glutathione S-transferase mu, omega, pi, and theta class variants and smoking in Parkinson's disease. *Neurosci.Lett.* 2007; 413(3):274–278. [PubMed: 17194543]
94. Winkler S, Konig IR, Lohmann-Hedrich K, Vieregge P, Kostic V, Klein C. Role of ethnicity on the association of MAPT H1 haplotypes and subhaplotypes in Parkinson's disease. *Eur.J Hum.Genet.* 2007; 15(11):1163–1168. [PubMed: 17637803]
95. Woodward, M. *Epidemiology Study Design and Data Analysis.* 2 ed. Boca Raton: Chapman & Hall/CRC; 2005.
96. Wright KO, Messing EM, Reeder JE. DBCCR1 mediates death in cultured bladder tumor cells. *Oncogene.* 2004; 23(1):82–90. [PubMed: 14712213]
97. Zabetian CP, Hutter CM, Factor SA, Nutt JG, Higgins DS, Griffith A, Roberts JW, Leis BC, Kay DM, Yearout D, Montimurro JS, Edwards KL, Samii A, Payami H. Association analysis of MAPT H1 haplotype and subhaplotypes in Parkinson's disease. *Ann.Neurol.* 2007; 62(2):137–144. [PubMed: 17514749]
98. Zappia M, Annesi G, Nicoletti G, Serra P, Arabia G, Pugliese P, Messina D, Caracciolo M, Romeo N, Annesi F, Pasqua AA, Spadafora P, Civitelli D, Romeo N, Epifanio A, Morgante L, Quattrone A. Association of tau gene polymorphism with Parkinson's disease. *Neurol.Sci.* 2003; 24(3):223–224. [PubMed: 14600827]
99. Zarranz JJ, Alegre J, Gomez-Esteban JC, Lezcano E, Ros R, Ampuero I, Vidal L, Hoenicka J, Rodriguez O, Atares B, Llorens V, Gomez TE, del ST, Munoz DG, de Yebenes JG. The new mutation, E46K, of alpha-synuclein causes Parkinson and Lewy body dementia. *Ann.Neurol.* 2004; 55(2):164–173. [PubMed: 14755719]
100. Zhang H, Jia Y, Cooper JJ, Hale T, Zhang Z, Elbein SC. Common variants in glutamine:fructose-6-phosphate amidotransferase 2 (GFPT2) gene are associated with type 2 diabetes, diabetic nephropathy, and increased GFPT2 mRNA levels. *J Clin.Endocrinol.Metab.* 2004; 89(2):748–755. [PubMed: 14764791]
101. Zimprich A, Biskup S, Leitner P, Lichtner P, Farrer M, Lincoln S, Kachergus J, Hulihan M, Uitti RJ, Calne DB, Stoessl AJ, Pfeiffer RF, Patenge N, Carbajal IC, Vieregge P, Asmus F, Muller-Myhsok B, Dickson DW, Meitinger T, Strom TM, Wszolek ZK, Gasser T. Mutations in LRRK2 cause autosomal-dominant parkinsonism with pleomorphic pathology. *Neuron.* 2004; 44(4):601–607. [PubMed: 15541309]

Table 1

Demographic properties of the MIHG, CIDR and NINDS samples.

	MIHG			CIDR			NINDS		
	All MIHG	Cases	Controls	All CIDR	Cases	Controls	All NINDS	Cases	Controls
Male (freq)	595(0.49)	381(0.63)	214(0.34)	875(0.50)	358(0.41)	517(0.60)	339(0.54)	162(0.61)	127(0.48)
Female (freq)	631(0.51)	224(0.37)	407(0.66)	869(0.50)	523(0.59)	346(0.40)	289(0.46)	104(0.39)	136(0.52)
AAO(SD)	--	56.39(12.95)	--	--	62.34(10.32)	--	--	65.33(7.86)	--
AAE(SD)	66.97(11.31)	64.56(12.18)	69.32(9.86)	70.99(9.32)	71.21(8.46)	54.83(13.08)	71.73(7.82)	71.99(6.53)	69.69(8.77)

AAO – Age at onset of PD
AAE – Age at examination
SD – Standard deviation
freq – Relative frequency

Table 2

Trend test of association with PD results from the MIHG GWAS sample and exact Hardy-Weinberg test results for all SNPs associated with PD at $p < 5 \times 10^{-5}$ or less.

SNP rs#	Trend p	HWE p	Chromosome	Gene
rs356220	2.67×10^{-6}	0.930	4q22.1	<i>SNCA</i>
rs1543467	2.97×10^{-6}	0.117	1p22.3	<i>CLCA4</i>
rs12063142	5.02×10^{-6}	0.468	1p36.13	--
rs9513249	5.92×10^{-6}	0.011	13q32.1	--
rs12870589	9.45×10^{-6}	0.002	13q32.2	--
rs976683	1.04×10^{-5}	0.529	3q26.31	<i>NLGNI</i>
rs1816879	1.16×10^{-5}	0.784	15q22.1	--
rs1406968	1.16×10^{-5}	0.705	20p11.23	<i>SLC24A3</i>
rs4813368	1.22×10^{-5}	0.701	20p11.23	<i>SLC24A3</i>
rs7646773	1.30×10^{-5}	0.865	3p11.2	--
rs1992695	1.34×10^{-5}	0.422	4q28.3	--
rs7322222	1.35×10^{-5}	0.061	13q32.1	--
rs10851073	1.44×10^{-5}	0.001	13q32.2	--
rs4238458	1.52×10^{-5}	0.407	15q24.1	--
rs11625012	1.61×10^{-5}	0.507	14q22.1	--
rs7698161	1.76×10^{-5}	0.738	4q28.3	--
rs7496513	1.85×10^{-5}	0.402	15q24.1	--
rs135066	1.91×10^{-5}	1.000	22q13.2	<i>MPPED1</i>
rs13032621	1.95×10^{-5}	1.000	2q37.2	--
rs9457743	2.03×10^{-5}	0.931	6q25.3	--
rs1159278	2.21×10^{-5}	0.025	13q32.1	<i>RAP2A</i>
rs12142266	2.31×10^{-5}	0.406	1p22.3	--
rs358079	2.99×10^{-5}	0.794	3p14.3	<i>CACNA2D3</i>
rs13411180	3.09×10^{-5}	0.832	2q31.2	<i>ZNF533</i>
rs13009601	3.13×10^{-5}	0.848	2q31.2	<i>ZNF533</i>
rs6930229	3.14×10^{-5}	1.000	6q25.3	--
rs1515274	3.39×10^{-5}	0.850	2q31.2	<i>ZNF533</i>
rs13157	3.52×10^{-5}	1.000	1p36.11	<i>RUNX3</i>
rs929708	3.70×10^{-5}	0.378	3p25.3	<i>ATP2B2</i>
rs10882088	3.89×10^{-5}	0.612	10q23.33	<i>KIF11</i>
rs11833635	4.09×10^{-5}	0.743	12q15	<i>PTPRR</i>
rs6798732	4.51×10^{-5}	0.736	3p11.2	--
rs4777585	4.57×10^{-5}	0.496	15q24.1	<i>NEO1</i>
rs12938031	4.61×10^{-5}	1.000	17q21.31	<i>CRHR1</i>

SNP rs#	Trend p	HWE p	Chromosome	Gene
rs6598020	4.63×10^{-5}	0.270	11p15.5	<i>TMEM16J</i>
rs4927602	4.79×10^{-5}	0.936	2p25.3	<i>SNTG2</i>

Table 3

Results for SNPs with $p < 1 \times 10^{-6}$ from the joint analysis with imputation of the MIHG, CIDR and NINDS GWAS data.

SNP	Chromosome	Gene	Type	P_M	P_C	P_N	P_3	Empirical p
rs11012	17q21.31	<i>PLEKHM1</i>	GCG	6.58×10^{-4}	4.40×10^{-5}	0.188	5.65×10^{-8}	0.011
rs2736990	4q22.1	<i>SNCA</i>	GCG	5.08×10^{-5}	8.67×10^{-4}	0.102	6.74×10^{-8}	0.014
rs12063142	1p36.13	<i>TAS1R2</i>	GCG	6.41×10^{-6}	0.032	0.033	4.83×10^{-7}	0.159
rs4837628	9q33.1	<i>DBC1</i>	GCG	4.83×10^{-3}	5.66×10^{-5}	0.984	1.03×10^{-6}	0.281
rs11248060	4p16.3	<i>DGKQ</i>	GCG	0.244	2.05×10^{-5}	0.005	2.26×10^{-6}	0.509
rs1981997	17q21.31	<i>MAPT</i>	GCG	4.96×10^{-3}	1.37×10^{-4}	0.733	2.45×10^{-6}	0.539
rs12373139	17q21.31	<i>IMP5</i>	GCI	4.96×10^{-3}	1.92×10^{-4}	0.349	2.85×10^{-6}	0.602
rs10464059	5q35.3	--	III	1.86×10^{-4}	4.78×10^{-4}	0.939	2.91×10^{-6}	0.610
rs13355682	5q35.3	--	III	1.86×10^{-4}	4.78×10^{-4}	0.888	3.16×10^{-6}	0.640
rs393152	17q21.31	<i>C17orf69</i>	GCG	5.86×10^{-3}	2.58×10^{-4}	0.281	3.46×10^{-6}	0.674
rs12185268	17q21.31	<i>IMP5</i>	GCG	4.96×10^{-3}	2.52×10^{-4}	0.349	3.62×10^{-6}	0.689
rs8070723	17q21.31	<i>MAPT</i>	GCG	5.86×10^{-3}	2.30×10^{-4}	0.392	4.49×10^{-6}	0.723
rs611199	9q33.1	<i>DBC1</i>	GCG	5.43×10^{-3}	9.61×10^{-4}	0.212	7.65×10^{-6}	0.908
rs17080196	5q35.3	<i>GFPT2</i>	III	8.35×10^{-4}	1.52×10^{-3}	0.633	7.73×10^{-6}	0.911
rs1635291	17q21.31	--	GCG	7.27×10^{-3}	1.89×10^{-4}	0.517	7.76×10^{-6}	0.912
rs7703402	5q35.3	<i>GFPT2</i>	III	1.57×10^{-4}	1.06×10^{-3}	0.192	8.41×10^{-6}	0.936
rs6864729	5q35.3	<i>GFPT2</i>	III	8.35×10^{-4}	1.52×10^{-3}	0.696	8.87×10^{-6}	0.945
rs2303012	5q35.3	<i>GFPT2</i>	III	9.09×10^{-4}	1.52×10^{-3}	0.696	9.49×10^{-6}	0.951
rs974002	14q13.1	<i>NPAS3</i>	GCG	6.29×10^{-3}	7.37×10^{-3}	0.017	9.85×10^{-6}	0.954

P_M – p-values from the MIHG GWAS

P_C – p-values from the CIDR GWAS

P_N – p-values from the NINDS GWAS

P_3 – p-values from the joint analysis of the three GWAS

G – Genotyped SNP identifier for each sample in order MIHG, CIDR, NINDS

I – Imputed SNP identifier for each sample in order MIHG, CIDR, NINDS

Association results that replicate in the MIHG, CIDR and NINDS studies at the $p<0.05$ level with effects in the same direction. All SNPs were genotyped.

Table 4

SNP	Chromosome	Gene	P _M	P _C	P _N	P ₃
rs12063142	1p36.13	<i>TAS1R2</i>	6.41×10^{-6}	0.032	0.033	4.83×10^{-7}
rs974002	14q13.1	<i>NPAS3</i>	6.32×10^{-3}	7.39×10^{-3}	0.017	9.85×10^{-6}
rs3851539	8q21.3	<i>MMP16</i>	7.86×10^{-3}	0.015	0.034	3.81×10^{-5}
rs7960736	12q24.23	<i>KSR2</i>	0.026	0.022	6.94×10^{-3}	6.26×10^{-5}
rs11159221	14q24.3	--	0.024	0.045	5.21×10^{-3}	1.32×10^{-4}
rs1991601	2q31.1	<i>WIPF1</i>	0.019	0.043	0.043	3.26×10^{-4}
rs1080074	11q23.2	<i>FAM55A</i>	0.038	0.036	0.035	3.77×10^{-4}
rs1863270	15q22.2	<i>RORA</i>	0.047	0.044	0.032	5.33×10^{-4}

PM – p-values from the MIHG GWAS
PC – p-values from the CIDR GWAS
PN – p-values from the NINDS GWAS
P3 – p-values from the joint analysis of the three GWAS

Table 5

Summary of results for the top 10 genes in order from PDgene (June 2009) for each GWAS dataset for the SNP in each gene with the smallest joint analysis p-value.

Gene	SNP	P _M	P _C	P _N	P ₃	#SNPs in gene	Gene Size (bp)
<i>GBA</i>	rs9628662	0.694	0.165	0.581	0.153	1	10,245
<i>LRRK2</i>	rs1907632	0.423	0.255	0.023	0.032	42	144,273
<i>SNCA</i>	rs2736990	5.08×10^{-5}	8.67×10^{-4}	0.102	6.70×10^{-8}	15	112,876
<i>PINK1</i>	rs2296223	0.546	0.031	0.247	0.212	2	18,056
<i>MAPT</i>	rs1981997	4.96×10^{-3}	1.37×10^{-4}	0.733	2.50×10^{-6}	15	134,002
<i>USP24</i>	rs12065953	8.96×10^{-3}	0.195	0.129	7.81×10^{-3}	2	149,006
<i>CYP2D6</i>	rs769258	0.572	0.848	0.033	0.574	1	4,382
<i>MAOB</i>	rs209766	0.031	0.172	0.056	0.003	31	115,864
<i>ELAVL4</i>	rs17105974	3.39×10^{-3}	0.059	0.142	0.042	10	153,854
<i>APOE</i>	rs439401	0.568	0.411	0.829	0.41	2	3,611

P_M – p-values from the MIHG GWAS

P_C – p-values from the CIDR GWAS

P_N – p-values from the NINDS GWAS

P₃ – p-values from the joint analysis of the three GWAS