Published in final edited form as:

J Toxicol Environ Health A. 2010 January ; 73(10): 684–700. doi:10.1080/15287390903578539.

Muramic Acid, Endotoxin, 3-Hydroxy Fatty Acids, and Ergosterol Content Explain Monocyte and Epithelial Cell Inflammatory Responses to Agricultural Dusts

Jill A. Poole, MD^{1,2}, Gregory P. Dooley, PhD^{3,4}, Rena Saito, PhD³, Angela M. Burrell, MA^{1,2}, Kristina L. Bailey, MD², Debra J. Romberger, MD^{1,2}, John Mehaffy³, and Stephen J. Reynolds, PhD, CIH³

¹Omaha Veterans Administration Medical Center, Omaha, NE 68105

²Pulmonary, Critical Care, Sleep & Allergy Section; Department of Internal Medicine, University of Nebraska Medical Center, 985300 The Nebraska Medical Center, Omaha, NE 68198-5300

³High Plains Intermountain Center for Agricultural Health and Safety; 154B EHB Department of Environmental and Radiological Health Sciences, Colorado State University, Ft. Collins, CO 80523-1681

⁴Center for Environmental Medicine, Department of Environmental and Radiological Health Sciences, Colorado State University, Ft. Collins, CO 80523-1681

Abstract

In agricultural and other environments, inhalation of airborne microorganisms is linked to respiratory disease development. Bacterial endotoxins, peptidoglycans, and fungi are potential causative agents, but relative microbial characterization and inflammatory comparisons amongst agricultural dusts are not well described. The aim of this study was to determine the distribution of microbial endotoxin, 3-hydroxy fatty acids (3-OHFA), muramic acid, and ergosterol and evaluate inflammatory responses in human monocytes and bronchial epithelial cells with various dust samples. Settled surface dust from 5 environments was obtained: swine facility, dairy barn, grain elevator, domestic home (no pets) and domestic home with dog. Endotoxin concentration was determined by recombinant Factor C (rFC). 3-OHFA, muramic acid and ergosterol were measured using gas chromatography-mass spectrometry. Dust-induced inflammatory cytokine secretion in human monocytes and bronchial epithelial cells was evaluated. Endotoxin-independent dust-induced inflammatory responses were evaluated. Endotoxin and 3-OHFA levels were highest in agricultural dusts. Muramic acid, endotoxin, 3-OHFA and ergosterol were detected in dusts samples. Muramic acid was highest in animal farming dusts. Ergosterol was most significant in grain elevator dust. Agricultural dusts induced monocyte TNF α , IL-6, IL-8 and epithelial cell IL-6 and IL-8 secretion. Monocyte and epithelial IL-6 and IL-8 secretion was not dependent on endotoxin. House dust(s)-induced monocyte TNF α , IL-6, IL-8 secretion. Swine facility dust generally produced elevated responses compared to other dusts. Agricultural dusts are complex with significant microbial component contribution. Large animal farming dust(s)-induced inflammation is not entirely dependent on endotoxin. Addition of muramic acid to endotoxin in large animal farming environment monitoring is warranted.

Send Correspondence to: Jill A. Poole, MD, Pulmonary, Critical Care, Sleep & Allergy Section, University of Nebraska Medical Center, 985300 The Nebraska Medical Center, Omaha, NE 68198-5300, japoole@unmc.edu, Fax: 402-559-8210, Phone: 402-559-4087.

Disclosure: none

Keywords

Agricultural dust; cytokines; endotoxin; muramic acid; ergosterol; monocytes; epithelial cells

INTRODUCTION

Inhalation of components of airborne microorganisms as found in organic dusts from various environmental settings, particularly agricultural environments, may lead to adverse chronic respiratory health effects including asthma, chronic bronchitis, chronic obstructive disease, and hypersensitivity pneumonitis (Von Essen et al. 1999). Microbial agents from Gram-negative bacteria (endotoxins, lipopolysaccharide; LPS), Gram-positive bacteria (peptidoglycan; PGN), and fungi are among potential causative agents. Endotoxin is routinely measured and is linked to inflammatory outcomes, but there is disagreement as to the association of endotoxin and disease in exposed workers. Several studies attempted to establish dose-response relationships between endotoxins and respiratory diseases (Schwartz et al. 1995; Reynolds et al. 1996; Donham et al. 2000; Vogelzang et al. 1998; Milton et al. 1996; Park et al. 2001), but a universal dose-response relationship has not been developed since some studies reported high exposure without symptoms (Rask-Andersen et al. 1989) or low exposure with a possible dose-response relationship (Kateman et al. 1990).

PGN is a microbial agent also recognized to induce inflammatory responses. Muramic acid (chemical marker of PGN) was shown to correlate with inflammatory outcomes in exposed workers in European swine barns (Zhiping et al. 1996; Larsson et al. 1999). However, it has not been systematically studied in US agricultural environments. Fungi (chemical marker is ergosterol) also exist in agricultural environments. The relative distribution and combination of these agents may explain disease manifestation.

In the Midwest United States, exposure to organic aerosols in grain elevators, swine confinement facilities and dairy farms was associated with chronic obstructive pulmonary disease (Bailey et al. 2007). Our study focusing on organic dust from swine confinement facilities showed that non-endotoxin components in the dust mediate bronchial epithelial cell inflammatory responses (Romberger et al. 2002), and that this dust regulates the expression of the Gram-positive ligand receptor, Toll-like receptor-2 (TLR2; (Bailey et al. 2008)). Although endotoxin induces inflammation in mononuclear phagocytes, the non-endotoxin components in the swine facility dust appear to be driving monocyte, macrophage, and dendritic cell innate immune inflammatory responses (Poole et al. 2007, 2008, 2009). Because the swine facility dust has high levels of muramic acid, it has been speculated that Gram-positive microbial components are important in the development of swine facility dust-induced chronic respiratory diseases (Poole et al. 2008). However, distribution of muramic acid levels in other US farming environments is not well known.

In the US, there are currently no established occupational guidelines and controls for endotoxin exposure. However, among workers in swine and poultry production facilities the suggested recommended limits for endotoxin exposure are 614 EU/m³ and 900 EU/m³, respectively (Donham et al. 1995, 2000, 2002; Reynolds et al. 1996). The levels of muramic acid and ergosterol in various environmental settings (livestock, animal confinement facilities, grain elevators) have not been well-determined. This information may be critical in establishing future guidelines and controls for reducing disease. As traditional culture methods to quantify microbial exposures have proven to be of limited use, non-culture methods and assessment for microbial constituents are recommended (Douwes et al. 2003). Applied gas chromatography-mass spectrometry (GC/MS) methods that provide high

detection selectivity are one such way to characterize these complex dust samples (Sebastian et al. 2003).

This study aimed to 1) investigate dust samples from differing agricultural environments (swine confinement facility, dairy barn, grain elevator) as compared to domestic homes for their respective inflammatory effects in human monocytes and bronchial epithelial cells, and 2) measure endotoxin, 3-OHFA, muramic acid, and ergosterol utilizing a chemical-analytical approach. It was hypothesized that these dusts may differ in microbial community chemical composition and ability to induce cellular inflammatory responses.

METHODS

Chemicals and Equipment

All organic solvents were high performance liquid chromatography (HPLC) grade. Acetone, methyl tert-butyl ether, diethyl ether, and hexane were purchased from Fisher Scientific (Fair Lawn, NJ). Methanol and heptane were obtained from Mallinckrodt (Phillipsburg, NJ). Potassium hydroxide, ammonium hydroxide, paraffin oil, pentadecanol, and pyridine were purchased from Sigma Aldrich (St. Louis, MO). Bis(trimethylsilyl)-trifluoroacetamide (BSTFA) was purchased from Regis (Morton Grove, IL). Lyophilized C₁₃ labeled algal cells were purchased from Isotec (Miami, OH). Endotoxin free water was obtained from Lonza (Basel, Switzerland). Strata-X and Strata-XC solid phase cartridges were purchased from Phenomenex (Torrance, CA).

3-OHFA, muramic acid, and ergosterol were separately analyzed with a Waters Quattro Micro GC-MS/MS system operated in the electron ionization (EI) positive mode. The carrier gas was ultra high purity helium at a head column pressure of 69kPa. Injections were made using an Agilent 7683B autosampler in the splitless mode onto a DB-5ms capillary column (30mm × 0.25mm I.D., 0.25μm film thickness) (J&W Scientific, Santa Clara, CA).

Dust Sample Collection and Preparation

Settled surface dust was collected using the Omega HEPA Vacuum with a Dust Collection Filter Sock (Midwest Filtration Company, Cincinnati, OH) from 5 different locations within the month of September 2008: floor of the operating facility of a swine confinement facility (housing approximately 400-600 hogs), floor of the feeding barn of a dairy farm (approximately 50 in the building sampled, 2500 total on the facility), floor of the storage facility of a grain elevator (feed grain/feedmill), domestic home without pets (pets never allowed during home ownership), and domestic home with pet, dog (~60 pound dog). House dust was collected in the carpeted living room. In each facility three 1m² areas were sampled vacuuming the whole area in two orthogonal passes. Dust samples in the Filter Sock were immediately placed in sterile Whirl-Pak bags (Nasco, Ft. Atkinson, WI). Samples were stored at -20°C.

For chemical marker analysis studies, dust samples (5 mg) were extracted with 10ml of 0.05% TWEEN-20 by vortexing for 1 min followed by 1 hr at 100 rpm on a rotary mixer at 20°C. The Tween extraction solution was then aliquoted for separate analysis of endotoxin (rFC), 3-OHFA (GC/MS/MS), muramic acid (GC/MS/MS), and ergosterol (GC/MS). Samples for mass spectrometry were not centrifuged before aliquoting. Individual aliquots were stored at -80°C, then lyophilized prior to sample digestion for mass spectrometry. Although Tween extraction is standard practice for mass spectrometry analysis, Tween is cellular toxic, and thus, not recommended for cell culture studies. All samples were analyzed in triplicate (i.e. three 5 mg extracts were used from each dust sample).

For cell culture studies, the dust samples were placed into solution and sterile filtered by a standard published procedure (Romberger et al. 2002). Briefly, 100 mg of each dust sample was placed in 1 ml of Hanks' balanced salt solution (HBSS) without calcium (Biofluids). The mixture was vortexed for 1 min and allowed to stand at room temperature for 1 hour. The mixture was centrifuged at x g for 10 min, and the supernatant was recovered and centrifuged again. The final supernatant was filter (0.22 μ m) sterilized. Individual aliquots were stored at -80°C prior to cell culture studies. HBSS extractions were also analyzed by rFC and mass spectrometry.

Endotoxin Bioassay

Analysis of endotoxin was performed using the recombinant factor C (rFC) assay developed by Lonza (Walkersville, MD) using a reference of endotoxin activity in Endotoxin Units relative to a standard reference LPS (*E. coli* O55:B5). The activation of rFC is determined by the fluorescence generated by the enzymatic cleavage of a peptide-coumarin substrate. Endotoxin concentrations of samples were calculated according to the standard curve. Lower limit of quantification of this assay is 0.01 EU/ml. Both Tween and HBSS extractions were analyzed using the rFC bioassay. Endotoxin was not detected in HBSS and Tween blanks.

3-OHFA Analysis

GC/MS/MS analysis was modified from previous studies (Saito et al. 2009; Reynolds et al. 2005). To prepare samples for GC/MS/MS analysis of 3-hydroxy fatty acids (3-OHFAs; marker of LPS), lyophilized dust samples and standards (C_8 - C_{10} and C_{12} - C_{18} at 0, 1, 2, 5, 20, 100, 500 ng) were spiked with 25ng of C_{11} as a surrogate and digested in 0.5ml of methanolic HCl overnight at 85°C . Samples were diluted with 1ml of Lonza pyrogen-free water and spiked with 10 μ L of 100ug/ml pentadecanol prior to solid phase extraction (SPE). Strata-X 60mg/3ml polymeric reversed phase columns were conditioned with 1ml of diethyl ether and 1ml of water. Samples were loaded to the column and pulled through drop-wise with vacuum. Cartridges were dried under full vacuum for 20 minutes. 3-OHFA was eluted from the column with diethyl ether and the eluent dried under a gentle stream of nitrogen. Dried samples were incubated with 50 μ L BSTFA/1% TMCS and 5 μ L pyridine at 85°C for 30 min to form trimethylsilyl derivatives. Following derivatization, cooled samples were diluted to 100 μ L with heptane for GC/MS/MS analysis.

3-OHFA were separated by gas chromatography with an inlet temperature of 280°C and an oven temperature profile of 90 to 250°C at $5^{\circ}\text{C}/\text{min}$, 250 to 290°C at $20^{\circ}\text{C}/\text{min}$, holding 290°C for 5 min. The GC-mass spectrometer interface temperature was 300°C . The mass spectrometer was operated in multiple reactions monitoring (MRM) mode with fragment ions generated with collision energy of 10eV. The two MRM transitions (quantitation and confirmation) monitored for each 3OHFA in their respective retention time windows were as follows: C_8 m/z 231>189, 131, C_9 m/z 245>203, 131, C_{10} m/z 259.02>217, 131, C_{11} m/z 273>241, 131, C_{12} m/z 287>255, 131, C_{13} m/z 301>269, 131, C_{14} m/z 315>283, 131, C_{15} m/z 329>297, 131, C_{16} m/z 343>311, 131, C_{17} m/z 357>325, 131, and C_{18} m/z 371>339, 131. Results are presented in pmol 3-OHFA per mg dust or ml of HBSS extract. 3-OHFAs were not detected in Tween and HBSS blanks.

Muramic Acid Analysis

To prepare samples for GC/MS/MS analysis of muramic acid (marker of PGN), lyophilized samples and standards (0, 2, 5, 10, 50, 100, or 500 ng) were digested in 1 ml of methanolic HCl overnight at 100°C . A solution of C_{13} muramic acid as an isotope dilution internal standard was obtained by digestion of 4mg of C_{13} -labeled algal cells (99% C_{13}) as for the samples. Prior to SPE, 30 μ L of the C_{13} muramic acid solution was spiked into each sample

and standard as an internal standard. SPE of the sample was done with Strata-XC 60mg/3mL strong cation exchange columns conditioned with 2 ml of methanol and 2 ml of aqueous 0.1% H₃PO₄. Samples were loaded to the column with vacuum assistance and then dried under full vacuum for 20 min. Muramic acid was eluted from the column with 5% NH₄OH in acetone. Ten μ l of 100 μ g/ml pentadecanol was added to the eluent and then dried under a stream of nitrogen. Dried samples were incubated with 50 μ l BSTFA/1%TMCS and 5 μ l pyridine at 85°C for 30 min to form trimethylsilyl derivatives. Following derivatization, cooled samples were diluted to 100 μ l with heptane for GC/MS/MS analysis.

Muramic acid was analyzed with an oven temperature profile of 120 to 290°C at 20°C/min, holding 280°C for 4 min. The inlet temperature was 260°C and the GC-mass spectrometer interface temperature was set at 300°C. The mass spectrometer was operated in MRM mode with fragment ions generated with collision energy of 6eV. The two MRM transitions monitored for muramic acid were m/z 185>142 and 185>130 and m/z 190>145 for C₁₃ muramic acid. Results are presented in ng muramic acid per mg dust or ml of HBSS extract. Muramic acid was not detected in Tween and HBSS blanks.

Ergosterol Analysis

To prepare samples for GC/MS analysis of ergosterol (marker of fungal biomass), lyophilized samples and standards (0, 1, 4, 10, or 40 ng) were spiked with 10 ng of D2-ergosterol as an internal standard and digested in 3 ml of 10% methanolic KOH for 90 min at 80°C. SPE was done with Strata-X 60mg/3ml polymeric reversed phase columns conditioned with 2 ml of methanol and 2 mL of water. Samples were loaded to the column, pulled through drop-wise with vacuum and then dried under full vacuum for 20 minutes. Ergosterol was eluted from the column with 10% methanol in methyl tert-butyl ether. Twenty μ l of 0.1% paraffin oil in acetone was added to the eluent and then dried under a stream of nitrogen. Dried samples were incubated with 50 μ L of 1:1 BSTFA/1%TMCS and hexane at 80°C for 30 min to form trimethylsilyl derivatives.

Ergosterol was analyzed with an oven temperature profile of 90 to 280°C at 20°C/min, holding 280°C for 15 min. The inlet temperature was 280°C and the GC-mass spectrometer interface temperature was set at 300°C. The mass spectrometer was operated in single ion monitoring (SIM) mode targeting m/z 363, 337, and 378. For quantification, m/z 363 and 365 were monitored for ergosterol and D2-ergosterol respectively. Results are presented in ng ergosterol per mg dust or ml of HBSS extract. Ergosterol was not detected in Tween and HBSS blanks.

Cell populations

To represent the inflammatory cytokine/chemokine response of lung mononuclear phagocytes (monocytes/macrophages), human peripheral blood monocytes were chosen. Populations of human peripheral blood monocytes were obtained from the UNMC Elutriation Core Facility. Briefly, monocytes were isolated by countercurrent centrifugal elutriation of mononuclear leukocyte-rich fractions of blood cells from healthy donors undergoing leukopheresis (Wahl et al. 1984) and were rested overnight prior to experimental assays. Elutriated monocytes were >99% pure as determined by flow cytometry of cell surface antigen expression of CD14. Peripheral blood was taken with written informed consent and studies were approved by the Institutional Review Board. Human monocytes were maintained in complete RPMI, which consisted of L-glutamine-RPMI 1640 (Life Technologies, Grand Island, NY) supplemented with 10% heat-inactivated fetal bovine serum (Biofluids), 2-mercaptoethanol (5×10^{-5} M), 50 μ g/ml of streptomycin (Invitrogen, Carlsbad, CA), and 80 U/ml of amphotericin B (Invitrogen).

To represent airway epithelial cell inflammatory responses, the human bronchial epithelial cell line, BEAS-2B (simian virus 40-immortalized; American Type Culture Collection, Manassas, VA), was chosen as it also allowed enough cells to conduct the described studies. Cells were cultured under serum-free conditions using a 1:1 medium mixture of RPMI 1640 medium (Invitrogen, Carlsbad, CA) and LHC-9 (prepared from LHC basal; Biofluids, Rockville, MD) on 1% type I collagen-coated 100 mm plates as previously described (Romberger et al. 2002). LHC-9 media contains LHC basal media (Biofluids), 0.5 μ M phosphoethanolamine-ethanolamine (Sigma, St. Louis, MO), 0.11 mM calcium (Fisher, Springfield, NJ), 50 U/ml penicillin and streptomycin (Life Technologies, Grand Island, NY), 2 μ g/ml fungizone (Life Technologies), trace elements, 5 μ g/ml bovine insulin (Sigma), 5 ng/ml epidermal growth factor (Sigma), 10 μ g/ml bovine transferrin (Sigma), 10 nM 3,3',5-triiodothyronine (Biofluids), bovine pituitary extract (50 μ g protein/ml; Pel Freeze, Rogers, AR), 0.2 μ M hydrocortisone (Biofluids), 0.5 μ g/ml epinephrine (Sigma), and 0.1 μ g/ml retinoic acid (Sigma). Cells were maintained in culture at 37°C in humidified 95% air/5% CO₂ for 48-72 hr before each experiment.

Cytokine/chemokine assays

Human peripheral blood monocytes and the human bronchial epithelial cell line were stimulated with the various HBSS dust extracts to determine inflammatory cytokine responsiveness. Separate dose and time course studies determined that a 1% concentration of swine facility dust extract (generally most potent inflammatory agent) at 5 hours post exposure elicited optimal monocyte TNF α production with no evidence of cellular toxicity (Poole et al. 2007). Thus, all HBSS dust extracts were diluted in the respective monocyte or epithelial culture medium to a final concentration of 1%. Although difficult to directly compare cell culture studies with real world exposure, using endotoxin as a biomarker of exposure, a 1% concentration of swine facility dust extract equates to approximately 37.28 EU/mg of endotoxin (Table 1), and we suggest that this corresponds to a “low” swine barn exposure condition (mean endotoxin $\pm$ SEM over 5 hours, 452.3 $\pm$ 65.8 EU/m³; Dosman et al. 2006). Cells were also incubated with polymyxin B (50 mg/ml; Sigma) plus HBSS dust extracts diluted to a final concentration of 1% to determine endotoxin-independent dust-induced inflammatory cytokine responsiveness (referred to as endotoxin-depleted HBSS dust extracts 1%). Human peripheral blood monocytes (1 $\times$ 10⁶ cells/mL) in duplicate were challenged with swine confinement facility, dairy barn, grain elevator, house with dog, house without pets HBSS dust extracts (1%) and media (control) for 5 hr (optimal time) in adjacent experiments. Cell-free culture supernatant was harvested and stored at -20°C for later cytokine/chemokine assays. For airway epithelial cell studies, confluent monolayers of epithelial cells in 6-well cell culture plates were exposed to the HBSS dust extracts (1%) for 24 hr, optimal time to elicit cytokine production in adjacent experiments. Cell-free supernatant was then harvested and stored at -20°C for later cytokine/chemokine assays. Cell viability was assessed by trypan blue exclusion method after stimulation.

TNF α , IL-6 and CXCL8/IL-8 protein levels were assayed by sandwich ELISA as previously published (Poole et al. 2007). Monocyte cytokine/chemokine secretion was reported as concentration (pg/mL) per 1 $\times$ 10⁶ viable cells. Epithelial cytokine/chemokine secretion was normalized to total protein (mg) of the cell layer using Nanodrop spectrophotometer (Applied Biosciences) and reported as pg/mg/mL.

Statistical Analysis

Endotoxin levels determined by rFC bioassay were normalized to the total amount of dust sampled (EU/mg). Endotoxin's 3-OHFA (pmol/mg or pmol/mL), muramic acid (ng/mg or ng/mL), and ergosterol (ng/mg or ng/mL) concentrations were normalized to the total amount of dust sampled or to concentration in a 1% HBSS extract solution. The sum of all

3-OHFA constituents measured is reported and the % total 3-OHFA for carbon chain length is reported. Cytokine/chemokine data are presented as the mean $\pm$ standard deviation (SD). Differences in dust-stimulated inflammatory mediator secretion were compared to no stimulation (media control) and to each other. Differences in inflammatory responses between complete (untreated) dust and endotoxin-depleted (polymyxin B treated) dusts were also made. Statistics were performed using one-way parametric ANOVA with post-hoc Bonferonni for multiple comparisons. To determine difference between complete (untreated) and endotoxin depleted (polymyxin-B treated), a two-tailed, paired *t*-test was utilized using SPSS (16.0) software. Due to sample size, correlation testing was not conducted. Significance was set at $p < 0.05$.

RESULTS

Endotoxin measurements by recombinant factor C (rFC)

In these studies, endotoxin concentration and distribution of endotoxin among the various dusts was measured using the recombinant factor C (rFC) to compare results with GC/MS/MS. Table 1 shows the relative endotoxin activity of the dusts as measured by the bioassay (EU/mg of dust). In all dust extracts, significant endotoxin activity was detected. Endotoxin was significantly higher in the swine facility and grain elevator dust extracts as compared to the other dust extracts, but not different from each other. Dairy barn Tween extract was significantly higher than and domestic homes. Higher endotoxin values were measured in the Tween extractions compared to the HBSS extractions. As expected, endotoxin was significantly depleted to the lower limit of detection with polymyxin-B treatment (Table 1).

Endotoxin and Proportions of 3-OHFAs in dust samples and dust extracts

In these studies, total and specific hydroxylated fatty acids of varying carbon chain lengths (3-OHFA; endotoxin lipid A component) were measured by GC/MS/MS as described above. Figure 1A depicts the total concentration (pmol/mg dust) of 3-OHFA measured in aliquots of the dust sample Tween extractions (average of 3 dust extractions). The swine facility dust demonstrated the highest concentration, with a predominance of long carbon chain length 3-OHFA (Figure 1B). There was no statistical difference in total 3-OHFA concentration between the dairy barn, grain elevator and domestic homes. Figure 1B depicts the distribution of the 3-OHFA (% total 3-OHFAs measured in the dust samples from the various agricultural and domestic home dusts. Even numbered carbon chain length 3-OHFA was consistently elevated in the dairy barn, grain elevator, and domestic homes as compared to odd numbered 3-OHFA. The variability of 3-OHFA concentration within the replicate (triplicate) dust extractions had an average CV of 25% over all dust types. As anticipated, 3-OHFAs were undetectable in dust samples + polymyxin B by GC/MS/MS consistent with the property of polymyxin B to bind/inhibit endotoxin (data not shown).

Next, dust samples were extracted with HBSS as described in the methods section for utilization in cell culture studies at a 1% concentration. These HBSS extracts were also analyzed by mass spectrometry for 3-OHFA. Figure 2A depicts the total 3-OHFA concentration (pmol/mL) measured in an HBSS dust extract preparation of each dust, and Figure 2B depicts the distribution of 3-OHFA (% total 3-OHFAs) measured in each dust extract. The concentration of HBSS dust extracts was overall decreased respective to the Tween dust extracts. There were some differences in the fingerprint pattern of the 3-OHFAs between the HBSS dust extracts as compared to the Tween dust extracts. There was relatively higher swine facility C_8 , C_{14} , C_{16} , dairy barn C_8 , and grain elevator C_{14} . Grain elevator C_8 , C_{10} , C_{12} , C_{16} , C_{18} , house with pet C_8 , C_{10} , C_{14} , C_{18} , and house without pet C_8 , C_{12} , C_{14} , C_{16} , C_{18} were below the limit of detection for the HBSS dust extracts.

Together, these studies demonstrate that each dust has a unique 3-OHFA fingerprint, and that overall swine facility dust 3-OHFA were significantly increased with a notably different carbon chain pattern (long-chains) as compared to the other agricultural and domestic house dusts, which were more evenly distributed. Differences in extraction methods for the chemical-analytical approach (Tween) versus HBSS for the cell culture approach also resulted in a decrease in recovery of total 3-OHFA and quantitative differences in the 3-OHFA carbon chain patterns.

Muramic acid in dust samples and dust extracts

In these studies, muramic acid was measured by GC/MS/MS as described above to determine relative distribution of the Gram-positive microbial component load in the dusts. As shown in Figure 3A, muramic acid was detected primarily in large animal farming environments, with swine facility dust greater than the dairy barn dust samples (Figure 3A; Tween extraction). In the HBSS dust extract utilized in cell culture studies, muramic acid detection was significantly elevated in the swine facility and dairy barn extracts, with minimal detection in the grain elevator and domestic house dusts (Figure 3B). These data demonstrate that large animal farming environments are a rich source of peptidoglycans (muramic acid), which are primarily found in Gram positive bacteria.

Ergosterol in dust samples and dust extracts

Ergosterol was next measured by GC/MS in the dust samples and cell culture dust extracts to determine relative distribution of fungal burden. Whereas ergosterol was detected in all dust samples, grain elevator dusts contained markedly increased levels of ergosterol as compared to the other dusts (Figure 4A). In the 1% dust extracts utilized for the cell culture studies, ergosterol was only detected in the grain elevator dust extracts (Figure 4B). These results are consistent with other studies demonstrating a strong presence of fungi in grain farming/grain elevators (Swan et al. 1998).

Dust extracts stimulate inflammatory cytokine secretion in human monocytes

In these studies, the effect of the various HBSS dust extracts on inflammatory cytokine responsiveness in human monocytes was investigated. As described in the methods section, human monocytes were stimulated with 1% dilutions of HBSS dust extracts and media control for 5 hr in adjacent experiments. All dust extracts resulted in significant monocyte secretion of TNF α , IL-6, and CXCL8/IL-8 as compared to control (Figure 5A-C; N=3 separate studies). To determine endotoxin-independent responses, 1% HBSS dust extract cultures were treated with polymyxin B (endotoxin-depleted HBSS dust extracts), and differences in the ability to induce cytokine/chemokine secretion were observed. Endotoxin-depleted agricultural HBSS 1% dust extracts resulted in significant secretion of TNF α , IL-6, and IL-8 as compared to media plus polymyxin-B control (Figure 5A-C, N=3 separate studies). With the all dust extracts, there was a significant dampening of TNF α secretion with endotoxin-depletion, but there was no significant change in IL-6 and IL-8 secretion. However, there was a quantitative reduction in IL-8 in the endotoxin-depleted grain elevator dust. There was no marked difference in cell count or cell viability between treatment conditions to explain these results.

In comparing dust extract-induced cytokine secretion among the dust groups, the endotoxin-depleted swine facility dust extract demonstrated significantly increased TNF α as compared to all other endotoxin-depleted dust extracts (Figure 5A; N=3). There was also a significant increase in IL-8 with the endotoxin-depleted swine facility dust extract as compared to the endotoxin-depleted domestic house dust extracts (Figure 5C; N=3).

Dust extracts stimulate inflammatory cytokine secretion in bronchial epithelial cells

In these studies, the effect of the various dust extracts on inflammatory cytokine responsiveness of human bronchial epithelial cells was investigated. Bronchial epithelial cells were stimulated with and without a 1% dilution of various HBSS dust extracts for 24 hr in adjacent experiments. The agricultural dust extracts, particularly swine facility and dairy barn, induced significant secretion of epithelial cell IL-6 and IL-8 as compared to control (Figure 6A-B; N=3 separate studies). There was no significant change in IL-6 and IL-8 secretion in epithelial cells stimulated with endotoxin-depleted swine confinement facility or dairy barn dust extract; although, with endotoxin-depleted dairy barn dust extract, a quantitative reduction in IL-6 secretion was observed (Figure 6A; N=3). The grain elevator and domestic house dust extracts-induced inflammatory responses were overall weak. There was no marked difference in cell viability to account for these differences. TNF α was not detected in epithelial cell culture supernatants.

In analyzing complete dust extract-induced cytokine secretion among the groups, the swine facility dust extract demonstrated significantly increased IL-6 as compared to grain elevator and domestic house dust extracts (Figure 6A; N=3). In analyzing endotoxin-depleted (polymyxin-B treated) dust extract-induced cytokine secretion among the groups, the swine confinement dust extracts induced significantly elevated IL-6 as compared to all other endotoxin-depleted dust extracts. There was significantly increased IL-8 secretion with endotoxin-depleted swine facility dust extracts as compared to endotoxin-depleted grain elevator and domestic house dust extracts with a quantitative observed rise when compared to endotoxin-depleted dairy barn dust extracts.

Discussion

Organic dust exposure is an important environmental factor that has been implicated in increased morbidity among repetitively exposed subjects (Von Essen et al. 2003). Whereas it is recognized that airway inflammatory immune effects occur with agricultural work exposures, the relative contribution of different constituents of the dust and the effect of these dusts on innate immune cells remains to be fully elucidated. In this study, agricultural dusts associated with respiratory disease (Bailey et al. 2007) in Colorado and Nebraska were compared to domestic house dusts to determine the relationship between endotoxin's 3-OHFA, muramic acid, and ergosterol. The dusts' ability to elicit inflammatory cytokine responsiveness in human monocytes and bronchial epithelial cells, cells important in respiratory disease, were also investigated.

Endotoxin, derived from Gram-negative bacteria, is a well-described inflammatory agent that has been inconsistently linked to inflammatory respiratory outcomes in exposed workers (Mayeux 1997). Endotoxin in agricultural settings can be measured by the limulus amoebocyte lysate bioassay, but also by the possibly more sensitive rFC bioassay. Unlike the bioassays that fail to detect cell-bound endotoxin, mass spectrometry analysis allows determination of total amounts of both cell-bound and non-cell-bound endotoxin; chemical marker: 3-OHFAs (Sonesson et al. 1990; Reynolds et al. 2005). Whereas correlations between the bioassay and mass spectrometry were previously reported for 3-OHFA in house dust (Hines et al. 2000; Park et al. 2004; Saraf et al. 1997, 1999), patterns of 3-OHFA distribution vary by sampling locations (Hines et al. 2000; Szponar et al. 2001). Bacterial composition varies by environment (Walters et al. 1994) as several studies suggested that the presence of house pets significantly increases the concentration of endotoxins in indoor household air samples (Walters et al. 1994; Zucker et al. 2000). Similarly, livestock dusts contain much higher levels and more variable 3-OHFA distributions than grain dusts (Reynolds et al. 2005; Sebastian et al. 2003). Species differences in livestock may contribute to differences in 3-OHFA distributions, but few studies compared 3-OHFA compositions

among various agricultural dusts (Reynolds et al. 2005). This is one of the first studies to use mass spectrometry in the US to investigate the specific 3-OHFA associated with specific bacterial sources in various agriculture settings and utilize these various dust extracts in cell culture studies.

Whereas endotoxin and 3-OHFA were detected in all dust samples, 3-OHFA was strikingly increased in swine facility dust as compared to the other dust samples. The chemical structure of the swine facility dust was also markedly different with a predominance of longer-chain 3-OHFA (14 to 18 carbons in length). Previous reports indicated that shorter-chain (C_{10} - C_{14}) are positively correlated with endotoxin activity in the *Limulus* bioassay, while longer-chain (C_{16} - C_{18}) 3-OHFAs tend to have lower or even no correlation in house dust samples (Saraf et al. 1999; Park et al. 2004). Saito et al. found the opposite trend in a recent study of agricultural dusts demonstrating that longer chain (C_{15} - C_{18}) 3-OHFA were strongly positively correlated with endotoxin activity in livestock dusts. It was also suggested that longer chain (C_{12} - C_{14}) 3-OHFA may elicit more significant potent immunological effects (Dehus et al. 2006). It is well-recognized that workers of swine confinement facilities have a high prevalence of chronic respiratory disease affecting up to one-third of exposed workers (Spurzem et al. 2002).

In contrast to the swine facility dust, the dairy barn, grain elevator, and domestic house dusts had a predominance of even numbered carbon chain length 3-OHFA with a lesser predominance of odd numbered 3-OHFA. These results are consistent with our recent investigation that found that even numbered carbon chain length 3-OHFA were more likely to be associated with nasal airway inflammation in agricultural workers than odd numbered carbon chain length 3-OHFAs (Burch et al. 2009). The domestic house dusts utilized here were utilized solely as a comparative domestic dust sample to that of agricultural dusts. However, it was observed that the total endotoxin concentration determined by mass spectrometry was no different with the house dusts as compared to the dairy barn and grain elevator endotoxin. This suggests that non-endotoxin components in the agricultural dust may be critical in eliciting disease.

One such non-endotoxin component is peptidoglycan. Peptidoglycan, derived primarily from Gram-positive bacteria but also to a lesser degree from Gram-negative bacteria, can not be directly measured, but its chemical marker, muramic acid, can be measured by mass spectrometry. Muramic acid was abundantly detected primarily in the large animal farming environments, swine facility and dairy barn dusts, and only minimally detected in the grain elevator and domestic house dusts. Using the described methods, our detection of muramic acid in the dusts was overall lower (10 fold) than the amounts of muramic acid detected by others in Sweden (Sebastian et al. 2003). The reasons for this are not clear, but it may limit the ability of this assay to discriminate subtle differences between environments where muramic acid levels are expected to be lower, such as domestic homes. The relative trend of high muramic acid in US swine confinement facilities as compared to domestic homes has been previously reported (Poole et al. 2008). This study confirms those findings and adds dairy barn farming to the list of work environments enriched with Gram-positive bacteria. The finding of elevated ergosterol in grain elevators is consistent with the known increased risk of fungal exposure with handling and storing grains (Swan et al. 1998). Our study underscores the predominance of fungal elements in grain elevator dust.

To understand the relative contribution of these various dust constituents to inflammatory outcomes, the ability of the dust extracts (complete and endotoxin depleted dust extracts) to elicit cytokine/chemokine secretion from representative airway inflammatory cells was investigated. In human monocytes, agricultural and domestic house dust extracts induced significant TNF α , IL-6, and CXCL8/IL-8 secretion. Removal of endotoxin significantly

reduced TNF α responsiveness of all the dusts, but had no significant effect on IL-6 and CXCL8 responsiveness, suggesting a selective role for endotoxin in inducing inflammation. IL-6 and CXCL-8 are two mediators that have been implicated in agricultural-induced respiratory disease (Larsson et al. 1994; Jagielo et al. 1996; Clapp et al. 1994). CXCL8 is a potent neutrophil chemoattractant, and the finding of increased neutrophils in bronchial alveolar lavage fluid and nasal lavage fluid is a consistent finding among workers exposed to agricultural environments (Cormier et al. 2000; Larsson et al. 1994; Clapp et al. 1994). In large animal farming environments, peptidoglycan likely drives the innate immune inflammatory response with possible synergy with endotoxin. In contrast, in the grain elevator environment, endotoxin plus ergosterol appear to be important.

In contrast to the human monocytes, the bronchial epithelial cells displayed significant cytokine responsiveness to the agricultural dusts, primarily swine confinement facility and dairy barn, whereas the grain elevator and domestic house dusts were weakly or not IL-6 and IL-8 responsive. Depletion of endotoxin did not alter swine facility dust extract-induced cytokine responsiveness, but endotoxin depletion did modulate IL-6, but not CXCL8, responsiveness of the dairy barn extract. Although the swine facility had significantly increased levels of endotoxin as compared to the other dusts, elevated muramic acid levels predominated in the swine confinement facility and dairy barn dusts. Consistent with the notion that endotoxin is not the critical player in the agricultural dust-induced bronchial epithelial cell responses, under normal conditions, epithelial cells lack significant expression of the TLR4-MD-2 complex (LPS receptor; (Elson et al. 2007). However, bronchial epithelial cells weakly express TLR2, a pattern recognition receptor that recognizes the structurally broadest range of different bacterial compounds such as PGN, lipoproteins, lipoarabinomannans, lipomannans, glycosylphosphatidylinositol, lipoteichoic acids, and zymosan (Zahringer et al. 2008). Uehara, 2007 also found in epithelial cells are nucleotide-binding oligomerization domain (NOD) proteins including NOD1 and NOD2, which are intracellular sensors for bacterial molecules and mediate inflammation of specific motifs in PGN. Taken together, data suggest that in bronchial epithelial cells, the inflammatory cytokine response to large animal farming dusts is being driven by Gram-positive microbial components such as peptidoglycan.

A limitation of this study is that air samples were not utilized. Settled dust samples were utilized in order to obtain sufficient mass to determine detection limits in this initial study, and also to obtain sufficient mass for cell culture studies. In this study there were differences in the microbial constituents detected in the dust sample using Tween extracts and the cell culture HBSS dust extracts. In addition to simple dilution effects, the HBSS cell culture extracts were also sterile filtered (0.2 μ M pore), which eliminates large particulates, but could result in an under-prediction of inflammatory potential. This standard process prevents cell culture contamination and is argued to represent the potential lung deposition of inhaled airborne agents. In addition, HBSS extracts, namely swine facility dust extracts, has been shown to demonstrate *in vivo* biological activity in a murine model (Poole et al. 2009). It is also possible that other unknown constituents are involved in mediating the organic dust inflammatory consequences, such as particles and/or metals. However, Poole et al. previously reported that swine facility dust heat-inactivated at 120°C (a process that inactivates the biologics and leaves the metals intact) did not result in inflammatory cytokine secretion (Poole et al. 2008). Finally, the inflammatory outcome of this study was investigation of various dust-induced cytokines/chemokines, other inflammatory indices such as free radical or oxidant production were not determined, but could be investigated in future studies.

In conclusion, utilization of this chemical-analytical approach can next be applied to human population environmental studies in the United States to determine the relative contribution

of microbial components and to determine if single or synergy of multiple microbial components is mediating chronic inflammatory respiratory disease in exposed workers. The addition of muramic acid and possibly ergosterol chemical analysis to that of endotoxin is warranted when performing agricultural environmental samplings for developing health hazard control standards or guidelines and for developing effective interventions.

Acknowledgments

The authors would like to thank Lisa Chudomelka for assistance with manuscript preparation. We would also like to thank Howard E. Gendelman, MD, of the UNMC Pathology Elutriation Facility for isolating monocytes using centrifugal elutriation. Study supported by grants from the High Plains Intermountain Center for Agricultural Health and Safety (CDC/NIOSH-5-U50-OH008085; JAP, RS); National Institute of Environmental Health Sciences (K08: ES015522-01; JAP)

Abbreviations

COPD	Chronic obstructive pulmonary disease
CXCL8	Interleukin-8 (IL-8)
GC/MS	Gas chromatography-mass spectrometry
HBSS	Hanks' balanced salt solution
HPLC	High performance liquid chromatography
IL-6	Interleukin-6
LPS	Lipopolysaccharide
3-OHFA	Three hydroxy fatty acid
PBS	Phosphate buffered saline
PGN	Peptidoglycan
rFC	Recombinant Factor C endotoxin assay
TNF- α	Tumor necrosis factor-alpha

References

- Bailey KL, Meza JL, Smith LM, Von Essen SG, Romberger DJ. Agricultural exposures in patients with COPD in health systems serving rural areas. *J Agromedicine* 2007;12(3):71–6. [PubMed: 19042672]
- Bailey KL, Poole JA, Mathisen TL, Wyatt TA, Von Essen SG, Romberger DJ. Toll-like receptor 2 is upregulated by hog confinement dust in an IL-6-dependent manner in the airway epithelium. *Am J Physiol Lung Cell Mol Physiol* 2008;294(6):L1049–54. [PubMed: 18359883]
- Burch J, Svendsen E, Siegel P, Wagner S, Von Essen S, Keefe T, Mehaffy J, Serrano Martinez A, Bradford M, Baker L, Cranmer B, Saito R, Tessari J, Prinz L, Andersen C, Christensen O, Koehncke N, Reynolds S. Endotoxin exposure and inflammation markers among agricultural workers in Colorado and Nebraska. *J Toxicol and Environ Health*. 2009 In press.
- Clapp WD, Becker S, Quay J, Watt JL, Thorne PS, Frees KL, Zhang X, Koren HS, Lux CR, Schwartz DA. Grain dust-induced airflow obstruction and inflammation of the lower respiratory tract. *Am J Respir Crit Care Med* 1994;150(3):611–7. [PubMed: 8087327]
- Cormier Y, Israel-Assayag E, Racine G, Duchaine C. Farming practices and the respiratory health risks of swine confinement buildings. *Eur Respir J* 2000;15(3):560–5. [PubMed: 10759453]
- Dehus O, Hartung T, Hermann C. Endotoxin evaluation of eleven lipopolysaccharides by whole blood assay does not always correlate with limulus amebocyte lysate assay. *J Endotoxin Res* 2006;12(3):171–80. [PubMed: 16719988]

- Donham KJ, Cumro D, Reynolds S. Synergistic effects of dust and ammonia on the occupational health effects of poultry production workers. *J Agromedicine* 2002;8(2):57–76. [PubMed: 12853272]
- Donham KJ, Cumro D, Reynolds SJ, Merchant JA. Dose-response relationships between occupational aerosol exposures and cross-shift declines of lung function in poultry workers: Recommendations for exposure limits. *J Occup Environ Med* 2000;42(3):260–9. [PubMed: 10738705]
- Donham KJ, Reynolds SJ, Whitten P, Merchant JA, Burmeister L, Pependorf WJ. Respiratory dysfunction in swine production facility workers: Dose-response relationships of environmental exposures and pulmonary function. *Am J Ind Med* 1995;27(3):405–18. [PubMed: 7747746]
- Dosman JA, Fukushima Y, Senthilselvan A, Kirychuk SP, Lawson JA, Pahwa P, Cormier Y, Hurst T, Barber EM, Rhodes CS. Respiratory response to endotoxin and dust predicts evidence of inflammatory response in volunteers in a swine barn. *Am J Ind Med* 2006;49(9):761–6. [PubMed: 16917830]
- Douwes J, Thorne P, Pearce N, Heederik D. Bioaerosol health effects and exposure assessment: Progress and prospects. *Ann Occup Hyg* 2003;47(3):187–200. [PubMed: 12639832]
- Elson G, Dunn-Siegrist I, Daubeuf B, Pugin J. Contribution of toll-like receptors to the innate immune response to gram-negative and gram-positive bacteria. *Blood* 2007;109(4):1574–83. [PubMed: 17038528]
- Hines CJ, Milton DK, Larsson L, Petersen MR, Fisk WJ, Mendell MJ. Characterization and variability of endotoxin and 3-hydroxy fatty acids in an office building during a particle intervention study. *Indoor Air* 2000;10(1):2–12. [PubMed: 10842455]
- Jagiello PJ, Thorne PS, Watt JL, Frees KL, Quinn TJ, Schwartz DA. Grain dust and endotoxin inhalation challenges produce similar inflammatory responses in normal subjects. *Chest* 1996;110(1):263–70. [PubMed: 8681637]
- Kateman E, Heederik D, Pal TM, Smeets M, Smid T, Spitteler M. Relationship of airborne microorganisms with the lung function and leucocyte levels of workers with a history of humidifier fever. *Scand J Work Environ Health* 1990;16(6):428–33. [PubMed: 2284591]
- Larsson BM, Larsson K, Malmberg P, Palmberg L. Gram positive bacteria induce IL-6 and IL-8 production in human alveolar macrophages and epithelial cells. *Inflammation* 1999;23(3):217–30. [PubMed: 10392756]
- Larsson K, Malmberg P, Eklund A. Acute exposure to swine dust causes airway inflammation and bronchial hyperresponsiveness. *Am J Ind Med* 1994;25(1):57–8. [PubMed: 8116654]
- Larsson KA, Eklund AG, Hansson LO, Isaksson BM, Malmberg PO. Swine dust causes intense airways inflammation in healthy subjects. *Am J Respir Crit Care Med* 1994;150(4):973–7. [PubMed: 7921472]
- Mayeux PR. Pathobiology of lipopolysaccharide. *J Toxicol Environ Health* 1997;51(5):415–35. [PubMed: 9233377]
- Milton DK, Wypij D, Kriebel D, Walters MD, Hammond SK, Evans JS. Endotoxin exposure-response in a fiberglass manufacturing facility. *Am J Ind Med* 1996;29(1):3–13. [PubMed: 8808037]
- Park JH, Gold DR, Spiegelman DL, Burge HA, Milton DK. House dust endotoxin and wheeze in the first year of life. *Am J Respir Crit Care Med* 2001;163(2):322–8. [PubMed: 11179100]
- Park JH, Szponar B, Larsson L, Gold DR, Milton DK. Characterization of lipopolysaccharides present in settled house dust. *Appl Environ Microbiol* 2004;70(1):262–7. [PubMed: 14711650]
- Poole JA, Alexis NE, Parks C, MacInnes AK, Gentry-Nielsen MJ, Fey PD, Larsson L, Allen-Gipson D, Von Essen SG, Romberger DJ. Repetitive organic dust exposure in vitro impairs macrophage differentiation and function. *J Allergy Clin Immunol* 2008;122(2):375–82. 382 e1–4. [PubMed: 18585769]
- Poole JA, Thiele GM, Alexis NE, Burrell AM, Parks C, Romberger DJ. Organic dust exposure alters monocyte-derived dendritic cell differentiation and maturation. *Am J Physiol Lung Cell Mol Physiol*. 2009
- Poole JA, Wyatt TA, Oldenburg PJ, Elliott MK, West WW, Sisson JH, Von Essen SG, Romberger DJ. Intranasal organic dust exposure-induced airway adaptation response marked by persistent lung inflammation and pathology in mice. *Am J Physiol Lung Cell Mol Physiol*. 2009

- Poole JA, Wyatt TA, Von Essen SG, Hervert J, Parks C, Mathisen T, Romberger DJ. Repeat organic dust exposure-induced monocyte inflammation is associated with protein kinase C activity. *J Allergy Clin Immunol* 2007;120(2):366–73. [PubMed: 17555806]
- Rask-Andersen A, Malmberg P, Lundholm M. Endotoxin levels in farming: Absence of symptoms despite high exposure levels. *Br J Ind Med* 1989;46(6):412–6. [PubMed: 2818976]
- Reynolds SJ, Donham KJ, Whitten P, Merchant JA, Burmeister LF, Pependorf WJ. Longitudinal evaluation of dose-response relationships for environmental exposures and pulmonary function in swine production workers. *Am J Ind Med* 1996;29(1):33–40. [PubMed: 8808040]
- Reynolds SJ, Milton DK, Heederik D, Thorne PS, Donham KJ, Croteau EA, Kelly KM, Douwes J, Lewis D, Whitmer M, Connaughton I, Koch S, Malmberg P, Larsson BM, Deddens J, Saraf A, Larsson L. Interlaboratory evaluation of endotoxin analyses in agricultural dusts—comparison of LAL assay and mass spectrometry. *J Environ Monit* 2005;7(12):1371–7. [PubMed: 16307099]
- Romberger DJ, Bodlak V, Von Essen SG, Mathisen T, Wyatt TA. Hog barn dust extract stimulates IL-8 and IL-6 release in human bronchial epithelial cells via PKC activation. *J Appl Physiol* 2002;93(1):289–96. [PubMed: 12070216]
- Saito R, Cranmer BK, Tessari JD, Larsson L, Mehaffy JM, Keefe TJ, Reynolds SJ. Recombinant factor c (RFC) assay and gas chromatography/mass spectrometry (GC/MS) analysis of endotoxin variability in four agricultural dusts. *Ann Occup Hyg* 2009;53(7):713–22. [PubMed: 19638393]
- Saraf A, Larsson L, Burge H, Milton D. Quantification of ergosterol and 3-hydroxy fatty acids in settled house dust by gas chromatography-mass spectrometry: Comparison with fungal culture and determination of endotoxin by a limulus amoebocyte lysate assay. *Appl Environ Microbiol* 1997;63(7):2554–9. [PubMed: 9212406]
- Saraf A, Park JH, Milton DK, Larsson L. Use of quadrupole gc-ms and ion trap GC-MS-MS for determining 3-hydroxy fatty acids in settled house dust: Relation to endotoxin activity. *J Environ Monit* 1999;1(2):163–8. [PubMed: 11529095]
- Schwartz DA, Donham KJ, Olenchok SA, Pependorf WJ, Van Fossen DS, Burmeister LF, Merchant JA. Determinants of longitudinal changes in spirometric function among swine confinement operators and farmers. *Am J Respir Crit Care Med* 1995;151(1):47–53. [PubMed: 7812571]
- Sebastian A, Larsson L. Characterization of the microbial community in indoor environments: A chemical-analytical approach. *Appl Environ Microbiol* 2003;69(6):3103–9. [PubMed: 12788704]
- Sonesson A, Larsson L, Schutz A, Hagmar L, Hallberg T. Comparison of the limulus amoebocyte lysate test and gas chromatography-mass spectrometry for measuring lipopolysaccharides (endotoxins) in airborne dust from poultry-processing industries. *Appl Environ Microbiol* 1990;56(5):1271–8. [PubMed: 2187411]
- Spurzem JR, Romberger DJ, Von Essen SG. Agricultural lung disease. *Clin Chest Med* 2002;23(4):795–810. [PubMed: 12512166]
- Swan JR, Crook B. Airborne microorganisms associated with grain handling. *Ann Agric Environ Med* 1998;5(1):7–15. [PubMed: 9852487]
- Szponar B, Larsson L. Use of mass spectrometry for characterising microbial communities in bioaerosols. *Ann Agric Environ Med* 2001;8(2):111–7. [PubMed: 11748866]
- Vogelzang PF, van der Gulden JW, Folgering H, Kolk JJ, Heederik D, Preller L, Tielen MJ, van Schayck CP. Endotoxin exposure as a major determinant of lung function decline in pig farmers. *Am J Respir Crit Care Med* 1998;157(1):15–8. [PubMed: 9445272]
- Von Essen S, Donham K. Illness and injury in animal confinement workers. *Occup Med* 1999;14(2):337–50. [PubMed: 10329909]
- Von Essen S, Romberger D. The respiratory inflammatory response to the swine confinement building environment: The adaptation to respiratory exposures in the chronically exposed worker. *J Agric Saf Health* 2003;9(3):185–96. [PubMed: 12970949]
- Wahl SM, Katona IM, Stadler BM, Wilder RL, Helsel WE, Wahl LM. Isolation of human mononuclear cell subsets by counterflow centrifugal elutriation (CCE). I. Functional properties of B-lymphocyte-, T-lymphocyte-, and monocyte-enriched fractions. *Cell Immunol* 1984;85(2):384–95. [PubMed: 6232003]

- Walters M, Milton D, Larsson L, Ford T. Airborne environmental endotoxin: A cross-validation of sampling and analysis techniques. *Appl Environ Microbiol* 1994;60(3):996–1005. [PubMed: 8161191]
- Zahringer U, Lindner B, Inamura S, Heine H, Alexander C. TLR2 - promiscuous or specific? A critical re-evaluation of a receptor expressing apparent broad specificity. *Immunobiology* 2008;213(3-4): 205–24. [PubMed: 18406368]
- Zhiping W, Malmberg P, Larsson BM, Larsson K, Larsson L, Saraf A. Exposure to bacteria in swine-house dust and acute inflammatory reactions in humans. *Am J Respir Crit Care Med* 1996;154(5): 1261–6. [PubMed: 8912733]
- Zucker BA, Trojan S, Muller W. Airborne Gram-negative bacterial flora in animal houses. *J Vet Med B Infect Dis Vet Public Health* 2000;47(1):37–46. [PubMed: 10780171]

Figure 1.

(A-B). Endotoxin's 3-hydroxy fatty acid (3-OHFA) analysis of various agricultural and domestic house dusts after Tween extraction. (A) Total 3-OHFA normalized to the total amount of dust (pmol/mg). (B) The percent of the total 3-OHFAs for each carbon chain length reported for each individual dust extract. Mean results $\pm$ SEM are presented (N=3). *Statistically different than other dust extracts ($p < 0.001$).

Figure 2. (A-B). Endotoxin's 3-hydroxy fatty acid (3-OHFA) analysis of various agricultural and domestic house dusts with HBSS extraction, which was utilized in cell culture studies at a 1% concentration. (A) Total 3-OHFA normalized to the total amount of dust in a 1% concentration (pmol/mL). (B) The percent of the total 3-OHFAs for each carbon chain length reported for each individual dust extract. Mean results $\pm$ SEM are presented (N=3). *Statistically different from all other dust extracts ($p < 0.001$).

Figure 3.

(**A-B**). Muramic acid analysis of the various agricultural and domestic house dust extracts. (**A**) Muramic acid concentration normalized to the total amount of dust extracted in Tween (ng/mg). (**B**) Muramic acid concentration normalized to total volume in the 1% HBSS dust extract (ng/mL). Mean results $\pm$ SEM are presented (N=3). *Statistically different from all other dust extracts ($p<0.001$). #Statistically different from all other dust extracts ($p<0.05$).

Figure 4.

(**A-B**). Ergosterol analysis of the various agricultural and domestic house dust samples. (**A**) Ergosterol concentration normalized to the total amount of dust extracted in Tween (ng/mg). (**B**) Ergosterol concentration normalized to total volume in the 1% HBSS dust extract (ng/mL). Mean results $\pm$ SEM are presented (N=3). *Statistically different from all other dust extracts ($p < 0.001$). #Statistically different from all other dust extracts ($p < 0.05$)

Figure 5. (A-C). Secretion of TNF α (A), IL-6 (B), and CXCL8/IL-8 (C) in human peripheral blood monocytes stimulated with complete (untreated) and polymyxin B-treated (endotoxin-depleted) various agricultural and domestic house 1% HBSS dust extracts for 5 hours. *Significantly different from respective media control ($p < 0.05$). #Significantly different from matched endotoxin-depleted dust extract (polymyxin B treated vs. untreated dust extract; $p < 0.05$). N=3 separate experiments. Mean results are presented per 1×10^6 cells $\pm$ SD. ^aStatistically different from all other endotoxin-depleted dust extracts. ^bStatistically different from endotoxin-depleted domestic house dust extracts.

Figure 6.
(A-B). Secretion of IL-6 **(A)** and CXCL8/IL-8 **(B)** in human bronchial epithelial cells (BEAS-2B) stimulated with complete (untreated) and polymyxin B-treated (endotoxin-depleted) various agricultural and domestic house 1% HBSS dust extracts for 24 hours. *Significantly different from respective media control ($p < 0.05$). $N = 3$ separate experiments. Mean results $\pm$ SD are presented. ^aStatistically different as compared to grain elevator and domestic house dust extracts. ^bStatistically different as compared to all other endotoxin-depleted dust extracts. ^cStatistically different as compared to endotoxin-depleted grain elevator and domestic house dust extracts.

Table 1

Recombinant factor C (rFC) bioassay results of endotoxin measurements of the various dusts (EU/mg). Mean (standard deviation) for each dust type and extraction type (N=3)

	rFC (EU/mg)		
	Tween Extract	HBSS Extract	Polymyxin-B Treated HBSS Extract
Swine Facility	1338.7 (176.4)*	372.8 (30.5)*	3.1 (1.1)
Dairy Barn	433.5 (26.1)#	22.9 (4.0)	<LOD
Grain Elevator	1130.3 (635.2)*	295.2 (29.3)*	13.8 (2.3)
House, Dog	64.0 (19.0)	20.9 (2.1)	0.1 (0)
House, No Pets	45.9 (7.3)	12.5 (1.1)	0.1 (0)

* Statistically difference (p<0.05): Tween and HBSS Swine facility and grain elevator extracts significantly greater endotoxin as compared to other dusts, but not different from each other.

Statistical difference (p<0.05): Tween dairy barn extract significantly greater than domestic house dust extracts.