Published in final edited form as:

Tetrahedron Lett. 2010 May 5; 51(18): 2536–2538. doi:10.1016/j.tetlet.2010.03.009.

An Alkoxide Anion Triggered *tert*-Butyloxycarbonyl Group Migration. Mechanism and Application

Fengtian Xue[†] and Richard B. Silverman^{*}

Department of Chemistry, Department of Biochemistry, Molecular Biology, and Cell Biology, Center for Molecular Innovation and Drug Discovery, Chemistry of Life Processes Institute, Northwestern University, 2145 Sheridan Road, Evanston, Illinois 60208-3113, USA

Abstract

We report a fast N→O *tert*-butyloxycarbonyl (Boc) migration of the imide (3*R*,4*R*)-*tert*-butyl 3-(((6-(bis(*tert*-butoxycarbonyl)amino)-4-methylpyridin-2-yl)methyl)-4-hydroxypyrrolidine-1-carboxylate (**2**) via a base-generated alkoxide. The mechanism of the migration is intramolecular, involving an unusual nine-membered cyclic transition state.

Keywords

N to O migration; Boc group migration; intramolecular migration; crossover experiment

In the course of research on the development of chiral pyrrolidine-based inhibitors for neuronal nitric oxide synthase (nNOS), we evaluated various methods for the synthesis of compounds **1a** and **1b**, the key intermediates to prepare drug candidates for different kinds of neurodegenerative diseases.^{1–3} Given the poor efficiency of benzyl deprotection during the reported synthetic route to **1a**,² we attempted to use bis-Boc protection of the amino group on the pyridine ring (**1b**); these groups could be removed in one step during the late stage deprotection of the synthesis in excellent yields, providing a much more practical preparation of the final products on a multigram scale.⁴

© 2010 Elsevier Ltd. All rights reserved.

*Correspondence to: Professor Richard B. Silverman at the Department of Chemistry, Phone: 847-491-5653, Fax: 847-491-7713, Agman@chem.northwestern.edu.

[†]Present Address: ¹Department of Chemistry, University of Louisiana at Lafayette, P.O. Box 44370, Lafayette, Louisiana 70504, USA

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

To generate **1b**, allylation of alcohol **2** was attempted by treating a solution of **2** in DMF with NaH (2 equiv) at room temperature, followed by the addition of allyl bromide (2 equiv). The reaction was quenched with H₂O to form the product with 93% isolated yield (Scheme 1). To our surprise, mass spectrum and ¹H NMR data for this product did not match the anticipated product (**1b**). The isolated product has only an (M + H⁺) peak at 448, which is 100 less than the calculated molecular weight of **1b**, implying a possible loss of one Boc group from the desired product. This was further confirmed by the fact that there were two distinctive singlets at 1.46 and 1.52 ppm (each integrating to nine protons) in the ¹H NMR spectrum (in CDCl₃) of the product. The instability of the Boc protecting group under strong basic conditions has been documented.^{5–6} Interestingly however, one broad singlet in the ¹H NMR spectrum was found from 2.30 to 2.40 ppm, indicating the presence of a hydroxyl group in the product. Further NOSEY NMR data showed that the allyl group was connected through the nitrogen atom of the amino functionality to the pyridine ring.⁷ On the basis of these results, we assigned the product as (3*R*,4*R*)-*tert*-butyl 3-((6-(allyl(*tert*-butoxycarbonyl)amino)-4-methylpyridin-2-yl)methyl)-4-hydroxypyrrolidine-1-carboxylate (**3**).⁸ It was also noted that: 1) compound **2** showed significant stability in aqueous NaOH even at accelerated temperature,⁹ and 2) no O-allylation product was detected in the reaction process.

To elucidate the origin of *N*-allyl alcohol **3**, the reaction was repeated and monitored closely by thin layer chromatography (TLC) and LC/MS analysis. Time course studies clearly showed the disappearance of the starting material (**2**), and the buildup of a new compound with significantly less polarity when **2** was treated with NaH in DMF. After the addition of H₂O, the product (**3**), with similar polarity to that of **2**, was formed quickly (Scheme 1). Accordingly, we speculated that the basic environment generated during the quenching step catalyzed a hydrolysis reaction of the initial product, leading to the formation of alcohol **3**. To test this hypothesis, the same reaction was repeated and quenched with saturated aqueous NH₄Cl, to avoid the potential base-catalyzed hydrolysis step. As a result, compound **4** was isolated in a 96% yield (Scheme 2).¹⁰ This result implies that a carbonate derivative was an intermediate involved in the reaction course, which explains why there was no O-allylation product formed from the reaction.

On the basis of this collected evidence, we propose that deprotonation of **2** by treatment with NaH forms **5** at the beginning of the reaction (Scheme 3). Alkoxide **5** initiates the migration of one of the two Boc groups on the aminopyridine through a nine-membered ring transition state to generate amide anion (**6**), which reacts with allyl bromide to generate **4**. Several examples of anion triggered migration reactions have been described in literature.^{11–15} The carbonate linkage of **4** is unstable to the strong basic environment (e.g., aqueous NaOH, generated during the quench step), and is hydrolyzed quickly to give alcohol **3** as the only product. To prove the presence of **6**, the reaction was quenched with saturated aqueous NH₄Cl before the addition of allyl bromide. As seen in Scheme 4, compound **7** was isolated in quantitative yields and characterized.¹⁶

To further investigate the reaction mechanism, we carried out a crossover experiment using a mixture of compounds **2** and **8** as starting material. The mixture was treated with NaH, and after 5 min the reaction mixture was quenched with saturated aqueous NH₄Cl. The reaction did not give any crossover product (i.e., *O*-Boc-**8**) as shown in Scheme 5: compound **7** was isolated in almost quantitative yield, and **8** was fully recovered from the reaction mixture. These results clearly indicate that the Boc migration reaction takes place by an intramolecular pathway.

The unusual Boc-protecting group migration also provides a facile approach to selectively add an allyl or a benzyl group onto the amide N atom without hydroxyl group protection.

The previous method for this reaction suffered from competition of *O*-allylation or *O*-benzylation reactions, leading to unsatisfactory yields of products unless the hydroxyl group was protected (Scheme 6).² Using di-Boc-protected **2** as the starting material, however, the reaction (e.g., allylation or benzylation) produces the desired compound as the only product in excellent yields without interference of side reactions at the hydroxyl group.

In summary, we demonstrated a novel alkoxide anion triggered N→O Boc migration, involving an unusually large nine-membered ring transition state. The Boc migration process provides a new high yield method to perform selective amide allylation/benzylation over hydroxyl allylation/benzylation reactions.

Acknowledgments

The authors are grateful for financial support from the National Institutes of Health (GM049725).

References

1. For a comprehensive review of the chemistry and biology related to the present work, see: Silverman RB. Acc. Chem. Res 2009;42:439. [PubMed: 19154146]
2. Lawton GR, Ranaivo HR, Wing LK, Ji H, Xue F, Martasek P, Roman LJ, Watterson DM, Silverman RB. Bioorg. Med. Chem 2009;17:2371. [PubMed: 19268602]
3. Xue F, Fang J, Lewis WW, Martasek P, Roman LJ, Silverman RB. Bioorg. Med. Chem. Lett 2010;20:554. [PubMed: 19963381]
4. A synthetic route using **2b** as key intermediate for the preparation of multi-gram scale drug candidates is complete.
5. Foillard S, Rasmussen MO, Razkin J, Boturn D, Dumy P. J. Org. Chem 2008;73:983. [PubMed: 18173281]
6. Tom NJ, Simon WM, Frost HN, Ewing M. Tetrahedron Lett 2004;45:905.
7. Xue F, Silverman RB. unpublished result.
8. Synthesis and characterization of (3*R*,4*R*)-*tert*-butyl 3-(allyloxy)-4-((6-(bis(*tert*-butoxycarbonyl)amino)-4-methylpyridin-2-yl)methyl)pyrrolidine-1-carboxylate (**3**). To a solution of **2** (1.15 g, 2.0 mmol) in DMF (15 mL) was added NaH (60% in mineral oil, 160 mg, 4.0 mmol) at room temperature. After five min, allyl bromide (480 μ L, 4.0 mmol) was added. The reaction mixture was allowed to stir at room temperature for 30 min, then quenched with H₂O (5.0 mL). The solvents were removed by rotary evaporation at 50 °C, and the resulting oil was purified by flash column chromatography (silica gel, EtOAc/hexanes, 1:1) to yield **3** as a colorless oil (1.05 g, 93%): ¹H NMR (500 MHz, CDCl₃) δ 1.44-1.45 (m, 9H), 1.50 (m, 9H), 2.33 (s, 3H), 2.33-2.38 (br s, 1H), 2.75-2.85 (dt, *J* = 5.0, 13.5 Hz, 1H), 2.85-2.93 (m, 1H), 3.16-3.22 (dt, *J* = 5.0, 10.5 Hz, 1H), 3.40-3.55 (m, 2H), 11 3.55-3.59 (t, *J* = 9.0 Hz, 0.5H), 3.62-3.66 (t, *J* = 10.0 Hz, 1H), 4.05 (s, 1H), 4.20 (s, 0.5H), 4.38 (s, 0.5H), 4.43 (s, 2H), 5.07-5.12 (m, 2H), 5.85-6.00 (ddd, *J* = 5.0, 10.0, 15.5 Hz, 1H), 6.70-6.80 (m, 1H), 7.34 (s, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 21.1, 21.2, 28.2, 28.51, 28.54, 35.3, 35.4, 44.7, 45.3, 49.1, 49.48, 49.52, 53.3, 53.6, 60.4, 70.2, 71.0, 79.1, 81.4, 115.7, 118.2, 120.2, 120.3, 134.35, 134.37, 149.5, 149.6, 154.0, 154.1, 154.4, 157.4, 157.5; LCTOF (M +H⁺) calcd for C₂₄H₃₈N₃O₅ 448.2811, found 448.2801.
9. Compound **2** was allowed to stir in a mixture of 1 N NaOH and MeOH (1:1) at room temperature. After 30 min, only 15% of the hydrolyzed product (3*R*,4*R*)-*tert*-butyl 3-((6-(*tert*-butoxycarbonylamino)-4-methylpyridin-2-yl)methyl)-4-hydroxypyrrolidine-1-carboxylate was generated: ¹H NMR (500 MHz, CDCl₃) δ 1.45-1.46 (m, 9H), 1.52 (s, 9H), 2.33 (s, 3H), 2.33-2.35 (br s, 1H), 2.70-2.80 (dt, *J* = 6.0, 9.5 Hz, 1H), 2.80-2.85 (m, 1H), 3.15-3.20 (dt, *J* = 5.5, 10.5 Hz, 1H), 3.40-3.55 (m, 2H), 3.58-3.62 (dd, *J* = 8.5, 10.0 Hz, 0.5H), 3.63-3.67 (dd, *J* = 8.5, 10.0 Hz, 0.5H), 4.09 (s, 1H), 4.75 (br s, 1H), 6.68-6.69 (d, *J* = 7.0 Hz, 1H), 7.19 (s, 1H), 7.68 (s, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 21.1, 21.4, 28.2, 28.52, 28.54, 35.16, 35.24, 44.5, 45.2, 49.0, 49.5, 53.2, 53.6, 60.4, 70.3, 71.3, 79.2, 81.2, 110.6, 119.1, 119.2, 151.0, 151.1, 151.2, 152.2, 154.4, 157.4, 157.5; LC-TOF (M+H⁺) calcd for C₂₁H₃₄N₃O₅ 408.2498, found 408.2499.

10. Synthesis and characterization of (3*R*,4*R*)-*tert*-butyl 3-(allyloxy)-4-((6-(bis(*tert*-butoxycarbonyl)amino)-4-methylpyridin-2-yl)methyl)pyrrolidine-1-carboxylate (**4**). To a solution of **2** (505 mg, 1.0 mmol) in DMF (15 mL) was added NaH (60% in mineral oil, 80 mg, 2.0 mmol) at room temperature. After 5 min, allyl bromide (240 μ L, 2.0 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature for 30 min then quenched with ice cold saturated NH₄Cl (5.0 mL). The solvent was removed *in vacuo*, and the resulting residue was purified by flash column chromatography (silica gel, EtOAc/hexanes, 1:4) to yield **4** as a colorless oil (525 mg, 96%): ¹H NMR (500 MHz, CDCl₃) δ 1.42 (s, 9H), 1.47-1.49 (m, 18H), 2.25-2.27 (m, 3H), 2.70-2.85 (m, 2H), 2.89-2.93 (dd, *J* = 5.5, 13.0 Hz, 1H), 3.14-3.21 (dd, *J* = 11.0, 21.5 Hz, 1H), 3.46-3.59 (m, 3H), 4.40-4.60 (m, 2H), 5.03-5.12 (m, 3H), 5.86-5.92 (ddd, *J* = 5.5, 12.0, 17.5 Hz, 1H), 6.64 (s, 1H), 7.32 (s, 0.5H), 7.36 (s, 0.5H); ¹³C NMR (125 MHz, CDCl₃) δ 21.11, 21.13, 27.8, 28.3, 28.47, 28.50, 34.7, 41.8, 42.3, 48.8, 49.06, 49.10, 49.2, 52.4, 52.8, 76.7, 76.8, 79.4, 80.9, 81.0, 82.4, 82.5, 115.8, 117.3, 117.4, 119.8, 134.91, 134.93, 148.5, 148.6, 153.1, 153.2, 153.9, 154.0, 154.1, 154.2, 154.4, 156.8; LC-TOF (M+H⁺) calcd for C₂₉H₄₆N₃O₇ 548.3336, found 548.3339.
11. Liu Y, Ding Q, Wu X. J. Org. Chem 2008;73:6025. [PubMed: 18610981]
12. Gérard S, Marchand-Brynaert J. Tetrahedron Lett 2003;44:6339.
13. Hara O, Ito M, Hamada Y. Tetrahedron Lett 1998;39:5537.
14. Valentekovich RJ, Schreiber SL. J. Am. Chem. Soc 1995;117:9069.
15. Bunch L, Norrby P-O, Frydenvang K, Krogsgaard-Larsen P, Madsen U. Org. Lett 2001;3:433. [PubMed: 11428032]
16. Synthesis and characterization of (3*R*,4*R*)-*tert*-butyl 3-((6-(*tert*-butoxycarbonylamino)-4-methylpyridin-2-yl)methyl)-4-(*tert*-butoxycarbonyloxy)pyrrolidine-1-carboxylate (**7**). To a solution of **2** (250 mg, 0.5 mmol) in DMF (10 mL) was added NaH (60% in mineral oil, 40 mg, 1.0 mmol) at room temperature. After 5 min, the reaction mixture was quenched with saturated NH₄Cl (5.0 mL). The solvent was removed *in vacuo*, and the resulting residue was purified by flash column chromatography (silica gel, EtOAc/hexanes, 1:4) to yield **7** as a colorless oil (505 mg, 99%): ¹H NMR (500 MHz, CDCl₃) δ 1.43-1.44 (m, 9H), 1.48-1.50 (m, 18H), 2.27-2.30 (m, 3H), 2.60-2.75 (m, 2H), 2.85-2.92 (m, 1H), 3.15-3.22 (dt, *J* = 4.0, 11.0 Hz, 1H), 3.45-3.65 (m, 3H), 5.10 (br s, 1H), 6.60 (s, 1H), 7.13-7.17 (d, *J* = 15.5 Hz, 1H), 7.58-7.60 (d, *J* = 9.0 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 21.28, 21.30, 24.7, 27.79, 27.82, 28.3, 28.47, 28.50, 34.4, 34.5, 36.6, 42.0, 42.5, 48.9, 49.2, 52.5, 52.8, 76.2, 79.4, 79.5, 80.75, 80.80, 82.5, 110.18, 110.23, 115.8, 118.9, 150.0, 151.4, 151.5, 152.37, 152.41, 153.2, 153.3, 154.2, 154.4, 157.2, 157.3; LC-TOF (M+H⁺) calcd for C₂₆H₄₂N₃O₅ 508.3023, found 508.3037.

Scheme 1.
Formation of **3** from **2**

Scheme 2.
Formation of **4** from **2**

Scheme 3.
Proposed mechanism for the formation of **3**

Scheme 5.
Crossover experiment for Boc migration

Scheme 6.
Alkylation of the aminopyridine without hydroxyl protection