Published in final edited form as:

J Nat Prod. 2010 April 23; 73(4): 679–682. doi:10.1021/np1000297.

Structure Elucidation at the Nanomole-Scale. 3. Phorbasides G-I from *Phorbas* sp.

Doralyn S. Dalisay^{†,‡} and Tadeusz F. Molinski^{*,†,§}

Department of Chemistry and Biochemistry, and Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California, San Diego, 9500 Gilman Drive MC0358, La Jolla, California 92093-0358

Abstract

Three new phorbasides (G-I), chlorocyclopropyl-ene-yne macrolide glycosides, were isolated from the sponge *Phorbas* sp. in yields of 7–9.5 μ g and fully characterized by MS, CD and microcryoprobe NMR. The structures of the new compounds differ only in the nature of the sugar residues. The absolute configurations of the new compounds were correlated by ROESY and CD with the parent compounds phorbasides A and B.

The use of cryo-microprobe NMR spectroscopy, integrated with other ‘nanomole-scale’ techniques (CD, FTIR, FTMS) for natural products discovery has expanded our view to extremely minute amounts of compound.¹ Nanomole-scale methods exploit technical improvements in signal-to-noise of spectroscopic methods, but more importantly they open new vistas for discovery of new compounds from rare organisms and uncovering chemical complexity and diversity within single specimens.² The marine sponge *Phorbas* sp. collected off the Muiron Island in Western Australia, has been shown to contain highly cytotoxic macrolides including phorboxazoles A (**1**) and B (**2**),³ and phorbasides A (**3**) and B (**4**).⁴ Recently, the structures of very minor natural products from this one specimen of *Phorbas* were elucidated to reveal appreciably broader chemical diversity within the sponge than encountered in earlier studies. The new compounds, which were fully characterized by MS, CD and NMR (COSY, ROESY, HSQC, HMBC), include phorbaside F (**5**, 8 μ g, 13 nmol),⁵ hemi-phorboxazole A (16.5 μ g)⁶ and muironolide A (**6**, 90 μ g).⁷ The latter represents the first member of an entirely new carbon skeleton based on a hexahydro-1*H*-isindolin-1-one. We now report additional phorbasides G-I (**7–9**) in total yields of 7–9.5 μ g. The structures were solved by integrated application of microcryoprobe NMR, including quantitative analysis, MS and CD analysis.

Results and Discussion

Reexamination of the CCl₄-soluble fraction of sponge *Phorbas* sp., which previously afforded **3** and **4**,⁴ revealed very minor components that were further purified by reversed phase HPLC to give *total yields* of pure compounds in amounts ranging from 7 to 9.5 μ g (quantitation by comparative NMR ‘¹³C satellite’ integration)⁵: phorbaside F⁵ (**5**), 8 μ g (3.8

*To whom correspondence should be addressed: Tel: +1 (858) 534-7115. Fax: +1 (858) 822-0386. tmolinski@ucsd.edu.

[†]UC San Diego, Department of Chemistry and Biochemistry

[‡]Present address: Department of Chemistry, University of British Columbia, Vancouver, BC, Canada V6T 1Z1. EWR

[§]UC San Diego, Skaggs School of Pharmacy and Pharmaceutical Sciences

Supporting Information Available. ¹H NMR, 2D NMR spectra of **7**, **8** and **9**. These materials are available free of charge via the Internet at <http://pubs.acs.org>.

$\times 10^{-6} \%$), phorbaside G (**7**), $9.5 \mu\text{g}$ ($4.0 \times 10^{-6} \%$), phorbaside H (**8**), $7.0 \mu\text{g}$ ($2.9 \times 10^{-6} \%$) and phorbaside I (**9**), $9.4 \mu\text{g}$ ($3.9 \times 10^{-6} \%$ dry wt).

The molecular formula of phorbaside G (**7**), $\text{C}_{41}\text{H}_{61}\text{ClO}_{14}$, derived from HRESITOFMS (m/z $[\text{M}+\text{Na}]^+$, 835.2631), is two H atoms less than that of phorbaside B (**4**). The ^1H , COSY, NOESY, HSQC and HMBC experiments revealed the same macrolide carbon framework present in **4**, but differing in the disaccharide group attached at C-5 (Tables 1 and 2). The second sugar unit of **7** is oxidized with respect to **4** and has a keto group at C-4'' (δ 208.0). In addition, the ^{13}C NMR chemical shifts of C-3' (δ 80.5) and C-4' (δ 70.4) in the first sugar unit appeared downfield of those in **4** by ~ 10 ppm. The configuration of the sugar unit was established by ROESY as a 6-deoxy-2,3-*C,O*-dimethyl- α -L-*manno*-pyranoside (*O*-methyl-L-*evalose*). Careful analysis of the HMBC spectrum showed that the anomeric proton signal H-1'' (δ 5.82, s) was correlated with the quaternary C-3', thus indicating a 1'' $\rightarrow$ 3' disaccharide linkage. The downfield ^{13}C NMR chemical shift of C-3' at δ 80.5 is characteristic of a 1'' $\rightarrow$ 3' linkage when the tertiary OH of *C*-methyl sugars is glycosylated and matches ^{13}C NMR chemical shifts of L-*evalose* residues with similar glycosylation patterns found in glycolipid antigens from *Mycobacterium avium*.⁸

Phorbaside H (**8**) has a molecular formula of $\text{C}_{41}\text{H}_{63}\text{ClO}_{14}$ as established by HRESITOFMS (m/z 837.3780 $[\text{M}+\text{Na}]^+$) and is isomeric with phorbaside B (**4**). Significant differences in NMR chemical shifts were again seen in the sugar units as revealed by the ^1H , COSY, ROESY, HSQC and HMBC experiments (Tables 1 and 2). The configuration of the first sugar unit in **8** is 6-deoxy-2,3-*C,O*-dimethyl- α -L-*manno*-pyranoside, the same as in **4**, while the second sugar unit is a 6-deoxy-2,3-*C,O*-dimethyl- α -L-*talo*-pyranoside on the basis of interpretation of a ROESY experiment (600 MHz, t_m = 400 ms). Key ROESY correlations were observed from the anomeric signal H-1'' to H-2'', H-2'' to H-7'' and H-4'' to H-5''. Furthermore, the ^1H NMR signal due to the methyl group at C-7'' (δ 1.38) showed ROESY correlations to H-2'' and H-5'' (Figure 1). The C4'' configuration in **8**, therefore, is epimeric to C-4 in *evalose* residues that occur in **3** and **4**, and is the first example of a *talo*-hexose in the callipeltoside-phorbaside family, although the 4-amino-2,3-*O,C*-dimethyl-4,6-deoxypyranosides in callipeltosides A (**10**) and B (**11**),⁹ are also *talo*. The linkage of the sugar units was established from cross peak assignments of HMBC and HSQC data. As with **7**, a 1'' $\rightarrow$ 3' disaccharide linkage was identified by characteristic changes in ^{13}C chemical shifts and heteronuclear correlation; the signal of the anomeric proton H-1'' (δ 5.76, s) in the second sugar of **8** shared an HMBC cross peak with the quaternary C-3' (δ 80.1) of the first sugar, which was also shifted downfield by about 10 ppm compared to **3**.

Phorbaside I (**9**) has a molecular formula of $\text{C}_{34}\text{H}_{50}\text{ClNNaO}_{10}$, as determined by HRESITOFMS (m/z $[\text{M}+\text{Na}]^+$, 690.3020). Analysis of ^1H , COSY, HSQC and HMBC experiments indicated that **9** has the same aglycone as phorbaside A (**3**), and deductive reasoning which accounted for the differences in the NMR signals and the presence of N in the formula, led to an assignment of amino sugar glycone (Tables 1 and 2). The ^1H NMR spectrum of **9** showed several signals that were doubled in the ratio of 2:1, characteristic of interconverting rotamers of a formamide NH-CH=O group. The associated ^1H NMR signals were split into two sets; a formyl signal H-8' (δ 8.37, s, major and 7.93, d, J = 12 Hz, minor) and an NH signal (6.32, d, J = 10.6 Hz, major and 6.14, t, J = 11.4 Hz, minor). Signal doubling was also observed for H-7' (δ 1.43, s, major and 1.45, s, minor) and H-2'-OMe (δ 3.47, s, major and 3.46, s, minor). A *talo* configuration for the sugar was supported by ROESY (600 MHz, t_m = 400 ms) data, cross peaks between *syn*-facial hydrogens across the pyranose ring; H-1' to H-2', H2' to H-7', H'-4 to H-6' and H-7'. These lines of evidence pointed to a *N*-formyl 4-amino-4,6-dideoxy-2,3-*C,O*-dimethyl- α -L-*talo*-pyranoside residue, first seen in callipeltoside B (**11**), but assigned as β -L.⁹ Thus the configuration at each anomeric carbon in **9** is α -L, the same as in **3** and **4** derived from similar evidence.¹⁰ Note,

the disaccharides **7** and **8** have 1''→3' disaccharide linkages instead of the 1''→4' assigned to phorbaside C and the 1''→2' linkage for phorbaside E.⁴

The anomeric configuration of evalose residues and related sugars in the callipeltose-phorbaside family of glycosides has been a matter of some equivocation. The ¹H NMR signals for both H-1 and H-2 in evalose residues are singlets that lack vicinal couplings that would otherwise resolve axial or equatorial dispositions from simple Karplus relationships. D'Auria and coworkers assigned a β-L configuration to callipeltosides B and C,⁹ however, we assigned α-L to the anomeric carbons C-1' and C-1'' in **3**, **4** and other phorbasides⁴ based on consistent observation of a weak ROESY cross peak between each acetal proton (H-1' or H-1'') and the adjacent *syn*-C-2'-OMe group. Similar ROE's were observed for phorbasides G (**7**) and phorbaside H (**8**, see Figure 1). Here, with the enhanced mass-sensitivity of the microcryoprobe NMR spectroscopy, we were able to provide independent evidence of the axial configuration of the C-1' and C-1'' OR groups in **7** and in **8** the form of ¹J_{CH}'s of the anomeric CH signals measured from ¹H coupled HSQC spectra (Table 2). The value of ¹J_{CH} of the anomeric CH in aldopyranoses is sensitive to the axial or equatorial disposition of the C-1 alkoxy substituent and fall into the ranges of 169–171 Hz and 158–162 Hz, respectively.¹¹ The measured values for C-1' (*J* = 167 Hz) and C1'' (*J* = 178 Hz) for **7** and **8** are essentially the same and fall into the higher range, consistent with axial alkoxy substituents, O-C-5 and O-C-3', respectively.

The absolute configurations of all the new compounds were readily assigned by comparisons with **3** and **4** after quantitation,⁵ quantitative recovery from the 1.7 mm NMR tubes, and measurement of their CD spectra (Figure 2). The Cotton effects observed at λ 231 and 241 nm in **7–9** arise from π-π* transitions of the hyperconjugated cyclopropyl-en-yne chromophore.⁴ These Cotton effects are identical in sign and magnitude compared to those of **3** and **4** that we had earlier assigned⁴ by correlation with synthetic models of known configuration, prepared from diastereomers of *O*-menthyl 2,2-dichlorocyclopropanecarboxylate.¹² Independently, the sugar configurations of **3** and **4** were assigned as L- by methanolysis (HCl, MeOH), conversion of the resultant *O*-methylglycosides to bi-chromophoric derivatives, and CD comparison with synthetic standards prepared in several steps from L-rhamnose.¹³ Since all phorbasides A-I were obtained from the same single specimen of *Phorbas* sp., it is highly likely that the sugars in **7–9** are also L.

Unlike phorbasides A-E, where differential cytotoxicities against the colon tumor cell line HCT116 were reported,^{4b} the amounts of **6–9** available were insufficient to assess cytotoxic activity. Nanomole-scale natural products methods provide powerful tools for structure identification,^{1,2} however fuller investigation of their properties is a challenge that is presently met only by synthesis and well-designed SAR studies. This was nicely illustrated recently by SAR studies of the antifungal hemiphorboxazole A,⁶ a compound that was obtained in a yield of only 16.5 μg, but subsequently prepared in milligram quantities by directed synthesis.¹⁴

In conclusion, the structures of the new phorbasides G-I (**7–9**) were elucidated from vanishingly small samples (7–9.5 μg) using CD, MS and NMR data obtained from a 600 MHz NMR spectrometer equipped with a 1.7 mm microcryoprobe. The outstanding quality of NMR data obtained from this instrument allowed full assignments and revealed differences in the glycone units. Phorbaside G (**7**) contained a novel C-4 oxidized pyranoside, and **8** and **9** were variations on the C-methyl pyranoside sugars first reported by Zampella and coworkers in callipeltosides A (**10**), B (**11**) and C (**12**).

Experimental Section

General Experimental Procedures

UV-Vis spectra were recorded on a Jeol V-630 dual beam spectrometer in 0.2 or 1 cm quartz cells CD spectra were recorded on a Jasco 810 spectropolarimeter in 0.2 cm quartz cells at 23 °C unless otherwise stated. ^1H and ^{13}C NMR spectra were recorded in CDCl_3 on a 600 MHz spectrometer equipped with a Bruker 1.7 mm $\{^{13}\text{C}\}^1\text{H}$ CPTCI microcryoprobe. NMR spectra were measured in CDCl_3 and referenced to residual solvent signals (^1H , δ 7.26 ppm; ^{13}C , δ 77.16 ppm). HRMS measurements were measured under ESI conditions with an Agilent 6000 series TOF mass spectrometer, or a ThermoFinnigan Orbitrap mass spectrometer. Semi-preparative HPLC was carried out with a Varian high capacity dual-pump system coupled to a Rainin UV-1 high-dynamic UV detector under specified conditions. All solvents for HPLC purification were redistilled from commercial 'HPLC grade' solvent.

Extraction and Isolation

The sponge *Phorbas* sp. (sample ID: 93-054) was collected from Muiron Island, Western Australia in 1993. A taxonomic description of the specimen has appeared elsewhere.¹⁵ The sample was immediately frozen and stored at -20°C until extraction (~2 months). The CCl_4 -soluble fraction (350 mg) of the MeOH extract of *Phorbas* sp.³ was separated by flash chromatography (silica cartridge, Analogix RS-12, 12 g, 2 cm $\times$ 7.5 cm) using mixtures of *n*-hexane and EtOAc in increasing polarity (0–100%) to yield seven fractions. Fraction 5 (93-054-B1–5, 12.5 mg) was further purified twice by reversed-phase HPLC (Phenylhexyl column, 250 $\times$ 10 mm, 90:10 MeOH:H₂O, 2 mL.min⁻¹, then Phenylhexyl column, 250 $\times$ 4.6 mm, 60:40 CH₃CN:H₂O, 2 mL.min⁻¹) to yield pure phorbaside I (**9**), t_R = 9 min, 9.4 μg (3.9×10^{-6} % dry wt), phorbaside F⁵ (**5**), t_R = 12 min, 8 μg (3.8×10^{-6} % dry wt), phorbaside G (**7**), t_R = 18 min, 9.5 μg (4.0×10^{-6} % dry wt) and phorbaside H (**8**) t_R = 19 min, 7.0 μg (2.9×10^{-6} % dry wt).

Phorbaside G (7)—colorless film; UV (MeOH) λ_{max} (log ϵ) 233 nm (4.07), 241 (4.07); CD (MeOH) λ_{max} ($\Delta\epsilon$) 231 nm (+11.07), 241 (+9.60); ^1H and ^{13}C NMR (see Tables 1 and 2); HRESITOFMS m/z 835.3631 [$\text{M}+\text{Na}$]⁺ (calcd for C₄₁H₆₁ClNaO₁₄, 835.3642).

Phorbaside H (8)—colorless film; UV (MeOH) λ_{max} (log ϵ) 233 nm (4.07), 241 (4.07); CD (MeOH) λ_{max} ($\Delta\epsilon$) 231 nm (+11.6), 241 (+10.3); ^1H and ^{13}C NMR (see Tables 1 and 2); HRESITOFMS m/z 837.3780 [$\text{M}+\text{Na}$]⁺ (calcd for C₄₁H₆₃ClNaO₁₄, 837.3799).

Phorbaside I (9)—colorless film; UV (MeOH) λ_{max} (log ϵ) 233 nm (4.07), 241 (4.07); CD (MeOH) λ_{max} ($\Delta\epsilon$) 231 nm (+11.4), 241 (+10.5); ^1H and ^{13}C NMR (see Table 1 and 2); HRESITOFMS m/z 690.3020 [$\text{M}+\text{Na}$]⁺ (calcd for C₃₄H₅₀ClNNaO₁₀, 690.3021).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank B. Morinaka and E. Rogers for helpful discussions and A. Jansma for assistance with NMR experiments, and Y. Su (UCSD) for HRMS measurements. This research was supported by grants from the NIH (RO1 AI 039987, CA122256).

References and Notes

1. (a) Molinski TF. *Curr Opin Drug Discov Devel* 2009;12:197–206. (b) Schroeder FC, Gronquist M. *Angew Chem Intl Ed* 2006;45:7122–7131.
2. Molinski TF. *Nat Prod Rep*. 2010 in press.
3. (a) Searle PA, Molinski TF. *J Am Chem Soc* 1995;117:8126. (b) Searle PA, Molinski TF, Brzezinski LJ, Leahy JW. *J Am Chem Soc* 1996;118:9422–9423. (c) Molinski TF. *Tetrahedron Lett* 1996;37:7879–7880.
4. (a) Skepper CK, MacMillan JB, Zhou GX, Masuno MN, Molinski TF. *J Am Chem Soc* 2007;129:4150–1. [PubMed: 17373800] (b) MacMillan JB, Xiong-Zhou G, Skepper CK, Molinski TF. *J Org Chem* 2008;73:3699–706. [PubMed: 18412387]
5. Dalisay DS, Molinski TF. *J Nat Prod* 2009;72:739–744. [PubMed: 19399996]
6. Dalisay DS, Molinski TF. *Org Lett* 2009;11:1967–1970. [PubMed: 19341267]
7. Dalisay DS, Morinaka BI, Skepper CK, Molinski TF. *J Am Chem Soc* 2009;131:7552–7553. [PubMed: 19453148]
8. Chatterjee D, Bozic C, Aspinall GO, Brennan PJ. *J Biol Chem* 1988;263:4092–7. [PubMed: 3346237]
9. Zampella A, D'Auria MV, Minale L, Debitus C. *Tetrahedron* 1997;53:3243–3248.
10. A survey of the structures of natural products with *O*-evalosyl sugar residues (Ref. ⁸ and references cited within) reveals a preference for the α -L linkage that happens to be the more thermodynamically stable isomer (anomeric effect. For an X-ray structure of methyl *O*-methyl evalose, see Giuliano RM, Kasperowicz S, Boyko WJ, Rheingold AL. *Carbohydr Res* 1989;185:61–67. [PubMed: 2713872]). Contrary to this, the structures of callipeltosides B and C were assigned (Ref. 9) or depicted (Ref. 13) with the inverted β -L anomeric configuration.
11. Pretsch, E.; Bühlmann, P.; Badertscher, M. *Structure Determination of Organic Compounds. Tables of Spectra Data*. 4. Springer; Heidelberg: 2009. p. 149
12. Masuno MN, Young DM, Hoepker AC, Skepper CK, Molinski TF. *J Org Chem* 2005;70:4162–4165. [PubMed: 15876110]
13. A total synthesis of callipeltoside C (**12**) was reported recently (Carpenter J, Northrup AB, Chung D, Wiener JJM, Kim S-G, MacMillan DWC. *Angew Chem Intl Ed* 2008;47:3568–3572.), with a tentative assignment of the anomeric carbon as β -L, with the caveat, “At the present time we cannot state definitively the stereochemistry of the C-1' anomeric position of callipeltoside C”. Although the title of this paper refers to a “revision” of configuration of callipeltoside C, the configurations of sugars in callipeltosides B (**11**) and C (**12**) in the original isolation paper were, in fact, arbitrarily assigned to the D- series (Ref. 9).
14. Smith AB III, Liu Z, Hogan A-ML, Dalisay DS, Molinski TF. *Org Lett* 2009;11:3766–3769. [PubMed: 19637861]
15. See Supporting Information for Reference 3a.

Figure 1. ROESY correlations in the sugar residues of **8**. (a) 6-deoxy-2,3-C,O-dimethyl- α -L-manno-pyranoside (first sugar unit) (b) 6-deoxy-2,3-C,O-dimethyl- α -L-talo-pyranoside (second sugar unit). Correlations marked with blue arrows are key to assignment of the anomeric configuration (see text).

Figure 2. Circular dichroism (CD) spectra of phorbosides G-I (MeOH, 23 °C). (a) **7** ($c = 1.0 \times 10^{-4}$ M). (b) **8** ($c = 7.8 \times 10^{-5}$ M). (c) **9** ($c = 1.78 \times 10^{-4}$ M).

Table 1NMR data of phorbasides G-I, **7–9** (600 MHz, CDCl₃)

	7	8	9^a
#	^δ H, mult. (J in Hz)	^δ H, mult. (J in Hz)	^δ H, mult. (J in Hz)
1			
2	2.52, q (7.3)	2.52, q (7.3)	2.51, m
3			
4	1.10, m 2.31, dd (12.3, 5.2)	1.10, m 2.29, dd (12.0, 4.7)	1.15, m 2.31, dd (12.0, 4.5)
5	3.63, td (10.6, 4.4)	3.61, td (10.5, 4.7)	3.64, td (10.6, 4.5)
6	1.46, m	1.46, m	1.45, m
7	3.66, dd (10.6, 2.6)	3.66, dd (10.5, 2.4)	3.67, m
8	2.18, m	2.19, m	2.18, m
9	3.77, dd (9.5, 2.6)	3.77, dd (9.5, 2.4)	3.78, dd (9.5, 2.2)
10	5.28, d (9.5)	5.28, d (9.5)	5.28, d (9.5)
11			
12	2.25, d (7.3)	2.25, d (7.3)	2.25, d (7.0)
13	5.72, dd (14.0, 6.5)	5.72, dd (16.0, 6.2)	5.72, dd (16.0, 6.2)
14	6.05, dd (15.8, 6.5)	6.05, dd (16.0, 6.2)	6.05, dd (16.0, 6.2)
15	5.66, d (15.8)	5.66, d (16.0)	5.66, d (16.0)
18	1.78, m	1.78, m	1.77, m
19	3.17, ddd (9.0, 6.6, 3.0)	3.17, ddd (9.0, 6.6, 3.0)	3.17, ddd (9.0, 6.6, 3.0)
20	1.29, m	1.27, m	1.27, m
21	1.73, s	1.73, s	1.73, s
22	0.98, d (6.7)	0.97, d (7.2)	0.98, d (7.0)
23	0.92, d (6.7)	0.91, d (6.4)	0.92, d (6.4)
24	1.13, d (7.0)	1.13, d (7.2)	1.13, d (7.2)
OMe	3.23, s	3.23, s	3.23, s
OH	4.79, s	4.78, s	
1'	4.88, s	4.85, s	4.96, s
2'	3.26, s	3.18, s	2.97, s
3'			
4'	3.94, d (5.4)	3.94, dd (9.4, 3.2)	3.97, d (10.5)
5'	3.76, m	3.75, dd (9.4, 6.5)	4.11, q
6'	1.30, d (6.6)	1.31, d (6.5)	1.11, d (6.4)
7'	1.45, s	1.38, s	1.43, s [1.45, s ^b]
8'			8.37, s [(7.93, d (12.0)) ^b]
3'-OH			
2'-OMe	3.43, s	3.41, s	3.47, s [3.46, s ^b]
1''	5.82, s	5.76, s	
2''	3.44, s	3.00, s	
3''			

	7	8	9 ^a
#	δH , mult. (<i>J</i> in Hz)	δH , mult. (<i>J</i> in Hz)	δH , mult. (<i>J</i> in Hz)
4''		3.14, d (12.0)	
5''	4.70, q (6.8)	4.22, q (6.5)	
6''	1.31, d (6.0)	1.27, d (6.2)	
7''	1.63, s	1.38, s	
2''-OMe	3.49, s	3.48, s	
-NH			6.32, d (10.6) ^a [6.14, t, (11.4) ^b]

^a Mixture of rotamers (2:1), some signals doubled.

^b minor rotamer.

Table 2¹³C NMR data^a of phorbasides G-I, **7–9** (600 MHz, CDCl₃)

	7	8	9
#	δC, mult.	δC, mult.	δC mult.
1	176.1, C	176.2, C	176.6
2	47.0, CH	47.4, CH	47.0, CH
3	97.4, C	97.4, C	97.0, C
4	46.3, CH ₂	46.9, CH ₂	46.3, CH ₂
5	79.3, CH	79.1, CH	80.0, CH
6	38.7, CH	38.8, CH	38.5, CH
7	74.9, CH	74.8, CH	74.7, CH
8	37.0, CH	37.1, CH	36.7, CH
9	79.5, CH	79.8, CH	79.5, CH
10	127.9, CH	127.8, CH	127.5, CH
11	133.0, C	132.4, C	132.6, C
12	46.0, CH ₂	46.9, CH ₂	46.4, CH ₂
13	70.5, CH	70.8, CH	71.0, CH
14	139.7, CH	139.9, CH	140.0, CH
15	111.2, CH	111.3, CH	111.2, CH
16	<i>b</i>	90.6, C	90.5, C
17	<i>b</i>	<i>b</i>	<i>b</i>
18	<i>b</i>	<i>b</i>	<i>b</i>
19	<i>b</i>	35.0, CH	<i>b</i>
20	18.0, CH ₂	19.6, CH ₂	<i>b</i>
21	16.8, CH	16.7, CH	16.5, CH
22	6.5, CH ₃	6.7, CH ₃	6.2, CH ₃
23	12.4, CH ₃ ^c	12.8, CH ₃	12.4, CH ₃
24	12.4, CH ₃ ^c	12.6, CH ₃	12.6, CH ₃
OMe	55.4	55.3	55.1
1'	99.0, CH ^d	99.5, CH ^d	98.6, CH
2'	86.4, CH	87.0, CH	83.0, CH
3'	80.5, C	80.1, C	67.8, C
4'	70.4, CH	69.8, CH	55.4, CH
5'	67.0, CH	67.0, CH	64.4, CH
6'	18.2, CH ₃	18.3, CH ₃	17.3, CH
7'	19.4, CH ₃	19.5, CH ₃	23.5 [38.5] ^c CH ₃
8'			149.1, C
2'-OMe	58.8	59.3	55.3 [59.2] ^c
1''	92.4, CH ^e	92.9, CH ^e	
2''	89.3, CH	84.6, CH	

	7	8	9
#	δ^{C} , mult.	δ^{C} , mult.	δ^{C} mult.
3"	76.3, C	68.0, C	
4"	208.0, C	76.5, CH	
5"	68.2, CH	65.8, CH	
6"	14.7, CH ₃	17.3, CH ₃	
7"	23.3, CH ₃	22.4, CH ₃	
2"-OMe	59.2	59.4	

^a δ 's measured by indirect detection, HSQC and HMBC ($^1J_{\text{CH}} = 8$ Hz).

^b not detected.

^c minor rotamer.

^d $^1J_{\text{CH}} = 167$ Hz.

^e $^1J_{\text{CH}} = 178$ Hz.