Published in final edited form as:

Neuroimage. 2009 October 15; 48(1): 191–199. doi:10.1016/j.neuroimage.2009.06.028.

Perceptual load interacts with involuntary attention at early processing stages: event-related potential studies

Shimin Fu¹, Yuxia Huang², Yuejia Luo², Yan Wang³, John Fedota¹, Pamela M. Greenwood¹, and Raja Parasuraman¹

¹ARCH Laboratory, Department of Psychology, George Mason University, Fairfax, VA 22030, USA

²National Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, Beijing, 100875, CHINA

³Key Laboratory of Mental Health, Institute of Psychology, Chinese Academy of Sciences, Beijing, 100101, CHINA

Abstract

Perceptual load is known to influence the locus of attentional selection in the brain but through an unknown underlying mechanism. We used event-related potentials (ERPs) to investigate how perceptual load interacts with cue-driven involuntary attention. Perceptual load was manipulated in a line orientation discrimination task in which target location was cued involuntarily by means of peripheral cues. Attentional modulation was observed for P1m (the posterior midline P1 component with peak latency between 108 and 140 ms) with invalid trials eliciting larger P1m than valid trials. This attentional effect on P1m increased as a function of perceptual load, suggesting an early temporal locus for the interaction of perceptual load and involuntary attention. Attentional modulation for the C1 component (peak latency at approximately 80 ms) was also observed, but only for high load stimuli that were presented intermixed with low load stimuli. Results suggest that (a) perceptual load affects attentional selection at early processing stages; (b) perceptual load interacts with *involuntary* attention earlier and with different brain mechanisms relative to *voluntary* attention; and (c) attentional modulation in the C1 time range is possible under optimal experimental conditions.

Keywords

perceptual load; involuntary attention; event-related potentials (ERPs); peripheral cueing; C1; P1m

Introduction

The question of how early in visual processing selective attention exerts its effects has long been debated in cognitive psychology (for a review, see Johnson and Dark, 1986). There is consistent support that early selection occurs in extrastriate cortex, based on work with the

© 2009 Elsevier Inc. All rights reserved.

Please send correspondence to: Dr Shimin Fu, ARCH Laboratory, MS 3F5, George Mason University, Fairfax, VA 22030-4444, Phone: 703-993-4268, Fax: 703-993-1330, shimin.fu@gmail.com; sfu@gmu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

P1 component of the visual event-related potential (ERP) (Clark and Hillyard, 1996; Heinze et al., 1994; Mangun et al., 1998, 2001; Woldorff et al., 1997). Yet, other techniques provide evidence of even earlier selection: single-unit (Ito and Gilbert, 1999; McAdams and Reid, 2005; Motter, 1993; Roelfsema et al., 1998) and brain imaging studies (Gandhi et al., 1999; Hopf et al., 2004; Somers et al., 1999) have shown that attention can modulate activity in the striate cortex. However, the visual ERP C1 component --- the first visual cortical component (peak latency at approximately 80 ms after stimulus onset) arising from visual information processing and reflecting the initial cortical volley (Clark et al., 1995; Clark and Hillyard, 1996) -- is not consistently found to be modulated by attention. Several authors have argued that the attentional effects observed in the striate cortex reflect top-down processes from higher visual cortex back onto the striate cortex (Lamme and Spekreijse, 2000; Mehta et al., 2000a; Mehta et al., 2000b; Martinez et al., 1999, 2001; Noesselt et al., 2002; Super et al., 2001; see also Juan and Walsh, 2003; and Pascual-Leone and Walsh, 2001, for confirmatory evidence obtained from transcranial magnetic stimulation studies). On the other hand, recent ERP and MEG studies have shown C1 modulation by perception and/or attention (Kelly and Foxe, 2008; Khoe et al., 2005; Wu et al., 2005; see also Poghosyan et al., 2005 for MEG evidence).

However, the story is more complex than whether or not C1 is modulated by attention. Perceptual load also appears to influence the temporal locus of attentional selection in the brain. There is behavioral evidence that interference from task-irrelevant distractors with a target is reduced when the target is under high perceptual load (Lavie and Tsai, 1994; Lavie 1995). Early selection only occurs when the perceptual load is high. However, electrophysiological evidence is needed to determine how early in processing these influences are exerted. A direct translation of Lavie's behavioral conception of perceptual load into the ERP paradigm poses problems because simultaneously presented target and distractor stimuli produce overlapping ERP components. Nevertheless, by using operational definitions of perceptual load that vary somewhat from Lavie's, several researchers have explored the relationship between perceptual load and attention using ERPs (Fu et al., 2008; Handy and Mangun, 2000). Under voluntary attention conditions, different load by attention interactions have been observed for the visual P1 (Experiment 1 of Handy and Mangun, 2000) or N1 components (Fu et al., 2008; Experiment 2 of Handy and Mangun, 2000). The paradigms used in these studies differed in some respects, so the reasons for the different load by voluntary attention interactions for P1 or N1 are not clear.

Another important factor concerns how attention is deployed -- voluntarily or involuntarily -- as that affects the temporal locus of selection as well. It has been known for some time that involuntary attention affects stimulus processing in a manner distinguishable from voluntary attention. Cue validity effects have a different time course for the two types of attention shifts (Muller and Rabbitt, 1989; Yantis and Jonides, 1990). This is also seen in different characteristics of the amplitudes and latencies of ERP P1 and/or N1 components for involuntary relative to voluntary attention (Doallo et al., 2004; Fu et al., 2001; Hopfinger and Mangun, 1998; Hopfinger and West, 2006).

There are several reasons why an involuntary paradigm is advantageous for examining the locus of perceptual load effects. As attentional shifting is faster when attention is involuntarily captured by peripheral onset of cues (Eimer, 2000), involuntary attention can be expected to exert effects earlier in processing. Further, the timing of feedback modulation in V1 is different for voluntary and involuntary shifting of attention, as suggested by a recent study using the involuntary attention paradigm (Fu et al., 2005b). In that study, the initial C1 component (approximately 80 ms) was not modulated by attention; however, re-activation of V1 (approximately 110 ms) was evident in the P1 time range, suggesting a relatively early feedback mechanism for tasks involving involuntary attention (as compared

with approximately 150 ms using voluntary attention). Despite substantial evidence that involuntary attention is fast and automatic (e.g., Mueller and Rabbitt, 1989), previous ERP studies using involuntary attention tasks have observed that the earliest attentional modulation occurs on the later P1 and not the earlier C1 component (Doallo et al., 2004; Fu et al., 2005a; Hopfinger and Mangun, 1998; Hopfinger and Maxwell, 2005; Van de Lubbe et al., 1997). However, those studies did not manipulate perceptual load and most of the above-mentioned ERP studies on attention and the perceptual load-attention interaction (e.g., Fu et al., 2008; Handy and Mangun, 2000) have studied the locus of attention using voluntary attention paradigms with trial-by-trial cuing using central, symbolic cues. Therefore, to better understand the locus of attention, involuntary orienting should be assessed as perceptual load is manipulated.

Use of ERPs allows investigation into the temporal (and perhaps anatomic) locus of attentional selection and the interaction between perceptual load and attention. Previous work has shown that perceptual load interacts with voluntary attention within the P1 or N1 time range (Handy and Mangun, 2000) and the effect of this interaction on N1 has been localized to the temporal parieto-occipital region by using dipole modeling (Fu et al., 2008). The present study sought to extend our previous work to involuntary attention, investigating when the earliest attentional selection could occur and when perceptual load interacted with involuntary attention. We hypothesized that attentional selection might be evident earlier under conditions of high perceptual load and involuntary attention. We predicted earlier selection for high-load stimuli relative to low-load stimuli. We also expected that the interaction between perceptual load and attention would occur earlier under involuntary relative to the voluntary attention conditions that have been the focus of previous studies.

Method

Participants

Sixteen healthy college students participated as paid volunteers. Data from one participant were excluded because of technical problems, and data from two participants were also excluded from further analyses because of excessive eye blinks. The remaining 13 participants (7 male) were between 18 and 23 years of age (mean age 20.8 years). All the participants were right-handed and had normal or corrected-to-normal vision. They reported no history of neurological problems. Informed consent was obtained from all the participants.

Stimuli

A black fixation cross ($0.31^\circ \times 0.31^\circ$) was presented on a white background at the center of the monitor throughout the entire recording session. On each trial, each target array ($2.45^\circ \times 2.54^\circ$) appeared randomly on the left or right side, with its center 6.36° to the side and 2.05° above the fixation cross. The perceptual load of the target array was manipulated as low, medium, and high (Figure 1a, 1b, and 1c, respectively). The target arrays classified as medium load were identical to those used in our previous study (Fu et al., 2005b) and consisted of two horizontal, one vertical, and one diagonal line, with one line in each of the four quadrants. For each of the medium-load target arrays, the length of all the element lines was 0.9° and all the lines could appear at any of the quadrant with equal probability, with a restriction that the two horizontal lines were always displayed in the diagonal quadrants. The low-load target arrays were created by cutting the 3 non-target lines of the medium arrays into 10 small dot-like pieces, and then scrambling these small pieces at their respective quadrant, so that the target element (the diagonal line) could “pop-out” from the dot-like background. The high-load target arrays were created by adding a background of nine “X”s (each subtending 0.37°) to the medium arrays, so that the target line element needed to be

disentangled from confounding elements with same orientation in these background “X”s, before participants could perform the orientation judgment of the target diagonal line. For all the target arrays, the diagonal line could appear in any of the four quadrants with equal probability, and its orientation could be backward (“\”) or forward (“/”) with equal probability. Prior to the presentation of each target array, a peripheral Kanizsa box ($2.95^\circ \times 3.11^\circ$) served as a location cue, which flashed randomly in the left or right visual field. The center of the cue overlapped with the center of the target array in the left or right visual field. The cue was non-informative about the location of the target array; that is, the target array could equally appear at the same location as the cue or at the opposite side. All the three types of target arrays were mixed in every experiment block with equal probability, and the stimulus sequence was randomized for every subject. In all the experimental trials, 3/7 trials were valid (with 1/7 for low, medium and high load each), and 3/7 trials were invalid (with 1/7 for low, medium and high load each). In the remaining 1/7 of the trials, the peripheral cue was presented without a following target (the cue-only condition). The purpose of this cue-only condition was to remove the ERP overlap from the cues (because of short and fixed cue-to-target SOA), as in our previous study (Fu et al., 2005b). The durations of cue and target were 50 and 100 ms, respectively. The cue-to-target stimulus onset asynchrony (SOA) was fixed at 150 ms, and the inter-trial interval (ITI, offset to onset) ranged randomly between 1200 and 1600 ms. The purpose of using a fixed cue-to-target SOA was to have a relatively constant cue validity effect, so that if there was any small but relatively constant attentional effect on early ERP components (such as C1), it would be more likely recorded relative to that varying across trials.

Procedure

Participants were required to fixate the central cross and minimize eye blinks and body movements during all the experimental blocks. Their task was to discriminate the orientation of the diagonal line. They were instructed to respond to the backward diagonal (\) by pressing key 4 on a NeuroScan (NeuroScan, Texas, USA) Stimpad using their right thumb, and to respond to the forward diagonal (/) by pressing key 1 using their left thumb. To help participants memorize the stimulus-response mapping, they were told that their hand position matched the orientation of the stimuli, if they were to raise the Stimpad in front of them. Participants were instructed to respond as quickly as possible while maintaining higher than 90% of accuracy, and to withhold a response on the “cue only” trials. Each participant was given 80 practice trials before the EEG recording; further practice was given if they did not reach 90% accuracy in target (line orientation) discrimination. During the EEG recording, short breaks were allowed after every 40 trials. To mitigate fatigue effects, all participants were required to take part in the EEG recording session twice within three days, with exactly the same task and procedure. There were 1344 trials in total for each EEG recording session, ensuring a theoretical sample size of 192 for each experimental condition across recording sessions.

Data acquisition

The STIM (NeuroScan, Texas, USA) software packages were used to present stimuli, and SCAN (NeuroScan, Texas, USA) software packages were used to record and analyze the EEG data. Sixty-four channels of EEG and EOG were recorded from the scalp with a NeuroScan quick-cap (AgCl electrodes). Standard 10 – 20 sites were FPZ, FP1, FP2, FZ, F3, F4, F7, F8, CZ, C3, C4, PZ, P3, P4, OZ, O1, O2, T3, T4, T5, and T6. Additional intermediate sites were AF3, AF4, AF7, AF8, F1, F2, F5, F6, FCZ, FC1, FC2, FC3, FC4, FC5, FC6, FT7, FT8, C1, C2, C5, C6, CPZ, CP1, CP2, CP3, CP4, CP5, CP6, TP7, TP8, P1, P2, P5, P6, PO3, PO4, POZ, PO5, PO6, PO7, and PO8. Physically linked left and right mastoid, rather than a commonly used mathematically linked mastoid (Picton et al., 2000), served as reference. Horizontal eye movements (HEOG) were monitored by placing two

electrodes lateral to the left and right orbits. The impedances of all electrodes were kept below 5 K Ω throughout the recording session. Vertical eye movements (VEOG) and eye blinks were measured by placing two electrodes, with one 1.5 cm below and the other 1.5 cm above the left eye. The EEG from each electrode site was digitized at 500 Hz and was filtered with a band-pass of 0.1 to 40 Hz. The EEG for the 200 ms preceding target onset was used as baseline.

Data Analysis

Prior to averaging the EEG, artifact rejection was performed to discard epochs contaminated by eye blinks, horizontal eye movements, body movements, and muscle activity. The rejection threshold was $\pm 75 \mu\text{V}$ for eye blinks/muscle movements and was $\pm 40 \mu\text{V}$ for horizontal eye movements. EEGs for the cue-plus-target conditions were averaged from the onset of the target stimulus (valid, invalid), and EEGs for the cue-only trials were averaged from the onset of the “blank” stimulus. All the data were collapsed over the two orientations of the diagonal line. An ERP subtraction procedure (Fu et al., 2005b) was applied to remove the ERPs of the cues. After ERP subtraction, the peak amplitudes and latencies were used for statistical analyses. For behavioral data analysis, RTs exceeding ± 3 standard deviations were excluded from data analysis (less than 2% data excluded). Behavioral data were analyzed by means of repeated measures analysis of variances (ANOVAs) with three factors: load (low, medium, or high), validity (valid or invalid), and visual field (left or right). For electrophysiological data analysis, data were analyzed from electrode exhibiting the largest amplitude waveform of each component of interest. The hemisphere factor (left or right) was included when analyzing data from symmetrical electrodes (e.g., non-midline electrodes). The time window for obtaining peak latency and amplitude data was 50–110 ms for the C1 component, 70–160 ms for the P1 component, and 150–220 ms for the N1 component. The relatively wide ranges for searching for C1 and P1 peaks were set to accommodate individual differences in the C1 and P1 components; however, the overlap between these search ranges should not cause any confusion on the sequence of these components because of their polarities. For example, the peak latency of P1 for one participant (e.g., 80 ms) might be earlier than the peak latency of C1 for another participant (e.g., 110 ms), but for each participant, C1 should always be earlier than P1. Peak-baseline differences were calculated for peak amplitudes of each component. For statistical significance, Greenhouse-Geisser correction for non-sphericity was applied where appropriate. To be conservative in making statistic inference after multiple comparisons, the most stringent Bonferroni correction ($p = 0.05/3 = 0.017$, for comparisons under 3 load levels), was used to evaluate the critical comparisons between different load levels for each electrode of interest.

Results

Behavioral measures

Figure 2 shows mean reaction times (RTs) and error rates to targets averaged across 13 participants. Participants responded faster on valid relative to invalid trials [$F(1, 12) = 38.792$, $p < 0.0005$; see Figure 2a]. The main effect of load was significant [$F(2, 24) = 77.539$, $p < 0.0005$, $\epsilon = 0.835$]. Separate analyses showed that the participants responded fastest to low-load stimuli, slower to medium-load stimuli, and slowest to high-load stimuli (all $p < 0.0005$). The validity effect on RTs (invalid minus valid) was smallest for the high-load stimuli relative to both the low- and medium-load stimuli (both $p < 0.005$). Participants responded more accurately to valid relative to invalid trials [$F(1, 12) = 17.598$, $p < 0.001$; see Figure 2b]. The main effect of load was significant [$F(2, 24) = 48.743$, $p < 0.0005$, $\epsilon = 0.971$]. Separate analyses showed that the participants had the fewest errors for low-load stimuli, more errors for the medium-load stimuli, and the highest errors for the high-load

stimuli (all $p < 0.005$). The validity effect on error rates was larger for high-load relative to low-load stimuli [$F(1, 12) = 7.59, p < 0.017$], which explains the smaller validity effect on RTs for high-load relative to low- and medium-load stimuli—participants made more errors to invalid trials with high load to achieve faster responses.

ERP measures

In order to isolate the ERPs to the target from the ERPs to the cue-plus-target, ERPs from the cue-only condition were subtracted from ERPs from the cue-plus-target condition, a method used in our previous study (Fu et al., 2005b). By doing so, the target elicited ERPs can be compared between valid and invalid conditions. Figure 3 and Figure 4 show the target ERPs from valid and invalid trials for low-, medium-, and high-load stimuli after ERP subtraction. There was a clear negative C1 component over the parietal area about 80 ms after target onset, and a midline P1 component (P1m, Figure 3); the lateral P1 and N1 components in posterior areas were also distinguishable (Figure 4). There were effects of cue validity and perceptual load on these early ERP components.

C1—The C1 component was seen over parietal cortex with its largest amplitude at the PZ site (Figure 3). The load $\times$ validity interaction was significant [PZ: $F(2, 24) = 8.452, p < 0.003, \epsilon = 0.833$]. Separate analysis showed that valid trials elicited larger C1 for high-load stimuli [$F(1, 12) = 8.63, p < 0.012$]. This validity effect only approached significance for medium-load stimuli [$F(1, 12) = 2.297, p < 0.155$], and was not significant for low-load stimuli [$F(1, 12) = 0.165, p < 0.692$]. Figure 5a and 5b show the scalp voltage maps of the difference waves obtained by subtracting ERPs on invalid trials from those on valid trials. At 80 ms after the onset of the target array, there were no attentional effects on the C1 component under the low-load condition, but a C1 attentional effect was evident over parietal areas under the high-load condition (Figure 5a). For the C1 peak latency, no significant main effects or interactions were observed.

Midline P1 (P1m)—The amplitudes of the midline P1 component (P1m) were larger for invalid relative to valid trials at the central parietal area [CPZ: $F(1, 12) = 9.607, p < 0.009$], and this attentional effect increased with perceptual load, as suggested by a significant load $\times$ validity interaction [$F(2, 24) = 6.551, p < 0.009, \epsilon = 0.830$] (Figure 3; Figure 5a and 5b, 100–140 ms). For the latency of P1m, invalid trials elicited an earlier P1m relative to valid trials [$F(1, 12) = 15.781, p < 0.002$].

Lateral P1 (P1l)—For the amplitude of the lateral P1 (P1l), the load $\times$ validity interaction was significant [P3/P4: $F(2, 24) = 4.329, p < 0.031, \epsilon = 0.872$], suggesting that the validity effect on lateral P1 amplitude was more pronounced with increased load (Figure 4). No other main effects or interactions related to load or validity were significant. Separate analyses showed no significant validity main effect or related interactions for all stimuli (all $p > 0.15$). For the latency of P1l, invalid trials elicited an earlier lateral P1 [$F(1, 12) = 7.043, p < 0.021$] relative to valid trials.

Lateral N1 (N1l)—For the amplitude of the lateral posterior N1 (N1l), the validity $\times$ visual field $\times$ hemisphere interaction was significant [P3/P4: $F(1, 12) = 18.924, p < 0.001$], suggesting that invalid trials elicited a larger contralateral N1 relative to the valid trials. The main effect of load approached significance [$F(2, 24) = 3.411, p < 0.057, \epsilon = 0.888$], and the load $\times$ visual field $\times$ hemisphere interaction was significant [$F(2, 24) = 11.416, p < 0.0005, \epsilon = 0.945$], suggesting that the effects of the perceptual load were more pronounced at sites contralateral to the stimulus visual field (Figure 4). The scalp voltage distributions of ERPs at the P1 and N1 time range are shown in Figure 5a (100–200 ms) and Figure 5b (100–180

ms). For the latency of N11, invalid trials elicited an earlier posterior N1 relative to valid trials [$F(1, 12) = 8.508, p < 0.013$].

Discussion

One challenge in attention research is to design a task that can reveal the full power of selectivity in attention (Desimone and Duncan, 1995). Such a task must be sufficiently sensitive to detect the earliest selective neural processing associated with attentional modulation of task-relevant visual stimuli under demanding conditions with irrelevant distractors (Maunsell, 1995). We approached this challenge by using peripheral cueing to elicit involuntary attention (Yantis and Jonides, 1990) and by manipulating the perceptual load of the stimuli by varying distractors (Lavie, 1995). In the present study we hypothesized that observing the earliest cortical sign of attentional selection requires certain optimal experimental conditions, which include at a minimum, high perceptual load and involuntarily driven attention. Consistent with this hypothesis, the P1m (the posterior midline P1 component) was reliably modulated by involuntary attention in the main study and as well as in two follow-up studies^{1, 2}. Moreover, the earliest visual C1 component (approximately 80 ms) was also modulated by attention under these optimal experimental conditions. Whether these conditions are *sufficient* to modulate C1 will require further investigation. Importantly, our novel finding of a P1m interaction between perceptual load and involuntary attention suggests that attention-type (involuntary vs. voluntary attention, via indirect comparison) involves different brain mechanisms.

¹A C1 that is generated in striate cortex should show polarity inversion for upper vs. lower visual field stimuli (e.g., Di Russo et al., 2003). To test this, 15 healthy participants were asked to maintain central fixation while the cues and targets were presented in the upper and lower visual fields. Only high-load stimuli were used (Figure 1c). The cue-to-target SOA was jittered between 100–250 ms. The cues predicted the location of the subsequent target on 50% of the trials. The participants were required to respond to the orientation of the diagonal line. The ADJAR algorithm (Talsma and Woldorff, 2005; Woldorff, 1993) was applied to remove the ERP overlap between the cues and the targets. The attentional effect on C1 was not found, although the reversed polarity of C1 for stimuli in LoVF vs. UVF fits well with the cruciform model of the striate cortex (Jeffreys and Axford, 1972; Mangun, 1995). Notably, the jittered cue-to-target SOA in this follow-up study could have smeared a potentially small C1 attentional effect. Perhaps more importantly, arousal might have played a role in eliciting C1 attentional effects differentially between these two studies. Previous studies have shown an arousal-like effect on a component in the C1 latency range (80–90 ms after stimulus onset) from emotional expressions (Eger et al., 2003; Pourtois et al., 2004) and threatening stimuli (Stolarova et al., 2006). Based on this, the mixed load levels in the main study might have increased the arousal level for the high-load stimuli and thus contributed to the C1 effect for high-load stimuli. Consistent with this arousal hypothesis, participants responded faster to the high-load stimuli when they were intermixed with low and medium load stimuli (main study) relative to when they were present alone (follow-up study). However, some researchers have argued that it is impossible—or only possible to a very limited extent—to observe visual ERPs derived from sub-cortical structures such as the thalamus (Proverbio and Zani, 2003), where manifestations of arousal might be generated. Thus, the role of arousal for the C1 effects remains unclear.

²One limitation of our main study is that the cue-target sensory interactions might have co-varied with the stimulus load, thereby contributing to the present attentional effects on C1 and P1m. That is, the sensory interactions under the high-load condition could have differed from those under medium- and low-load conditions as there were physical differences (such as orientation, luminance, and spatial frequency, etc.) among the stimulus arrays associated with different load levels. We addressed this issue by using a new set of stimuli that better controlled for the difference between stimulus physical properties and cue-target interaction between load levels. Nineteen healthy participants (2 participants' data were excluded because of a data collection issue, 5 male, mean age 20.3 years) were tested. The location of the cues was valid, invalid, or neutral (appeared simultaneously in both the left and right visual field) with equal probability. The durations of the cue and the target array were 50 ms and 100 ms, respectively. The SOAs between the cue and the target array was randomized between 100 and 300 ms, and the ITI (inter-trial interval) between the offset of the target array and the next cue was ranged between 1100 and 1600 ms. The participants' task was to discriminate the orientation of the bar in the center of the target array. The perceptual load of the stimuli was high or low with equal probability. In the high load condition, distractors sharing characteristic features surrounded the central bar -- a central horizontal or vertical bar surrounded by crosses, or a central forward or backward bar surrounded by Xs. In the low load condition, the central bar did not share any features with the surrounding distractor -- a central horizontal or vertical bar surrounded by Xs, or a central forward or backward bar surrounded by crosses. Note that the location of the four distractors was designed to avoid collinearity between the central bar and the distractors. The sequence of trials was randomized for each participant. In total 1920 trials were presented. The ADJAR algorithm (Talsma and Woldorff, 2005; Woldorff, 1993) was used to remove ERP overlap due to short SOAs (100–300 ms) between the cues and the targets. The attentional effect on P1m was found and the perceptual load by attention interaction was replicated for P1m but not for C1. The C1 attentional effect was observed, but with a more parietal distribution. Thus, this study basically replicated the major findings on C1 and P1m of our main study, but did so in the absence of target stimulus confounds and cue-target sensory interactions.

P1m, attentional modulation, and the early perceptual load by attention interaction

The major finding of the present study was that for P1m perceptual load reliably interacted with involuntary attention. This was evident in our main study, whose findings were replicated in a follow-up study under conditions with better control over cue-target sensory interactions². In our previous study using a voluntary (sustained) attention task, perceptual load interacted with voluntary attention during the lateral N1 time range (peak latency 190 ms) but not for P1m (Fu et al., 2008). Indirect comparisons between these two studies which used similar stimuli suggest that attention type affects the locus of interaction and possibly involves different underlying brain mechanisms. This claim is based on the earlier onset and more medially distributed scalp locus of interaction we observed under involuntary relative to voluntary attention conditions.

To our knowledge, these are the first observations of perceptual load by attention interactions for P1m. These novel observations are important in light of previous findings of lateralized attentional effects on P1 and N1, i.e., P1l and N1l (e.g., Fu et al., 2008; Handy and Mangun, 2000). The load by *involuntary* attention interaction on P1m (approximately 140 ms for invalid trials) in the present study occurred earlier and the effects were distributed more medially compared to the load by *voluntary* attention interaction on posterior N1 (approximately 190 ms) in a previous sustained attention study (Fu et al., 2008) which used similar stimuli. The generators of the P1m are not known; presumably they are a part of the dorsal parieto-frontal network (e.g., superior parietal lobule and frontal eye field) that is involved in involuntary attention (Corbetta and Shulman, 2002). Moreover, the scalp distribution maps (Figure 5a and Figure 5b, 100–140 ms) showed that the attentional effect on P1m started at posterior and then spread to more central and frontal areas over its latency range, regardless of perceptual load. This suggests that it is likely to be a feed-forward process from early visual cortex to parietal and frontal areas (Lamme, 2003). Therefore, both the underlying attentional processes involved and the scalp distribution suggested different generators between the P1m and the conventional lateral P1 component.

Why does perceptual load interact with involuntary attention at an early processing stage as we saw with P1m (and C1, under optimal experimental conditions)? One explanation is that attentional selection involves a competitive process between the target and the distractors in the stimulus display, akin to the competitive neural interactions in visual areas with large receptive field (Desimone and Duncan, 1995). Perceptual load may influence the timing of different competitive process. That is, if there is little or no competition between the target line and the distractors, as in the low-load condition, the target line “wins” without the need for resolving the competition. In contrast, in the high-load condition, more attentional resources are required to resolve the competition so that the target line can “win out”. It is possible that the obligatory (or automatic) nature of involuntary attention may have involved early brain areas as shown on P1m, and perceptual load may have caused the flexible timing of attentional selection as shown on P1m and C1.

The attentional effect on P1m was robust. It was observed in our main study and two follow-up studies^{1, 2}, with valid trials eliciting smaller P1m than invalid trials. In fact, this P1m attentional effect under involuntary attentional orienting has been clearly illustrated -- although without statistical confirmation -- in a previous study (Hopfinger and West, 2006, Figure 3b, electrode Pz). Interestingly, the direction of the P1m attentional effect was reversed (i.e., P1m was larger for valid relative to invalid trials) in a recent study using voluntary attention task (Fu et al., 2008). There appears to be no simple explanation for this difference. Perhaps similar to the reversal of P1 attentional effect that have been observed in previous studies (e.g., Hopfinger and Mangun, 1998; McDonald et al., 1999), this P1m attentional effect may be affected by stimulus and task parameters. Furthermore, as we discussed previously, the neural sources of the P1m may be different from that of the

extrastriate cortex-generated P11. Specifically, we speculated that the generators of the P1m are a part of the dorsal parieto-frontal network (e.g., superior parietal lobule and frontal eye field) that is involved in involuntary attention (Corbetta and Shulman, 2002).

Attentional modulation and the C1 component

The combination of involuntary attention, a cluttered display, and line orientation targets in the present study may have contributed to the earliest C1 modulation of attention by involving predominantly neural activity in striate cortex, as we previously discussed (Fu et al., 2005b). The line orientation discrimination task is consistent with the neural specificity hypothesis of visual selective attention that attentional selection is based on “neural channels,” aggregations of neurons with identical receptive field properties which are responsible for encoding a given feature (orientation in the present study) of a stimulus” (Harter and Aine, 1984). The reflexive and fast nature of involuntary attention may have facilitated this attentional selection and the perceptual load by attention interaction, thereby modulating C1. This is also consistent with a model which proposes that the bottom-up saliency map is located in the primary visual cortex (Zhaoping, 2002) and that attentional capture could occur at the V1 level even without awareness (Zhaoping, 2008).

Nevertheless, the C1 component was not consistently modulated by attention. In our main study an interaction between perceptual load and attention on C1 was observed, but the attentional effect was seen only under the high-load conditions. However, when only high-load stimuli were presented and cue-to-target SOAs were jittered in a follow-up study ¹, this C1 attentional effect was absent. In another follow-up study ² with better control of cue-target sensory interaction and again a jittered SOA, the C1 attentional effect for the high-load stimuli was not significant at the site where C1 was largest but was significant at more parieto-centrally distributed sites. Thus, consistent with the literature that attention-related C1 effects have not been consistently observed across studies (e.g., Clark and Hillyard, 1996; Martinez et al., 1999; Noesselt et al., 2002; and Khoe et al., 2005; Kelly and Foxe, 2008), the optimal conditions for eliciting such effects remain uncertain. The contributory factors may include perceptual load, arousal, cue-target sensory interaction, and cue-to-target SOA. The present study suggests that high perceptual load may be necessary for eliciting a C1 attentional effect, as seen in both the main study and the second follow-up study ². Moreover, arousal might also have played a role in the present study, as suggested by our first follow-up study ¹, which showed no C1 attentional effect for the high-load stimuli in the absence of lower load stimuli (see note 1 for more discussion). Consistently, previous studies have shown an arousal-like effect at the C1 time range with emotional facial expression (Eger et al., 2003; Pourtois et al., 2004) and threatening/arousing stimuli (such as attack scenes, mutilated bodies) (Stolarova et al., 2006). Nevertheless, an explanation based on a possible confound of cue-target sensory interaction in the main study is less likely. This is because our follow-up study ² greatly reduced the difference on such cue-target interactions between the low-and high-load conditions yet still found a C1 attentional effect. In addition, individual differences in striate cortical structure might have also played a role in the present study, as suggested by a recent study (Kelly and Foxe, 2008). Kelly and Foxe pre-tested two stimulus locations and selected one location where C1 was largest to elicit C1 components from each participant. By doing so, individual differences in V1 architecture between participants was to some extent mitigated and thus not only the statistical power but also the chance to find a tiny C1 attentional effect was improved. Therefore, while the present approach of using high-load stimuli and involuntary attention is a promising one for eliciting the C1 attentional effect, the question of whether attention typically modulates the amplitude of C1 is still unanswered.

Although the relationship between the P1m and C1 attentional effects is unclear, one possibility is that the attention effect in striate cortex (indexed by C1) is “amplified” along

the visual processing streams to higher cortex (indexed by P1m), similar to the progressively enhanced fMRI signal strength of the attentional effect from V1 to V4 (O'Connor et al., 2002). Alternatively, it is possible that the C1 attentional effect may be affected by the P1m attentional effect because of ERP overlap between these two components (see Fu et al., 2008 for similar overlap and possible account). This latter explanation seems less likely because the C1 attentional effect was not associated with a latency change for the high-load stimuli as it would be if it were contaminated with P1m.

Attentional modulation and the contralateral P1 and N1 components (i.e., P1l and N1l)

The attentional effects we observed on the contralateral P1l and N1l components essentially replicated our previous findings in a peripheral cueing paradigm (Fu et al., 2001, 2005a, 2005b). Compared to invalid trials, valid trials elicited a later P1l, a later N1l, and a smaller N1l. Compared to the enhanced N1l for attended stimuli observed in previous voluntary attention tasks, this reversed N1l effect under involuntary conditions indicates that there may be different mechanisms underlying involuntary and voluntary attention. Moreover, a simple sensory gain control mechanism (for a review, see Hillyard et al. 1998) which proposes that attention amplifies processing of the attended stimuli at an early sensory processing stage without changing the timing of the processing, cannot account for the presently observed attentional modulation of P1l and N1l latencies.

It should be noted that while previous studies (e.g., Fu et al., 2005b; Hopfinger and Mangun, 1998) observed P1l enhancement by attention under peripheral cueing conditions, the present study found no main effect of attention on contralateral P1l but a significant interaction between attention and stimulus load, such that the higher the load, the larger the P1l for invalid relative to valid trials (see also our follow-up studies^{1, 2}). This lack of P1l attentional effect differs from our previous studies showing a smaller (and earlier) P1l for invalid relative to valid trials (Fu et al., 2001; Fu et al., 2005a, 2005b). This is noteworthy but not very surprising, because results from other research groups have also shown different patterns of attentional effects on P1l with peripheral cueing (e.g., Doallo et al., 2004; Eimer, 1994; Hopfinger and Mangun, 1998; Hopfinger and Mangun, 2001; McDonald et al., 1999; Prime and Ward, 2004, 2006). The P1l attentional effect elicited by peripheral cueing has been reported as positive (i.e., larger P1l for valid relative to invalid trials, Hopfinger and Mangun, 1998, under short cue-to-target SOA condition; Hopfinger and Mangun, 2001), or negative (i.e., larger P1l for invalid relative to valid trials, Eimer, 1994; McDonald et al., 1999; Prime and Ward, 2004, 2006), depending on stimulus properties and cue-to-target SOA. Similarly, the attentional effect on N1l has been shown to be positive (i.e., larger N1l for valid relative to invalid trials, see Hopfinger and Mangun, 1998, long SOA condition, but without statistical analysis on this component), negative (i.e., larger N1l for invalid relative to valid trials, Fu et al., 2001, 2005a, 2005b; Hopfinger and Mangun, 2001, short SOA condition), or null (Hopfinger and Mangun, 1998, short SOA condition; Hopfinger and Mangun, 2001, long SOA condition) under involuntary attention tasks. Given that almost all the visual areas and also feedback mechanisms have been activated approximately 100 ms after stimulus onset (Lamme, 2003), it is not surprising that the P1l (approximately 70–150 ms) and N1l (approximately 160–200 ms) components vary across different stimuli and tasks. That is, the attentional effect on P1l and N1l may have overlapped with other processes occurring at the same time range in the present study.

Summary

The present work shows that perceptual load affects the locus of attentional selection at early processing stages as manifested in effects on P1m (the posterior midline P1 component). Additionally, perceptual load interacts with involuntary attention at the P1m time range (peak latency varied between 108 to 140 ms), with invalid trials eliciting a larger

attentional effect which increased with perceptual load. The scalp distribution of P1m suggests that the attentional effects on it differ from conventional attentional effects on the lateral P1 and N1 components. Considered in the context of previous studies, these results suggest that the interaction between perceptual load and involuntary attention involves neural mechanisms that differ from those involved in the interaction between perceptual load and voluntary attention. Results also suggest that attentional modulation of the C1 component (presumably originating in striate cortex) may only be observed under optimal experimental conditions, and, further, that high and/or varying perceptual load may be required to elicit attentional effects on C1. If that is correct, it would suggest that under high load, the filter is set earlier, thereby modulating early components.

Acknowledgments

This research was supported by NIH grant AG19653 to RP, and MSTC grant 2002CCA01000 and CAS grant KSCX2-SW-221 to YL. The authors thank Daniel Caggiano, Silu Fan, Steve Hillyard, Xiong Jiang, Yang Jiang, Andreas Kleinschmidt and anonymous reviewers for helpful comments on these studies and Autumn Brown for assistance in data collection.

References

- Anllo-Vento L, Hillyard SA. Selective attention to the color and direction of moving stimuli: electrophysiological correlates of hierarchical feature selection. *Perception & Psychophysics* 1996;58:191–206. [PubMed: 8838164]
- Clark VP, Fan S, Hillyard SA. Identification of early visual evoked potential generators by retinotopic and topographic analyses. *Human Brain Mapping* 1995;2:170–187.
- Clark VP, Hillyard SA. Spatial selective attention affects early extrastriate but not striate components of the visual evoked potential. *Journal of Cognitive Neuroscience* 1996;8:387–402.
- Corbetta M, Shulman GL. Control of goal-directed and stimulus-driven attention in the brain. *Nature Reviews Neuroscience* 2002;3:201–215.
- Desimone R, Duncan J. Neural mechanisms of selective visual attention. *Annual Review of Neuroscience* 1995;18:193–222.
- Di Russo F, Martinez A, Hillyard SA. Source analysis of event-related cortical activity during visuospatial attention. *Cerebral cortex* 2003;13:486–499. [PubMed: 12679295]
- Doallo S, Lorenzo-Lopez L, Vizoso C, Holguin SR, Amenedo E, Bara S, et al. The time course of the effects of central and peripheral cues on visual processing: an event-related potential study. *Clinical Neurophysiology* 2004;115:199–210. [PubMed: 14706489]
- Eger E, Jedynak A, Iwaki T, Skrandies W. Rapid extraction of emotional expression: evidence from evoked potential fields during brief presentation of face stimuli. *Neuropsychologia* 2003;41:808–817. [PubMed: 12631531]
- Eimer M. The time course of spatial orienting elicited by central and peripheral cues: evidence from event-related brain potentials. *Biological Psychology* 2000;53:253–258. [PubMed: 10967235]
- Fu S, Caggiano DM, Greenwood PM, Parasuraman R. Event-related potentials reveal dissociable mechanisms for orienting and focusing visuospatial attention. *Cognitive Brain Research* 2005a; 23:341–353. [PubMed: 15820641]
- Fu S, Fan S, Chen L, Zhuo Y. The attentional effects of peripheral cueing as revealed by two event-related potential studies. *Clinical Neurophysiology* 2001;112:172–185. [PubMed: 11137676]
- Fu S, Greenwood PM, Parasuraman R. Brain mechanisms of involuntary visuospatial attention: an event-related potential study. *Human Brain Mapping* 2005b;25:378–390. [PubMed: 15852465]
- Fu S, Zinni M, Squire P, Kumar R, Caggiano DM, Parasuraman R. When and where perceptual load interacts with voluntary visuospatial attention: an event-related potential and dipole modeling study. *NeuroImage* 2008;39:1345–1355. [PubMed: 18006335]
- Gandhi SP, Heeger DJ, Boynton GM. Spatial attention affects brain activity in human primary visual cortex. *Proceedings of the National Academy of Sciences of the United States of America* 1999;96:3314–3319. [PubMed: 10077681]

- Handy TC, Mangun GR. Attention and spatial selection: electrophysiological evidence for modulation by perceptual load. *Perception & Psychophysics* 2000;62:175–186. [PubMed: 10703265]
- Harter, MR.; Aine, CJ. Brain mechanisms of visual selective attention. In *Varieties of Attention*. Parasuraman, R.; Davies, DR., editors. Orlando, FL: Academic Press; 1984. p. 293–321.
- Heinze HJ, Mangun GR, Burchert W, Hinrichs H, Scholz M, Munte TF, et al. Combined spatial and temporal imaging of brain activity during visual selective attention in humans. *Nature* 1994;372:543–546. [PubMed: 7990926]
- Hillyard SA, Anillo-Vento L. Event-related brain potentials in the study of visual selective attention. *Proceedings of the National Academy of Sciences of the United States of America* 1998;95:781–787. [PubMed: 9448241]
- Hillyard SA, Vogel EK, Luck SJ. Sensory gain control (amplification) as a mechanism of selective attention: electrophysiological and neuroimaging evidence. *Philosophical Transactions of the Royal Society of London - Series B: Biological Sciences* 1998;353:1257–1270.
- Hopf JM, Noesselt T, Tempelmann C, Braun J, Schoenfeld A, Heinze HJ. Popout modulates focal attention in the primary visual cortex. *NeuroImage* 2004;22:574–582. [PubMed: 15193585]
- Hopfinger JB, Mangun GR. Reflexive attention modulates processing of visual stimuli in human extrastriate cortex. *Psychological Science* 1998;9:441–447.
- Hopfinger JB, Maxwell JS. Appearing and disappearing stimuli trigger a reflexive modulation of visual cortical activity. *Cognitive Brain Research* 2005;25:48–56. [PubMed: 15907377]
- Hopfinger JB, West VM. Interactions between endogenous and exogenous attention on cortical visual processing. *NeuroImage* 2006;31:774–789. [PubMed: 16490366]
- Ito M, Gilbert CD. Attention modulates contextual influences in the primary visual cortex of alert monkeys. *Neuron* 1999;22:593–604. [PubMed: 10197538]
- Jeffreys D, Axford J. Source locations of pattern-specific component of human visual evoked potentials. I: Component of striate cortical origin. *Experimental Brain Research* 1972;16:1–21.
- Johnson WA, Dark VJ. Selective attention. *Annual Review of Psychology* 1986;37:43–75.
- Juan CH, Walsh V. Feedback to V1: a reverse hierarchy in vision. *Experimental Brain Research* 2003;150:259–263.
- Kelly SP, Gomez-Ramirez M, Foxe JJ. Spatial attention modulates initial afferent activity in human primary visual cortex. *Cerebral cortex* 2008;2008:2629–2636. [PubMed: 18321874]
- Khoe W, Mitchell J, Reynolds J, Hillyard SA. Exogenous attentional selection of transparent superimposed surfaces modulates early event-related potentials. *Vision Research* 2005;45:3004–3014. [PubMed: 16153678]
- Lamme VA. Why visual attention and awareness are different. *Trends in Cognitive Sciences* 2003;7:12–18. [PubMed: 12517353]
- Lamme, VA.; Spekreijse, H. Contextual modulation in primary visual cortex and scene perception. In: Gazzaniga, MS., editor. *The new cognitive neuroscience*. Cambridge, MA: MIT press; 2000. p. 279–290.
- Lavie N. Perceptual load as a necessary condition for selective attention. *Journal of Experimental Psychology: Human Perception & Performance* 1995;21:451–468. [PubMed: 7790827]
- Lavie N. Distracted and confused?: Selective attention under load. *Trends in Cognitive Sciences* 2005;9:75–82. [PubMed: 15668100]
- Lavie N, Tsal Y. Perceptual load as a major determinant of the locus of selection in visual attention. *Perception & Psychophysics* 1994;56:183–197. [PubMed: 7971119]
- Luck SJ, Hillyard SA, Mouloua M, Woldorff MG, Clark VP, Hawkins HL. Effects of spatial cuing on luminance detectability: psychophysical and electrophysiological evidence for early selection. *Journal of Experimental Psychology: Human Perception & Performance* 1994;20:887–904. [PubMed: 8083642]
- Mangun GR. Neural mechanisms of visual selective attention. *Psychophysiology* 1995;32:4–18. [PubMed: 7878167]
- Mangun GR, Buonocore M, Girelli M, Jha A. ERP and fMRI measures of visual spatial selective attention. *Human Brain Mapping* 1998;6:383–389. [PubMed: 9788077]

- Mangun GR, Hinrichs H, Scholz M, Mueller-Gaertner HW, Herzog H, Krause BJ, et al. Integrating electrophysiology and neuroimaging of spatial selective attention to simple isolated visual stimuli. *Vision Research* 2001;41:1423–1435. [PubMed: 11322984]
- Martinez A, Anllo-Vento L, Sereno MI, Frank LR, Buxton RB, Dubowitz DJ, et al. Involvement of striate and extrastriate visual cortical areas in spatial attention. *Nature Neuroscience* 1999;2:364–369.
- Martinez A, DiRusso F, Anllo-Vento L, Sereno MI, Buxton RB, Hillyard SA. Putting spatial attention on the map: timing and localization of stimulus selection processes in striate and extrastriate visual areas. *Vision Research* 2001;41:1437–1457. [PubMed: 11322985]
- McAdams CJ, Reid RC. Attention modulates the responses of simple cells in monkey primary visual cortex. *The Journal of Neuroscience* 2005;25:11023–11033. [PubMed: 16306415]
- McDonald JJ, Ward LM, Kiehl KA. An event-related brain potential study of inhibition of return. *Perception & Psychophysics* 1999;61:1411–1423. [PubMed: 10572468]
- Mehta AD, Ulbert I, Schroeder CE. Intermodal selective attention in monkeys. I: distribution and timing of effects across visual areas. *Cerebral Cortex* 2000;10:343–358. [PubMed: 10769247]
- Mehta AD, Ulbert I, Schroeder CE. Intermodal selective attention in monkeys. II: physiological mechanisms of modulation. *Cerebral Cortex* 2000;10:359–370. [PubMed: 10769248]
- Motter BC. Focal attention produces spatially selective processing in visual cortical areas V1, V2, and V4 in the presence of competing stimuli. *Journal of Neurophysiology* 1993;70:909–919. [PubMed: 8229178]
- Muller HJ, Rabbitt PM. Reflexive and voluntary orienting of visual attention: time course of activation and resistance to interruption. *Journal of Experimental Psychology: Human Perception and Performance* 1989;15:315–330. [PubMed: 2525601]
- Noesselt T, Hillyard SA, Woldorff MG, Schoenfeld A, Hagner T, Jancke L, et al. Delayed striate cortical activation during spatial attention. *Neuron* 2002;35:575–587. [PubMed: 12165478]
- O'Connor DH, Fukui MM, Pinsk MA, Kastner S. Attention modulates responses in the human lateral geniculate nucleus. *Nature Neuroscience* 2002;5:1203–1209.
- Pascual-Leone A, Walsh V. Fast backprojections from the motion to the primary visual area necessary for visual awareness. *Science* 2001;292:510–512. [PubMed: 11313497]
- Picton TW, Bentin S, Berg P, Donchin E, Hillyard SA, Johnson R Jr, et al. Guidelines for using human event-related potentials to study cognition: recording standards and publication criteria. *Psychophysiology* 2000;37:127–152. [PubMed: 10731765]
- Poghosyan V, Shibata T, Loannides AA. Effects of attention and arousal on early responses in striate cortex. *European Journal of Neuroscience* 2005;22:225–234. [PubMed: 16029212]
- Pourtois G, Grandjean D, Sander D, Vuilleumier P. Electrophysiological correlates of rapid spatial orienting towards fearful faces. *Cerebral Cortex* 2004;14:619–633. [PubMed: 15054077]
- Prime DJ. Cortical expressions of inhibition of return. *Brain Research* 2006;1072:161–174. [PubMed: 16445889]
- Prime DJ, Ward LM. Inhibition of return from stimulus to response. *Psychological Science* 2004;15:272–276. [PubMed: 15043647]
- Proverbio, AM.; Zani, A. Electromagnetic manifestations of mind and brain. In: Zani, A.; Proverbio, A., editors. *The Cognitive Electrophysiology of Mind and Brain*. San Diego, CA: Academic Press; 2003. p. 13–40.
- Roelfsema PR, Lamme VA, Spekreijse H. Object-based attention in the primary visual cortex of the macaque monkey. *Nature* 1998;395:376–381. [PubMed: 9759726]
- Somers DC, Dale AM, Seiffert AE, Tootell RB. Functional MRI reveals spatially specific attentional modulation in human primary visual cortex. *Proceedings of the National Academy of Sciences of the United States of America* 1999;96:1663–1668. [PubMed: 9990081]
- Stolarova M, Keil A, Moratti S. Modulation of the C1 Visual Event-related component by Conditioned Stimuli: Evidence for Sensory Plasticity in Early Affective Perception. *Cerebral cortex* 2006;16:876–887. [PubMed: 16151178]
- Super H, Lamme VA. Two distinct modes of sensory processing observed in monkey primary visual cortex (V1). [see comment]. *Nature Neuroscience* 2001;4:304–310.

- Talsma, D.; Woldorff, MG. Methods for the estimation and removal of artifacts and overlap in ERP waveforms. In: Handy, TC., editor. *Event-related Potentials: a Method Handbook*. Cambridge, MA: MIT Press; 2005. p. 115-148.
- Van de Lubbe RHJ, Woestenburg JC. Modulation of early ERP components with peripheral precues: a trend analysis. *Biological Psychology* 1997;45:143–158. [PubMed: 9083648]
- Vogel EK, Luck SJ. The visual N1 component as an index of a discrimination process. *Psychophysiology* 2000;37:190–203. [PubMed: 10731769]
- Vogel EK, Woodman GF, Luck SJ. Pushing around the locus of selection: evidence for the flexible-selection hypothesis. *Journal of Cognitive Neuroscience* 2005;17:1907–1922. [PubMed: 16356328]
- Wijers AA, Lange JJ, Mulder G, Mulder LJ. An ERP study of visual spatial attention and letter target detection for isoluminant and nonisoluminant stimuli. *Psychophysiology* 1997;34:553–565. [PubMed: 9299910]
- Woldorff MG, Fox PT, Matzke M, Lancaster JL, Veeraswamy S, Zamarripa F, et al. Retinotopic organization of early visual spatial attentional effects as revealed by PET and ERPs. *Human Brain Mapping* 1997;5:280–286. [PubMed: 20408229]
- Wu Y, Chen J, Han S. Neural mechanisms of attentional modulation of perceptual grouping by collinearity. *Neuroreport* 2005;16:567–570. [PubMed: 15812309]
- Yantis S, Jonides J. Abrupt visual onsets and selective attention: voluntary versus automatic allocation. *Journal of Experimental Psychology: Human Perception and Performance* 1990;16:121–124. [PubMed: 2137514]
- Zhaoping L. A saliency map in primary visual cortex. *Trends in Cognitive Sciences* 2002;6:9–16. [PubMed: 11849610]
- Zhaoping L. Attention capture by eye of origin singletons even without awareness-A hallmark of a bottom-up saliency map in the primary visual cortex. *Journal of Vision* 2008;8:1–18.

Figure 1.

Illustrations of the stimuli with low, medium and high perceptual load. Low-load stimuli (Figure 1a) were developed by cutting the three non-target lines of the medium-load stimuli (Figure 1b) into small parts and scrambling them in their respective quadrants. High-load stimuli (Figure 1c) were developed by putting the medium-load stimulus on a background that consisted of “x”s.

(a)

(b)

Figure 2.
Reaction times (RTs) and error rates as a function of perceptual load and cue validity.

Figure 3.

The grand averaged ERPs across all participants for the valid and invalid trials under low-, medium- and high-load conditions for three midline electrodes (CPZ, PZ, and POZ). Data were averaged across visual fields. The attentional effect on P1m (the midline P1 component) increased as a function of perceptual load. The amplitude of C1 was larger for valid relative to invalid trials under the high-load condition.

Figure 4.

The grand averaged ERPs across all participants for the valid and invalid trials under low-, medium- and high-load conditions at the parietal (P3/P4), temporal (T5/T6) and occipital (O1/O2) regions contralateral to the target location. Data were averaged across visual field and hemisphere.

Figure 5.

(a) The 3D back view of the scalp voltage distributions of the attentional effects between 80 and 200 ms (every 20 ms) after target array onset. (b) The 2D top view of the mean scalp voltage distributions of the attentional effects between 60 and 180 ms (every 20 ms) after target array onset. ERP data were obtained from subtracting the averaged ERPs of the invalid trials from that of the valid trials. Data were collapsed across stimulus visual field. The attentional effects on the C1 (80 ms) and P1m (100–140 ms, midline area) increased with the increase of the perceptual load (Figure 5a). The attentional effects at the P1 and N1 time range (120–180 ms) showed strong activities on the occipito-temporal area (Figure 5a and 5b). Note that the bilateral occipito-temporal activities at the 120–180 ms time range

was the results of averaging across left and right visual fields; these attentional effects were actually more pronounced on the hemisphere contralateral to the stimulus location. Moreover, the attentional effect on P1m over its latency range (100–140 ms) started on the parietal area, and then spread to the central and frontal areas, regardless of perceptual load (Figure 5b).