Published in final edited form as:

Mech Ageing Dev. 2010 April ; 131(4): 270–275. doi:10.1016/j.mad.2010.02.008.

A potential link between phosphate and aging – lessons from Klotho-deficient mice

Makoto Kuro-o, M.D., Ph.D.

Department of Pathology, The University of Texas Southwestern Medical Center at Dallas, 5323 Harry Hines Blvd., Dallas, Texas 75390-9072, Phone: (214) 648-4018, Fax: (214) 648-4070

Makoto Kuro-o: Makoto.Kuro-o@UTSouthwestern.edu

Abstract

Phosphate homeostasis is maintained primarily by a bone-kidney endocrine axis. When phosphate is in excess, fibroblast growth factor-23 (FGF23) is secreted from bone and acts on kidney to promote phosphate excretion into urine. FGF23 also reduces serum vitamin D levels to suppress phosphate absorption from intestine. Thus, FGF23 functions as a hormone that induces negative phosphate balance. One critical feature of FGF23 is that it requires Klotho, a single-pass transmembrane protein expressed in renal tubules, as an obligate co-receptor to bind and activate cognate FGF receptors. Importantly, defects in either FGF23 or Klotho not only cause phosphate retention but also a premature-aging syndrome in mice, which can be rescued by resolving hyperphosphatemia. In addition, changes in extracellular and intracellular phosphate concentration affect glucose metabolism, insulin sensitivity, and oxidative stress *in vivo* and *in vitro*, which potentially affect aging processes. These findings suggest an unexpected link between inorganic phosphate and aging in mammals.

Keywords

Klotho; FGF23; phosphate; vitamin D; CKD

1. Introduction

Aging is one of the most potent risk factors for any age-related diseases, including cancer, cardiovascular disease, stroke, chronic kidney disease, and neurodegenerative disorders, which are leading causes of death and disability in aging society. Suppression of aging, if at all possible, would be the most effective way to prevent these diseases, reduce mortality, and improve quality of life. Aging processes can be accelerated or suppressed by various environmental factors. Diet restriction (a dietary regimen of 40–50% less food intake than the average diet) has been known to suppress aging and extend life span in diverse experimental organisms including nematodes, fruit flies, rodents, and non-human primates (Masoro, 2006). Thus, diet restriction very likely suppresses aging in humans as well. However, it is extremely difficult for anyone to reduce daily meals down to about one-half and continue this eating habit throughout life. The major challenge is to find a strategy for suppressing aging that is feasible in daily life. If one can identify a particular nutrient(s) responsible for the benefit of diet restriction, reducing intake of such nutrient(s) alone may be as effective as and more feasible

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

than diet restriction. In addition, identification of such nutrient(s) is expected to provide new insights into the molecular mechanism of aging. As discussed below, recent animal studies and epidemiological studies have raised the possibility that phosphate may be one of such nutrients. The purpose of this review is to discuss potential link between aging and phosphate.

2. Endocrine regulation of phosphate metabolism

Serum mineral levels are maintained within narrow ranges. In humans, serum calcium and phosphate levels are normally 2.2–2.5 mM (8.8–10.2 mg/dL) and 0.81–1.45 mM (2.5–4.5 mg/dL), respectively (Ravel, 1994; Tonelli et al., 2005). Serum calcium and phosphate levels are determined by counterbalance between absorption from intestine, mobilization from bone (the major reservoir of calcium phosphate in the body), and excretion from kidney into urine (Berndt and Kumar, 2009; Schiavi and Kumar, 2004). These processes are coordinately regulated by several endocrine factors. Vitamin D and parathyroid hormone (PTH) have been extensively studied as hormones that regulate calcium and phosphate metabolism (Dusso et al., 2005). The active form of vitamin D (1,25-dihydroxyvitamin D₃) is synthesized in the kidney and acts on intestine to increase absorption of dietary calcium and phosphate. It also acts on bone to stimulate osteoclastogenesis and promote mobilization of calcium and phosphate from the reservoir, leading to increase in both calcium and phosphate levels in blood. PTH is secreted from parathyroid gland in response to low serum calcium levels and acts on kidney to promote vitamin D synthesis, thereby increasing intestinal absorption of calcium and phosphate. Unlike vitamin D, PTH has an activity that promotes phosphate excretion into urine (phosphaturia). Thus, PTH selectively increases blood calcium levels without concomitant increase in blood phosphate levels (Berndt and Kumar, 2007).

Recent studies have identified fibroblast growth factor-23 (FGF23) as a novel hormone that selectively lowers blood phosphate levels (Kuro-o, 2006; Quarles, 2003; White et al., 2000). When phosphate is in excess, FGF23 is secreted from bone and acts on kidney to induce phosphaturia and suppress vitamin D synthesis to induce a negative phosphate balance and maintain phosphate homeostasis (Liu et al., 2007; Liu et al., 2006; Quarles, 2003). FGF23 was originally identified as a gene mutated in patients with autosomal dominant hypophosphatemic rickets (ADHR), one of the rare hereditary disorders that exhibit renal phosphate wasting (White et al., 2000). Patients with ADHR carry missense mutations in the *FGF23* gene that confer resistance to proteolytic degradation of the FGF23 protein. Wild-type FGF23 protein is cleaved by an unknown protease at the ¹⁷⁶RXXR¹⁷⁹ motif and inactivated (White et al., 2001). Missense mutations in this critical motif (R176Q, R179Q/W) make the protein resistant to the protease (Shimada et al., 2002). As a result, ADHR patients have high serum FGF23 levels and phosphate-wasting phenotypes including hypophosphatemia and defects in bone mineralization (rickets). In contrast, mice lacking FGF23 (*Fgf23*^{-/-} mice) develop severe phosphate-retention phenotypes characterized by extensive soft tissue calcification and hyperphosphatemia (Shimada et al., 2004b). Furthermore, several mutations that affect expression and/or proteolytic degeneration of FGF23 have been identified in mice and humans, in which phosphate wasting phenotypes or phosphate retention phenotypes are associated with increased or decreased serum FGF23 levels, respectively (Kurosaki and Kuro-o, 2009a; Quarles, 2008). These observations have established that FGF23 is a hormone indispensable for maintaining phosphate homeostasis.

3. Molecular basis of FGF23 action

Although FGF23 belongs to the FGF ligand superfamily (Yamashita et al., 2000), phylogenetic and sequence analyses have segregated FGF23 and two additional FGFs (FGF19 and FGF21) from the other FGF family members (Itoh and Ornitz, 2008). These three atypical FGFs, namely FGF19, FGF21, and FGF23, are also known as endocrine FGFs, because they function as

endocrine factors unlike the other classical FGFs that primarily function as paracrine and/or autocrine factors (Kuro-o, 2008). The molecular basis behind the endocrine mode of action may lie in the fact that these endocrine FGFs have low affinity to heparan sulfate (HS). In general, FGF ligands have high affinity to heparin and HS, because they share a conserved HS-binding domain (Mohammadi et al., 2005a; Mohammadi et al., 2005b; Murzin et al., 1992). However, the HS-binding domain of endocrine FGFs deviates from that of the other paracrine-acting FGFs and prohibits formation of hydrogen bonding between HS and amino acid residues in the HS-binding domain, which is the basis of affinity to HS (Goetz et al., 2007; Harmer et al., 2004). This unique structural feature reduces affinity of endocrine FGFs to HS and allows them to escape from HS-rich extracellular matrices and enter systemic circulation. Although the low affinity to HS may help endocrine FGFs function as hormones, it may attenuate their ability to activate FGF receptors (FGFRs), because HS is required for high affinity binding of FGFs to FGFRs. FGFRs are receptor tyrosin kinases that dimerize upon binding FGFs. HS participates in the FGF-FGFR interaction and promotes formation of a 2:2:2 FGF-FGFR-HS signaling complex, which is essential for efficient activation of FGFR tyrosine kinase (Schlessinger et al., 2000). Thus, endocrine FGFs may require a co-factor(s) other than and/or in addition to HS to secure efficient dimerization and activation of FGFR. In fact, FGF23 cannot activate FGF signaling in most cultured cells even when they express FGFRs endogenously, whereas classical FGFs like FGF2 can activate FGF signaling in these cells (Kurosu et al., 2006).

Recent studies identified Klotho, a single-pass transmembrane protein expressed in the kidney, as a co-factor required for FGF23 to bind to FGFRs (Kurosu et al., 2006; Urakawa et al., 2006). Klotho protein forms a constitutive binary complex with several FGFRs (FGFR1c, 3c, and 4) and increases their affinity specifically to FGF23. In other words, Klotho functions as an obligate co-receptor for FGF23. In fact, Klotho-deficient mice are totally resistant to FGF23 and exhibit phosphate-retention phenotypes like FGF23-deficient mice (Tsujikawa et al., 2003; Yoshida et al., 2002). Kidney-specific expression of Klotho explains why FGF23 recognizes kidney as its target organ among many organs that express FGFRs. Klotho protein may have evolved to compensate for the low affinity of FGF23 to heparan sulfate and specifically support FGFR activation with FGF23, which represents a novel mechanism for confining target organs in redundant ligand-receptor interactions. Thus, Klotho and FGF23 have emerged as important components of the bone-kidney endocrine axis that regulates phosphate metabolism (Kuro-o, 2006, 2008; Kurosu and Kuro-o, 2008, 2009b; Liu et al., 2007; Liu and Quarles, 2007).

The phosphaturic activity of FGF23 stems from its ability to suppress sodium-phosphate co-transporter type-2a (NaPi-2a) that mediates transepithelial phosphate reabsorption in renal tubules (Segawa et al., 2003; Segawa et al., 2007; Shimada et al., 2004c). Injection of recombinant FGF23 protein reduces the number of NaPi-2a expressed on the apical brush border membrane of proximal tubules and induces phosphaturia within hours, although the precise signaling pathway through which FGF23 regulates NaPi-2a trafficking and/or expression in a Klotho-dependent manner remains to be determined. Importantly, negative phosphate balance reduces serum FGF23 levels, indicating the existence of a negative feedback loop in the regulation of phosphate excretion (Yu et al., 2005) (Fig. 1). This negative feedback loop is essential for phosphate homeostasis, because disruption of this loop by ablating FGF23 or Klotho expression results in hyperphosphatemia and severe phosphate retention in mice and humans. On the other hand, the activity of FGF23 as a counter-regulatory hormone for vitamin D originates from its ability to suppress synthesis and promote degradation of 1,25-dihydroxyvitamin D₃ in the kidney (Shimada et al., 2004a). FGF23 down-regulates expression of the *Cyp27b1* gene that encodes 1 α -hydroxylase, the enzyme that synthesizes the active form of vitamin D (1,25-dihydroxyvitamin D₃) from its inactive precursor (25-hydroxyvitamin D₃). In addition, FGF23 up-regulates expression of the *Cyp24* gene that encodes 24-

hydroxylase, the enzyme that hydrolyzes and inactivates 1,25-dihydroxyvitamin D₃ (Liu et al., 2006). Importantly, 1,25-dihydroxyvitamin D₃ up-regulates expression of the *FGF23* gene (Shimada et al., 2004a) and closes a negative feedback loop (Fig. 1). This negative feedback loop is essential for maintaining vitamin D homeostasis, because disruption of this loop results in high serum 1,25-dihydroxyvitamin D₃ levels as observed in *Klotho*-deficient mice (Tsujikawa et al., 2003) and *FGF23*-deficient mice (Shimada et al., 2004b).

4. Genetic evidence for phosphate toxicity

Defects in either *Klotho* or *FGF23* disrupt the negative feedback loops that maintain phosphate and vitamin D homeostasis, resulting in high serum phosphate and vitamin D levels. High serum vitamin D promotes intestinal absorption of calcium and induces hypercalcemia as well. Of note, this metabolic state characterized by high serum phosphate, calcium, and vitamin D levels is associated with a premature aging syndrome. In fact, the *klotho* gene was originally identified as a gene mutated in a mouse strain that inherited a premature aging syndrome (Kuro-o et al., 1997). Mice lacking *Klotho* develop multiple aging-like phenotypes, including a shortened life span, growth retardation, hypogonadotropic hypogonadism, rapid thymus atrophy (Min et al., 2007), skin atrophy, sarcopenia, vascular calcification, osteopenia (Kawaguchi et al., 1999), pulmonary emphysema (Ishii et al., 2008; Sato et al., 2007; Suga et al., 2000), cognition impairment (Nagai et al., 2003), hearing disturbance (Kamemori et al., 2002), and motor neuron degeneration (Anamizu et al., 2005). These phenotypes are also observed in mice lacking *FGF23* (Razzaque et al., 2006).

These observations imply that phosphate, calcium, and/or vitamin D may be toxic when retained and accelerate aging. Several studies have supported this notion. First, vitamin D-deficient diet not only restored serum phosphate and calcium levels to normal but also rescued several aging-like phenotypes in both *Klotho*-deficient mice and *FGF23*-deficient mice (Stubbs et al., 2007; Tsujikawa et al., 2003). Second, ablation of vitamin D action in *Klotho*-deficient mice and *FGF23*-deficient mice by disrupting the *Cyp27b1* gene (Ohnishi et al., 2009; Razzaque et al., 2006) or vitamin D receptor gene (Hesse et al., 2007) also rescued hyperphosphatemia, hypercalcemia, and the premature aging syndrome.

Lastly, low phosphate diet rescued shortened life span and vascular calcification in both *FGF23*-deficient mice and *Klotho*-deficient mice (Morishita et al., 2001; Stubbs et al., 2007). These studies have provided unequivocal evidence that the premature aging syndrome caused by defects in the *Klotho*-*FGF23* endocrine axis is due to retention of phosphate, calcium, and/or vitamin D. It should be noted that low phosphate diet rescued multiple phenotypes of *FGF23*-deficient mice despite the fact that it further increased already high serum calcium and vitamin D levels (Stubbs et al., 2007), suggesting that phosphate, but not calcium or vitamin D, is primarily responsible for the aging-like phenotypes. It is likely that low vitamin D diet and ablation of vitamin D activity rescued accelerated aging through reducing serum phosphate levels, although it remains to be determined whether high serum vitamin D and/or calcium levels are required for phosphate to induce the premature aging syndrome.

5. Phosphate and aging

Inorganic phosphate is not only an essential component of cell structure (DNA and membrane phospholipids) but also a key mediator of numerous cellular activities, including energy metabolism (ATP production) and kinase-mediated signal transduction. It is also involved in pathophysiology of various disorders such as bone diseases, vascular calcification, and chronic kidney disease among others. Although little is known about direct effects of phosphate on aging, accumulating evidence has demonstrated that inorganic phosphate has significant impacts on glucose metabolism and oxidative stress, which potentially affect aging processes of any organism from yeast to human.

In *Saccharomyces cerevisiae*, phosphate starvation, as well as glucose starvation, induces cell cycle arrest and extension of chronological life span (Boer et al., 2008; Brauer et al., 2008). Several phosphate-responsive genes have been identified, including the *PHO85* gene that encodes a cyclin-dependent kinase whose activity is correlated with intracellular phosphate concentration. Pho85 is inactivated under low phosphate conditions, leading to a quiescent G₀-like state and prolonged survival (Wanke et al., 2005). Pho85 also functions as a negative regulator for a set of genes that are typically induced under glucose starvation (DeRisi et al., 1997; Mouillon and Persson, 2006). Thus, phosphate restriction causes metabolic changes similar to those induced by caloric (glucose) restriction in yeast.

In mammals, low phosphate diet causes changes in metabolism similar to those induced by diet restriction as discussed below. Animals under diet restriction reduce blood insulin levels to adapt reduced carbohydrate availability and alter expression of insulin-responsive genes, which leads to changes in glucose metabolism including increased gluconeogenesis and decreased glycolysis among others (Cao et al., 2001; Kayo et al., 2001; Lee et al., 1999; Masoro, 2006; Wetter et al., 1999). Although low phosphate diet does not reduce blood insulin levels, it indeed alters expression of insulin-responsive genes in a way similar to that induced by diet restriction (Xie et al., 1999; Xie et al., 2000), resulting in increased gluconeogenesis and decreased glycolysis as well. This may be partly explained by the fact that low phosphate diet induces moderate insulin resistance by unknown mechanisms (Haap et al., 2006; Paula et al., 1998). Thus, moderate insulin resistance induced by phosphate restriction, as well as hypoinsulinemia induced by diet restriction, attenuate intracellular insulin signaling activity and induce similar changes in insulin-responsive gene expression, resulting in a similar metabolic state.

Recent studies have shown that increased insulin resistance does not necessarily mean diabetes and short life span. Rather, it has become increasingly clear that adequate suppression of insulin-like signaling pathway is an evolutionarily conserved mechanism for anti-aging and life span extension. Reduction-of-function mutations in the genes encoding orthologs of insulin receptor, insulin receptor substrates (IRS), and PI3-kinase has been known to extend life span in *C. elegans* and *Drosophila* (Clancy et al., 2001; Kenyon, 2005; Kenyon et al., 1993; Morris et al., 1996; Tatar et al., 2001). In mammals, increased longevity is reported in mice lacking insulin receptor in adipose tissues (Bluher et al., 2003), mice heterozygous for a null allele of the insulin-like growth factor-1 (IGF-1) receptor gene (Holzenberger et al., 2003), mice lacking IRS-1 (Selman et al., 2008), mice lacking IRS-2 in the brain (Taguchi et al., 2007), and *dwarf* mice with impaired growth hormone (GH)-IGF-1 endocrine axis (Bartke and Brown-Borg, 2004; Brown-Borg et al., 1996; Flurkey et al., 2002). In humans, some centenarians show resistance to IGF-1, short stature, and high serum IGF-1 associated with loss-of-function mutations in the IGF-1 receptor gene (Suh et al., 2008). Importantly, some long-lived animals exhibit insulin resistance (Kurosu et al., 2005; Selman et al., 2008), indicating that increased insulin sensitivity is not a prerequisite for long life span and anti-aging. Although many long-lived animals indeed exhibit increased insulin sensitivity, it is always associated with hypoinsulinemia and attenuated insulin/IGF-1 signaling activity in tissues. Thus, attenuated insulin/IGF-1 signaling activity in tissues shows closer association with life span extension than increased insulin sensitivity.

In addition to its involvement in glucose metabolism and insulin sensitivity, inorganic phosphate is shown to increase oxidative stress both *in vivo* and *in vitro*. Phosphate retention caused by *Klotho* deficiency in mice results in cognition impairment due to increased oxidative damage and apoptosis in hippocampus neurons, which can be rescued by administration of an antioxidant (Nagai et al., 2003). Furthermore, human vascular endothelial cells cultured in high phosphate medium (2.5 mM) have higher levels of cellular reactive oxygen species (ROS) than those cultured in normal phosphate medium (1.0 mM) (Di Marco et al., 2008). These

observations have raised the possibility that hyperphosphatemia by itself may play a causative role in pathogenesis of endothelial dysfunction and cardiovascular disease in chronic kidney disease (CKD) patients, because phosphate retention is universally observed in CKD patients and blood phosphate levels higher than 2.5 mM are often observed in this population. In fact, recent studies have identified hyperphosphatemia as a potent risk of death in CKD patients, even when the blood phosphate levels are within the normal range (Goodman et al., 2000; Tonelli et al., 2005). Thus, controlling blood phosphate levels below 2.0 mM by prescribing low phosphate diet and/or phosphate binders has become an important therapeutic goal for CKD patients (Ganesh et al., 2001). Of note, the National Kidney Foundation task force has indicated that the cardiovascular mortality of a 35-year-old patient on dialysis is equivalent to that of an 80-year-old “healthy” individual, rendering CKD to be the most potent accelerator of vascular senescence (Meyer and Levey, 1998).

Furthermore, the American Heart Association announced that CKD should be included in the highest-risk group for cardiovascular disease (Sarnak et al., 2003). Importantly, Klotho expression is significantly reduced in kidneys of CKD patients (Koh et al., 2001). Thus, CKD may be viewed as a segmental progeroid syndrome associated with a state of Klotho deficiency and phosphate retention. Interestingly, patients with Hutchinson-Gilford progeria syndrome (HGPS) exhibit hyperphosphatemia and reduced fractional excretion of phosphate (Merideth et al., 2008; Ortiz, 2008), which are characteristic features of mice lacking Klotho or FGF23. Although hyperphosphatemia is not primarily responsible for the HGPS phenotypes, it may increase the cardiovascular mortality in HGPS patients.

Inorganic phosphate entered into the cell is transported into mitochondria, where it functions not only as a substrate of ATP synthase but also as a key regulator of oxidative phosphorylation. In fact, mitochondrial membrane potential ($\Delta\psi$), NADH concentration, and oxygen consumption in isolated mitochondria increase in a hyperbolic manner as extra-mitochondrial phosphate concentration increases (Bose et al., 2003). $\Delta\psi$ is known to positively correlate ROS production in the electron transport chain (Papa and Skulachev, 1997). In addition, high phosphate concentration enhances delivery of reducing equivalent to cytochrome c in Complex III in the electron transport chain, which also increases ROS production (Bose et al., 2003). Thus, changes in mitochondrial function induced by high phosphate can increase ROS generation in mitochondria. The effects of inorganic phosphate on mitochondrial ROS generation likely have physiological significance, because cytoplasmic phosphate levels in mouse and human tissues as determined by ^{31}P nuclear magnetic resonance (NMR) indeed change from <1 mM to >10 mM depending on tissue types and their metabolic states (Katz et al., 1988, 1989). Furthermore, inorganic phosphate induces mitochondrial permeability transition (the inner membrane becomes non-selectively permeable to small solutes) in Ca^{2+} -loaded mitochondria, which occurs under pathological settings including ischemia/reperfusion and triggers cell death (Kowaltowski et al., 2001). Thus, inorganic phosphate plays multiple roles in the regulation of oxidative stress and mitochondrial function in health and disease, which potentially affect aging processes.

Another piece of evidence in support of the notion that phosphate affects aging processes is an inverse correlation between longevity and serum phosphate levels in mammals (Fig. 2). Although there are difficulties in precisely estimating the longevity and “normal” serum phosphate levels, a close relationship is clearly observed between longevity and serum phosphate throughout diverse species.

6. Perspective

When compared with endocrine regulation of calcium metabolism, that of phosphate metabolism was poorly understood until quite recently. Identification of the bone-kidney

endocrine axis mediated by FGF23 and Klotho has greatly promoted our understanding on phosphate metabolism. Since disruption of phosphate homeostasis is universally observed in patients with chronic kidney disease and identified as a potent risk for death, the FGF23-Klotho endocrine axis has emerged as a novel therapeutic target of chronic kidney disease (CKD), which nearly 26 million adult Americans, or 13% of the US population, suffer from and is increasingly recognized as a global public health problem in aging society (Coresh et al., 2007). In addition, animal studies on FGF23- and Klotho-deficient mice have revealed a potential link between phosphate and aging in mammals. If phosphate retention not only increases mortality of CKD patients but also accelerates aging processes in general, several important hypotheses may be drawn. First, CKD may be viewed as a state of accelerated aging or a segmental progeroid syndrome caused by phosphate retention. Second, prescription of low phosphate diet and/or phosphate binders may be considered not only for CKD patients but also for various common age-related diseases. Lastly, prescription of low phosphate diet and/or phosphate binders may be considered even for healthy individuals to prevent aging and age-related diseases. Further studies are required to prove or disprove a novel concept that phosphate accelerates aging.

References

- Anamizu Y, Kawaguchi H, Seichi A, Yamaguchi S, Kawakami E, Kanda N, Matsubara S, Kuro-o M, Nabeshima Y, Nakamura K, et al. Klotho insufficiency causes decrease of ribosomal RNA gene transcription activity, cytoplasmic RNA and rough ER in the spinal anterior horn cells. *Acta Neuropathol (Berl)* 2005;109:457–466. [PubMed: 15834732]
- Asadi F, Rostami A, Asadian P, Pourkabar M. Serum biochemistry and hematology values and hemoglobin electrophoresis in Persian squirrels (*Sciurus anomalus*). *Vet Clin Pathol* 2007;36:188–191. [PubMed: 17523094]
- Bartke A, Brown-Borg H. Life extension in the dwarf mouse. *Curr Top Dev Biol* 2004;63:189–225. [PubMed: 15536017]
- Berndt T, Kumar R. Phosphatonins and the regulation of phosphate homeostasis. *Annu Rev Physiol* 2007;69:341–359. [PubMed: 17002592]
- Berndt T, Kumar R. Novel mechanisms in the regulation of phosphorus homeostasis. *Physiology (Bethesda)* 2009;24:17–25. [PubMed: 19196648]
- Bluher M, Kahn BB, Kahn CR. Extended longevity in mice lacking the insulin receptor in adipose tissue. *Science* 2003;299:572–574. [PubMed: 12543978]
- Boer VM, Amini S, Botstein D. Influence of genotype and nutrition on survival and metabolism of starving yeast. *Proc Natl Acad Sci U S A* 2008;105:6930–6935. [PubMed: 18456835]
- Bose S, French S, Evans FJ, Joubert F, Balaban RS. Metabolic network control of oxidative phosphorylation: multiple roles of inorganic phosphate. *J Biol Chem* 2003;278:39155–39165. [PubMed: 12871940]
- Brauer MJ, Huttenhower C, Airoidi EM, Rosenstein R, Matese JC, Gresham D, Boer VM, Troyanskaya OG, Botstein D. Coordination of growth rate, cell cycle, stress response, and metabolic activity in yeast. *Mol Biol Cell* 2008;19:352–367. [PubMed: 17959824]
- Brown-Borg HM, Borg KE, Meliska CJ, Bartke A. Dwarf mice and the ageing process [letter]. *Nature* 1996;384:33. [PubMed: 8900272]
- Cao SX, Dhahbi JM, Mote PL, Spindler SR. Genomic profiling of short- and long-term caloric restriction effects in the liver of aging mice. *Proc Natl Acad Sci U S A* 2001;98:10630–10635. [PubMed: 11535822]
- Clancy DJ, Gems D, Harshman LG, Oldham S, Stocker H, Hafen E, Leivers SJ, Partridge L. Extension of Life-Span by Loss of CHICO, a *Drosophila* Insulin Receptor Substrate Protein. *Science* 2001;292:104–106. [PubMed: 11292874]
- Coresh J, Selvin E, Stevens LA, Manzi J, Kusek JW, Eggers P, Van Lente F, Levey AS. Prevalence of chronic kidney disease in the United States. *JAMA* 2007;298:2038–2047. [PubMed: 17986697]

- DeRisi JL, Iyer VR, Brown PO. Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science* 1997;278:680–686. [PubMed: 9381177]
- Di Marco GS, Hausberg M, Hillebrand U, Rustemeyer P, Wittkowski W, Lang D, Pavenstadt H. Increased inorganic phosphate induces human endothelial cell apoptosis in vitro. *Am J Physiol Renal Physiol* 2008;294:F1381–F1387. [PubMed: 18385273]
- Dusso AS, Brown AJ, Slatopolsky E. Vitamin D. *Am J Physiol Renal Physiol* 2005;289:F8–28. [PubMed: 15951480]
- Feldhamer, GA.; Thompson, BC.; Chapman, JA. *Wild Mammals of North America: Biology, Management, and Conservation*. 2. Baltimore: JHU Press; 2003.
- Field, KJ.; Sibold, AL.; Suckow, MA. *The laboratory hamster & gerbil*. CRC Press; 1998.
- Flurkey K, Papaconstantinou J, Harrison DE. The Snell dwarf mutation *Pit1(dw)* can increase life span in mice. *Mech Ageing Dev* 2002;123:121–130. [PubMed: 11718806]
- Ganesh SK, Stack AG, Levin NW, Hulbert-Shearon T, Port FK. Association of elevated serum PO(4), Ca x PO(4) product, and parathyroid hormone with cardiac mortality risk in chronic hemodialysis patients. *J Am Soc Nephrol* 2001;12:2131–2138. [PubMed: 11562412]
- Goetz R, Beenken A, Ibrahimi OA, Kalinina J, Olsen SK, Eliseenkova AV, Xu C, Neubert T, Zhang F, Linhardt RJ, et al. Molecular Insights into the Klotho-Dependent, Endocrine Mode of Action of FGF19 Subfamily Members. *Mol Cell Biol* 2007;27:3417–3428. [PubMed: 17339340]
- Goodman WG, Goldin J, Kuizon BD, Yoon C, Gales B, Sider D, Wang Y, Chung J, Emerick A, Greaser L, et al. Coronary-artery calcification in young adults with end-stage renal disease who are undergoing dialysis. *N Engl J Med* 2000;342:1478–1483. [PubMed: 10816185]
- Gorbunova V, Bozzella MJ, Seluanov A. Rodents for comparative aging studies: from mice to beavers. *Age (Dordr)* 2008;30:111–119. [PubMed: 19424861]
- Haap M, Heller E, Thamer C, Tschritter O, Stefan N, Fritsche A. Association of serum phosphate levels with glucose tolerance, insulin sensitivity and insulin secretion in non-diabetic subjects. *Eur J Clin Nutr* 2006;60:734–739. [PubMed: 16391583]
- Harmer NJ, Pellegrini L, Chirgadze D, Fernandez-Recio J, Blundell TL. The crystal structure of fibroblast growth factor (FGF) 19 reveals novel features of the FGF family and offers a structural basis for its unusual receptor affinity. *Biochemistry* 2004;43:629–640. [PubMed: 14730967]
- Heard DJ, Ruiz MM, Harr KE. Comparison of serum and plasma for determination of blood biochemical values in Malaysian flying foxes (*Pteropus vampyrus*). *J Zoo Wildl Med* 2006;37:245–248. [PubMed: 17319121]
- Hesse M, Frohlich LF, Zeitz U, Lanske B, Erben RG. Ablation of vitamin D signaling rescues bone, mineral, and glucose homeostasis in *Fgf-23* deficient mice. *Matrix Biol* 2007;26:75–84. [PubMed: 17123805]
- Holliday, R. *Understanding of ageing*. Cambridge: Cambridge University Press; 1995.
- Holzenberger M, Dupont J, Ducos B, Leneuve P, Geloën A, Even PC, Cervera P, Le Bouc Y. IGF-1 receptor regulates lifespan and resistance to oxidative stress in mice. *Nature* 2003;421:182–187. [PubMed: 12483226]
- Ishii M, Yamaguchi Y, Yamamoto H, Hanaoka Y, Ouchi Y. Airspace enlargement with airway cell apoptosis in klotho mice: a model of aging lung. *J Gerontol A Biol Sci Med Sci* 2008;63:1289–1298. [PubMed: 19126841]
- Itoh N, Ornitz DM. Functional evolutionary history of the mouse *Fgf* gene family. *Dev Dyn* 2008;237:18–27. [PubMed: 18058912]
- Kamemori M, Ohyama Y, Kurabayashi M, Takahashi K, Nagai R, Furuya N. Expression of Klotho protein in the inner ear. *Hear Res* 2002;171:103–110. [PubMed: 12204354]
- Katz LA, Swain JA, Portman MA, Balaban RS. Intracellular pH and inorganic phosphate content of heart in vivo: a ³¹P-NMR study. *Am J Physiol* 1988;255:H189–196. [PubMed: 3394819]
- Katz LA, Swain JA, Portman MA, Balaban RS. Relation between phosphate metabolites and oxygen consumption of heart in vivo. *Am J Physiol* 1989;256:H265–274. [PubMed: 2912189]
- Kawaguchi H, Manabe N, Miyaura C, Chikuda H, Nakamura K, Kuro-o M. Independent impairment of osteoblast and osteoclast differentiation in klotho mouse exhibiting low-turnover osteopenia. *J Clin Invest* 1999;104:229–237. [PubMed: 10430604]

- Kayo T, Allison DB, Weindruch R, Prolla TA. Influences of aging and caloric restriction on the transcriptional profile of skeletal muscle from rhesus monkeys. *Proc Natl Acad Sci U S A* 2001;98:5093–5098. [PubMed: 11309484]
- Kenyon C. The Plasticity of Aging: Insights from Long-Lived Mutants. *Cell* 2005;120:449–460. [PubMed: 15734678]
- Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A C. elegans mutant that lives twice as long as wild type. *Nature* 1993;366:461–464. [PubMed: 8247153]
- Koh N, Fujimori T, Nishiguchi S, Tamori A, Shiomi S, Nakatani T, Sugimura K, Kishimoto T, Kinoshita S, Kuroki T, et al. Severely reduced production of klotho in human chronic renal failure kidney. *Biochem Biophys Res Commun* 2001;280:1015–1020. [PubMed: 11162628]
- Kowaltowski AJ, Castilho RF, Vercesi AE. Mitochondrial permeability transition and oxidative stress. *FEBS Lett* 2001;495:12–15. [PubMed: 11322939]
- Kuro-o M. Klotho as a regulator of fibroblast growth factor signaling and phosphate/calcium metabolism. *Curr Opin Nephrol Hypertens* 2006;15:437–441. [PubMed: 16775459]
- Kuro-o M. Endocrine FGFs and Klothos: emerging concepts. *Trends Endocrinol Metab* 2008;19:239–245. [PubMed: 18692401]
- Kuro-o M, Matsumura Y, Aizawa H, Kawaguchi H, Suga T, Utsugi T, Ohyama Y, Kurabayashi M, Kaname T, Kume E, et al. Mutation of the mouse klotho gene leads to a syndrome resembling ageing. *Nature* 1997;390:45–51. [PubMed: 9363890]
- Kurosu H, Kuro-o M. The Klotho gene family and the endocrine fibroblast growth factors. *Curr Opin Nephrol Hypertens* 2008;17:368–372. [PubMed: 18660672]
- Kurosu H, Kuro-o M. Endocrine fibroblast growth factors as regulators of metabolic homeostasis. *Biofactors* 2009a;35:52–60. [PubMed: 19319846]
- Kurosu H, Kuro-o M. The Klotho gene family as a regulator of endocrine fibroblast growth factors. *Mol Cell Endocrinol* 2009b;299:72–78. [PubMed: 19063940]
- Kurosu H, Ogawa Y, Miyoshi M, Yamamoto M, Nandi A, Rosenblatt KP, Baum MG, Schiavi S, Hu MC, Moe OW, et al. Regulation of fibroblast growth factor-23 signaling by klotho. *J Biol Chem* 2006;281:6120–6123. [PubMed: 16436388]
- Kurosu H, Yamamoto M, Clark JD, Pastor JV, Nandi A, Gurnani P, McGuinness OP, Chikuda H, Yamaguchi M, Kawaguchi H, et al. Suppression of Aging in Mice by the Hormone Klotho. *Science* 2005;309:1829–1833. [PubMed: 16123266]
- Lee CK, Klopp RG, Weindruch R, Prolla TA. Gene Expression Profile of Aging and Its Retardation by Caloric Restriction. *Science* 1999;285:1390–1393. [PubMed: 10464095]
- Liu S, Gupta A, Quarles LD. Emerging role of fibroblast growth factor 23 in a bone-kidney axis regulating systemic phosphate homeostasis and extracellular matrix mineralization. *Curr Opin Nephrol Hypertens* 2007;16:329–335. [PubMed: 17565275]
- Liu S, Quarles LD. How fibroblast growth factor 23 works. *J Am Soc Nephrol* 2007;18:1637–1647. [PubMed: 17494882]
- Liu S, Tang W, Zhou J, Stubbs JR, Luo Q, Pi M, Quarles LD. Fibroblast growth factor 23 is a counter-regulatory phosphaturic hormone for vitamin D. *J Am Soc Nephrol* 2006;17:1305–1315. [PubMed: 16597685]
- Masoro EJ. Dietary restriction-induced life extension: a broadly based biological phenomenon. *Biogerontology* 2006;7:153–155. [PubMed: 16732403]
- Merideth MA, Gordon LB, Clauss S, Sachdev V, Smith AC, Perry MB, Brewer CC, Zalewski C, Kim HJ, Solomon B, et al. Phenotype and course of Hutchinson-Gilford progeria syndrome. *N Engl J Med* 2008;358:592–604. [PubMed: 18256394]
- Meyer KB, Levey AS. Controlling the epidemic of cardiovascular disease in chronic renal disease: report from the National Kidney Foundation Task Force on cardiovascular disease. *J Am Soc Nephrol* 1998;9:S31–42. [PubMed: 11443766]
- Min D, Panoskaltsis-Mortari A, Kuro-o M, Hollander GA, Blazar BR, Weinberg KI. Sustained thymopoiesis and improvement in functional immunity induced by exogenous KGF administration in murine models of aging. *Blood* 2007;109:2529–2537. [PubMed: 17138819]

- Mohammadi M, Olsen SK, Goetz R. A protein canyon in the FGF-FGF receptor dimer selects from an a la carte menu of heparan sulfate motifs. *Curr Opin Struct Biol* 2005a;15:506–516. [PubMed: 16154740]
- Mohammadi M, Olsen SK, Ibrahimi OA. Structural basis for fibroblast growth factor receptor activation. *Cytokine Growth Factor Rev* 2005b;16:107–137. [PubMed: 15863029]
- Moreau B, Vie JC, Cotellon P, De Thoisy I, Motard A, Raccourt CP. Hematologic and serum biochemistry values in two species of free-ranging porcupines (*Coendou prehensilis*, *Coendou melanurus*) in French Guiana. *J Zoo Wildl Med* 2003;34:159–162. [PubMed: 12885133]
- Morishita K, Shirai A, Kubota M, Katakura Y, Nabeshima Y, Takeshige K, Kamiya T. The progression of aging in *klotho* mutant mice can be modified by dietary phosphorus and zinc. *J Nutr* 2001;131:3182–3188. [PubMed: 11739863]
- Morris JZ, Tissenbaum HA, Ruvkun G. A phosphatidylinositol-3-OH kinase family member regulating longevity and diapause in *Caenorhabditis elegans*. *nature* 1996;382:536–539. [PubMed: 8700226]
- Mouillon JM, Persson BL. New aspects on phosphate sensing and signalling in *Saccharomyces cerevisiae*. *FEMS Yeast Res* 2006;6:171–176. [PubMed: 16487340]
- Munson L, Koehler JW, Wilkinson JE, Miller RE. Vesicular and ulcerative dermatopathy resembling superficial necrolytic dermatitis in captive black rhinoceroses (*Diceros bicornis*). *Vet Pathol* 1998;35:31–42. [PubMed: 9545133]
- Murzin AG, Lesk AM, Chothia C. beta-Trefoil fold. Patterns of structure and sequence in the Kunitz inhibitors interleukins-1 beta and 1 alpha and fibroblast growth factors. *J Mol Biol* 1992;223:531–543. [PubMed: 1738162]
- Nagai T, Yamada K, Kim HC, Kim YS, Noda Y, Imura A, Nabeshima Y, Nabeshima T. Cognition impairment in the genetic model of aging *klotho* gene mutant mice: a role of oxidative stress. *Faseb J* 2003;17:50–52. [PubMed: 12475907]
- Ohnishi M, Nakatani T, Lanske B, Razzaque MS. Reversal of mineral ion homeostasis and soft-tissue calcification of *klotho* knockout mice by deletion of vitamin D 1alpha-hydroxylase. *Kidney Int* 2009;75:1166–1172. [PubMed: 19225558]
- Ortiz A. Hutchinson-Gilford progeria syndrome. *N Engl J Med* 2008;358:2410. author reply 2410–2411. [PubMed: 18512277]
- Papa S, Skulachev VP. Reactive oxygen species, mitochondria, apoptosis and aging. *Mol Cell Biochem* 1997;174:305–319. [PubMed: 9309704]
- Passeri G, Vescovini R, Sansoni P, Galli C, Franceschi C, Passeri M. Calcium metabolism and vitamin D in the extreme longevity. *Exp Gerontol* 2008;43:79–87. [PubMed: 17698310]
- Paula FJ, Plens AE, Foss MC. Effects of hypophosphatemia on glucose tolerance and insulin secretion. *Hormone and Metabolic Research* 1998;30:281–284. [PubMed: 9660090]
- Pugh, DG. *Sheep & goat medicine*. Elsevier Health Sciences; 2002.
- Quarles LD. FGF23, PHEX, and MEPE regulation of phosphate homeostasis and skeletal mineralization. *Am J Physiol Endocrinol Metab* 2003;285:E1–9. [PubMed: 12791601]
- Quarles LD. Endocrine functions of bone in mineral metabolism regulation. *J Clin Invest* 2008;118:3820–3828. [PubMed: 19033649]
- Ramsay, EC. *Ursidae and Hyanidae*. In: Fowler, ME.; Miller, RE., editors. *Zoo and Wild Animal Medicine*. St Louis: Saunders; 2003. p. 523-538.
- Ravel, R. *Clinical laboratory medicine: clinical application of laboratory data*. 6. Elsevier Health Sciences; 1994.
- Razzaque MS, Sitara D, Taguchi T, St-Arnaud R, Lanske B. Premature aging-like phenotype in fibroblast growth factor 23 null mice is a vitamin D-mediated process. *FASEB J* 2006;20:720–722. [PubMed: 16436465]
- Sarnak MJ, Levey AS, Schoolwerth AC, Coresh J, Culleton B, Hamm LL, McCullough PA, Kasiske BL, Kelepouris E, Klag MJ, et al. Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. *Circulation* 2003;108:2154–2169. [PubMed: 14581387]

- Sato A, Hirai T, Imura A, Kita N, Iwano A, Muro S, Nabeshima Y, Suki B, Mishima M. Morphological mechanism of the development of pulmonary emphysema in klotho mice. *Proc Natl Acad Sci U S A* 2007;104:2361–2365. [PubMed: 17284608]
- Schiavi SC, Kumar R. The phosphatonin pathway: new insights in phosphate homeostasis. *Kidney Int* 2004;65:1–14. [PubMed: 14675031]
- Schlessinger J, Plotnikov AN, Ibrahimi OA, Eliseenkova AV, Yeh BK, Yayon A, Linhardt RJ, Mohammadi M. Crystal structure of a ternary FGF-FGFR-heparin complex reveals a dual role for heparin in FGFR binding and dimerization. *Mol Cell* 2000;6:743–750. [PubMed: 11030354]
- Segawa H, Kawakami E, Kaneko I, Kuwahata M, Ito M, Kusano K, Saito H, Fukushima N, Miyamoto K. Effect of hydrolysis-resistant FGF23-R179Q on dietary phosphate regulation of the renal type-II Na/Pi transporter. *Pflugers Arch* 2003;446:585–592. [PubMed: 12851820]
- Segawa H, Yamanaka S, Ohno Y, Onitsuka A, Shiozawa K, Aranami F, Furutani J, Tomoe Y, Ito M, Kuwahata M, et al. Correlation between hyperphosphatemia and type II Na-Pi cotransporter activity in klotho mice. *Am J Physiol Renal Physiol* 2007;292:F769–779. [PubMed: 16985213]
- Selman C, Lingard S, Choudhury AI, Batterham RL, Claret M, Clements M, Ramadani F, Okkenhaug K, Schuster E, Blanc E, et al. Evidence for lifespan extension and delayed age-related biomarkers in insulin receptor substrate 1 null mice. *FASEB J* 2008;22:807–818. [PubMed: 17928362]
- Shimada T, Hasegawa H, Yamazaki Y, Muto T, Hino R, Takeuchi Y, Fujita T, Nakahara K, Fukumoto S, Yamashita T. FGF-23 is a potent regulator of vitamin D metabolism and phosphate homeostasis. *J Bone Miner Res* 2004a;19:429–435. Epub 2003 Dec 2029. [PubMed: 15040831]
- Shimada T, Kakitani M, Yamazaki Y, Hasegawa H, Takeuchi Y, Fujita T, Fukumoto S, Tomizuka K, Yamashita T. Targeted ablation of Fgf23 demonstrates an essential physiological role of FGF23 in phosphate and vitamin D metabolism. *J Clin Invest* 2004b;113:561–568. [PubMed: 14966565]
- Shimada T, Muto T, Urakawa I, Yoneya T, Yamazaki Y, Okawa K, Takeuchi Y, Fujita T, Fukumoto S, Yamashita T. Mutant FGF-23 responsible for autosomal dominant hypophosphatemic rickets is resistant to proteolytic cleavage and causes hypophosphatemia in vivo. *Endocrinology* 2002;143:3179–3182. [PubMed: 12130585]
- Shimada T, Urakawa I, Yamazaki Y, Hasegawa H, Hino R, Yoneya T, Takeuchi Y, Fujita T, Fukumoto S, Yamashita T. FGF-23 transgenic mice demonstrate hypophosphatemic rickets with reduced expression of sodium phosphate cotransporter type IIa. *Biochem Biophys Res Commun* 2004c;314:409–414. [PubMed: 14733920]
- Stubbs JR, Liu S, Tang W, Zhou J, Wang Y, Yao X, Quarles LD. Role of Hyperphosphatemia and 1,25-Dihydroxyvitamin D in Vascular Calcification and Mortality in Fibroblastic Growth Factor 23 Null Mice. *J Am Soc Nephrol* 2007;18:2116–2124. [PubMed: 17554146]
- Suga T, Kurabayashi M, Sando Y, Ohyama Y, Maeno T, Maeno Y, Aizawa H, Matsumura Y, Kuwaki T, Kuro-o M, et al. Disruption of the klotho gene causes pulmonary emphysema in mice. Defect in maintenance of pulmonary integrity during postnatal life. *Am J Respir Cell Mol Biol* 2000;22:26–33. [PubMed: 10615062]
- Suh Y, Atzmon G, Cho MO, Hwang D, Liu B, Leahy DJ, Barzilai N, Cohen P. Functionally significant insulin-like growth factor I receptor mutations in centenarians. *Proc Natl Acad Sci U S A* 2008;105:3438–3442. [PubMed: 18316725]
- Taguchi A, Wartschow LM, White MF. Brain IRS2 signaling coordinates life span and nutrient homeostasis. *Science* 2007;317:369–372. [PubMed: 17641201]
- Tatar M, Kopelman A, Epstein D, Tu MP, Yin CM, Garofalo RS. A Mutant Drosophila Insulin Receptor Homolog That Extends Life-Span and Impairs Neuroendocrine Function. *Science* 2001;292:107–110. [PubMed: 11292875]
- Thrall, MA.; Baker, DC.; Campbell, TW.; Lassen, ED.; DeNicola, DB.; Rebar, A.; Fettman, MJ.; Weiser, G. *Veterinary Hematology and Clinical Chemistry: Text and Clinical Case Presentations Set*. Hoboken: Wiley-Blackwell; 2004.
- Tonelli M, Sacks F, Pfeffer M, Gao Z, Curhan G. Relation between serum phosphate level and cardiovascular event rate in people with coronary disease. *Circulation* 2005;112:2627–2633. [PubMed: 16246962]

- Tsujikawa H, Kurotaki Y, Fujimori T, Fukuda K, Nabeshima Y. Klotho, a gene related to a syndrome resembling human premature aging, functions in a negative regulatory circuit of vitamin D endocrine system. *Mol Endocrinol* 2003;17:2393–2403. [PubMed: 14528024]
- Tuntasuvan, D.; Theeraphan, A.; Phoengpong, N.; Jitnupong, W.; Lungka, G. In: Baker, I; Kashio, M., editors. Comparison of serum chemistry values and serum mineral values between captive and free-ranging elephants in Thailand; Giants on Our Hands: Proceedings of the International Workshop on the Domesticated Asian Elephant; Bangkok: RAP publication; 2002. p. 278
- Urakawa I, Yamazaki Y, Shimada T, Iijima K, Hasegawa H, Okawa K, Fujita T, Fukumoto S, Yamashita T. Klotho converts canonical FGF receptor into a specific receptor for FGF23. *Nature* 2006;444:770–774. [PubMed: 17086194]
- Wanke V, Pedruzzi I, Cameroni E, Dubouloz F, De Virgilio C. Regulation of G0 entry by the Pho80-Pho85 cyclin-CDK complex. *EMBO J* 2005;24:4271–4278. [PubMed: 16308562]
- Wetter TJ, Gazdag AC, Dean DJ, Cartee GD. Effect of calorie restriction on in vivo glucose metabolism by individual tissues in rats. *American Journal Of Physiology* 1999;276:E728–738. [PubMed: 10198310]
- White KE, Carn G, Lorenz-Depiereux B, Benet-Pages A, Strom TM, Econs MJ. Autosomal-dominant hypophosphatemic rickets (ADHR) mutations stabilize FGF-23. *Kidney Int* 2001;60:2079–2086. [PubMed: 11737582]
- White KE, Evans WE, O'Riordan JLH, Speer MC, Econs MJ, Lorenz-Deplereux B, Grabowski M, Meitinger T, Storm TM. Autosomal dominant hypophosphatemic rickets is associated with mutations in FGF23. *Nat Genet* 2000;26:345–348. [PubMed: 11062477]
- Xie W, Li Y, Mechin MC, Van De Werve G. Up-regulation of liver glucose-6-phosphatase in rats fed with a P(i)-deficient diet. *Biochem J* 1999;343(Pt 2):393–396. [PubMed: 10510305]
- Xie W, Tran TL, Finegood DT, van de Werve G. Dietary P(i) deprivation in rats affects liver cAMP, glycogen, key steps of gluconeogenesis and glucose production. *Biochem J* 2000;352(Pt 1):227–232. [PubMed: 11062077]
- Yahav S, Buffenstein R, Pettifor JM. Calcium and inorganic phosphorus metabolism in naked mole rats *Heterocephalus glaber* is only indirectly affected by cholecalciferol. *Gen Comp Endocrinol* 1993;89:161–166. [PubMed: 8381370]
- Yamashita T, Yoshioka M, Itoh N. Identification of a novel fibroblast growth factor, FGF-23, preferentially expressed in the ventrolateral thalamic nucleus of the brain. *Biochem Biophys Res Commun* 2000;277:494–498. [PubMed: 11032749]
- Yoshida T, Fujimori T, Nabeshima Y. Mediation of unusually high concentrations of 1,25-dihydroxyvitamin D in homozygous klotho mutant mice by increased expression of renal 1 α -hydroxylase gene. *Endocrinology* 2002;143:683–689. [PubMed: 11796525]
- Yu X, Sabbagh Y, Davis SI, Demay MB, White KE. Genetic dissection of phosphate- and vitamin D-mediated regulation of circulating Fgf23 concentrations. *Bone* 2005;36:971–977. [PubMed: 15869926]

Figure 1.

The bone-kidney endocrine axes mediated by FGF23 and Klotho. In osteocytes, active form of vitamin D (1,25-dihydroxyvitamin D₃) binds to vitamin D receptor (VDR) and forms heterodimers with another nuclear receptor (RXR) to transactivate transcription of FGF23 gene. Phosphate (P) also increases expression of FGF23 in bone, but its mechanism remains to be determined. FGF23 secreted from bone acts on the Klotho-FGFR complex expressed in renal epithelium in the kidney. As a phosphaturic hormone, FGF23 inhibits transepithelial phosphate reabsorption by suppressing sodium-phosphate co-transporter type-IIa (NaPi-2a) on the apical brush-border membrane of renal tubules, thereby closing a negative feedback loop for phosphate homeostasis. As a counter-regulatory hormone for vitamin D, FGF23 reduces

serum levels of 1,25-dihydroxyvitamin D₃ through suppressing its synthesis by down-regulating expression of the *Cyp27b1* gene and promoting its inactivation by up-regulating expression of the *Cyp24* gene, thereby closing a negative feedback loop for vitamin D homeostasis.

Figure 2.

Relation between longevity and serum phosphate in mammals. 1: *Klotho*^{-/-} mouse, 2: Mouse, 3: Rat, 4: Hamster, 5: Gerbil, 6: Nutria, 7: Rabbit, 8: Guinea pig, 9: Sheep, 10: Squirrel, 11: Porcupine, 12: Naked mole rat, 13: Flying fox, 14: Bear, 15: Rhinoceros, 16: Elephant, 17: Human, 18: Human (centenarian). Serum phosphate levels are average or median values, whichever available in literatures (Asadi et al., 2007; Feldhamer et al., 2003; Field et al., 1998; Gorbunova et al., 2008; Heard et al., 2006; Holliday, 1995; Kuro-o et al., 1997; Moreau et al., 2003; Munson et al., 1998; Passeri et al., 2008; Pugh, 2002; Ramsay, 2003; Segawa et al., 2007; Thrall et al., 2004; Tuntasuvan et al., 2002; Yahav et al., 1993).