Published in final edited form as:

Dev Neuropsychol. 2010 May ; 35(3): 233–256. doi:10.1080/87565641003689556.

Longitudinal Study of Callosal Microstructure in the Normal Adult Aging Brain Using Quantitative DTI Fiber Tracking

Edith V. Sullivan¹, Torsten Rohlfing², and Adolf Pfefferbaum^{1,2}

¹Department of Psychiatry and Behavioral Sciences, Stanford University School of Medicine, Stanford, CA

²Neuroscience Program, SRI International, Menlo Park, CA

Abstract

We present a review of neuroimaging studies of normal adult aging conducted with Diffusion Tensor Imaging (DTI) and data from one of the first longitudinal studies using DTI to study normal aging. To date, virtually all DTI studies of normal adult aging have been cross-sectional and have identified several patterns of white matter microstructural sparing and compromise that differentiate regional effects, fiber type, and diffusivity characteristics: 1) fractional anisotropy (FA) is lower and mean diffusivity is higher in older than younger adults, 2) aging is characterized by an anterior-to-posterior gradient of greater-to-lesser compromise also seen in superior-to-inferior fiber systems, and 3) association fibers connecting cortical sites appear to be more vulnerable to aging than projection fibers. The results of this longitudinal study of the macrostructure and microstructure of the corpus callosum yielded a consistent pattern of differences between healthy, young (20s to 30s) and elderly (60s to 70s) men and women without change over 2 years. We then divided the fibers of the corpus callosum into the midsagittal strip and the lateral distal fibers in an attempt to identify the locus of the age-related differences. The results indicated that, on average, mean values of FA and longitudinal diffusivity (λ_L) were lower in the distal than midsagittal fibers in both groups, but the age effects and the anterior-to-posterior gradients were more pronounced for the distal than midsagittal fibers and extended more posteriorly in the distal than midsagittal fibers. Despite lack of evidence for callosal aging over 2 years, ventricular enlargement occurred and was disproportionately greater in the elderly relative to the young group, being 8.2% in the elderly but only 1.2% in the young group. Thus, different brain regions can express different rates of change with aging. Our longitudinal DTI data indicate that normal aging is associated with declining FA and increasing diffusivity in both λ_L (longitudinal diffusivity) and λ_T (transverse diffusivity), perhaps defining the normal ontological condition rather than a pathological one, which can be marked by low FA and low diffusivity.

There is little doubt that even healthy aging is marked by decline in sensory, motor, and selective cognitive functions, well delineated by hundreds of careful, quantitative, cross-sectional and longitudinal assessments (for volumes of reviews, Birren & Schaie, 2001; Craik & Salthouse, 2008). Less known are the neural mechanisms responsible for involutional changes in function, but quantitative in vivo neuroimaging studies conducted over the last quarter century provide critical leads to identify the selective brain structures and systems that contribute to functional decline.

We and others have recently reviewed the neuroimaging findings in normal adult aging (Madden, Bennet & Song, 2009; Sullivan & Pfefferbaum, 2007, 2009; Pfefferbaum & Sullivan, 2009, 2010; Zahr, Pfefferbaum & Sullivan, 2009) based on magnetic resonance imaging (MRI),

capable of yielding quantification of macrostructural characteristics (notably, size and shape) of brain tissue, and diffusion tensor imaging (DTI), designed to yield quantification of microstructural characteristics of brain tissue, typically white matter axonal and myelin integrity. Many other reviews on the radiologically-identified structural characteristics of normal aging brain are also available, and most are cross-sectional in design. The goal of identifying specific brain mechanisms determining age-related declines in selective cognitive, sensory, and motor processes is a worthy neuropsychological goal in its own right and also forms an essential context for characterizing abnormalities of neurodegenerative diseases and other conditions affecting brain tissue integrity.

The purpose of this paper is to provide an overview of our current knowledge of white matter changes detected with neuroimaging across the adult age span. We focus on microstructural changes observed with DTI but start our review with a brief summary of macrostructural changes observed with MRI in order to give a context for both levels of analysis of neuromorphological ontological change. In addition, we present new data from a longitudinal DTI study of normal aging.

Differential Effects of Age on Brain Volume of Gray Matter and White Matter: MRI

Throughout development and aging, the brain undergoes extensive volume growth measurable with MRI. Quantitative cross-sectional MRI study of brain ontology in humans, ages 3 months to 70 years, implied growth of the cortical gray matter compartment until about age 5 years, followed by a linear volume decline (Pfefferbaum et al., 1994) until very old age when an accelerated decline occurs (Fotenos, Mintun, Snyder, Morris, & Buckner, 2008). A different pattern emerged for cortically subjacent white matter volume, which showed growth acceleration during adolescence to asymptote in the third decade; during the same time, CSF spaces, including sulci and ventricles, expanded continually (Pfefferbaum et al., 1994).

A majority of cross-sectional and longitudinal MRI studies in the adult age range report systematic age-related volume enlargement in CSF-filled spaces that occurs at the expense of cortical gray matter and with little volume change in white matter (cross-sectional Blatter et al., 1995; Courchesne et al., 2000; Good et al., 2001; Pfefferbaum et al., 1994; Raz et al., 1997; Smith, Chebrolu, Wekstein, Schmitt, & Markesbery, 2007; Sowell, Thompson, & Toga, 2004; Sullivan, Deshmukh, Desmond, Lim, & Pfefferbaum, 2000; Sullivan, Rosenbloom, Serventi, & Pfefferbaum, 2004; Taki et al., 2006; D. Tisserand, Van Boxtel, Gronenschild, & Jolles, 2001); (longitudinal Liu et al., 2003; Pfefferbaum, Sullivan, Rosenbloom, Mathalon, & Lim, 1998; Raz, Rodrigue, Kennedy, & Acker, 2007; Resnick, Pham, Kraut, Zonderman, & Davatzikos, 2003). A minority of studies has reported the opposite pattern of tissue shrinkage, with greater age-related volume decline in white matter than gray matter (Guttmann et al., 1998; Jernigan et al., 2001). When regional white matter volume does show age-related loss, it is typically small, estimated at 1% per year decline in midsagittal area of the corpus callosum of elderly men examined over a 4-year span (Sullivan, Pfefferbaum, Adalsteinsson, Swan, & Carmelli, 2002) and 2% per decade in a neuropathology study (Miller, Alston, & Corsellis, 1980); (but see Jernigan et al., 2001; Walhovd et al., 2005)]. In contrast to volumetric results, microstructural study of white matter with DTI appears to be more sensitive to aging's degenerative effects, as evidenced in the consistency of findings reported and reviewed below.

On MRI, neither the pons nor the corpus callosum show substantial age-related volume decline with age (Driesen & Raz, 1995; Raz, Gunning-Dixon, Head, Williamson, & Acker, 2001; Sullivan et al., 2002; Sullivan et al., 2004). Use of DTI to measure regional white matter fiber bundle volume revealed differential effects of age, with significant linear declines detected in the corona radiata, anterior cingulum, regions of the fornix and cerebellar peduncle and

nonlinear decline in the genu in healthy individuals, age 13 to 70 years (Pagani, Agosta, Rocca, Caputo, & Filippi, 2008). Visual inspection of corpora callosa of healthy elderly individuals can suggest excessive thinning, especially of the isthmus, in the elderly, that is actually the result of ventricular expansion and not necessarily callosal atrophy (Pfefferbaum, Sullivan, & Carmelli, 2001, 2004). Thus, caution must be exercised when interpreting callosal thinning in conditions, such as normal pressure hydrocephalus, marked by ventriculomegaly. Regional volume shrinkage, whether in gray matter or white matter, tends to accelerate in very old age (Raz et al., 2005; Salat, Kaye, & Janowsky, 1999) and may reflect heterogeneity arising from common occult conditions, such as preclinical or undetected dementia, hypertension, metabolic disorders, or alcoholism. Indeed, the Leukoaraiosis And DISability (LANDIS) study identified 569 elderly men and women with mild to severe subcortical white matter hyperintensities (Fazekas, Chawluk, Alavi, Hurtig, & Zimmerman, 1987) and found significant atrophy of the corpus callosum that was associated with poor scores on the Mini-Mental State Examination, a short physical performance battery, and walking speed (Ryberg et al., 2007) and selective relations between anterior but not posterior callosal atrophy and deficits in attention and executive functions (Jokinen et al., 2007).

The dynamic nature of aging makes it particularly desirable to track its course with longitudinal examination. Such reports based on conventional MRI have been accumulating over the past decade (e.g., Fotenos et al., 2008; Kramer et al., 2007; Raz et al., 2005; Resnick et al., 2003); some suggest that cross-sectional study underestimates the toll of age on the brain (for review, Raz & Rodrigue, 2006). One longitudinal study showed that accelerated decline in whole brain volume correlated with higher socioeconomic status, which was interpreted as an indication that individuals with higher status have greater “reserve” than those with lower status and therefore express cognitive decline later despite accelerating signs of brain shrinkage (Fotenos et al., 2008). Yet another longitudinal study found no protective effects of educational attainment on preservation of brain volumes (Raz et al., 2005). Although the loci and extent of age-related tissue volume decline may differ from study to study, whether due to differences in study cohort, image acquisition parameters, or image data analysis, the most consistent finding is a salient, rapid decline of prefrontal tissue volume (Liu et al., 2003; Pfefferbaum et al., 1998; Raz et al., 2004; Resnick et al., 2003; Tang, Whitman, Lopez, & Baloh, 2001; Tisserand et al., 2002), possibly contributing to (if not accounting for) parallel declines in component processes of executive functioning, such as working memory (Leung, Gore, & Goldman-Rakic, 2002; Park et al., 1996), sequencing (Allain et al., 2007) and temporal ordering (Fuster, 2000), response inhibition (Müller-Oehring, Schulte, Raassi, Pfefferbaum, & Sullivan, 2007), error monitoring (Wang, Ulbert, Schomer, Marinkovic, & Halgren, 2005), stimulus evaluation response latency (Pfefferbaum, Ford, Wenegrat, Roth, & Kopell, 1984), and attention allocation (Madden, Pierce, & Allen, 1992), each drawing on adequate prefrontal functioning (sample of reviews, Buckner, 2004; Fuster, 2000; Tisserand & Jolles, 2003; Y. Wang et al., 2006). In contrast with the MRI literature, longitudinal DTI studies of microstructural integrity in normal aging are only recently emerging, and this report adds to this new area of study.

Differential Effects of Age on Regional White Matter Microstructure: DTI

MR diffusion weighted imaging (DWI) and diffusion tensor imaging (DTI) allow quantification of microscopic movement of water molecules modeled as Brownian motion *within* each voxel of an image (Basser, 1995; Moseley et al., 1990). Because of the linear organization of the brain’s fiber bundles, fasciculi, and commissures, DTI has been useful in characterizing the microstructural condition and constituents of white matter. The principles of DTI are presented in a number of overviews (e.g., Bammer, Acar, & Moseley, 2003; Le Bihan, 2003, 2007; Mori & Zhang, 2006; Pfefferbaum & Sullivan, 2005a) and are summarized next.

In regions with few or no constraints imposed by physical boundaries, such as cerebrospinal fluid (CSF) in the ventricles, water movement is random, that is, freely diffusing, and is therefore isotropic. By contrast, the water molecule path, for example, in a white matter fiber, is constrained by the physical boundaries, such as the axon sheath, causing the movement to be greater along the long axis of the fiber than across it and is anisotropic, typically measured as fractional anisotropy (FA) and ranging between 0 and 1 on a normalized scale (Pierpaoli & Basser, 1996). Thus, DTI is selectively sensitive to the detection of tightly packed fibers in parallel orientation of white matter systems.

The tensor is associated with three eigenvalues ($\lambda_1, \lambda_2, \lambda_3$), each corresponding to one of three mutually orthogonal orientational eigenvectors, describing the diffusion ellipsoid by its major axes (Figure 1). The eigenvalue average, which is equal to the trace of the tensor, reflects the magnitude of diffusion. The largest eigenvalue, λ_1 , is the longitudinal diffusivity, λ_L , whereas λ_2 and λ_3 quantify transverse diffusivity, $\lambda_T = (\lambda_2 + \lambda_3)/2$. The extent to which one eigenvalue, λ_1 , dominates the other two, λ_2 and λ_3 , determines the degree of anisotropy within a voxel. DTI data sets are commonly reduced to an anisotropy image and a mean diffusivity image, but the utility of decomposing mean diffusivity into λ_L and λ_T has been shown in studies of the developing neonate and children and normal aging as well as in tracking neurological conditions, including stroke (C. Wang et al., 2006), head injury (Sidaros et al., 2008; Wilde et al., 2006), multiple sclerosis (Bonzano et al., 2008), and alcohol (Pfefferbaum, Adalsteinsson, & Sullivan, 2006a, 2006b) and drug dependence (Moeller et al., 2007).

Fractional anisotropy (FA)

In less than a decade, quantitative DTI has expanded our understanding of the toll aging takes on the brain's white matter microstructure. Quantitative DTI has revealed degradation of white matter microstructure (in terms of diffusion coherence on an intravoxel basis) undetectable with bulk volume (in terms of the average of many voxels of a specific tissue type) measures from conventional MRI in normal aging (Hugenschmidt et al., 2008; Pfefferbaum & Sullivan, 2003; Sullivan et al., 2001) and neuropathology (Herve et al., 2005; multiple sclerosis: Kolind et al., 2008; alcoholism Pfefferbaum & Sullivan, 2002; Sidaros et al., 2008; stroke: C. Wang et al., 2006; head injury Wilde et al., 2006). Most consistently observed has been a decline in FA and complementary increase in diffusivity in white matter with advancing age (Chun, Filippi, Zimmerman, & Ulug, 2000; Head et al., 2004; Madden et al., 2004; Nusbaum, Tang, Buchsbaum, Wei, & Atlas, 2001; O'Sullivan et al., 2001; Ota et al., 2006; Pfefferbaum, Sullivan, Hedehus, Lim et al., 2000; Salat et al., 2005; Stebbins et al., 2001). This aging pattern is similar in men and women (Ota et al., 2006; Sullivan et al., 2001), although regional variation may occur. Hsu et al. (Hsu et al., 2008) reported differential age-related declines in FA, where men showed a steeper decline in a global measure of FA and in FA of the right anterior limb of the internal capsule than women, whereas the FA decline was steeper in women than men in the right deep temporal white matter. Although the FA decline with age is linear from about 20 years onwards, the rise in diffusivity is not and accelerates in older age (Pfefferbaum, Rosenbloom, Adalsteinsson, & Sullivan, 2007).

One of the most robust findings describing age-related differences in regional FA has been a distribution of low FA selective to frontal white matter in the elderly (Ardekani, Kumar, Bartzokis, & Sinha, 2007a; Bhagat & Beaulieu, 2004; Bucur et al., 2007; Foong et al., 2000; Grieve, Williams, Paul, Clark, & Gordon, 2007; Head et al., 2004; Madden et al., 2007; Madden et al., 2004; Nusbaum et al., 2001; O'Sullivan et al., 2001; Pfefferbaum, Adalsteinsson, & Sullivan, 2005; Pfefferbaum & Sullivan, 2003; Pfefferbaum, Sullivan, Hedehus, Lim et al., 2000; D. H. Salat et al., 2005; Sullivan et al., 2001; Takahashi et al., 2004) that was confirmed in a monkey model of aging (Makris et al., 2007). This anterior-posterior gradient, where anterior fiber systems are more susceptible to age-related compromise than are posterior

systems, has been repeatedly replicated in the corpus callosum but also holds true for lateralized fiber bundles and endures whether measured with region-of-interest analysis (O'Sullivan et al., 2001; Pfefferbaum et al., 2005; Pfefferbaum & Sullivan, 2003; Sullivan et al., 2001), voxel-based approach (Ardekani, Kumar, Bartzokis, & Sinha, 2007b; Hsu et al., 2008; Salat et al., 2005), or quantitative fiber tracking (Pfefferbaum et al., 2007; Sullivan et al., 2001; Sullivan, Rohlfing, & Pfefferbaum, 2009).

Diffusivity

The typical aging pattern of white matter microstructure is characterized by a decrease in intravoxel anisotropy (FA) accompanied by an increase in diffusivity (Chen, Li, & Hindmarsh, 2001; Engelter, Provenzale, Petrella, DeLong, & MacFall, 2000; Head et al., 2004; Helenius et al., 2002; Naganawa, Sato, Katagiri, Mimura, & Ishigaki, 2003; Pfefferbaum et al., 2005; Pfefferbaum & Sullivan, 2003). Modeling age's effect on FA and diffusivity across 120 adults, 20 to 80 years, revealed a linear decline in FA and nonlinear increase in diffusivity with age that was greater in the genu than splenium of the corpus callosum (Pfefferbaum et al., 2007). Diffusivity can be inflated by partial voluming from non-white matter tissue, such as gray matter and cerebrospinal fluid, both characterized by lower FA and higher diffusivity than white matter (Bhagat & Beaulieu, 2004). Yet, even when controlling for partial voluming by eroding peripheral voxels of a particular region of interest, that is, by removing voxels in the periphery of white matter regions most likely to contain signal from non-white matter tissue, the complementary aging functions—FA decrease with diffusivity increase—are observable (Pfefferbaum & Sullivan, 2002; Pfefferbaum & Sullivan, 2003).

The magnitude of the FA-diffusivity relationship varies across brain regions and is greater in older than younger individuals (Pfefferbaum & Sullivan, 2003). This relationship suggests that decreased brain white matter intravoxel coherence is attributable, at least in part, to the accumulation of interstitial or intracellular fluid, or both fluid compartments (e.g., Norris, Niendorf, & Leibfritz, 1994; Pfefferbaum & Sullivan, 2005b; Rumpel, Ferrini, & Martin, 1998; Sehy, Ackerman, & Neil, 2002; Silva et al., 2002) and may reflect age-related loosening of myelin, dense cytoplasm, and formation of fluid-filled balloons, which were observed in area 46 white matter in nonhuman primate models of normal aging (Peters & Sethares, 2003).

Tractography

FA is a measure of the magnitude and orientation of diffusion derived from the tensor's eigenvalues on an *intravoxel* basis. By contrast, coherence measures, including tractography, provide an orientational measure on an *intervoxel* basis, that is, the degree to which the diffusion orientation of a voxel is similar to its neighbors. The use of voxel-to-voxel coherence measures serves the conceptual basis for quantitative fiber tracking (Fillard & Gerig, 2003; Gerig, Corouge, Vachet, Krishnan, & MacFall, 2005). Although the connectivity and coherence between different brain regions on vector and fiber tracking maps is readily apparent on visual inspection, these displays have only recently been subjected to quantification (Gerig et al., 2005; Mori et al., 2002; Xu, Mori, Solaiyappan, van Zijl, & Davatzikos, 2002; Xue, van Zijl, Crain, Solaiyappan, & Mori, 1999). Methods for quantitative analysis of structural connectivity of white matter include fiber-tract trajectories (Basser & Pierpaoli, 1998; Masutani, Aoki, Abe, Hayashi, & Otomo, 2003; Mori & van Zijl, 2002) and maps of the degree of "alignment" among neighboring vectors, that is, on a voxel-to-voxel basis, resulting in a measure of intervoxel coherence (Jones, Simmons, Williams, & Horsfield, 1999; Pfefferbaum, Sullivan, Hedehus, Adalsteinsson et al., 2000). In addition to providing useful visual depiction of empirically derived fiber tracts (Schmahmann et al., 2007), advantages of quantitative fiber tracking over focal region of interest analysis include the ability to measure the entire extent of a fiber tract and to characterize its integrity along its full extent.

In an initial study using quantitative fiber tracking in normal aging, we observed higher diffusivity and fewer imaging-defined fiber bundles in the anterior but not posterior segments of the corpus callosum, based on known interhemispheric projection sites (Pandya & Seltzer, 1986) in elderly compared with young, healthy men and women (Sullivan, Adalsteinsson, & Pfefferbaum, 2006). Examination of regionally distinct fiber bundles identified throughout the supratentorium and infratentorium replicated the anterior-posterior gradient of age-related degradation of white matter quality in a cohort of 120 healthy men and women, who spanned the adult age range from 20 to 81 years and extended the pattern to a superior-inferior gradient (Sullivan et al., 2009). In this case, quantitative fiber tracking revealed lower anisotropy and higher diffusivity in older than younger healthy individuals and in superior than inferior bundles (longitudinal fasciculi, cingulate bundles), but no age effect in pontine or cerebellar fiber systems. Robust sex differences were not identified in this study of adult aging nor were they forthcoming in a developmental study of regional fiber systems using quantitative tractography in children, age 6 to 17 years (Eluvathingal, Hasan, Kramer, Fletcher, & Ewing-Cobbs, 2007); however, sex differences have been reported in voxel-based analysis (Schneiderman et al., 2007).

Stadlbauer et al. (Stadlbauer, Salomonowitz, Strunk, Hammen, & Ganslandt, 2008) used fiber tracking to examine three different categories of fiber systems: association fibers, which included the superior longitudinal, inferior longitudinal, and inferior fronto-occipital fasciculi; callosal fibers; and projection fibers, which included corticobulbar and corticospinal tracts and thalamic fibers. The age-related declines in FA were graded from greatest in the association fibers, less so in the callosal fibers, and least in the projection fibers. Nonetheless, significant increases in mean diffusivity were present in all three, fiber systems, indicating that none was fully immune to involutional changes. Another fiber tracking study examined 38 healthy men and women, age 18 to 88 years, and found age-related FA decline and diffusivity increase in ADC and each of the three separate lambdas (eigenvalues) in the fornix but not the cingulate bundles (Stadlbauer, Salomonowitz, Strunk, Hammen, & Ganslandt, 2007). Using quantitative fiber tracking in groups of healthy young and elderly men and women, Zahr et al. (Zahr, Rohlfing, Pfefferbaum, & Sullivan, 2009) identified an anterior-posterior and superior-inferior gradient of age-related degradation in fiber bundles, notable in the genu, fornix, and uncinate fibers; functional relations were observed between Working Memory Problem Solving, and Motor factor scores and DTI metrics indicating regionally compromised fiber tracts.

Brain structure-function relationships

The functional ramifications of the DTI metrics have been regularly verified with observations of correlations between low FA or high diffusivity and poor cognitive (Bucur et al., 2007; Charlton et al., 2007; Grieve et al., 2007; Madden et al., 2007; O'Sullivan et al., 2001; Shenkin et al., 2003; Stebbins et al., 2001) or motor (Sullivan et al., 2001) test performance in healthy aging men and women. Some studies provide evidence for selectivity of DTI-behavioral relationships. Low FA in frontal white matter correlated with low scores on tests of executive functions assessed with a visual odd-ball task (Madden et al., 2004). This study also revealed an age-related difference in regional FA relationships with reaction time, where the relationship in the older but not younger adults was selective to the internal capsule but not the splenium, whereas the younger adults showed the opposite relationship. We found a selective relation between performance on alternated finger tapping (but not the control condition of unimanual finger tapping) and FA in the splenium and perisplenial white matter in healthy men and women (Sullivan et al., 2001). A cross-sectional developmental DTI study of 92 children to young adults, age 9 to 24 years, found an age-related increase in FA of the splenium and that this increase, possibly reflecting further myelination, was also predictive of speed in the alternated finger tapping, which requires interhemispheric coordination of bimanual movements (Muetzel et al., 2008).

Other studies report nonspecific relationships. For example, Grieve et al. (Grieve et al., 2007) showed that low FA in three separate regions—frontal, temporal, and parietal white matter—correlated with faster maze completion time in a group of 87 healthy volunteers, age 20 to 73 years; attentional shifting accuracy was correlated with FA from frontal to occipital regions. Lower whole brain FA and mean diffusivity were predictive of working memory performance (Charlton et al., 2007; Charlton et al., 2006). Significant correlations were observed between high anterior white matter mean diffusivity and prolonged time to complete the Trail Making test, whereas high diffusivity was related to verbal fluency (O’Sullivan et al., 2001); unfortunately, selectivity of these relationships was not tested. A monkey model of aging provided evidence for age-related decrease in FA of association fibers, namely, the superior longitudinal fasciculus and cingulate bundle, and also the anterior corpus callosum, that also correlated with a cognitive set shifting task (Makris et al., 2007).

Brain structure-function relationships using quantitative fiber tracking have also been established in healthy adults. One report provided evidence for a relationship between Stroop word-color naming and central callosal FA integrity (Sullivan et al., 2006). In another study, fine finger movement speed correlated with FA and transverse diffusivity measured in three lateral fiber bundles supporting motor movement (internal capsules, external capsules, and cerebellar hemisphere bundles) but no commissural fiber tract metrics (Sullivan et al., 2009).

Longitudinal Study of Microstructure vs. Macrostructure of the Corpus Callosum

DTI studies of normal aging have largely relied on cross-sectional examination of healthy men and women drawn from either contrasting age groups (young vs. elderly adults) or a continuous age distribution. The lack of longitudinal results limits generalization of available studies (c.f., Raz et al., 2007; Rohlfing, Sullivan, & Pfefferbaum, 2006). The few studies reporting longitudinal results in normal aging do so secondarily, in that the target study groups were individuals with neurological conditions, including head injury (Sidaros et al., 2008) and amyotrophic lateral sclerosis (ALS) (Blain et al., 2007). In neither control group were age-related declines in FA or increases in diffusivity detected.

Here, we present new longitudinal DTI and MRI data collected over a 2-year interval in a sample of healthy young and elderly men and women. In follow-up to our initial study on the cross-sectional comparison of quantitative fiber tracking in this cohort (Sullivan et al., 2006), we predicted that DTI would reveal an anterior-posterior gradient of microstructural decline with age that would be even greater in the follow-up session in the older group. We also expected that DTI would be more sensitive to degenerative changes with age than size measures derived from MRI. We also examined anisotropy and diffusivity after dividing the callosal fibers into midsagittal and bilateral distal components. Given the consistency of ventricular expansion with age, we measured the ventricular system as a positive comparison brain structure at both times on MRI, anticipating that expansion over the inter-scanning interval would occur even if aging effects were not evident macroscopically on MRI or microscopically on DTI.

METHOD

Participants

The 10 men and 10 women who served in our earlier aging study were contacted, on average, 2 years later and invited to participate in a follow-up study. Re-contacting these 20 volunteers revealed that one man and one woman from the young group had moved too far away for a return visit and that two elderly volunteers had died; one man died of cancer unknown at MRI

1, and one woman was fatally hit as a pedestrian by a car. The remaining participants formed a balanced, two-group design of healthy, highly educated adults of 4 young and 4 elderly men and 4 young and 4 elderly women: 8 young (mean $\pm$ SD=28.6 $\pm$ 5.2, range=22 to 37 years at MRI 1; mean $\pm$ SD=31.1 $\pm$ 5.3, range=24 to 40 years at MRI 2; 17.9 $\pm$ 1.6 years of education) and 8 older (mean $\pm$ SD=72.7 $\pm$ 5.2, range=65 to 79 years at MRI 1; mean $\pm$ SD=74.6 $\pm$ 5.1, range=67 to 81 years at MRI 2; 16.0 $\pm$ 1.5 years of education). The younger subjects included laboratory members and men and women recruited from the local community. All older subjects were recruited from a larger ongoing study of normal aging and scored well within the normal range (above 25 out of 30) on the Mini-Mental State Examination (Folstein, Folstein, & McHugh, 1975) at both scanning sessions: mean $\pm$ SD=29.0 $\pm$ 1.3, range=27 to 30 at MRI 1; mean $\pm$ SD=28.6 $\pm$ 1.1, range=27 to 30 at MRI 2. The time between MRI 1 and MRI 2 was 2.5 $\pm$.22 years for the young group and 1.9 $\pm$.25 years for the elderly group.

DTI and MRI Acquisition

The DTI and structural data were acquired on a 3T MRI scanner: 1) structural Fast Spin Echo (FSE); 2) an Inversion Recovery Prepared SPOiled Gradient Recalled echo (IRPrepSPGR); 3) Diffusion Tensor Images (DTI) with 6 non-collinear diffusion directions repeated with opposite gradient polarity, and 4) a field map used for correction of spatial distortion due to main field (B_0) inhomogeneity. The SPGR data were aligned such that adjacent pairs of 1.25 mm thick SPGR slices subtended each 2.5 mm thick FSE and DTI slice.

DTI Analysis

DTI quantification was preceded by eddy current correction on a slice-by-slice basis by within-slice registration, which took advantage of the symmetry of the opposing polarity acquisition (Bodammer, Kaufmann, Kanowski, & Tempelmann, 2004) and also allowed for compensation of the diffusion effect created by the imaging gradients (Neeman, Freyer, & Sillerud, 1991), reducing the data to 6 non-collinear diffusion-weighted images per slice. Using the field maps, B_0 -field inhomogeneity-induced geometric distortion in the eddy current-corrected images was corrected with PRELUDE (Phase Region Expanding Labeller for Unwrapping Discrete Estimates, (Jenkinson, 2003)) and FUGUE (FMRIB's Utility for Geometrically Unwarping EPIs (Jenkinson, 2001)). These "native" DTI data were used for fiber tracking. From the $b=0$ and 6 diffusion weighted images, 6 maps of the apparent diffusion coefficient (ADC) were calculated. Solving 6 simultaneous equations with respect to ADC_{xx}, ADC_{xy}, etc. yielded the elements of the diffusion tensor. The diffusion tensor was then diagonalized, yielding eigenvalues $\lambda_1, \lambda_2, \lambda_3$, as well as eigenvectors that define the predominant diffusion orientation. Based on the eigenvalues, FA and ADC were calculated on a voxel-by-voxel basis (Basser & Jones, 2002; Basser & Pierpaoli, 1998; Pierpaoli & Basser, 1996).

Fiber Tracking

To achieve common anatomical coordinates across subjects, FA data for each subject were aligned with a laboratory standard average brain FA template (Rohlfing, Zahr, Sullivan, & Pfefferbaum, 2008) using nonrigid registration (Rohlfing, Brandt, Menzel, & Maurer, 2004; Rohlfing & Maurer, 2003). The midsagittal corpus callosum (the target region of interest for fiber tracking) was identified with an interactive program on the laboratory standard, as were parallel planes 10mm bilaterally (the sources for fiber tracking). For fiber tracking, the target and sources were then warped to the corresponding locations on the native basis images for each subject with a numerical inversion of the subject-to-standard transformation. The tensor matrix, targets, and sources were passed to the fiber tracking routine in native space.

Fiber tracking was performed with the software distributed by Gerig et al. (Gerig et al., 2005) based on the method of Mori and colleagues (Mori & van Zijl, 2002; Xu et al., 2002; Xue et al., 1999). Fiber tracking parameters included white matter extraction threshold

(minimum FA) of .17, minimum fiber length of 37.5 mm, maximum fiber length of 187.5 mm, fiber tracking threshold of .125 (that is, .125 is the minimum FA of a voxel allowable in a fiber tract), and maximum voxel-to-voxel coherence minimum transition smoothness threshold of .80 (~37° maximum deviation between fiber segments from neighboring voxels), with no limit on the number of fibers. Identified fibers were required to pass through both sources to ensure identification of callosal fibers that extended to both hemispheres. The mean FA and ADC of all voxels comprising each fiber, for all fibers, were determined. After fiber detection the fiber locations were transformed back to common standard coordinates for display and further analysis. In common space, the midsagittal corpus callosum was divided geometrically into 6 regions of interest (Figure 2) but guided by the callosal anatomical projections described by (Pandya & Seltzer, 1986). For each callosal region the number of fibers, the mean fiber length, FA, ADC, λ_L , and λ_T were determined. We refer hereafter to the fibers coursing through each of the six callosal regions as “fiber bundles” following the Pandya and Selzer convention: prefrontal, premotor, precentral, postcentral, posterior parietal, and temporal-occipital (Figure 2).

MRI Quantification

The volume of the ventricular system and the midsagittal volume of the corpus callosum were determined by a parcellation method (Pfefferbaum, Rosenbloom et al., 2006). To achieve common anatomical coordinates for brain structure, the IRPrepSPGR data for each subject were aligned with a laboratory standard average structural template (Rohlfing et al., 2008) using nonrigid registration (Rohlfing et al., 2004; Rohlfing & Maurer, 2003). The entire ventricular system and the midsagittal plus two bilateral, 1 mm thick, immediately parasagittal slices of the corpus callosum were outlined with a semiautomated routine on the laboratory standard. The regions of interest were then warped to the native locations on the native IRPrepSPGR for each subject using numerical inversion of the initial transformation, and the volumes computed by voxel count.

Statistical Analysis

Primary analyses were based on group-by-brain region-by-time analysis of variance (ANOVA) and follow-up analyses used t-tests. For single-level variables, such as age, repeated measures with one within-group factor was used. For multiple-level variables, such as 6 callosal sectors for FA or 2 cerebral hemispheres for ventricular volume, repeated measures with multiple within-group factors was used with Greenhouse-Geiser (GG) correction where appropriate.

RESULTS

DTI Metrics of Fiber Tracking

Mean $\pm$ S.E. for FA, ADC, λ_L , and λ_T for the two age groups at each scanning session are presented in Figure 2.

Fractional anisotropy (FA)—The group-by-sector-by-time ANOVA examining group differences in FA of the 6 callosal sectors over time revealed a group-by-sector interaction ($F(5,70)=10.084$, $p=.0001$ GG). Follow-up analysis indicated significantly lower FA in the elderly than young group in the four anterior sectors ($p=.0001$ to $.0477$). The premotor sector only showed a modest decline over time in both groups ($p=.0501$).

Diffusivity—The group-by-sector-by-time ANOVA for the diffusivity (ADC) identified one significant interaction: group-by-sector ($F(5,70)=8.2254$, $p=.0002$ GG). Follow-up analyses indicated significantly higher ADC in the elderly than young group in five of the six sectors ($p=.0044$ to $.0008$), the exception being the posterior parietal sector. Change over time was not significant in either age group. The pattern of group differences observed for ADC, with the

elderly showing greater diffusivity than the young in five callosal sectors, was the same for λ_L ($F(5,70)=5.508$, $p=.0021$ GG) and λ_T ($F(5,70)=9.467$, $p=.0001$ GG). In follow-up testing, the p -values of the group differences for λ_L ranged from .0026 to .0429, and for λ_T ranged from .0049 to .0001.

Midsagittal vs. distal fibers—In a further analysis, we divided the fibers of the corpus callosum into the midsagittal strip and the lateral distal fibers in an attempt to identify the location of the age differences. Accordingly, we conducted group-by-sector-by-time ANOVA for each DTI metric of the midsagittal and distal fibers. For FA, the group-by-sector interactions were significant for both sets of fibers (midsagittal: ($F(5,70)=5.134$, $p=.0032$ GG; distal: ($F(5,70)=13.452$, $p=.0001$ GG). Follow-up analyses together with inspection of Figure 3 reveal a more extensive effect of age in the distal than midsagittal fibers, extending from prefrontal to postcentral sectors for the distal fibers but only to the two most anterior sectors for the midsagittal fibers. Further, mean FA of the midsagittal fibers ranged from .55 to .75 but only .40 to .55 for FA of the distal fibers.

As observed for FA, the age effects and the anterior-to-posterior gradients were more pronounced for the distal than midsagittal fibers, although a time effect was not forthcoming (Figure 3). Again, the group-by-sector-by-time ANOVAs for each diffusivity measure showed a similar set of effects for the midsagittal and distal fibers: ADC: midsagittal: ($F(5,70)=3.126$, $p=.04$ GG; distal: ($F(5,70)=6.101$, $p=.0017$ GG; λ_L : midsagittal: ($F(5,70)=2.954$, $p=.0419$ GG; distal: ($F(5,70)=2.606$, $p=.0682$ GG; and λ_T : midsagittal: ($F(5,70)=3.833$, $p=.0173$ GG; distal: ($F(5,70)=9.031$, $p=.0001$ GG. On average, mean values of FA, ADC, and λ_L were lower and mean λ_T values higher in the distal than midsagittal fibers in both groups.

MRI Volumes of Macrostructure

Mean $\pm$ S.E. of the volumes of the corpus callosum and lateral ventricles for the two age groups at each scanning session are presented in Figure 4.

Corpus Callosum—Unlike the callosal DTI measures, the MRI measures of callosal volume did not differ significantly by group ($F(1,14)=1.957$, $p=.1836$) nor time ($F(1,14)=.144$, $p=.7102$) and showed no group-by-time interaction ($F(1,14)=.0093$, $p=.9245$).

Lateral Ventricles—The group-by-hemisphere-by-time ANOVA revealed significant effects of group ($F(1,14)=48.681$, $p=.0001$) and hemisphere ($F(1,14)=9.515$, $p=.0081$). Examples of age differences in ventricular volumes are presented in Figure 5. A group-by-time interaction ($F(1,14)=8.921$, $p=.0098$) revealed a disproportionately greater expansion of the lateral ventricles in the elderly than young group. Even though the significant hemisphere effect indicated that the right ventricle was larger than the left, this pattern held similarly for young and elderly individuals, and an absence of a group-by-hemisphere-by-time interaction ($F(1,14)=1.115$, $p=.3090$) indicated lack of a laterality effect in aging.

DISCUSSION

The results of this longitudinal study of the macrostructure and microstructure of the corpus callosum yielded a consistent pattern of differences between healthy, young (20s to 30s) and elderly (60s to 70s) men and women without change over 2 years. The microstructural result is consistent with other longitudinal DTI observations in controls, showing no change over 12 months (Sidaros et al., 2008) or over an average of 8 months (Blain et al., 2007). Even in a group of 215 elderly men (age 70 to 82 years) examined over a 4-year interval, we detected <1% decrease per year in the midsagittal area of the corpus callosum, whereas the lateral ventricles expanded 2.9% annually (Sullivan et al., 2002). Although this time interval may

have been too brief for detection of structural features of callosal aging, the interval was adequate for detection of ventricular enlargement, which was disproportionately greater in the elderly relative to the young group: the increase in the lateral ventricles was 8.2% in the elderly compared with 1.2% in the young group. Thus different brain regions express different rates of change with aging.

The analyses based on dividing the corpus callosum into a midsagittal component and a bilateral distal component revealed a pattern of anisotropy and diffusivity age differences similar to those observed across the entire extent of fibers tracked but more dramatic for the distal than midsagittal fibers. Specifically, the age effects and the anterior-to-posterior gradients were more robust for the distal than midsagittal fibers and extended more posteriorly in the distal than midsagittal fibers. On average, FA and λ_L values were lower in the distal than midsagittal fibers in both groups. The higher FA of the midsagittal fibers may reflect a greater linearity of microstructure than in the distal fibers, but the consistency of group differences and small variance indicate surprising homogeneity of fibers identified as they emerge from the midsagittal corpus callosum. Although fibers that extend into the centrum semiovale should be more susceptible to partial voluming from boggy tissue characteristic of white matter hyperintensities, the low diffusivity in the distal relative to the midsagittal fibers dispels this possibility. Indeed, the consistency of the age effects and replicability of the measurements over 2 years provide predictive validity and reliability to quantitative fiber tracking in studies of normal aging.

The frontal microstructural effect of age observed in the DTI results at both scan session comports with the mainstay of cross-sectional reports on the effects of normal age, indicating greater age-related compromise of frontal relative to posterior brain white matter. This anterior-posterior gradient may reflect the normal developmental pattern, where frontal sites develop relatively late (Sowell, Thompson, Holmes, Jernigan, & Toga, 1999), show the greatest vulnerability to functional decline in normal aging (Gunning-Dixon & Raz, 2003; Raz, Gunning-Dixon, Head, Dupuis, & Acker, 1998), and are under proportionately greater environmental than genetic control than are posterior regions (Pfefferbaum et al., 2001).

The regionally differential effects of age in white matter FA and diffusivity across brain regions is likely attributable to the structure of the underlying fibers (Barkovich, 2000; Peters & Sethares, 2003), which influence the homogeneity of fiber orientations within each voxel. This variation with age may also contribute to some reports of fewer fiber representations identified with DTI (Stadlbauer et al., 2008; Sullivan et al., 2006). This speculation must be tempered, however, by the actual size of white matter fibers imaged. Given the diameters of axons, for example in the corpus callosum ranging from 0.4 to 5 μm , with the preponderance less than 1 μm , it is estimated that a cross-section of the genu of the human corpus callosum contains approximately 400,000 fibers per mm^2 (Aboitiz, Scheibel, Fisher, & Zaidel, 1992). A single voxel of a DTI study using relatively high resolution ($2 \times 2 \times 2 \text{ mm}^3$) for current standards could contain 1.6 million fibers in a cross-section. Additionally, the number of fibers identified can be influenced by the size of the brain from the simple fact that larger brains have more voxels from which fibers can be mathematically constructed.

Taken together, the last decade of DTI studies has identified several patterns of white matter microstructural sparing and compromise in normal adult aging that differentiate regional effects, fiber type, and diffusivity characteristics: FA is lower and diffusivity is higher in older than younger adults. These aging patterns are regional, characterized by an anterior-to-posterior gradient of greater-to-lesser compromise also seen in superior-to-inferior fiber systems. Association fibers connecting cortical sites may be more vulnerable to aging than projection fibers, which are corticospinal and corticothalamic systems; commissural fiber systems show an anterior-posterior gradient, which is paralleled by postmortem investigations. These studies

reveal degradation of white matter microstructure, including degradation of myelin and microtubules (Kemper, 1994) and axon deletion (Aboitiz, Rodriguez, Olivares, & Zaidel, 1996; Meier-Ruge, Ulrich, Bruhlmann, & Meier, 1992), especially of myelinated fibers of the precentral gyrus and small connecting fibers of the anterior corpus callosum. Whether the freely diffusing water molecules characterized with DTI are from the intracellular or extracellular compartments remains controversial (e.g., Sen & Basser, 2005). Although the general pattern is that FA declines with advancing age in healthy adults, it must also be recognized that surprisingly high FA is not necessarily a sign of exceptional health. Indeed, selective deletion of uniformly-orientated white matter fibers from a tissue sample of crossing fibers (as can occur with Wallerian degeneration) causes abnormally high anisotropy (Pierpaoli et al., 2001). Therefore, interpretation of FA results requires guidance by knowledge of the underlying regional architecture, especially white matter, given current applications (Pierpaoli et al., 2001; Shimony et al., 1999; Virta, Barnett, & Pierpaoli, 1999).

Further refinement of the observed aging patterns is revealed by consideration of the separate contributions from the ellipsoid components of the apparent diffusion coefficient (i.e., mean diffusivity). Animal models of stroke, fiber crushing, and dysmyelination indicate that decline in longitudinal diffusivity, λ_L , reflects axonal injury, whereas increase in transverse or radial diffusivity, λ_T , reflects damaged myelin. In particular, data from a shiver mouse model suggests that dysmyelination results in decreased FA because of increased λ_T , leaving λ_L unaffected (Song et al., 2002). By contrast, traumatic injury results in decreased FA typically because of both increased λ_T and decreased λ_L (Nevo et al., 2001). Human developmental studies of neonates through late adolescence or young adulthood report increasing FA and initially decreasing λ_T , together indicative of myelination. Curiously, studies of the normal human aging adult report declining FA and increasing diffusivity in both λ_L and λ_T , perhaps defining the normal ontological condition rather than a pathological one.

Given the assertion that white matter lesions measured on structural MRI “account for *all* age-related declines in speed but not in intelligence” (Rabbitt et al., 2007), neuropsychological studies of normal aging need to consider the condition of white matter supporting connectivity of gray matter structures, which function as a circuit and underlie complex cognitive, sensory, and motor abilities. Quantitative DTI and fiber tracking can contribute to the characterization of these white matter fiber systems, serve as correlates and predictors of selective functional abilities, and suggest mechanisms of compromise and decline with age and disease. Ideally, conclusions about brain structure-function relationships require support from formal tests of double or multiple dissociations to establish selectivity of such relationships (c.f., Bates, Appelbaum, Salcedo, Saygin, & Pizzamiglio, 2003; Teuber, 1955).

Acknowledgments

This work was supported by NIH grants AG017919, AA005965, AA010723, AA012388, AA17168

References Cited

- Aboitiz F, Rodriguez E, Olivares R, Zaidel E. Age-related changes in fibre composition of the human corpus callosum: sex differences. *Neuroreport* 1996;7(11):1761–1764. [PubMed: 8905659]
- Aboitiz F, Scheibel AB, Fisher RS, Zaidel E. Fiber composition of the human corpus callosum. *Brain Research* 1992;598:143–153. [PubMed: 1486477]
- Allain P, Berrut G, Etcharry-Bouyx F, Barre J, Dubas F, Le Gall D. Executive functions in normal aging: an examination of script sequencing, script sorting, and script monitoring. *Journal of Gerontol B, Psychological Sciences and Social Sciences* 2007;62(3):P187–190.
- Ardekani S, Kumar A, Bartzokis G, Sinha U. Exploratory voxel-based analysis of diffusion indices and hemispheric asymmetry in normal aging. *Magnetic Resonance Imaging* 2007a;25(2):154–167. [PubMed: 17275609]

- Ardekani S, Kumar A, Bartzokis G, Sinha U. Exploratory voxel-based analysis of diffusion indices and hemispheric asymmetry in normal aging. *Magnetic Resonance Imaging* 2007b;25(2):154–167. [PubMed: 17275609]
- Bammer R, Acar B, Moseley ME. In vivo MR tractography using diffusion imaging. *European Journal of Radiology* 2003;45(3):223–234. [PubMed: 12595107]
- Barkovich AJ. Concepts of myelin and myelination in neuroradiology. *American Journal of Neuroradiology* 2000;21(6):1099–1109.
- Basser PJ. Inferring microstructural features and the physiological state of tissues from diffusion-weighted images. *NMR in Biomedicine* 1995;8(7-8):333–344. [PubMed: 8739270]
- Basser PJ, Jones DK. Diffusion-tensor MRI: theory, experimental design and data analysis - a technical review. *NMR in Biomedicine* 2002;15(7-8):456–467. [PubMed: 12489095]
- Basser PJ, Pierpaoli C. A simplified method to measure the diffusion tensor from seven MR images. *Magnetic Resonance in Medicine* 1998;39(6):928–934. [PubMed: 9621916]
- Bates E, Appelbaum M, Salcedo J, Saygin AP, Pizzamiglio L. Quantifying dissociations in neuropsychological research. *Journal of Clinical and Experimental Neuropsychology* 2003;25(8):1128–1153. [PubMed: 14566586]
- Bhagat YA, Beaulieu C. Diffusion anisotropy in subcortical white matter and cortical gray matter: changes with aging and the role of CSF-suppression. *Journal of Magnetic Resonance Imaging* 2004;20(2):216–227. [PubMed: 15269946]
- Birren, JE.; Schaie, KW. *Handbook of the Psychology of Aging*. 5th Edition. Academic Press; San Diego: 2001.
- Blain CR, Williams VC, Johnston C, Stanton BR, Ganesalingam J, Jarosz JM, et al. A longitudinal study of diffusion tensor MRI in ALS. *Amyotrophic Lateral Sclerosis* 2007;8:348–355. [PubMed: 17924235]
- Blatter DD, Bigler ED, Gale SD, Johnson SC, Anderson C, Burnett BM, et al. Quantitative volumetric analysis of brain MRI: Normative database spanning five decades of life. *American Journal of Neuroradiology* 1995;16(2):241–245. [PubMed: 7726068]
- Bodammer N, Kaufmann J, Kanowski M, Tempelmann C. Eddy current correction in diffusion-weighted imaging using pairs of images acquired with opposite diffusion gradient polarity. *Magnetic Resonance in Medicine* 2004;51(1):188–193. [PubMed: 14705060]
- Bonzano L, Tacchino A, Roccatagliata L, Abbruzzese G, Mancardi GL, Bove M. Callosal contributions to simultaneous bimanual finger movements. *Journal of Neuroscience* 2008;28(12):3227–3233. [PubMed: 18354026]
- Buckner RL. Memory and executive function in aging and AD: multiple factors that cause decline and reserve factors that compensate. *Neuron* 2004;44(1):195–208. [PubMed: 15450170]
- Bucur B, Madden DJ, Spaniol J, Provenzale JM, Cabeza R, White LE, et al. Age-related slowing of memory retrieval: Contributions of perceptual speed and cerebral white matter integrity. *Neurobiology of Aging*. 2007
- Charlton R, Landau S, Schiavone F, Barrick TR, Clark CA, Markus HS, et al. A structural equation modeling investigation of age-related variance in executive function and DTI measured white matter damage. *Neurobiology of Aging* 2007;29(10):1547–1555. [PubMed: 17451845]
- Charlton RA, Barrick TR, McIntyre DJ, Shen Y, O'Sullivan M, Howe FA, et al. White matter damage on diffusion tensor imaging correlates with age-related cognitive decline. *Neurology* 2006;66(2):217–222. [PubMed: 16434657]
- Charlton RA, Landau S, Schiavone F, Barrick TR, Clark CA, Markus HS, et al. A structural equation modeling investigation of age-related variance in executive function and DTI measured white matter damage. *Neurobiology of Aging* 2007;29(10):1547–1555. [PubMed: 17451845]
- Chen ZG, Li TQ, Hindmarsh T. Diffusion tensor trace mapping in normal adult brain using single-shot EPI technique. A methodological study of the aging brain. *Acta Radiologica* 2001;42(5):447–458. [PubMed: 11552881]
- Chun T, Filippi CG, Zimmerman RD, Ulug AM. Diffusion changes in the aging human brain. *American Journal of Neuroradiology* 2000;21(6):1078–1083. [PubMed: 10871018]

- Courchesne E, Chisum HJ, Townsend J, Cowles A, Covington J, Egaas B, et al. Normal brain development and aging: quantitative analysis at in vivo MR imaging in healthy volunteers. *Radiology* 2000;216(3):672–682. [PubMed: 10966694]
- Craik, FIM.; Salthouse, TA. *The Handbook of Aging and Cognition*. Third ed.. Psychology Press; New York: 2008.
- Driesen NR, Raz N. The influence of sex, age, and handedness on corpus callosum morphology: a meta-analysis. *Psychobiology* 1995;23(3):240–247.
- Eluvathingal TJ, Hasan KM, Kramer L, Fletcher JM, Ewing-Cobbs L. Quantitative diffusion tensor tractography of association and projection fibers in normally developing children and adolescents. *Cerebral Cortex* 2007;17(12):2760–2768. [PubMed: 17307759]
- Engelter ST, Provenzale JM, Petrella JR, DeLong DM, MacFall JR. The effect of aging on the apparent diffusion coefficient of normal-appearing white matter. *American Journal of Roentgenology* 2000;175(2):425–430. [PubMed: 10915688]
- Fazekas F, Chawluk JB, Alavi A, Hurtig HI, Zimmerman RA. MR signal abnormalities at 1.5 T in Alzheimer's dementia and normal aging. *American Journal of Radiology* 1987;149:351–356.
- Fillard, P.; Gerig, G. Analysis Tool For Diffusion Tensor MRI; Paper presented at the Proceedings of Medical Image Computing and Computer-assisted Intervention, Lecture Notes in Computer Science; Saint-Malo, France. 2003;
- Folstein MF, Folstein SE, McHugh PR. Mini-mental state: A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research* 1975;12:189–198. [PubMed: 1202204]
- Foong J, Maier M, Clark C, Barker G, Miller D, Ron M. Neuropathological abnormalities of the corpus callosum in schizophrenia: a diffusion tensor imaging study. *Journal of Neurology Neurosurgery and Psychiatry* 2000;68(2):242–244.
- Fotenos AF, Mintun MA, Snyder AZ, Morris JC, Buckner RL. Brain volume decline in aging: evidence for a relation between socioeconomic status, preclinical Alzheimer disease, and reserve. *Archives of Neurology* 2008;65(1):113–120. [PubMed: 18195148]
- Fuster JM. Executive frontal functions. *Experimental Brain Research* 2000;133(1):66–70.
- Gerig, G.; Corouge, I.; Vachet, C.; Krishnan, KR.; MacFall, JR. Quantitative analysis of diffusion properties of white matter fiber tracts: a validation study; Paper presented at the 13th Proceedings of the International Society for Magnetic Resonance in Medicine; Miami, FL. 2005;
- Good CD, Johnsrude IS, Ashburner J, Henson RN, Friston KJ, Frackowiak RS. A voxel-based morphometric study of ageing in 465 normal adult brains. *NeuroImage* 2001;14:21–36. [PubMed: 11525331]
- Grieve SM, Williams LM, Paul RH, Clark CR, Gordon E. Cognitive aging, executive function, and fractional anisotropy: a diffusion tensor MR imaging study. *American Journal of Neuroradiology* 2007;28(2):226–235. [PubMed: 17296985]
- Gunning-Dixon FM, Raz N. Neuroanatomical correlates of selected executive functions in middle-aged and older adults: a prospective MRI study. *Neuropsychologia* 2003;41(14):1929–1941. [PubMed: 14572526]
- Guttmann CRG, Jolesz FA, Kikinis R, Killiany RJ, Moss MB, Sandor T, et al. White matter changes with normal aging. *Neurology* 1998;50:972–978. [PubMed: 9566381]
- Head D, Buckner RL, Shimony JS, Williams LE, Akbudak E, Conturo TE, et al. Differential vulnerability of anterior white matter in nondemented aging with minimal acceleration in dementia of the Alzheimer type: Evidence from Diffusion Tensor Imaging. *Cerebral Cortex* 2004;14(4):410–423. [PubMed: 15028645]
- Helenius J, Soinne L, Perkio J, Salonen O, Kangasmaki A, Kaste M, et al. Diffusion-weighted MR imaging in normal human brains in various age groups. *American Journal of Neuroradiology* 2002;23(2):194–199. [PubMed: 11847041]
- Herve D, Molko N, Pappata S, Buffon F, LeBihan D, Bousser MG, et al. Longitudinal thalamic diffusion changes after middle cerebral artery infarcts. *Journal of Neurology Neurosurgery and Psychiatry* 2005;76(2):200–205.

- Hsu JL, Leemans A, Bai CH, Lee CH, Tsai YF, Chiu HC, et al. Gender differences and age-related white matter changes of the human brain: A diffusion tensor imaging study. *Neuroimage* 2008;39(2):566–577. [PubMed: 17951075]
- Hugenschmidt CE, Peiffer AM, Kraft RA, Casanova R, Deibler AR, Burdette JH, et al. Relating imaging indices of white matter integrity and volume in healthy older adults. *Cerebral Cortex* 2008;18(2):433–442. [PubMed: 17575289]
- Jenkinson M. Improved unwarping of EPI volumes using regularised B₀ Maps (abs). *Human Brain Mapping - HBM200*. 2001
- Jenkinson M. A fast, automated, N-dimensional phase unwrapping algorithm. *Magnetic Resonance in Medicine* 2003;49:193–197. [PubMed: 12509838]
- Jernigan TL, Archibald SL, Fennema-Notestine C, Gamst AC, Stout JC, Bonner J, et al. Effects of age on tissues and regions of the cerebrum and cerebellum. *Neurobiology of Aging* 2001;22(4):581–594. [PubMed: 11445259]
- Jokinen H, Ryberg C, Kalska H, Ylikoski R, Rostrup E, Stegmann MB, et al. Corpus callosum atrophy is associated with mental slowing and executive deficits in subjects with age-related white matter hyperintensities: the LADIS Study. *Journal of Neurology, Neurosurgery, and Psychiatry* 2007;78(5):491–496.
- Jones D, Simmons A, Williams S, Horsfield M. Non-invasive assessment of axonal fiber connectivity in the human brain via diffusion tensor MRI. *Magnetic Resonance in Medicine* 1999;42:37–41. [PubMed: 10398948]
- Kemper, TL. Neuroanatomical and neuropathological changes during aging and dementia. In: Albert, ML.; Knoefel, JE., editors. *Clinical Neurology of Aging*. 2nd edition ed.. Oxford University Press; New York: 1994. p. 3-67.
- Kolind SH, Laule C, Vavasour IM, Li DK, Traboulsee AL, Madler B, et al. Complementary information from multi-exponential T(2) relaxation and diffusion tensor imaging reveals differences between multiple sclerosis lesions. *Neuroimage* 2008;40(1):77–85. [PubMed: 18226549]
- Kramer JH, Mungas D, Reed BR, Wetzel ME, Burnett MM, Miller BL, et al. Longitudinal MRI and cognitive change in healthy elderly. *Neuropsychology* 2007;21(4):412–418. [PubMed: 17605574]
- Le Bihan D. Looking into the functional architecture of the brain with diffusion MRI. *Nature Reviews Neuroscience* 2003;4(6):469–480.
- Le Bihan D. The ‘wet mind’: water and functional neuroimaging. *Physics and Medicine and Biology* 2007;52:R57–R90.
- Leung HC, Gore JC, Goldman-Rakic PS. Sustained mnemonic response in the human middle frontal gyrus during on-line storage of spatial memoranda. *J Cognitive Neuroscience* 2002;14(4):659–671.
- Liu RS, Lemieux L, Bell GS, Sisodiya SM, Shorvon SD, Sander JW, et al. A longitudinal study of brain morphometrics using quantitative magnetic resonance imaging and difference image analysis. *NeuroImage* 2003;20(1):22–33. [PubMed: 14527567]
- Madden DJ, Bennett IJ, Song AW. Cerebral white matter integrity and cognitive aging: Contributions from diffusion tensor imaging. *Neuropsychology Review*. 2009 in press.
- Madden DJ, Pierce TW, Allen PA. Adult age differences in attentional allocation during memory search. *Psychology of Aging* 1992;7(4):594–601.
- Madden DJ, Spaniol J, Whiting WL, Bucur B, Provenzale JM, Cabeza R, et al. Adult age differences in the functional neuroanatomy of visual attention: a combined fMRI and DTI study. *Neurobiology of Aging* 2007;28(3):459–476. [PubMed: 16500004]
- Madden DJ, Whiting WL, Huettel SA, White LE, MacFall JR, Provenzale JM. Diffusion tensor imaging of adult age differences in cerebral white matter: relation to response time. *NeuroImage* 2004;21(3):1174–1181. [PubMed: 15006684]
- Makris N, Papadimitriou GM, van der Kouwe A, Kennedy DN, Hodge SM, Dale AM, et al. Frontal connections and cognitive changes in normal aging rhesus monkeys: A DTI study. *Neurobiology of Aging* 2007;28(10):556–567.
- Masutani Y, Aoki S, Abe O, Hayashi N, Otomo K. MR diffusion tensor imaging: recent advance and new techniques for diffusion tensor visualization. *European Journal of Radiology* 2003;46(1):53–66. [PubMed: 12648802]

- Meier-Ruge W, Ulrich J, Bruhlmann M, Meier E. Age-related white matter atrophy in the human brain. *Annals of the New York Academy of Sciences* 1992;673:260–269. [PubMed: 1485724]
- Miller AKH, Alston RL, Corsellis JAN. Variations with age in the volumes of grey and white matter in the cerebral hemispheres of man: measurements with an image analyzer. *Neuropathology and Applied Neurobiology* 1980;6:119–132. [PubMed: 7374914]
- Moeller FG, Hasan KM, Steinberg JL, Kramer LA, Valdes I, Lai LY, et al. Diffusion tensor imaging eigenvalues: preliminary evidence for altered myelin in cocaine dependence. *Psychiatry Research Neuroimaging* 2007;154(3):253–258.
- Mori S, Kaufmann WE, Davatzikos C, Stieltjes B, Amodei L, Fredericksen K, et al. Imaging cortical association tracts in the human brain using diffusion-tensor-based axonal tracking. *Magnetic Resonance in Medicine* 2002;47(2):215–223. [PubMed: 11810663]
- Mori S, van Zijl PC. Fiber tracking: principles and strategies - a technical review. *NMR in Biomedicine* 2002;15(7-8):468–480. [PubMed: 12489096]
- Mori S, Zhang J. Principles of diffusion tensor imaging and its applications to basic neuroscience research. *Neuron* 2006;51:527–539. [PubMed: 16950152]
- Moseley ME, Mintonovitch J, Cohen Y, Asgari HS, Derugin N, Norman D, et al. Early detection of ischemic injury: comparison of spectroscopy, diffusion-, T2-, and magnetic susceptibility-weighted MRI in cats. *Acta Neurochirurgica Supplementa* 1990;51:207–209.
- Muetzel RL, Collins PF, Mueller BA, A MS, Lim KO, Luciana M. The development of corpus callosum microstructure and associations with bimanual task performance in healthy adolescents. *Neuroimage* 2008;39(4):1918–1925. [PubMed: 18060810]
- Müller-Oehring EM, Schulte T, Raassi C, Pfefferbaum A, Sullivan EV. Local-global interference is modulated by age, sex and anterior corpus callosum size. *Brain Research* 2007;1142:189–205. [PubMed: 17335783]
- Naganawa S, Sato K, Katagiri T, Mimura T, Ishigaki T. Regional ADC values of the normal brain: differences due to age, gender, and laterality. *European Radiology* 2003;13(1):6–11. [PubMed: 12541104]
- Neeman M, Freyer JP, Sillerud LO. A simple method for obtaining cross-term-free images for diffusion anisotropy studies in NMR microimaging. *Magnetic Resonance in Medicine* 1991;21(1):138–143. [PubMed: 1943671]
- Nevo U, Hauben E, Yoles E, Agranov E, Akselrod S, Schwartz M, et al. Diffusion anisotropy MRI for quantitative assessment of recovery in injured rat spinal cord. *Magnetic Resonance in Medicine* 2001;45(1):1–9. [PubMed: 11146478]
- Norris DG, Niendorf T, Leibfritz D. Healthy and infarcted brain tissues studied at short diffusion times: the origins of apparent restriction and the reduction in apparent diffusion coefficient. *NMR Biomedicine* 1994;7(7):304–310.
- Nusbaum AO, Tang CY, Buchsbaum MS, Wei TC, Atlas SW. Regional and global changes in cerebral diffusion with normal aging. *American Journal of Neuroradiology* 2001;22(1):136–142. [PubMed: 11158899]
- O'Sullivan M, Jones D, Summers P, Morris R, Williams S, Markus H. Evidence for cortical “disconnection” as a mechanism of age-related cognitive decline. *Neurology* 2001;57:632–638. [PubMed: 11524471]
- Ota M, Obata T, Akine Y, Ito H, Ikehira H, Asada T, et al. Age-related degeneration of corpus callosum measured with diffusion tensor imaging. *NeuroImage* 2006;31(4):1445–1452. [PubMed: 16563802]
- Pagani E, Agosta F, Rocca MA, Caputo D, Filippi M. Voxel-based analysis derived from fractional anisotropy images of white matter volume changes with aging. *Neuroimage* 2008;41(3):657–667. [PubMed: 18442927]
- Pandya, DN.; Seltzer, B. The topography of commissural fibers. In: Lepore, F.; Püto, M.; Jasper, HH., editors. *Two Hemispheres-One Brain: Functions of the Corpus Callosum*. Alan R. Liss, Inc.; New York: 1986. p. 47-74.
- Park DC, Smith AD, Lautenschlager G, Earles JL, Frieske D, Zwahr M, et al. Mediators of long-term memory performance across the life span. *Psychology of Aging* 1996;11(4):621–637.
- Peters A, Sethares C. Is there remyelination during aging of the primate central nervous system? *Journal of Comparative Neurology* 2003;460(2):238–254. [PubMed: 12687688]

- Pfefferbaum A, Adalsteinsson E, Sullivan EV. Frontal circuitry degradation marks healthy adult aging: Evidence from diffusion tensor imaging. *NeuroImage* 2005;26(3):891–899. [PubMed: 15955499]
- Pfefferbaum A, Adalsteinsson E, Sullivan EV. Dymorphology and microstructural degradation of the corpus callosum: Interaction of age and alcoholism. *Neurobiology of Aging* 2006a;27(7):994–1009. [PubMed: 15964101]
- Pfefferbaum A, Adalsteinsson E, Sullivan EV. Supratentorial profile of white matter microstructural integrity in recovering alcoholic men and women. *Biological Psychiatry* 2006b;59(4):364–372. [PubMed: 16125148]
- Pfefferbaum A, Ford J, Wenegrat B, Roth WT, Kopell BS. Clinical application of the P3 component of event-related potentials: I. Normal aging. *Electroencephalography and Clinical Neurophysiology* 1984;59:85–103. [PubMed: 6200311]
- Pfefferbaum A, Mathalon DH, Sullivan EV, Rawles JM, Zipursky RB, Lim KO. A quantitative magnetic resonance imaging study of changes in brain morphology from infancy to late adulthood. *Archives of Neurology* 1994;51:874–887. [PubMed: 8080387]
- Pfefferbaum A, Rosenbloom MJ, Adalsteinsson E, Sullivan EV. Diffusion tensor imaging with quantitative fiber tracking in HIV infection and alcoholism comorbidity: Synergistic white matter damage. *Brain* 2007;130:48–64. [PubMed: 16959813]
- Pfefferbaum A, Rosenbloom MJ, Rohlfing T, Adalsteinsson E, Kemper CA, Deresinski S, et al. Contribution of alcoholism to brain dymorphology in HIV infection: Effects on the ventricles and corpus callosum. *NeuroImage* 2006;33(1):239–251. [PubMed: 16877010]
- Pfefferbaum A, Sullivan EV. Microstructural but not macrostructural disruption of white matter in women with chronic alcoholism. *NeuroImage* 2002;15:708–718. [PubMed: 11848714]
- Pfefferbaum A, Sullivan EV. Increased brain white matter diffusivity in normal adult aging: Relationship to anisotropy and partial voluming. *Magnetic Resonance in Medicine* 2003;49:953–961. [PubMed: 12704779]
- Pfefferbaum A, Sullivan EV. Diffusion MR imaging in neuropsychiatry and aging. In: Gillard, J.; Waldman, A.; Barker, P., editors. *Clinical MR Neuroimaging: Diffusion, Perfusion and Spectroscopy*. Cambridge University Press; Cambridge: 2005a. p. 558–578.
- Pfefferbaum A, Sullivan EV. Disruption of brain white matter microstructure by excessive intracellular and extracellular fluid in alcoholism: Evidence from diffusion tensor imaging. *Neuropsychopharmacology* 2005b;30:423–432. [PubMed: 15562292]
- Pfefferbaum A, Sullivan EV. Diffusion MR imaging in neuropsychiatry and aging. In: Gillard, JH.; Waldman, AD.; Barker, PB., editors. *Clinical MR Neuroimaging: Diffusion, Perfusion and Spectroscopy*. 2nd edition. Cambridge University Press; Cambridge: 2010. in press
- Pfefferbaum A, Sullivan EV, Carmelli D. Genetic regulation of regional microstructure of the corpus callosum in late life. *Neuroreport* 2001;12:1677–1681. [PubMed: 11409738]
- Pfefferbaum A, Sullivan EV, Carmelli D. Morphological changes in aging brain structures are differentially affected by time-linked environmental influences despite strong genetic stability. *Neurobiology of Aging* 2004;25(2):175–183. [PubMed: 14749135]
- Pfefferbaum A, Sullivan EV, Hedehus M, Adalsteinsson E, Lim KO, Moseley M. In vivo detection and functional correlates of white matter microstructural disruption in chronic alcoholism. *Alcoholism: Clinical and Experimental Research* 2000;24(8):1214–1221.
- Pfefferbaum A, Sullivan EV, Hedehus M, Lim KO, Adalsteinsson E, Moseley M. Age-related decline in brain white matter anisotropy measured with spatially corrected echo-planar diffusion tensor imaging. *Magnetic Resonance in Medicine* 2000;44(2):259–268. [PubMed: 10918325]
- Pfefferbaum A, Sullivan EV, Rosenbloom MJ, Mathalon DH, Lim KO. A controlled study of cortical gray matter and ventricular changes in alcoholic men over a five year interval. *Archives of General Psychiatry* 1998;55(10):905–912. [PubMed: 9783561]
- Pierpaoli C, Barnett A, Pajevic S, Chen R, Penix L, Virta A, et al. Water diffusion changes in Wallerian degeneration and their dependence on white matter architecture. *NeuroImage* 2001;13:1174–1185. [PubMed: 11352623]
- Pierpaoli C, Basser PJ. Towards a quantitative assessment of diffusion anisotropy. *Magnetic Resonance in Medicine* 1996;36:893–906. [PubMed: 8946355]

- Rabbitt P, Scott M, Lunn M, Thacker N, Lowe C, Pendleton N, et al. White matter lesions account for all age-related declines in speed but not in intelligence. *Neuropsychology* 2007;21(3):363–370. [PubMed: 17484599]
- Raz N, Gunning FM, Head D, Dupuis JH, McQuain J, Briggs SD, et al. Selective aging of the human cerebral cortex observed in vivo: Differential vulnerability of the prefrontal gray matter. *Cerebral Cortex* 1997;7(3):268–282. [PubMed: 9143446]
- Raz N, Gunning-Dixon F, Head D, Rodrigue K, Williamson A, Acker J. Aging, sexual dimorphism, and hemispheric asymmetry of the cerebral cortex: replicability of regional differences in volume. *Neurobiology of Aging* 2004;25(3):377–396. [PubMed: 15123343]
- Raz N, Gunning-Dixon F, Head D, Williamson A, Acker JD. Age and sex differences in the cerebellum and the ventral pons: a prospective MR study of healthy adults. *American Journal of Neuroradiology* 2001;22(6):1161–1167. [PubMed: 11415913]
- Raz N, Gunning-Dixon FM, Head D, Dupuis JH, Acker JD. Neuroanatomical correlates of cognitive aging: evidence from structural magnetic resonance imaging. *Neuropsychology* 1998;12(1):95–114. [PubMed: 9460738]
- Raz N, Lindenberger U, Rodrigue KM, Kennedy KM, Head D, Williamson A, et al. Regional brain changes in aging healthy adults: General trends, individual differences, and modifiers. *Cerebral Cortex* 2005;15:1676–1689. [PubMed: 15703252]
- Raz N, Rodrigue KM. Differential aging of the brain: Patterns, cognitive correlates and modifiers. *Neuroscience and Biobehavioral Reviews* 2006;30:730–748. [PubMed: 16919333]
- Raz N, Rodrigue KM, Kennedy KM, Acker JD. Vascular health and longitudinal changes in brain and cognition in middle-aged and older adults. *Neuropsychology* 2007;21(2):149–157. [PubMed: 17402815]
- Resnick SM, Pham DL, Kraut MA, Zonderman AB, Davatzikos C. Longitudinal magnetic resonance imaging studies of older adults: a shrinking brain. *Journal of Neuroscience* 2003;23(8):3295–3301. [PubMed: 12716936]
- Rohlfing T, Brandt R, Menzel R, Maurer CR Jr. Evaluation of atlas selection strategies for atlas-based image segmentation with application to confocal microscopy images of bee brains. *NeuroImage* 2004;21(4):1428–1442. [PubMed: 15050568]
- Rohlfing T, Maurer CR. Nonrigid image registration in shared-memory multiprocessor environments with application to brains, breasts, and bees. *IEEE Transactions on Information Technology in Biomedicine* 2003;7(1):16–25. [PubMed: 12670015]
- Rohlfing T, Sullivan EV, Pfefferbaum A. Deformation-based brain morphometry to track the course of alcoholism: Differences between intra-subject and inter-subject analysis. *Psychiatry Research: NeuroImaging* 2006;146:157–170.
- Rohlfing, T.; Zahr, NM.; Sullivan, EV.; Pfefferbaum, A. The SRI24 multi-channel brain atlas: Construction and applications; *Medical Imaging 2008: Image Processing*, Proceedings of SPIE; 2008; EID 691409 (691412 pages)
- Rumpel H, Ferrini B, Martin E. Lasting cytotoxic edema as an indicator of irreversible brain damage: A case of neonatal stroke. *American Journal of Neuroradiology* 1998;19:1636–1638. [PubMed: 9802484]
- Ryberg C, Rostrup E, Stegmann MB, Barkhof F, Scheltens P, van Straaten EC, et al. Clinical significance of corpus callosum atrophy in a mixed elderly population. *Neurobiology of Aging* 2007;28(6):955–963. [PubMed: 16797787]
- Salat DH, Kaye JA, Janowsky JS. Prefrontal gray and white matter volumes in healthy aging and Alzheimer disease. *Archives of Neurology* 1999;56(3):338–344. [PubMed: 10190825]
- Salat DH, Tuch DS, Greve DN, van der Kouwe AJW, Hevelone ND, Zaleta AK, et al. Age-related alterations in white matter microstructure measured by diffusion tensor imaging. *Neurobiology of Aging* 2005;26:1215–1227. [PubMed: 15917106]
- Salat DH, Tuch DS, Hevelone ND, Fischl B, Corkin S, Rosas HD, et al. Age-related changes in prefrontal white matter measured by diffusion tensor imaging. *Annals of the New York Academy of Sciences* 2005;1064:37–49. [PubMed: 16394146]

- Schmahmann JD, Pandya DN, Wang R, Dai G, D'Arceuil HE, de Crespigny AJ, et al. Association fibre pathways of the brain: parallel observations from diffusion spectrum imaging and autoradiography. *Brain* 2007;130(Pt 3):630–653. [PubMed: 17293361]
- Schneiderman JS, Buchsbaum MS, Haznedar MM, Hazlett EA, Brickman AM, Shihabuddin L, et al. Diffusion tensor anisotropy in adolescents and adults. *Neuropsychobiology* 2007;55(2):96–111. [PubMed: 17587876]
- Sehy JV, Ackerman JJ, Neil JJ. Evidence that both fast and slow water ADC components arise from intracellular space. *Magnetic Resonance in Medicine* 2002;48(5):765–770. [PubMed: 12417990]
- Sen PN, Basser PJ. A model for diffusion in white matter in the brain. *Biophysical Journal* 2005;89(5):2927–2938. [PubMed: 16100258]
- Shenkin SD, Bastin ME, MacGillivray TJ, Deary IJ, Starr JM, Wardlaw JM. Childhood and current cognitive function in healthy 80-year-olds: a DT-MRI study. *Neuroreport* 2003;14(3):345–349. [PubMed: 12634481]
- Shimony JS, McKinstry RC, Akbudak E, Aronovitz JA, Snyder AZ, Lori NF, et al. Quantitative diffusion-tensor anisotropy brain MR imaging: normative human data and anatomic analysis. *Radiology* 1999;212(3):770–784. [PubMed: 10478246]
- Sidaros A, Engberg AW, Sidaros K, Liptrot MG, Herning M, Petersen P, et al. Diffusion tensor imaging during recovery from severe traumatic brain injury and relation to clinical outcome: a longitudinal study. *Brain* 2008;131(Pt 2):559–572. [PubMed: 18083753]
- Silva MD, Omae T, Helme r. K. G. Li F, Fisher M, Sotak CH. Separating changes in the intra- and extracellular water apparent diffusion coefficient following focal cerebral ischemia in the rat brain. *Magnetic Resonance in Medicine* 2002;48(5):826–837. [PubMed: 12417997]
- Smith CD, Chebrolu H, Wekstein DR, Schmitt FA, Markesbery WR. Age and gender effects on human brain anatomy: a voxel-based morphometric study in healthy elderly. *Neurobiology of Aging* 2007;28(7):1075–1087. [PubMed: 16774798]
- Song SK, Sun SW, Ramsbottom MJ, Chang C, Russell J, Cross AH. Dysmyelination revealed through MRI as increased radial (but unchanged axial) diffusion of water. *NeuroImage* 2002;17(3):1429–1436. [PubMed: 12414282]
- Sowell ER, Thompson PM, Holmes CJ, Jernigan TL, Toga AW. In vivo evidence for post-adolescent brain maturation in frontal and striatal regions [letter]. *Nature & Neuroscience* 1999;2(10):859–861.
- Sowell ER, Thompson PM, Toga AW. Mapping changes in the human cortex throughout the span of life. *Neuroscientist* 2004;10(4):372–392. [PubMed: 15271264]
- Stadlbauer A, Salomonowitz E, Strunk G, Hammen T, Ganslandt O. Quantitative diffusion tensor fiber tracking of age-related changes in the limbic system. *European Radiology*. 2007
- Stadlbauer A, Salomonowitz E, Strunk G, Hammen T, Ganslandt O. Age-related degradation in the central nervous system: assessment with diffusion-tensor imaging and quantitative fiber tracking. *Radiology* 2008;247(1):179–188. [PubMed: 18292477]
- Stebbins G, Carrillo MD, Medina D, de Toledo-Morrell L, Klingberg T, Poldrack RA, et al. Frontal white matter integrity in aging and its relation to reasoning performance: A diffusion tensor imaging study (abs 456.3). *Society for Neuroscience Abstracts* 2001;27:1204.
- Sullivan EV, Adalsteinsson E, Hedehus M, Ju C, Moseley M, Lim KO, et al. Equivalent disruption of regional white matter microstructure in aging healthy men and women. *Neuroreport* 2001;12(22):99–104. [PubMed: 11201100]
- Sullivan EV, Adalsteinsson E, Pfefferbaum A. Selective age-related degradation of anterior callosal fiber bundles quantified in vivo with fiber tracking. *Cerebral Cortex* 2006;16(7):1030–1039. [PubMed: 16207932]
- Sullivan EV, Deshmukh A, Desmond JE, Lim KO, Pfefferbaum A. Cerebellar volume decline in normal aging, alcoholism, and Korsakoff's syndrome: Relation to ataxia. *Neuropsychology* 2000;14(3):341–352. [PubMed: 10928737]
- Sullivan EV, Pfefferbaum A. Neuroradiological characterization of normal adult aging. *British Journal of Radiology* 2007;60:S99–S108. [PubMed: 18445750]

- Sullivan, EV.; Pfefferbaum, A. Diffusion tensor imaging in aging and age-related neurodegenerative disorders. In: Jones, DK., editor. Diffusion MRI. Cambridge University Press; Cambridge: 2009. in press
- Sullivan EV, Pfefferbaum A, Adalsteinsson E, Swan GE, Carmelli D. Differential rates of regional change in callosal and ventricular size: A 4-year longitudinal MRI study of elderly men. *Cerebral Cortex* 2002;12:438–445. [PubMed: 11884358]
- Sullivan EV, Rohlfing T, Pfefferbaum A. Quantitative fiber tracking of lateral and interhemispheric white matter systems in normal aging: Relations to timed performance. *Neurobiology of Aging*. 2009 online May 19 2008.
- Sullivan EV, Rosenbloom MJ, Serventi KL, Pfefferbaum A. Effects of age and sex on volumes of the thalamus, pons, and cortex. *Neurobiology of Aging* 2004;25:185–192. [PubMed: 14749136]
- Takahashi T, Murata T, Omori M, Kosaka H, Takahashi K, Yonekura Y, et al. Quantitative evaluation of age-related white matter microstructural changes on MRI by multifractal analysis. *Journal of the Neurological Sciences* 2004;225(1-2):33–37. [PubMed: 15465083]
- Taki Y, Kinomura S, Sato K, Goto R, Inoue K, Okada K, et al. Both global gray matter volume and regional gray matter volume negatively correlate with lifetime alcohol intake in non-alcohol-dependent Japanese men: a volumetric analysis and a voxel-based morphometry. *Alcoholism: Clinical and Experimental Research* 2006;30(6):1045–1050.
- Tang Y, Whitman GT, Lopez I, Baloh RW. Brain volume changes on longitudinal magnetic resonance imaging in normal older people. *Journal of Neuroimaging* 2001;11(4):393–400. [PubMed: 11677879]
- Teuber H-L. Physiological psychology. *Annual Review of Psychology* 1955;6:267–296.
- Tisserand D, Van Boxtel M, Gronenschild E, Jolles J. Age-related volume reductions of prefrontal regions in healthy individuals are differential. *Brain and Cognition* 2001;47:182–185.
- Tisserand DJ, Jolles J. On the involvement of prefrontal networks in cognitive ageing. *Cortex* 2003;39(4-5):1107–1128. [PubMed: 14584569]
- Tisserand DJ, Pruessner JC, Sanz Arigita EJ, van Boxtel MP, Evans AC, Jolles J, et al. Regional frontal cortical volumes decrease differentially in aging: an MRI study to compare volumetric approaches and voxel-based morphometry. *NeuroImage* 2002;17(2):657–669. [PubMed: 12377141]
- Virta A, Barnett A, Pierpaoli C. Visualizing and characterizing white matter fiber structure and architecture in the human pyramidal tract using diffusion tensor MR. *Magnetic Resonance Imaging* 1999;17(8):1121–1133. [PubMed: 10499674]
- Walhovd KB, Fjell AM, Reinvang I, Lundervold A, Dale AM, Eilertsen DE, et al. Effects of age on volumes of cortex, white matter and subcortical structures. *Neurobiology of Aging* 2005;26(9):1261–1270. discussion 1275–1268. [PubMed: 16005549]
- Wang C, Stebbins GT, Nyenhuis DL, deToledo-Morrell L, Freels S, Gencheva E, et al. Longitudinal changes in white matter following ischemic stroke: a three-year follow-up study. *Neurobiology of Aging* 2006;27(12):1827–1833. [PubMed: 16310892]
- Wang C, Ulbert I, Schomer DL, Marinkovic K, Halgren E. Responses of human anterior cingulate cortex microdomains to error detection, conflict monitoring, stimulus-response mapping, familiarity, and orienting. *Journal of Neuroscience* 2005;25(3):604–613. [PubMed: 15659596]
- Wang Y, Markram H, Goodman PH, Berger TK, Ma J, Goldman-Rakic PS. Heterogeneity in the pyramidal network of the medial prefrontal cortex. *Nature Neurosci* 2006;9(4):534–542. [PubMed: 16547512]
- Wilde EA, Chu Z, Bigler ED, Hunter JV, Fearing MA, Hanten G, et al. Diffusion tensor imaging in the corpus callosum in children after moderate to severe traumatic brain injury. *Journal of Neurotrauma* 2006;23(10):1412–1426. [PubMed: 17020479]
- Xu D, Mori S, Solaiyappan M, van Zijl PC, Davatzikos C. A framework for callosal fiber distribution analysis. *NeuroImage* 2002;17:1131–1143. [PubMed: 12414255]
- Xue R, van Zijl PC, Crain BJ, Solaiyappan M, Mori S. In vivo three-dimensional reconstruction of rat brain axonal projections by diffusion tensor imaging. *Magnetic Resonance in Medicine* 1999;42(6):1123–1127. [PubMed: 10571934]

- Zahr, NM.; Pfefferbaum, A.; Sullivan, EV. Changes in macrostructure and microstructure of the aging brain. In: Abou-Saleh, M.; Katona, C.; Kumar, A., editors. Principles and Practice of Geriatric Psychiatry. 3rd Edition. Wiley-Blackwell; London: 2009. in press
- Zahr NM, Rohlfing T, Pfefferbaum A, Sullivan EV. Problem solving, working memory, and motor correlates of association and commissural fiber bundles in normal aging: A quantitative fiber tracing study. *NeuroImage* 2009;44(3):1050–1062. [PubMed: 18977450]

Conventional Magnetic Resonance

Diffusion Tensor

Figure 1.

Axial images at the level of the lateral ventricles. Top two images. Examples of conventional magnetic resonance images from a fast spin-echo sequence (FSE). On the left is an early-echo image, which differentiates gray matter (light gray) and white matter (darker gray), both of which are relatively homogeneous in intensity. On the right is a late-echo image, on which CSF is brightest. Bottom two images: Examples of images from a diffusion tensor imaging sequence. On the left is a fractional anisotropy (FA) image, on which white matter is the brightest. On the right is a diffusivity image, on which CSF is brightest. The diffusion ellipsoid model displays in red the preferred orientation of the longitudinal diffusion, λ_L , and in blue the two minor axes of diffusion, the mean of which is λ_T .

Figure 2.

Top sagittal image: Example of fiber tracking of the corpus callosum of a 23 year-old healthy woman. The colors represent six different fiber bundles based on divisions described by Pandya and Selzer (Pandya & Seltzer, 1986) identified with fiber tracking based on the DTI-derived FA. From anterior (far left) to posterior (far right), the bundles are deemed prefrontal, premotor, precentral, postcentral, posterior parietal, and temporal-occipital. Bottom data figures: Mean $\pm$ S.E. of the six callosal sectors for the young and elderly groups at each MRI session of the four principal metrics of DTI: fractional anisotropy (upper left), apparent diffusion coefficient (upper right), λ_L (lower left), and λ_T (lower right). The elderly group had lower FA and higher

diffusivity than the young group at both scanning sessions, and the group differences were greatest in the anterior sectors.

Figure 3.

Top axial image on left: Example of fiber tracking of the midsagittal corpus callosum. The colors represent six different fiber bundles defined in Figure 2. Left column of line plots: Mean $\pm$ S.E. of FA, ADC, λ L, and λ T of the midsagittal callosal fibers. Top axial image on right: Example of fiber tracking of the distal fibers of the corpus callosum. Right column of line plots: Mean $\pm$ S.E. of FA, ADC, λ L, and λ T of the distal callosal fibers.

Figure 4.

Mean $\pm$ S.E. of the volumes of the corpus callosum and lateral ventricles for the young and elderly groups at each MRI session. While the callosal volumes showed no detectable change in either group over the 2-year follow-up period (group-by-time interaction ($F(1,14)=.0093$, $p=.9245$), the lateral ventricles expanded disproportionately in the elderly relative to the young group (group-by-time interaction ($F(1,14)=8.921$, $p=.0098$).

Figure 5.

Three-dimensional rendering of three views of the ventricular system: oblique sagittal view on left (frontal to occipital is right to left); axial view in middle (frontal to occipital is top to bottom); coronal view on right. All views reveal that the ventricles of the elderly women are substantially greater than those of the younger woman.