Published in final edited form as:

J Occup Environ Med. 2010 April ; 52(4): 407–414. doi:10.1097/JOM.0b013e3181d5e386.

Evaluation of cumulative lead dose and longitudinal changes in structural MRI in former organolead workers

Brian S. Schwartz, MD, MS^{1,2,5}, Brian Caffo, PhD³, Walter F. Stewart, PhD, MPH^{2,4}, Haley Hedlin³, Bryan D. James, MA, PhD^{8,9}, David Yousem, MD⁶, and Christos Davatzikos, PhD⁷

¹ Department of Environmental Health Sciences, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

² Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

³ Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

⁴ Center for Health Research, Geisinger Clinic, Danville, PA

⁵ Department of Medicine, Johns Hopkins School of Medicine, Baltimore, MD

⁶ The Russell H. Morgan Department of Radiology and Radiological Sciences, Johns Hopkins School of Medicine, Baltimore, MD

⁷ Department of Radiology, University of Pennsylvania School of Medicine, Philadelphia, PA

⁸ Rush University Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL

⁹ Department of Internal Medicine, Rush University Medical Center, Chicago, IL

Abstract

Objective—We evaluated whether tibia lead was associated with longitudinal change in brain volumes and white matter lesions in male former lead workers and population-based controls in whom we have previously reported on the cognitive and structural consequences of cumulative lead dose.

Methods—We used linear regression to identify predictors of change in brain volumes and white matter lesion grade scores, using two MRIs an average of five years apart.

Results—On average, total brain volume declined almost 30 cm³, predominantly in gray matter. Increasing age at the first MRI was strongly associated with larger declines in volumes and greater increases in white matter lesion scores. Tibia lead was not associated with change in brain volumes or white matter lesion scores.

Conclusions—In former lead workers in whom cumulative lead dose was associated with progressive declines in cognitive function decades after occupational exposure had ended, cumulative lead dose was associated with earlier persistent effects on brain structure, but not with additional worsening over five years.

We previously reported on relations of lifetime cumulative lead dose (estimated as the concentration of lead in tibia bone by X-ray fluorescence [XRF]) with cognitive function and brain structure measured by MRI in a cohort of older former workers with past exposure to organic and inorganic lead [1]. The long period between last occupational lead exposure and study follow-up (an average of 16 years at the first study visit) allows us to evaluate whether

lifetime lead dose was associated with reversible, persistent, or progressive effects on cognitive function and brain structure. Distinguishing among these various effects is an essential utility of longitudinal data, which is relatively rare in occupational epidemiology studies.

Understanding these relations is also directly relevant to the general population, since most older Americans were exposed to high levels of environmental lead exposure in the past and can have average tibia lead concentrations that are higher than in the former workers [2]. Past cumulative inorganic lead dose is adversely associated with cognitive function in older persons in the general population [3,4].

We reported that higher lifetime lead dose in former lead workers was associated with: 1) poorer cognitive test scores at cross-section [5] and progressive declines in cognitive function over time [6,7]; 2) smaller brain volumes in both regions of interest (ROIs) and voxel-wise analytic approaches [8]; 3) increased prevalence and severity of white matter lesions [8]; and 4) greater decrements in cognitive function from cumulative lead dose in subjects with the apolipoprotein E ϵ 4 allele (*APOE- ϵ 4*) [9]. In cross-sectional analyses smaller brain volumes were associated with worse cognitive function [10], and there was evidence that the associations of lead dose with worse cognitive function were mediated, at least in part, by changes in brain volumes [11].

With data from a second MRI, an average of five years later, we now report on relations of cumulative lead dose (tibia lead levels) with longitudinal changes in brain volumes and white matter lesions, to evaluate whether the effects of lead dose on brain structure are likely to be reversible, persistent, or progressive.

METHODS

Study Design and Overview

Subjects were initially recruited during two study phases between 1994 and 2003, as previously described [8]. In phase I (1994–7), former employees of a chemical manufacturing plant in the eastern United States were recruited. The first MRI was obtained in phase II (2001–3). During phase III (2005–8), summarized herein, subjects who completed the first MRI were invited for a second MRI. All phases of the study were reviewed and approved by the Johns Hopkins Bloomberg School of Public Health Committee on Human Research and written informed consent was obtained from all participants.

Selection and Recruitment of Study Subjects

The selection, recruitment, and enrollment over time of former lead workers and community-dwelling controls without occupational lead exposure (hereafter referred to as controls) have been previously reported [5,6,8,10,12]. During phase II, first MRIs were completed on 589 of 979 (60%) former lead workers and 67 of 131 (51%) controls. All participants in earlier phases of the study were eligible for this first MRI measurement. During phase III, a second MRI was obtained from 317 of 589 (54%) former lead workers and 45 of 67 (67%) controls. Second MRIs were not obtained because of death ($N = 52$), chronic illness ($N = 44$), discomfort (e.g., claustrophobia, inability to lie down) with the procedure ($N = 12$), contraindications (e.g., metal foreign body in eye) to MRI scanning ($N = 16$), loss to follow-up ($N = 47$), out-migration ($N = 9$), and refusal for unspecified reasons ($N = 99$).

Data Collection

Detailed data collection methods for the first two phases of the study have been previously described [8]. Following we describe only measures specifically used for the analysis presented herein.

Subject interview—In phase III, the subject interview was expanded to include a number of additional study variables, similar to the one used in the Baltimore Memory Study [13,14]. Health outcomes (e.g., diabetes, heart disease) were ascertained by interview response to the following question format for each condition, “Has a doctor ever told you that you had [name of condition]?” For educational attainment, information was obtained by interview on years of education, trade school, GED, and other educational certificates using previously published methods [14].

Tibia lead—Tibia lead, an estimate of lifetime cumulative lead dose, was available from earlier phases of the study on all former lead workers and all but one control with two MRIs. This was measured with ^{109}Cd -induced K-shell X-ray fluorescence (μg lead per gram bone mineral) and modeled as the estimated level at the end of employment (peak tibia lead), as previously described [5].

MRI acquisition—For the first MRI, all subjects were imaged at the same location on the same General Electric 1.5 T Signa model as previously described [8]. Eighteen of the first MRIs were not suitable for volumetric analysis due to image quality. For the second MRI, a 3 T General Electric scanner was utilized. T1-weighted images were acquired using an SPGR sequence (TE = 8 ms, TR = 21 ms, flip angle = 30° , FOV = 24 cm). Axial PD/T2 (TR/TE/TE2 = 2,200/27/120) and FLAIR (TR/TE/T1 = 8,000/100/2000) images were also acquired for lesion grading.

Clinical MRI review and white matter grading—All MRIs were reviewed to exclude urgent or emergent brain disease (subjects and their physicians were notified if present) [15]. MRIs were assigned a white matter lesion grade score by a trained neuroradiologist using the Cardiovascular Health Study (CHS) ten-point (0 to 9) scale [16,17], as previously reported [8], allowing analysis of change in ratings.

Image Analysis

The methods to obtain regional and voxel-wise volumes, including skull stripping, segmentation, registration, and transformation to regional analysis of volumes examined in normalized space (RAVENS), were completed using previously published methods [8,18–22]. Due to the inevitable changes in scanner technology and pulse sequences, we employed specialized image analysis methods that minimized the discontinuity between the two scans. We used the CLASSIC algorithm [23], which employs a 4-dimensional segmentation framework in which the baseline and follow-up scans are considered jointly during segmentation to minimize discrepancies between the two segmentations and better estimate longitudinal change. This algorithm has been previously validated [23].

Statistical Analysis

The purpose of the analysis was to determine if the effect of lead on brain structure was progressive in nature, an essential task which requires longitudinal data; that is, after lead exposure, lead gains access to the blood, then to the brain, causes an effect there, then leaves the brain, but the effect (e.g., volume loss) continues over time as a function of cumulative lead dose.

Multiple linear regression was used to evaluate associations of predictor variables with change in brain volumes (or volumes at the second MRI separately), using both ROI-based and voxel-wise approaches, and change in CHS scores. All regression models were adjusted for baseline age, duration of time between MRIs, APOE genotype, peak tibia lead (in analysis with former lead workers only), control status (i.e., former lead worker vs. control, in analysis with both

only), baseline ROI volume, height (cm), and education [14]. Model diagnostics were used to evaluate influence and normality for the ROI-based analysis.

ROI-based approach—To be consistent with the results of our prior published reports, we modeled change in 20 previously selected ROI volumes [8]. For bilateral structures, the volume represented the sum of right and left structures to minimize multiplicity concerns, but analyses were also performed separately for change in left- and right-sided ROI volumes (data not reported). Since we did not formally adjust for multiple comparisons in the ROI analysis we acknowledge that a p -value < 0.05 does not necessarily imply statistical significance.

Voxel-wise approach—Change in voxel volumes was modeled controlling for the aforementioned covariates using multivariate permutation testing in the R statistical programming language (www.cran.r-project.org). The SPM5 package (Statistical Parametric Software, Functional Imaging Laboratory, Wellcome Department of Imaging Neuroscience, University College London, 2003) was used to perform smoothing using a 3D isotropic Gaussian filter and MRIcro [24] to display results. Statistical significance was evaluated using a permutation approach that controlled for confounding variables. The maximum cluster size and cluster peak above the threshold was used to define a conservative permutation distribution on cluster sizes and peaks that, when compared to the observed cluster sizes and peaks, controls for multiple comparisons.

White matter lesions—Linear regression was used to model change in CHS white matter lesion grade scores.

RESULTS

Descriptive Summary of Study Subjects

Compared to those with no or one MRI, subjects with two MRIs were slightly younger, had a shorter time since last occupational exposure to lead, and lower peak tibia lead levels (Table 1). The mean (SD) duration from the first MRI to the second was 5.0 (0.4) years (range 3.6 to 6.1). The current age of the 317 former lead workers and 45 controls was 64.1 (7.6) and 71.9 (6.0) years, respectively ($p < 0.001$) (Table 2). Among all cohort members, controls had higher current tibia lead levels than did former workers (mean of 19.0 vs. 14.8 $\mu\text{g/g}$, Table 1), likely due to a cohort effect associated with the higher average age of controls.

Descriptive Summary of Change in ROI Volumes

There was no evidence that declines in volumes differed between former lead workers and controls (data not shown), so results are presented for all subjects combined. On average, total brain volume declined almost 30 cm^3 over a five-year period (Table 3 and Figure 1), with a more substantial decline in gray (24.4 cm^3) compared to white (5.4 cm^3) matter. All ROIs evidenced decline except for occipital white matter. The mean (SD) percent gray matter at the first MRI was 47.9% (1.8%) and at the second MRI it was 47.0% (1.9%). When change is expressed as a percent of the baseline total brain volume, the greatest decline was observed for total brain (2.55%), followed by total gray matter (2.09%), total white matter (0.46%), and medial structures (0.39%).

Predictors of Change in ROI Volumes

Among former lead workers, peak tibia lead was not associated with change in ROI volumes in adjusted models (Table 4). The remaining predictors were evaluated in all subjects to maximize power, given the lack of association for tibia lead and lack of differences between former lead workers and controls. As baseline age increased, ROI volumes declined (Table 4), with the expected exception of ventricle volume which increased in relation to baseline age.

A larger ROI volume at baseline was associated with a greater decline in volume, a finding expected from regression towards the mean (data not shown). Larger durations between MRIs were associated with larger declines in gray matter volumes, larger increases in white matter volumes, and larger increases in ventricle volume (Table 4).

Predictors of Change in Voxel Volumes

In a parallel analysis, results were substantively similar using a voxel-wise approach. For example, supra-threshold clusters for the association of lead with change in volume were well within the range expected by chance. In contrast, the adjusted association between baseline age and change in voxel volumes identified large supra-threshold clusters, whose sizes were well above the distribution of the maximum cluster size under the null hypothesis (Figure 2).

Predictors of Change in CHS White Matter Lesion Grade Score

For the change in CHS white matter lesion grade score (CHS2 minus CHS1), 6 persons (1.7%) improved by one category, 87 persons (24.0%) were unchanged, 134 persons (36.9%) worsened one category, 100 persons (27.6%) worsened by two categories, 24 persons (6.6%) by three categories, and 12 (3.3%) by four or five categories. Neither peak tibia lead nor control status were associated with change in CHS scores. Baseline age and increasing duration between MRIs were associated with increases in CHS scores ($\beta = 0.055$, $p < 0.001$ and $\beta = 0.286$, $p = 0.05$, respectively).

DISCUSSION

In this cohort of 45 to 75 year old men with past occupational exposure to organic and inorganic lead and population controls, we had previously observed that peak tibia lead concentration (an estimate of past cumulative lead dose) was associated with worse neurobehavioral test scores at cross-section [5], longitudinal decline in cognitive function [6], the prevalence and severity of white matter lesions, and with decreased volumes in both larger (e.g., total brain, lobar gray and white matter) and smaller (e.g., cingulate gyrus, insula, corpus callosum) ROIs [8], almost two decades after occupational lead exposure had ended. Because tibia lead was not associated with change in brain volumes over time using both ROI- and voxel-based methods, the current analysis suggests that the influence of lead on brain structure is persistent, but the results do not support progressive changes over the five years as measured by volumes in two MRIs. Our previous reports of progressive cognitive decline associated with past cumulative lead dose, which we termed “accelerated aging,” may be explained by a persistent lead-associated structural lesion combined with the effect of other risk factors associated with aging [1,6,7]. That is, cognitive decline in subjects without occupational lead exposure is age-dependent, but is more rapid when aging is combined with such exposure, even after exposure ceases. However, it should be noted that a portion of what has been previously termed age-related cognitive decline may be due, at least in part, to ubiquitous neurotoxicants such as lead or mercury [1,25,26].

More specifically, in the prior cross-sectional analysis [8], the association of tibia lead with brain volumes and white matter lesions was evidence of a persistent influence of lead on brain structure. In that analysis, the studied former workers had a mean (SD) lead exposure duration of 8.7 (9.8) years and a mean (SD) duration since last occupational exposure to lead of 18.0 (11.0) years. This implies that the lower brain volumes and increased prevalence and severity of white matter lesions associated with tibia lead levels in the cross-sectional analysis could reflect changes that had occurred over as much as the prior 26.7 years, since the beginning of occupational lead exposure. In the current analysis, we did not observe additional longitudinal change associated with cumulative lead dose over the next five years. Thus, we conclude that cumulative lead dose was associated with persistent, but not likely progressive structural

changes in the brain. These findings are also not inconsistent with our prior conclusion that at least part of the influence of cumulative lead dose on cognitive function is mediated through volume loss [11], for the same reasoning as above regarding the differing time periods of opportunity for change associated with lead dose to occur in the cross-sectional and longitudinal analyses.

Our data on the magnitude of changes in MRI volumes associated with age and aging are similar to those previously reported with some notable differences [27–33]. Our whole brain atrophy rate of approximately 0.5% per year is similar to values reported in some prior studies [27, 29,30] but not others [34,35]. Resnick et al. reported slightly larger losses in white matter than gray matter [27] in an older population of 50 men and 42 women; white matter losses were widespread, while gray matter losses were more localized [36]. In a study of 362 volunteers ranging in age from 18 to 93 years, whole-brain volume adjusted for head size declined by 0.22% per year between 20 and 80 years, then more rapidly after that [28]. In an earlier report of 370 adults ranging from 18 to 97 years of age, the rate of decline in old non-demented subjects was 0.45% per year [37], with the observation that gray matter volume loss began at age 20 and continued to very old age. For the latter study, the white matter loss appears to begin in the fifth or sixth decade, a finding consistent with our estimates of the relative amount of gray and white matter losses.

An important consideration is whether selection bias could account for the results we have reported. After the first MRI, we determined that average cognitive function did not differ by first MRI status and the relations of tibia lead with neurobehavioral test scores did not differ in those with and without MRIs [8]. We concluded after the first MRI that there was unlikely to be meaningful selection bias among those who completed the first MRI that could influence study results [8]. Former lead workers with two MRIs had lower tibia lead levels and were younger than those with only one (Table 1). We believe these differences are likely to mask, rather than spuriously create, associations.

A fundamental methodological challenge in longitudinal MRI studies is posed by changes in scanner hardware and software between scans. Initial analysis showed that applying standard 3D segmentation methods independently to each scan was insufficient and led to low longitudinal stability of the volumetric measurements. We therefore used an advanced 4D segmentation and atlas registration technique, which has been developed and validated specifically for longitudinal studies [38]. A potential pitfall of this approach is that it can over-smooth, and therefore underestimate longitudinal brain changes, if the parameters that control temporal smoothness are not properly set. However, previous validation studies of this approach carefully determined the appropriate parameter range.

In conclusion, in this cohort of former lead workers, cumulative lead dose was associated with persistent effects on brain volume, but recent changes in brain volume over five additional years were not associated with tibia lead. Advancing age is associated with annual declines in brain volumes of approximately 0.5% per year, primarily in gray matter in this age range.

Clinical Significance Statement

Cognitive difficulties and abnormalities on structural MRI can be due to cumulative lead dose in former lead workers. Although measurement of tibia lead in patients has clinical utility, this test is not widely available. Given the long-term nature of the exposure and the chronicity of the structural and functional changes, it is unlikely that chelation treatment would offer such patients any benefit.

Acknowledgments

This research was supported by grant R01 AG10785 from the National Institute on Aging (NIA). Its content is solely the responsibility of the authors. The authors thank Dr. Andrew Todd for bone lead measurements in earlier phases of the study.

References

1. Stewart WF, Schwartz BS. Effects of lead on the adult brain: A 15-year exploration. *Am J Ind Med* 2007;50(10):729–39. [PubMed: 17311281]
2. Theppeang K, et al. Gender and race/ethnicity differences in lead dose biomarkers. *Am J Public Health* 2008;98(7):1248–55. [PubMed: 18511728]
3. Shih RA, et al. Environmental lead exposure and cognitive function in community-dwelling older adults. *Neurology* 2006;67(9):1556–62. [PubMed: 16971698]
4. Shih RA, et al. Cumulative lead dose and cognitive function in adults: a review of studies that measured both blood lead and bone lead. *Environ Health Perspect* 2007;115(3):483–92. [PubMed: 17431502]
5. Stewart WF, et al. Neurobehavioral function and tibial and chelatable lead levels in 543 former organolead workers. *Neurology* 1999;52(8):1610–7. [PubMed: 10331686]
6. Schwartz BS, et al. Past adult lead exposure is associated with longitudinal decline in cognitive function. *Neurology* 2000;55(8):1144–50. [PubMed: 11071492]
7. Links JM, et al. Characterization of toxicokinetics and toxicodynamics with linear systems theory: application to lead-associated cognitive decline. *Environ Health Perspect* 2001;109(4):361–8. [PubMed: 11335184]
8. Stewart WF, et al. Past adult lead exposure is linked to neurodegeneration measured by brain MRI. *Neurology* 2006;66(10):1476–84. [PubMed: 16717205]
9. Stewart WF, et al. ApoE genotype, past adult lead exposure, and neurobehavioral function. *Environ Health Perspect* 2002;110(5):501–5. [PubMed: 12003753]
10. Schwartz BS, et al. Relations of brain volumes with cognitive function in males 45 years and older with past lead exposure. *Neuroimage* 2007;37(2):633–41. [PubMed: 17600728]
11. Caffo B, et al. Are brain volumes based on magnetic resonance imaging mediators of the associations of cumulative lead dose with cognitive function? *Am J Epidemiol* 2008;167(4):429–37. [PubMed: 18079133]
12. Schwartz BS, et al. Decrements in neurobehavioral performance associated with mixed exposure to organic and inorganic lead. *Am J Epidemiol* 1993;137(9):1006–21. [PubMed: 8317446]
13. Glass TA, Rasmussen MD, Schwartz BS. Neighborhoods and obesity in older adults: the Baltimore Memory Study. *Am J Prev Med* 2006;31(6):455–63. [PubMed: 17169707]
14. Schwartz BS, et al. Disparities in cognitive functioning by race/ethnicity in the Baltimore Memory Study. *Environ Health Perspect* 2004;112(3):314–20. [PubMed: 14998746]
15. Alphas HH, et al. Findings on brain MRI from research studies of occupational exposure to known neurotoxicants. *AJR Am J Roentgenol* 2006;187(4):1043–7. [PubMed: 16985155]
16. Fried LP, et al. The Cardiovascular Health Study: design and rationale. *Ann Epidemiol* 1991;1(3):263–76. [PubMed: 1669507]
17. Longstreth WT Jr, et al. Lacunar infarcts defined by magnetic resonance imaging of 3660 elderly people: the Cardiovascular Health Study. *Arch Neurol* 1998;55(9):1217–25. [PubMed: 9740116]
18. Goldszal AF, et al. An image-processing system for qualitative and quantitative volumetric analysis of brain images. *J Comput Assist Tomogr* 1998;22(5):827–37. [PubMed: 9754125]
19. Kabani N, et al. A 3D atlas of the human brain. *Neuroimage* 1998;7(4):S717.
20. Shen, D. Proceedings of the IEEE International Symposium on Biomedical Imaging, Arlington, Va: 2004. 4D Image Warping For Measurement of Longitudinal Brain Changes.
21. Andreasen NC, et al. Automatic atlas-based volume estimation of human brain regions from MR images. *Journal of Computer Assisted Tomography* 1996;20(1):98–106. [PubMed: 8576490]
22. Resnick SM, et al. One-year age changes in MRI brain volumes in older adults. *Cereb Cortex* 2000;10(5):464–72. [PubMed: 10847596]

23. Xue Z, Shen D, Davatzikos C. CLASSIC: Consistent Longitudinal Alignment and Segmentation for Serial Image Computing. *Neuroimage* 2006;30(2):388–399. [PubMed: 16275137]
24. Rorden C, Brett M. Stereotaxic display of brain lesions. *Behav Neurol* 2000;12(4):191–200. [PubMed: 11568431]
25. Schwartz BS, Stewart WF. Lead and cognitive function in adults: a questions and answers approach to a review of the evidence for cause, treatment, and prevention. *Int Rev Psychiatry* 2007;19(6):671–92. [PubMed: 18092244]
26. Rowland AS, McKinsty RC. Lead toxicity, white matter lesions, and aging. *Neurology* 2006;66(10):1464–5. [PubMed: 16717201]
27. Resnick SM, et al. Longitudinal magnetic resonance imaging studies of older adults: a shrinking brain. *J Neurosci* 2003;23(8):3295–301. [PubMed: 12716936]
28. Fotenos AF, et al. Brain volume decline in aging: evidence for a relation between socioeconomic status, preclinical Alzheimer disease, and reserve. *Arch Neurol* 2008;65(1):113–20. [PubMed: 18195148]
29. Chen K, et al. Correlations between apolipoprotein E epsilon4 gene dose and whole brain atrophy rates. *Am J Psychiatry* 2007;164(6):916–21. [PubMed: 17541051]
30. Enzinger C, et al. Risk factors for progression of brain atrophy in aging: six-year follow-up of normal subjects. *Neurology* 2005;64(10):1704–11. [PubMed: 15911795]
31. Schmidt R, et al. White matter lesion progression, brain atrophy, and cognitive decline: the Austrian stroke prevention study. *Ann Neurol* 2005;58(4):610–6. [PubMed: 16178017]
32. Liu RS, et al. A longitudinal study of brain morphometrics using quantitative magnetic resonance imaging and difference image analysis. *Neuroimage* 2003;20(1):22–33. [PubMed: 14527567]
33. Bernick C, et al. Statins and cognitive function in the elderly: the Cardiovascular Health Study. *Neurology* 2005;65(9):1388–94. [PubMed: 16275825]
34. Mueller EA, et al. Brain volume preserved in healthy elderly through the eleventh decade. *Neurology* 1998;51(6):1555–62. [PubMed: 9855501]
35. Tang Y, et al. Brain volume changes on longitudinal magnetic resonance imaging in normal older people. *J Neuroimaging* 2001;11(4):393–400. [PubMed: 11677879]
36. Braak H, Braak E. Frequency of stages of Alzheimer-related lesions in different age categories. *Neurobiol Aging* 1997;18(4):351–7. [PubMed: 9330961]
37. Fotenos AF, et al. Normative estimates of cross-sectional and longitudinal brain volume decline in aging and AD. *Neurology* 2005;64(6):1032–9. [PubMed: 15781822]
38. Xue Z, Shen D, Davatzikos C. CLASSIC: Consistent Longitudinal Alignment and Segmentation for Serial Image Computing. *Neuroimage*. 2005

Figure 1.

“Spaghetti plots” for relations of age with change (cm³) in four ROI volumes (panel A = total gray matter, panel B = total brain, panel C = total white matter, and panel D = bilateral hippocampus), including former lead workers and controls. Each line represents an individual’s change in age and change in ROI volume across the two MRIs.

Figure 2.

Transverse template brain slices with t-statistic maps of the adjusted association between age and change in brain volumes on a voxel-wise basis. Location of slice is identified by figure in lower right corner. The figure displays t-statistics < -3.1 with colors defined by key in lower right corner. Panel A is for gray matter and panel B is for white matter. For gray matter, the maps identified one large cluster that exhibited both maximum cluster size and peak value significance after controlling for multiplicity (p-values < 0.05). For white matter, there were 28 clusters that satisfied these two criteria.

Selected summary statistics for 1,110 former lead workers and controls who participated in any visit of the Former Lead Worker Study, 1994–2008. As former lead workers were enrolled over time, and tibia lead and MRIs were measured at different times, these data can be used to evaluate selection bias over time.

Table 1

Variable	Former Worker N = 979	Control N = 131	No MRI N = 439	One MRI N = 309	Two MRIs N = 362	By MRI Status p-value
Age at enrollment, years, mean (SD)	56.5 (8.0)	58.6 (7.0)	57.1 (8.0)	57.6 (8.4)	55.6 (7.1)	0.002
Age at first MRI, years, mean (SD)	60.2 (8.1)	66.7 (6.3)	--	61.6 (8.5)	60.2 (7.8)	0.03
Employment duration, years, mean (SD)	8.0 (9.6)	--	7.2 (9.3)	8.2 (9.8)	8.8 (9.6)	0.10
Duration since last lead exposure, years, mean (SD)	18.5 (11.1)	--	19.6 (11.5)	19.5 (11.4)	16.9 (10.5)	0.006
Controls, N (%)	0 (0%)	131 (100%)	64 (14.6%)	22 (7.1%)	45 (12.4%)	
Enrollment year, N (%)						< 0.001
P1-Y1	437 (44.6%)	113 (87.0%)	260 (59.2%)	119 (38.5%)	172 (47.5%)	
P1-Y2	218 (22.3%)	14 (11.5%)	111 (25.3%)	51 (16.5%)	71 (19.6%)	
P1-Y3	48 (4.9%)	2 (1.5%)	21 (4.8%)	10 (3.2%)	19 (5.3%)	
P2-Y5	107 (10.9%)	0 (0%)	22 (5.0%)	36 (11.7%)	49 (13.5%)	
P2-Y6	169 (17.3%)	0 (0%)	25 (5.7%)	93 (10.1%)	51 (14.1%)	
Current tibia lead, N (%)	820 (83.8%)	80 (61.1%)	264 (60.1%)	274 (88.7%)	362 (100.0%)	
Current tibia lead, µg/g, mean (SD)	14.8 (9.7)	19.0 (10.8)	16.0 (9.7)	15.5 (10.6)	14.2 (9.22)	0.06
Peak tibia lead, N (%)	795 (81.2%)	0 (0%)	248 (56.5%)	242 (78.3%)	305 (84.3%)	
Peak tibia lead, µg/g, mean (SD)	25.0 (18.6)	--	27.6 (19.5)	26.9 (20.6)	21.4 (15.5)	< 0.001
MRI P2, N (%)	589 (60.2%)	67 (51.2%)	--	294 (95.2%)	362 (100.0%)	
MRI P3, N (%)	332 (33.9%)	45 (34.4%)	--	15 (4.9%)	362 (100.0%)	
APOE genotype, N (%)						0.49
Not genotyped	97 (9.9%)	49 (37.4%)	134 (30.5%)	11 (12.3%)	1 (3.0%)	
ε2/ε2	3 (0.3%)	1 (1.2%)	1 (0.3%)	1 (0.3%)	2 (0.6%)	

Variable	Former Worker N = 979	Control N = 131	No MRI N = 439	One MRI N = 309	Two MRIs N = 362	By MRI Status p-value
ε2/3	90 (10.2%)	12 (14.6%)	30 (9.8%)	27 (9.1%)	45 (12.5%)	
ε3/3	565 (64.1%)	48 (58.5%)	209 (68.5%)	185 (62.1%)	219 (60.7%)	
ε2/4	26 (3.0 %)	4 (4.9%)	9 (3.0%)	10 (3.4%)	11 (3.1%)	
ε3/4	176 (20.0%)	16 (19.5%)	48 (15.7%)	66 (22.2%)	78 (21.6%)	
ε4/4	22 (2.5%)	1 (1.2%)	8 (2.6%)	9 (3.0%)	6 (1.7%)	
CHS score, P2, mean (SD)	0.9 (1.5)	1.1 (1.5)	--	1.0 (1.5)	0.8 (1.4)	0.09
CHS score, P3, mean (SD)	1.9 (1.7)	2.6 (1.4)	--	1.7 (1.3)	2.0 (1.7)	0.48
TBV, MRI1 4D fit, cm ³ , mean (SD)	1171.7 (100.8)	1140.8 (100.8)	--	--	1167.8 (101.1)	
TBV, MRI2 4D fit, cm ³ , mean (SD)	1143.0 (99.4)	1102.5 (99.6)	--	--	1138.0 (100.2)	

* Defined as SBP > 140 or DBP > 90 or self-reported use of anti-hypertensive medications at enrollment visit. Note that, as the study questionnaire changed for phase III, diabetes and cardiovascular disease status could not be determined using the same method for earlier phases of the study.

† Comparing those with one MRI to those with two MRIs.

Table 2

Comparing former lead workers and controls with two MRIs (N = 362) on selected variables from the Phase III visit

Variable	Former Lead Workers	Controls	p-value
	N = 317	N = 45	
Age, years, mean (SD) *	64.1 (7.6)	71.9 (6.0)	< 0.001
Education, high school graduate with or without additional trade school, N (%)	239 (75.4)	30 (66.7)	0.40 [†]
White race/ethnicity, N (%)	284 (89.6)	42 (93.3)	0.43
APOE genotype, at least one e4 allele, N (%)	81 (25.6)	14 (31.1)	0.48

* In Phase III

** There are small differences in sample sizes for some of the lipid panel measurements (among those with two MRIs, one former lead worker was missing HDL and eight former lead workers and one control were missing LDL)

[†] p-value from five education group comparison

Table 3

Summary statistics for change in selected region of interest volume measures for 353* former lead workers and population controls without a history of occupational exposure to lead

ROI [†]	Delta ROI** (cm ³)		Delta ROI/TBV1 (%)	
	Mean (SD)	MIN, MED, MAX	Mean (SD)	MIN, MED, MAX
TBV	-29.87 (24.34)	-98.08, -31.62, 72.19	-2.55 (2.09)	-8.48, -2.69, 6.88
VENTRICLES	-0.53 (2.75)	-8.71, -0.73, 21.1	-0.043 (0.24)	-0.72, -0.06, 1.74
TOTAL GM	-24.43 (17.45)	-77.35, -24.74, 47.50	-2.09 (1.50)	-6.67, -2.19, 4.53
FRONT GM	-2.95 (5.18)	-23.99, -2.75, 14.49	-0.25 (0.44)	-2.09, -0.24, 1.46
OCCIP GM	-3.30 (1.82)	-8.42, -3.37, 2.75	-0.28 (0.15)	-0.75, -0.29, 0.26
PARI GM	-3.81 (2.67)	-13.36, -3.83, 7.40	-0.33 (0.23)	-1.16, -0.33, 0.71
TEMP GM	-2.69 (4.42)	-14.42, -2.76, 15.55	-0.23 (0.38)	-1.25, -0.22, 1.48
TOTAL WM	-5.44 (13.48)	-48.33, -6.41, 42.80	-0.46 (1.16)	-4.23, -0.55, 3.49
FRONT WM	-4.37 (5.50)	-27.72, -4.21, 19.00	-0.37 (0.46)	-2.42, -0.36, 1.55
OCCIP WM	0.56 (1.67)	-5.23, 0.51, 5.93	0.05 (0.14)	-0.41, 0.05, 0.52
PARI WM	-1.02 (3.03)	-10.76, -0.88, 12.39	-0.09 (0.26)	-0.94, -0.08, 1.01
TEMP WM	-2.40 (3.18)	-12.80, -2.35, 6.61	-0.20 (0.27)	-1.12, -0.20, 0.61
ERC	-0.30 (0.24)	-1.20, -0.29, 0.43	-0.03 (0.02)	-0.10, -0.03, 0.04
AMYG	-0.25 (0.21)	-1.02, -0.25, 0.49	-0.02 (0.02)	-0.09, -0.02, 0.05
HIPPO	-0.48 (0.39)	-1.89, -0.47, 0.87	-0.04 (0.03)	-0.16, -0.04, 0.09
CEREB	-3.42 (4.40)	-14.73, -4.27, 18.74	-0.30 (0.37)	-1.52, -0.36, 1.41
MEDIAL	-4.53 (3.03)	-13.63, -4.51, 7.89	-0.39 (0.25)	-1.19, -0.39, 0.75
INSULA	-1.26 (0.76)	-3.90, -1.17, 1.08	-0.11 (0.06)	-0.34, -0.10, 0.10
CINGULATE	-0.40 (1.26)	-5.76, -0.43, 4.69	-0.03 (0.11)	-0.50, -0.04, 0.45
CORP CALL	-0.96 (0.48)	-2.81, -0.88, 0.47	-0.08 (0.04)	-0.25, -0.08, 0.05
INT CAPS	-0.28 (0.44)	-2.31, -0.25, 1.09	-0.02 (0.04)	-0.20, -0.02, 0.09

* Of the 362 persons with two MRIs, 8 former lead workers and 1 control had first MRIs that were of insufficient quality for analysis.

** Delta ROI = volume at second MRI minus volume at first MRI; all volumes combine bilateral structures.

[†] TBV = total brain volume (TBV1 = TBV at first MRI); GM = gray matter; FRONT = frontal; OCCIP = occipital; PARI = parietal; TEMP = temporal; WM = white matter; ERC = entorhinal cortex; AMYG = amygdala; HIPPO = hippocampus; CEREB = cerebellum; MEDIAL = medial structures (bilateral amygdala, cuneus, entorhinal cortex, hippocampal formation, lingual gyrus, medial front-orbital gyrus, medial frontal gyrus, medial occipito-temporal gyrus, parahippocampal gyrus, perirhinal cortex, precuneus, and uncus); CORP CALL = corpus callosum; INT CAPS = internal capsule.

Table 4

Linear regression^a results for delta ROI models for former lead workers and controls (N = 352), adjusting for confounding variables.

ROI	Baseline age Beta (SE)	Duration between MRIs Beta (SE)	Peak Tibia Lead Beta (SE)
TBV	-1.771 (0.154) ***	-2.533 (3.002)	0.1496 (0.0865)
VENTRICLES	0.102 (0.021) ***	1.230 (0.387) ***	-0.0001 (0.0111)
TOTAL GM	-1.072 (0.119) ***	-7.162 (2.260) ***	0.0906 (0.0646)
FRONT GM	-0.350 (0.035) ***	-1.473 (0.658) **	0.0217 (0.0186)
OCCIP GM	-0.072 (0.012) ***	-0.866 (0.245) ***	0.0184 (0.0071) **
PARI GM	-0.163 (0.018) ***	-0.523 (0.345)	0.0137 (0.0102)
TEMP GM	-0.252 (0.029) ***	-1.528 (0.561) ***	0.0245 (0.0161)
TOTAL WM	-0.758 (0.089) ***	4.904 (1.779) ***	0.0544 (0.0505)
FRONT WM	-0.185 (0.039) ***	1.201 (0.763)	0.0224 (0.0220)
OCCIP WM	-0.064 (0.012) ***	0.618 (0.232) ***	-0.0047 (0.0065)
PARI WM	-0.097 (0.021) ***	1.014 (0.429) **	0.0174 (0.0124)
TEMP WM	-0.107 (0.022) ***	1.436 (0.446) ***	0.0104 (0.0128)
ERC	-0.005 (0.001) ***	-0.0004 (0.028)	0.0013 (0.0008)
AMYG	-0.008 (0.001) ***	0.054 (0.028) *	0.0001 (0.0008)
HIPPO	-0.015 (0.003) ***	0.090 (0.050) *	-0.0002 (0.0015)
CEREB	-0.213 (0.027) ***	-0.749 (0.540)	-0.0125 (0.0157)
MEDIAL	-0.188 (0.020) ***	-0.217 (0.380)	0.0077 (0.0110)
INSULA	-0.029 (0.005) ***	0.084 (0.833)	0.0044 (0.0029)
CINGULATE	-0.081 (0.007) ***	0.001 (0.142)	0.0058 (0.0042)
CORP CALL	0.005 (0.003)	0.053 (0.065)	0.0002 (0.0019)
INT CAPS	-0.010 (0.003) ***	0.026 (0.060)	0.0001 (0.0018)

* 0.05 < p < 0.10;

** 0.01 < p < 0.05;

*** p < 0.01

^aRegressions also included *APOE* genotype (2-3, 2-4, and 3-4 plus 4-4, each compared to 3-3 as reference group), height, baseline ROI, control status, duration between MRIs, and education. The model with peak tibia lead was in former lead workers only.