It must be born in mind that this glucose solution is strongly hypertonic (1110 mosmol/L), so that it would be expected that this should irritate the wall of the vein. After repeated episodes of hypoglycemia and repeated glucose injections, all peripheral veins will be obliterated sooner or later. Rather than using high percentage glucose solutions, it is better to use isotonic 5% glucose solution, as even a slight increase in blood sugar is enough to recover consciousness. An alternative is treatment with IM glucagon (2).

DOI: 10.3238/arztebl.2010.0303a

REFERENCES

1. Meyburg J, Bernhard M, Hoffmann GF, Motsch J: Principles of Pediatric Emergency Care. Dtsch Arztebl Int 2009; 106: 739–48.
2. Seewi O, Land C: Metabolische und endokrine Entgleisungen. In: Kretz FJ (ed.) Kinder Notfall-Intensiv. 3rd Edition. Elsevier 2010; 281–3.

Dr. med. Ora Seewi

Kinder-Endokrinologie und Diabetologie
Klinik und Poliklinik für Allgemeine Kinderheilkunde der Universität zu Köln
Kerpener Str. 62
50924 Köln, Köln
ora.seewi@uk-koeln.de

Pay Attention to the Target Group

The paper discussed (3) was addressed to emergency physicians. The rate of severely ill children is low (less than 1% of all emergency cases). Because of the low incidence we suggest hands-on training possibilities for emergency physicians in pediatric and pediatric surgery departments. As a prerequisite in caring for paediatric emergency patients the equipment (drugs, material etc.) available plays a key role. The regular equipment in the German EMS is according to DIN.

The authors should address the following points:

- Non-invasive oscillometric blood pressure measurement only assesses the mean arterial pressure. Values like 98/64 mm Hg give a fiction of accuracy.
- Do the authors only use cuffed endotracheal tubes (sometimes inflated, sometimes not), or do they mean cuffed and uncuffed tubes?
- Intravenous piritramide can produce respiratory arrest in a neonate even at a dose of 50 µg/kg. A dose of 30 µg/kg would be safer and could be repeated in 10 minutes if necessary. Morphine is more potent than piritramide (1:0.75), not only for its primary effect, but for its side effects as well, and should NOT be given at the same doses. Our experience suggests that opioids are underused in pediatrics despite their immense clinical value. There are no nebulizers in first aid materials, although families often have them.
- Although EMS vehicles generally do not carry nebulizers, the families of the affected patients often already have one at home.
- In general, emergency physicians have more experience with midazolam than with diazepam. Midazolam has nearly only inactive metabolites. If a seizure continues until the emergency physician arrives (12–15 min), it is a mistake to administer

a rectal drug. Intravenous midazolam and propofol are to be preferred.

- According to the ERC guidelines after the third attempt to place an IV catheter line or after 120 sec an intraosseous needle should be placed. What are the relevant prehospital risks of IO-access seen by the authors? Furthermore, the 22 G peripheral venous catheter is the smallest catheter in the EMS; this could be used to deliver 7 mL/kg/min of fluid to a child weighing 5 kg. Fluid resuscitation using a 26 G peripheral venous catheter is impossible. The recommendations of the association of German pediatric anesthesiologists for fluid management (1) represent the evidence, as does the work of Murat and Dubois. Fluid replacement should be done with a balanced, acetate-based full electrolyte solution, in children <3 years with the addition 1% glucose.
- The authors propose a treatment of diabetic ketoacidosis with 0.9% NaCl solution or 0.45% NaCl solution with glucose, but at the same time they warn of a cerebral edema of unknown etiology. Under conditions of hyperglycemia and hyponatremia a cerebral edema is anticipated. The free water infused follows the concentration gradient of sodium ions into the cells (2). A balanced, acetate-based full electrolyte solution would be better.

DOI: 10.3238/arztebl.2010.0303b

REFERENCES

1. Sümpelmann R, et al.: Empfehlungen zur perioperativen Infusionstherapie bei Neugeborenen, Säuglingen und Kleinkindern. Anästh Intensivmed 2007; 48: 73–7.
2. Arieff A: Postoperativ hyponatraemische encephalopathy following elective surgery in children. Paediatric Anaesthesia 1998; (8): 1–4.
3. Meyburg J, Bernhard M, Hoffmann GF, Motsch J: Principles of Pediatric Emergency Care. Dtsch Arztebl Int 2009; 106: 739–48.

Dr. med. Thomas Mader DEAA

Rosenstr. 16
06114 Halle (Saale), Germany
maderth@gmx.de

Additions Are Needed

As advance analgesia is not possible, venipuncture must be performed on the conscious child.

If there is an indication for venipuncture, tedious and painful failed punctures must be avoided. A maximum of two attempts should be made at the classical and promising sites of puncture. If those fail, intra-osseous puncture should be performed. The possible complications are of secondary importance only. The best puncture site is at the transition between the proximal and mid third of the lower leg. If the Cook needle is used, the infant's lower leg must rest on the palm of the operator, in order to avoid greenstick fractures.

The sentence "Symptomatic hyperglycemia in childhood is almost always due to diabetic ketoacidosis" is totally unacceptable. Ketoacidosis is a symptom, not the cause of excessive hyperglycemia. Volume replacement is actually very important in the therapy of hyper-

glycemic ketotacidosis. If however 0.9% NaCl solution is infused, the acidosis may be enhanced, as there is a lack of metabolisable anions. A balanced electrolyte solution is to be preferred here.

Table 3 is not very helpful. It is hardly possible to distinguish between “dry” and “dried out” mucus membranes. Experience has shown that the clinical evaluation of the child should be based on the general condition, respiration and circulation, with the help of the Pediatric Assessment Triangle. This tool is a valuable didactic aid when the severity of a pediatric emergency is to be assessed.

DOI: 10.3238/arztebl.2010.0303c

REFERENCES

1. Müller-Lobeck L, Birnbaum J, Spies C: Kindernotfälle – Allgemeine Behandlungsstrategien, Anästhesiol Intensivmed Notfallmed Schmerzther 2009; (44)6: 422–7.
2. Heinzel O, Schwandt JC, Eppich WJ: Das ABC des lebensbedrohlichen Kindernotfalls. Monatsschr Kinderheilkd 2009; 157: 801–16.
3. Meyburg J, Bernhard M, Hoffmann GF, Motsch J: Principles of Pediatric Emergency Care. Dtsch Arztebl Int 2009; 106: 739–48.

Dr. med. Lutz Müller-Lobeck

Kinderanästhesie
Charité Universitätsmedizin Berlin
Klinik für Anästhesiologie mit Schwerpunkt operative Intensivmedizin
Campus Virchow Klinikum und Campus Charité Mitte
Augustenburger Platz 1
13353 Berlin, Germany
lutz.mueller-lobeck@charite.de

Use Familiar Material

The authors are quite right to state that, “Securing the airway [...] requires special knowledge and manual skills”. What is difficult to understand is the assertion that “oro-tracheal intubation [...] is to be preferred in emergencies”. The primary issue is really not intubation, but oxygenation.

If a child has a difficult airway, or if ventilation with a face mask turns out to be difficult or impossible, this is mostly due to errors in the positioning of the head or mask, or of the patient’s body. If positioning, the Esmarch maneuver, lifting the chin or Guedel’s tubes are all unsuccessful, the anesthesia must be immediately deepened and an alternative airway must be selected. As many physicians have had extensive experience with the laryngeal mask, the laryngeal mask should be the first choice in children.

Most emergency physicians are not pediatricians and are used to curved McIntosh’s spatulas. To avoid mistakes, they should use familiar material, particularly under stress or in emergencies. If Miller’s spatula fails to lift the epiglottis, intubation is impossible.

The cricoid cartilage (not “below the cricoid cartilage”) is the physiological bottleneck in a child’s respiratory tract and must be passed atraumatically. If there is damage, this may have happened because the selected tube is too wide. If intubation is necessary, a small but cuffed tube must be regarded as the gold standard. The resistance of the respiratory tract is irrelevant in ventilation. And why should an excessively small tube cause hyperexpansion of the lung?

Even under routine conditions, intubation of children requires enormous experience and regular training. Safeguarding a child’s respiratory tract is an enormous challenge for many emergency physicians. Although endotracheal intubation is established as the gold standard within the hospital, it can hardly be used preclinically. In this situation the supraglottic airway must be used and will guarantee a safe airway. The physician performing the intubation must master this flawlessly.

DOI: 10.3238/arztebl.2010.0304a

REFERENCES

1. Hoehne C, Haack M, Machotta A, Kaisers U: Atemwegsmanagement in der Kinderanästhesie, Anaesthesist 2006; 55: 809–20.
2. Holzki J: Laryngeal damage from tracheal intubation. Paediatric An-aesthesia 1997; 7: 435–7.
3. Müller-Lobeck L, Birnbaum J, Spies C: Kindernotfälle – Allgemeine Behandlungsstrategien, Anästhesiol Intensivmed Notfallmed Schmerzther 2009; (44) 6: 422–7.
4. Meyburg J, Bernhard M, Hoffmann GF, Motsch J: Principles of Pediatric Emergency Care. Dtsch Arztebl Int 2009; 106: 739–48.

Prof. Dr. med. Jochen Strauß

Dr. med. Karin Becke

Wissenschaftlicher Arbeitskreis Kinderanästhesie der DGAI
HELIOS-Klinikum Berlin Buch
Klinik für Anästhesie, Perioperative Medizin und Schmerztherapie
Schwanebecker Chaussee 50
13125 Berlin, Germany
jochen.strauss@helios-kliniken.de

In Reply:

We would like to thank our colleagues for the detailed discussion and for the valuable input from numerous letters. This has shown us that we have succeeded in waking the interest of many readers.

We have read the current review article referred to by Dr. Lukowski, but do not share his reservations about the administration of antipyretics. In most of the studies cited there, the fever could not be adequately reduced under the study medication (1). In one study, higher temperatures were even measured in children with recurrent febrile seizures than in the control group (2). We therefore do not regard it as justified to make a clear statement about the lack of reduction in the frequency of seizures. As Dr. Lukowski himself admits, the antipyretics help to control the fever and to safeguard the child’s well-being. If weight-adjusted doses are used and the contraindications respected, we see no evidence that the treatment is useless or even dangerous.

Genzwürker, Gernoth and Hinkelbein emphasize that the procedure of percutaneous coniotomy, as described by us, is not a standard procedure, for many reasons. They correctly point out that there is a DIN norm for first aid material. It is, however, common knowledge that, even for adult emergencies, desirable materials—or even material prescribed by the DIN norm—are often omitted. If medically approved products are not available in an emergency, the technique with simple materials which we describe (and use ourselves) can be employed to prevent potentially fatal hypoxia in the child, till the respiratory tract can be adequately safeguarded.