Published in final edited form as:

Synlett. 2009 December 1; 19: 3135–3138. doi:10.1055/s-0029-1218293.

Catalytic intermolecular hydroamination of vinyl ethers

Nirmal K. Pahadi and Jon A. Tunge

Department of Chemistry, University of Kansas, 1251 Wescoe Hall Drive, Lawrence, KS, 66045-7582

Abstract

This manuscript details the development of a palladium-catalyzed hydroamination of vinyl ethers. It is proposed that palladium catalyzes the hydroamination via Bronsted base catalysis, where palladium is protonated by the relatively acidic sulfonamide to generate a palladium hydride as well as the active anionic sulfonamide nucleophile. Thus, this process is distinct from known palladium-catalyzed hydroaminations of styrene derivatives that utilize less acidic amines.

Keywords

Hydroamination; vinyl ether; palladium; sulfonamide

Introduction

Transition metal-catalyzed intermolecular hydroamination of alkenes has attracted a great attention in recent years.¹ Much research in this line has centered on the *intramolecular* hydroamination of simple alkyl-substituted α -olefins.^{2,3} The analogous *intermolecular* hydroamination of alkenes has been most successful with electron deficient alkenes,⁴ dienes,⁵ and vinyarenes.⁶ In contrast to these reactions, the related transition metal-catalyzed hydroaminations of vinyl ethers have not been extensively investigated. Electron rich olefins such as vinyl ethers undergo palladium-catalyzed intramolecular hydroamination;⁷ intermolecular hydroaminations are catalyzed by acid⁸ or bifunctional rhenium-oxo catalysts.⁹

At the outset of our investigations, it was known that palladium can catalyze the hydroamination of olefins by several mechanisms. For instance, palladium(II) catalyzes the intramolecular hydroamination of vinyl ethers.⁷ This reaction is thought to occur via Lewis acid catalysis where palladium coordinates the olefin, thus polarizing it toward nucleophilic attack by the amine.^{7c,10} In addition, Hartwig has developed a hydroamination of styrenes that is proposed to go via acid-induced formation of π -benzyl complexes.⁶ Herein we report a catalytic intermolecular hydroamination of vinyl ethers that we propose utilizes low valent palladium as a specialized Bronsted base catalyst.

Results and Discussion

We began investigating the potential for hydroamination of 1-(vinyloxy)butane **2a** with a variety of aniline derivatives. Heating *N*-alkyl or *N*-acyl anilines in the presence of 5 mol% Pd (PPh₃)₄ at 100 °C resulted in no product formation. However, the more acidic *N*-tosyl aniline produced the corresponding addition product **3a**, which was obtained in excellent yield (Table

1, entry 4) when the reaction was performed in excess, neat vinyl ether. Lowering the concentration of vinyl ether to 1.6 M (4 equiv.), slightly lowers the yield (entry 5), while using just 2 equivalents considerably lowered the yield of aminol (entry 6) and increased time necessary for reaction completion. Moreover, substantial quantities of enamine **4a** were formed. Further lowering the concentration of **2** to 0.5 M, dramatically decreased the yield of **3** and only a trace amount of enamine **4** was observed. Dioxane, CH₃CN, (CH₂Cl)₂, and THF were also tested as potential solvents; only dioxane provided any product. However, when 1,4-dioxane was used as solvent, a 1:1 mixture of **3a** and **4a** was obtained as judged by ¹H NMR spectroscopy. In addition, a variety of other palladium catalysts were screened, but none was as effective at promoting the hydroamination as Pd(PPh₃)₄ (entries 7–10). Thus, reactions could be run in neat vinyl ether or in concentrated toluene solutions. For convenience, we chose to run reactions in neat vinyl ether for examination of the scope of the hydroamination.

Importantly, attempting to perform the same transformation without the addition of Pd(PPh₃)₄ produced < 5% yield of **3a** over 17 h at 100 °C (entry 11). Furthermore, since such a transformation could potentially be catalyzed by simple Bronsted acids,^{3,8} the reaction was run with 5 mol% TfOH with or without Pd(PPh₃)₄ and <5% product was observed (entries 12 and 13). Instead, TfOH promoted the decomposition of the butyl vinyl ether to acetaldehyde dibutyl acetal.¹¹ The reaction did proceed with Pd(PPh₃)₄ and 10 mol% benzoic acid (entry 14), however the acid additive inhibited the reaction and longer times were required for completion of the reaction. Importantly, the observed inhibition by acid contrasts with the Pd(PPh₃)₄-catalyzed hydroamination of vinylarenes which requires strong acid additives.⁶ Thus, we propose that the mechanism of the palladium-catalyzed hydroamination of vinyl ethers differs from that of vinyl arenes.

Given the inhibition of the reaction by added acid, we hypothesized that the Pd catalyst was behaving as a Bronsted base to facilitate the hydroamination. If this is the case, then the triphenylphosphine that accompanies the palladium may also be capable of acting as a base and catalyzing the reaction.⁸ Indeed, the reaction also proceeded in the presence of 20 mol % PPh₃ as a Lewis base, however the conversion was only 25 % after extended reaction time. Thus, phosphine catalysis does not account for our observed results.

Having the optimized conditions in hand, the scope and limitations of the hydroamination of butyl vinyl ether with a variety tosyl protected amines was explored (table 2). Both electron rich and electron poor anilines are good substrates for the reaction and groups like esters, nitro groups, and nitriles are compatible with the reaction conditions. Interestingly, *ortho*-substitution of the aniline lead to exclusive formation of the enamine product (entries 3 and 4, table 2) in good to moderate yield. Unfortunately, high concentrations of the vinyl ether were often necessary to promote formation of the aminol over the enamine. Lastly, when an aliphatic amine was exposed, the anticipated addition product **3k** was isolated in moderate yield along with enamine **4k**.

Next we turned our attention to investigating the vinyl ethers that were compatible reaction partners in the hydroamination. Simple aliphatic vinyl ethers react smoothly to afford the desired aminols except in the case of vinyl cyclohexyl ether, which gives only a low yield of aminol (table 3, entry 7). Interestingly, the bulkier *tert*-butyl ether forms the aminol in good yield (table 3, entry 5). Cyclic vinyl ethers undergo the palladium-catalyzed hydroamination as well. Both 5- and 6- membered vinyl ethers undergo hydroamination to provide the cyclic aminols (eq. 1). An aliphatic amine also participates in this transformation, however the product **3u** is formed in rather low yield.

(1)

A plausible mechanism for this transformation is outlined in Scheme 2. Protonation of Pd(0) with the tosyl aniline ($\text{pK}_a \sim 11.5$ in DMSO) produces a palladium hydride species^{12,13} which can undergo coordination of the vinyl ether followed by insertion to form an alkylpalladium intermediate **B**. Related carbopalladations of vinyl ethers have been proposed by Jordan.¹⁴ This intermediate can undergo nucleophilic substitution by the tosyl amide, probably via an $\text{S}_{\text{N}}1$ mechanism, to produce the observed product. The proposed intermediacy of the tosyl amide anion is supported by the fact that we observe acid inhibition. In addition, we observed near quantitative formation of $\text{TsNHCH}_2\text{CH}_2\text{Cl}$ when reactions were attempted in dichloroethane. Finally, the enamine byproduct may arise from elimination of alcohol from the aminol product, which is promoted by sterically bulky *ortho*-substituted anilines.

In conclusion, we have developed a Pd(0)-catalyzed hydroamination of vinyl ethers. The reaction occurs under formally neutral conditions and we propose that the Pd(0) catalyst acts as a Brønsted base to facilitate the hydroamination. This mechanism contrasts with that of the hydroamination of styrenes with less acidic amines.

Experimental Section

General procedure for the hydroamination of vinyl ethers

The preparation of **3a** is representative. To butyl vinyl ether (**2a**) (0.49 mL, 3.8 mmol) and Pd(PPh_3)₄ (11.55 mg, 0.01 mmol), was added *N*-tosylaniline (49.41 mg, 0.2 mmol) in a well-sealed pressure vial and heated to 100 °C in an aluminum block for the designated time. The completion of reaction was monitored via TLC and the reaction stopped when amine was completely consumed. The reaction mixture was passed through a short pad of silica gel and the product was isolated by flash column chromatography using silica gel as the stationary phase and 4:1 hexane+ethyl acetate as the eluent. 66 mg (94%) of the product *N*-(1-butoxyethyl)-4-methyl-*N*-phenylbenzene-sulfonamide (**3a**) was obtained as a slightly yellow oil. ¹H NMR (400 MHz, CDCl_3) δ ppm 7.56 (d, $J = 8.0$ Hz, 2H), 7.37 – 7.28 (m, 3H), 7.24 (d, $J = 8.0$ Hz, 2H), 7.05 – 7.01 (m, 2H), 5.69 (q, $J = 6.0$ Hz, 1H), 3.77 (dt, $J = 9.3, 6.7$ Hz, diastereotopic 1H), 3.56 (dt, $J = 9.3, 6.7$ Hz, diastereotopic 1H), 2.43 (s, Ts CH_3), 1.58 (tt, $J = 13.5, 4.0$ Hz, 2H), 1.37 (dq, $J = 14.9, 7.3$ Hz, 2H), 1.14 (d, $J = 6.0$ Hz, 3H), 0.94 (t, $J = 7.3$ Hz, 3H); ¹³C NMR (125 MHz, CDCl_3) δ ppm 143.17, 137.67, 134.44, 131.96, 129.18, 128.60, 128.54, 127.52, 85.06, 68.05, 31.51, 21.46, 20.62, 19.31, 13.88; IR (CH_2Cl_2): ν 3259, 2931, 1598, 1494, 1338, 1091, 919, 813, 754 cm^{-1} ; Calcd. HRMS for $\text{C}_{19}\text{H}_{25}\text{NO}_3\text{SNa}$ ($\text{M}+\text{Na}$), 370.1453; Found, 370.1450.

Acknowledgments

We thank the National Institute of General Medical Sciences (1R01GM079644).

References

- (1). (a) Mueller TE, Hultsch KC, Yus M, Foubelo F, Tada M. Chem. Rev 2008;108:3795–3892. [PubMed: 18729420] (b) Widenhoefer RA, Han X. Eur. J. Org. Chem 2006:4555–4563. (c) Hazari N, Mountford P. Acc. Chem. Res 2005;38:839–849. [PubMed: 16285707] (d) Hultsch KC. Org. Biomol. Chem 2005;3:1819–1824. [PubMed: 15889160] (e) Hong S, Marks TJ. Acc. Chem. Res 2004;37:673–686. [PubMed: 15379583] (f) Mueller TE, Beller M. Chem. Rev 1998;98:675–703. [PubMed: 11848912]
- (2). (a) Ohimiya H, Moriya T, Sawamura M. Org. Lett 2009;11:2145–2147. [PubMed: 19379007] (b) Hesp KD, Stradiotto M. Org. Lett 2009;11:1449–1452. [PubMed: 19231849] (c) Biyikal M, Lohnwitz K, Roesky PW, Bechert S. Synlett 2008:3106–3110. (d) Graebe K, Pohlki F, Doye S. Eur. J. Org. Chem 2008:4815–4823. (e) Majumder S, Odom AL. Organometallics 2008;27:1174–1177. (f) Liu Z, Hartwig JF. J. Am. Chem. Soc 2008;130:1570–1571. [PubMed: 18183986]
- (3). For recent acid-catalyzed hydroaminations see: (a) Zigang Li, Zhang J, Brouwer C, Yang C-G, Reich NW, He C. Org. Lett 2006;8:4175–4178. [PubMed: 16956180] . (b) Motokura K, Nakagiri N, Mori K, Mizugaki T, Ebitani K, Jitsukawa K, Kaneda K. Org. Lett 2006;8:4617–4620. [PubMed: 16986964] . (c) Rosenfeld DC, Shekhar S, Takemiya A, Utsunomiya M, Hartwig JF. Org. Lett 2006;8:4179–4182. [PubMed: 16956181] . (d) Schlummer B, Hartwig JF. Org. Lett 2002;4:1471–1474. [PubMed: 11975606] . (e) Colinas PA, Bravo RD. Org. Lett 2003;5:4509–4511. [PubMed: 14602037] . (f) Toshima K, Nagai H, Ushiki Y, Matsumura S. Synlett 1998:1007.. (g) Karshedt D, Bell AT, Tilly TD. J. Am. Chem. Soc 2005;127:12640–12646. [PubMed: 16144412] . (h) Widenhoefer R. Angew. Chem., Int. Ed 2006;45:1747–1749.. (i) Jiao P, Nakashima D, Yamamoto H. Angew. Chem., Int. Ed 2008;47:2411–2413..
- (4). (a) Kawatsura M, Hartwig JF. Organometallics 2001;20:1960–1964. (b) Fadini L, Togni A. Chem. Commun 2003:30–31. (c) Li K, Hii KK. Chem. Commun 2003:1132–1133. (d) Munro-Leighton C, Blue ED, Gunnoe TB. J. Am. Chem. Soc 2006;128:1446–1447. [PubMed: 16448104]
- (5). (a) Zhou J, Hartwig JF. J. Am. Chem. Soc 2008;130:12220–12221. [PubMed: 18715004] (b) Giner X, Najera C. Org. Lett 2008;10:2919–2822. [PubMed: 18553970] (c) Qin H, Yamagiwa N, Matsunaga S, Shibasaki M. J. Am. Chem. Soc 2006;128:1611–1614. [PubMed: 16448133]
- (6). (a) Johns AM, Utsunomiya M, Incarvito CD, Hartwig JF. J. Am. Chem. Soc 2006;128:1828–1839. [PubMed: 16464081] (b) Utsunomiya M, Hartwig JF. J. Am. Chem. Soc 2003;125:14286–14287. [PubMed: 14624571] (c) Nettekoven U, Hartwig JF. J. Am. Chem. Soc 2002;124:1166–1167. [PubMed: 11841274] (d) Kawatsura M, Hartwig JF. J. Am. Chem. Soc 2000;122:9546–9547. (e) Utsunomiya M, Hartwig JF. J. Am. Chem. Soc 2004;126:2702–2703. [PubMed: 14995178] (f) Utsunomiya M, Kuwano R, Motoi K, Hartwig JF. J. Am. Chem. Soc 2003;125:5608–5609. [PubMed: 12733880]
- (7). (a) Tada M, Shimamoto M, Sasaki T, Iwasawa Y. Chem. Commun 2004:2562–2563. (b) Penzien J, Su RQ, Muller TE. J. Mol. Catal. A 2002:182–183. 489–498. (c) Vogels CM, Hayes PG, Shaver MP, Westcott SA. Chem. Commun 2000:51–52.
- (8). Colinas PA, Bravo RD. Org. Lett 2003;5:4509–4511. [PubMed: 14602037]
- (9). Sherry BD, Loy RN, Toste FD. J. Am. Chem. Soc 2004;126:4510–4511. [PubMed: 15070356]
- (10). Cochran B, Michael FE. J. Am. Chem. Soc 2008;130:2786–2792. [PubMed: 18254623]
- (11). Shainyan BA, Grigor'eva A. A. Russ. J. Org. Chem 2001;37:1177–1178.
- (12). Knight JG, Ainge SW, Harm AM, Harwood SJ, Maughan HI, Armour DR, Hollinshead DM, Jaxa-Chamiec AA. J. Am. Chem. Soc 2000;122:2944–2945.
- (13). Trost BM. Chem. Eur. J 1998;4:2405–2412.
- (14). Luo S, Jordan RF. J. Am. Chem. Soc 2006;128:12072–12073. [PubMed: 16967949]

Scheme 1.

Table 1

Conditions for hydroamination^a

entry	catalyst	PG	time (h)	3a	4a
1 ^b	Pd(PPh ₃) ₄	Ac	24	<5%	<5%
2 ^b	Pd(PPh ₃) ₄	Me	24	<5%	<5%
3 ^b	Pd(PPh ₃) ₄	Bn	24	<5%	<5%
4	Pd(PPh ₃) ₄	Ts	15	94%	<5%
5 ^c	Pd(PPh ₃) ₄	Ts	12	82%	<5%
6 ^d	Pd(PPh ₃) ₄	Ts	24	50%	25%
7	PdCl ₂ (PhCN) ₂	Ts	24	10%	<5%
8	Pd ₂ dba ₃ /BINAP	Ts	24	80%	<5%
9	Pd ₂ dba ₃ /dppb	Ts	24	72%	<5%
10	Pd ₂ dba ₃ /dppf	Ts	24	75%	<5%
11	none	Ts	17	<5%	<5%
12	5 mol% TfOH	Ts	12	<5%	<5%
13	5 mol% TfOH/Pd(PPh ₃) ₄	Ts	12	<5%	<5%

entry	catalyst	PG	time (h)	3a	4a
14	10 mol% PhCO ₂ H/Pd(PPh ₃) ₄	Ts	41	85%	<5%

^a Isolated yields of reactions carried out in a sealed vial with **1a** (0.2 mmol) and neat **2a** at 100 °C.

^b Starting material was recovered.

^c 1.6 M olefin in toluene

^d 2 equiv **2a**

Table 2

Scope of hydroamination^a

entry	R ¹	2 (equiv)	time (h)	3 _s (%) ^b	4 _s (%) ^b
1	<i>p</i> -Br-C ₆ H ₄	19	14	3b : 67	-
2	<i>m</i> -Me-C ₆ H ₄	19	12	3c : 92	-
3	<i>o</i> -Me-C ₆ H ₄	19	20	-	4d : 90
4	<i>o</i> -Me ₂ -C ₆ H ₃	19	20	-	4e : 45
5	<i>p</i> -OMe-C ₆ H ₄	19	25	3f : 80	-
6	<i>p</i> -OMe-C ₆ H ₄	4b	25	3f : 70	-
7	<i>m</i> -OMe-C ₆ H ₄	19	24	3g : 80	-
8	<i>m</i> -OMe-C ₆ H ₄	4b	24	3g : 60	-
9	<i>p</i> -NO ₂ -C ₆ H ₄	4b	15	3h : 60	-
10	<i>p</i> -CN-C ₆ H ₄	19	26	3i : 93	-
11	<i>p</i> -CO ₂ Et-C ₆ H ₄	19	15	3j : 93	-
12	<i>p</i> -CO ₂ Et-C ₆ H ₄	4b	15	3j : 50	4j : (50)
13	CH ₂ (CH ₂) ₃ CH ₃	19	24	3k : 40	4k : <20

^a All reactions were carried out in a sealed vial with **1** (0.2 mmol) and **2** in the presence of 5 mol % Pd(PPh₃)₄ at 100 °C unless otherwise indicated.

b_1 in toluene solvent

Table 3

Scope of vinyl ethers^a

$$\begin{array}{c}
 \text{Ts} \\
 | \\
 \text{R}^1 \text{N}^+ \text{H} \\
 \text{1}
 \end{array}
 +
 \begin{array}{c}
 \text{OR}^2 \\
 | \\
 \text{CH}=\text{CH} \\
 \text{2}
 \end{array}
 \xrightarrow[100^\circ\text{C}]{5\text{ mol\% Pd(PPh}_3)_4}
 \begin{array}{c}
 \text{Ts} \\
 | \\
 \text{R}^1 \text{N}^+ \text{OR}^2 \\
 \text{3}
 \end{array}$$

entry	R ¹	R ²	equiv 2	time (h)	3, (%) ^b
1	C ₆ H ₅	<i>n</i> -Pr	19	36	3l : 75
2	<i>p</i> -F-C ₆ H ₄	<i>n</i> -Bu	8 ^b	14	3m : 92
3	<i>p</i> -F-C ₆ H ₄	<i>n</i> -Pr	4 ^b	14	3n : 87
4	<i>p</i> -F-C ₆ H ₄	Et	19	26	3o : 89
5	<i>p</i> -F-C ₆ H ₄	<i>t</i> -Bu	19	26	3p : 82
6	<i>p</i> -CF ₃ -C ₆ H ₄	<i>n</i> -Bu	8 ^b	14	3q : 82
7	<i>p</i> -CN-C ₆ H ₄	Cy	19	25	3r : 30

^a All reactions were carried out in a sealed vial with **1** (0.2 mmol) in the presence of 5 mol % Pd(PPh₃)₄ at 100 °C

^b in toluene