Published in final edited form as:

Synlett. 2010 February 1; 2010(3): 470–474. doi:10.1055/s-0029-1219186.

Synthesis of Chiral Nonracemic Tertiary α -Thio and α -Sulfonyl Acetic Esters via S_N2 reactions of Tertiary Mesylates

Jimmie D. Weaver, David K. Morris, and Jon A. Tunge

Department of Chemistry, University of Kansas, 1251 Wescoe Hall Drive, Lawrence, KS, 66045-7582 Fax: 785-864-5396

Jon A. Tunge: nge@ku.edu

Abstract

Syntheses of enantioenriched sulfides and sulfones via substitution of tertiary mesylate with thiolate nucleophile with modest to excellent success.

Keywords

sulfide; sulfone

Introduction

Recently, we have shown that the α -sulfonyl allyl esters cleanly undergo decarboxylative coupling to afford tertiary sulfones, making α -sulfonyl allyl esters a powerful precursor to tertiary sulfones that are not easily synthesized by other methods (Scheme 1).¹ In the course of this research we became interested in the synthesis of chiral nonracemic allyl α -sulfonylacetic esters.

To our knowledge, a general, simple, enantioselective method to fully substituted α -sulfonyl acid derivatives does not exist. Typically syntheses of optically active compounds have been racemic and rely on preparatory chiral HPLC to obtain individual stereoisomers.² Alternatively, modest success at the asymmetric synthesis has been achieved with the use of a chiral oxazolidine auxiliary to both quaternize the α -center and control the absolute stereochemistry of the sulfenylation.³ However, this methodology requires harsh and toxic reagents and is not general.

A scan of the literature revealed several less direct strategies for the synthesis of α -sulfonyl acid derivatives including chiral resolution of α -thio acids⁴ as well as several procedures for the activation and substitution of α -hydroxy esters.^{5,6} More specifically, several early reports obtained enantiomerically enriched α -thio and α -sulfonyl acids via resolution of the racemic α -thio acids with enantiopure amines via crystallization of diastereomeric salts.⁴ While this is a reliable method, it is not general and it requires stoichiometric amounts of chiral amine and requires repeated recrystallizations, resulting in very low overall yields (Scheme 2). Furthermore the optimal amine is substrate dependent; consequently selecting the appropriate amine requires a separate search and optimization for each substrate.

We felt that it would be possible to approach the synthesis of optically active α -thio and α -sulfonyl acids by a direct S_N2 displacement of an enantioenriched tertiary α -hydroxy ester derivative. Specifically, it was expected that the electron-withdrawing ester group would deactivate the requisite substrates toward S_N1 reactions. It was also expected that the planar nature of the ester group would better accommodate the sterically demanding transition state

for S_N2 displacement. Thus, we posited that combining these two features with the use of highly nucleophilic thiolates would allow S_N2 displacement at tertiary carbon centers (Scheme 3).^{5,7,8}

Searching the literature revealed two examples of S_N2 displacements of tertiary α -hydroxy acid derivatives with thiolates. Mukaiyama reported S_N2 displacement of diphenyl phosphinite derived from α -hydroxy acids in the presence of the appropriate quinuclidine and benzothiazole⁹ and Corey reported the displacement of a mesylate derived from the α -hydroxy atrolactic acid.⁶ With the goal of verifying these procedures with our substrates, we attempted to repeat the known Mitsunobu variant that has been used for the synthesis similar α -thio esters.⁹ Unfortunately this method did not work well in our hands (Scheme 4).

Consequently, it was decided to investigate the simpler, albeit less direct, two-step approach involving sequential activation and substitution of an α -hydroxy ester. Previously, Corey converted (+)-atrolactic acid to the corresponding sulfone by this protocol, although the experimental details were not provided and no reaction scope was demonstrated.⁶ Herein we provide a detailed examination of the methodology, provide expanded scope of the reaction, and outline the limits of the methodology.

Results and Discussion

While Corey was successful in his substitution of atrolactic acid, our attempts to displace the mesylate from the more sterically demanding butanoate led solely to elimination (Scheme 5). Consequently, further investigation was limited to propanoic acid derivatives.

In order to further explore this methodology we synthesized several nonracemic hydroxy esters as outlined in table 1. α -Methyl styrenes **2a–e** were subjected to Sharpless asymmetric dihydroxylation, which typically afford highly enantioenriched derivatives (90–94% ee).¹⁰ Next, the resulting diols underwent Pt/C-catalyzed oxidation under mild conditions utilizing air as the terminal oxidant.¹¹ Using this protocol, the ester substituted arene derivative **3a**, failed to give significant quantities of the desired product **4a**. However, the remaining substrates (**3b–3e**) produced the corresponding α -hydroxy acids in good yields using this practical and simple synthesis.^{12,13}

Next, the hydroxy acids **4b–e** were subjected to typical reaction conditions for esterification. Unfortunately, the tertiary benzylic hydroxy acids are not well-suited for common Fisher or DCC/DMAP-catalyzed esterifications. However, it was gratifying to find that esterifications via an S_N2 reaction of the potassium carboxylates and allyl bromides yielded the desired hydroxy esters in high yields and purities (Table 2).¹⁴

With the requisite α -hydroxy esters in hand, we turned to evaluating the mesylation/substitution procedure for the synthesis of α -thio esters.⁶ The esters, **5f–m** were subjected to mesylation followed by careful workup, then immediately subjected to the conditions for substitution with sodium phenyl thiolate (Scheme 6).

As can be seen from the data in table 3, the yields of α -thio esters (**6**) using this protocol are moderate, at best. In particular, the more electron rich *p*-Me arene substrates (entries 2 and 8, Table 2) formed α -thioesters in dismal yields. The low yields in these cases are attributed to the formation of significant quantities of β -sulfides (**7**) that likely arise via elimination followed by conjugate addition of the thiolate to the acrylate derivative (Scheme 6). The mixture of **6** and **7** was subjected to oxidation with *m*CPBA.^{1, 15} Upon oxidation the α -thioesters **6** readily formed the corresponding sulfones, while the β -thioesters **7** underwent elimination to form acrylates. The stereo-chemical fidelity of the mesylation/substitution was good in general, although several substrates gave substantial racemization (entries 1, 6, and 7). The delicate

nature of the mesylates is illustrated by the much lower ee of **8i** (entry 4) when the mesylation was inadvertently performed at room temperature; mesylation of an analogous substrate at -5°C , provided the sulfone in excellent ee (entry 3). Thus, we hypothesized that the observed racemization was due to ionization of the intermediate mesylate and that the methodology might be better suited for α , α -dialkyl sulfonyl esters, whose requisite mesylates would ionize less readily.

To test this hypothesis, Sharpless dihydroxylation was used to provide chiral nonracemic diol which was selectively deoxygenated to give hydroxy ester **9b** in 61% yield from **9a** (Scheme 7).⁷ As expected the mesylation of **9b** results in a stable mesylate that could be purified by column chromatography and stored at low temperatures without any observed racemization. As posited, exposure of the mesylate to sodium thiolate led to clean substitution to provide the α -thio ester which was oxidized to form the highly enantioenriched sulfone **9c**. Here, we assume that the reaction occurred with inversion of configuration in analogy to the observations of Corey.⁶

Similarly, enantioenriched dialkyl α -sulfonyl esters may also be synthesized from 1,1-disubstituted olefins such as **10a** (Scheme 8) via a straightforward dihydroxylation/oxidation sequence. Specifically, the PMP protected alcohol, **10a** underwent asymmetric dihydroxylation and Pt-catalyzed oxidation to provide the crystalline α -hydroxy acid **10b** in excellent yield.¹⁶ Esterification of the acid provided the requisite α -hydroxy ester for mesylation. Treatment with MsCl provided the mesylate, in 81% yield. Once again, this α , α -dialkyl mesylate was much more stable and less prone to undergo racemization, remaining unchanged after months at room temperature. Subsequent substitution of the mesylate cleanly afforded the sulfide in 96% yield. Oxidation to the sulfone provided **10c**, with excellent enantioenrichment (94% ee), indicating that the substitution of the tertiary alcohol derivative proceeded cleanly via an $\text{S}_{\text{N}}2$ mechanism. The overall yield of sulfone **10c** from **10a** was a satisfactory 54%.

In conclusion we have demonstrated the limits of the two step activation/substitution methodology on several benzylic hydroxy esters. Problematic ionization of the tertiary α -mesyl- α -aryl esters indicates that the best substituents are those that electronically destabilize carbocation formation. Additionally, we have shown that the two step mesylation and substitution protocol works satisfactorily for α , α -dialkyl- α -hydroxyl esters to provide chiral non racemic tertiary α -thio and α -sulfonyl esters in high yields and ee's.

Experimental Section

All reactions were run in flame-dried glassware under an Ar atmosphere unless otherwise noted. Commercially available reagents were used without additional purification unless otherwise stated. Compound purification was effected by flash chromatography using 230 $\times$ 400 mesh, 60 Å porosity, silica obtained from Sorbent Technologies. ^1H NMR and ^{13}C NMR spectra were obtained on either a Bruker Avance DRX 500 spectrometer equipped with a broadband observe probe or a Bruker Avance AVIII 500 spectrometer equipped with a $^{13}\text{C}/^1\text{H}$ cryoprobe and referenced to residual protio solvent signals. Structural assignments are based on ^1H , ^{13}C , DEPT-135, COSY, HSQC and IR spectroscopy. All ee's were determined using chiral HPLC with columns purchased from Daicel (Chiracel AD, OD-H, or AS) using 99:1 to 85:15 mixtures of hexanes:isopropanol.

Synthesis of hydroxy ester, **5h**

Hydroxy acid **4b** was synthesized via an adaptation of a known method.¹³ **4b** (200mg, 0.82 mmol) and dry K_2CO_3 (566 mg, 4.1 mmol) were added to a flame-dried flask followed by the addition of allylbromide (296 mg, 2.4 mmol). Next, acetone (2 mL, distilled from

MgSO₄) was added and the mixture stirred vigorously for 5 h. The mixture was then extracted with ethyl acetate (20 mL), washed with water (2 × 5 mL), dried over MgSO₄ and concentrated *in vacuo*. Azeotropic removal of excess allyl bromide and allyl alcohol provided pure **5h** (210 mg, 0.74 mmol).¹⁴ ¹H NMR (500 MHz, CDCl₃) δ 7.48 – 7.41 (m, 4H, ArCH's), 5.90 – 5.78 (m, *J* = 16.4, 11.3, 5.7 Hz, 1H, CH₂CHCH₂), 5.24 (dd, *J* = 7.5, 1.3 Hz, 1H, CH₂CHCHH), 5.21 (t, *J* = 1.3 Hz, 1H, CH₂CHCHH), 4.70 – 4.57 (m, *J* = 13.2, 11.8, 5.7 Hz, 2H, ROCH₂R), 3.76 (s, 1H, ROH), 1.75 (s, 3H, quatCCH₃). ¹³C NMR (126 MHz, CDCl₃) δ 175.13 (RCO₂R), 141.92 (qArC), 131.61 (ArC's), 131.24 (RCHCH₂), 127.40 (ArC's), 122.21 (qArC), 119.34 (RCHCH₂), 75.62 (qC), 67.12 (ROCH₂R), 27.02 (RCH₃).

Synthesis of sulfone **8h**

To a flame dried flask with septum was added **5h** (205 mg, 0.72 mmol), methane sulfonyl chloride (0.27 mL, 3.45 mmol), DMAP (13 mg, 0.11 mmol) and pyridine (0.72 mL) and the reaction was placed in a chiller where the temperature was maintained at –5 to 0 °C for 18 h. After 12 h, additional methane sulfonyl chloride (0.20 mL) was added. After 6 additional hours, the reaction mixture was extracted with ethyl acetate (20 mL) and washed with an ice/H₂O/HCl mixture (1:1 ice/1M HCl), dried over MgSO₄ and concentrated *in vacuo* with minimum heat. The crude mesylate was then added to a Schlenk flask and chilled to 0 °C. To the mesylate was added a prechilled solution of sodium phenyl thiolate and phenyl thiol in ethanol (NaH, 248 mg, 6.5 mmol and PhSH 0.73 mL 7.2 mmol, in EtOH 7.2 mL) and allowed to slowly warm to room temperature and stirred for 12 h. The reaction mixture was extracted with CH₂Cl₂ (40 mL) and washed with K₂CO₃ solution (2 × 10 mL). The organic layer was dried over MgSO₄, concentrated *in vacuo* and purified via flash column chromatography (95:5 to 50:50 Hexanes:CH₂Cl₂ to afford **6h** (110 mg, 41%). **6h** was oxidized to the sulfone using *m*CPBA to afford sulfone **8h** (41 mg, 38%).¹⁵ ¹H NMR (500 MHz, CDCl₃) δ 7.55 (t, *J* = 7.4 Hz, 1H), 7.52 – 7.49 (m, 2H), 7.43 – 7.40 (m, *J* = 5.8 Hz, 2H), 7.38 – 7.33 (m, 2H), 7.23 – 7.19 (m, 2H), 5.84 (ddt, *J* = 16.3, 10.5, 5.9 Hz, 1H), 5.29 (dq, *J* = 17.1, 1.4 Hz, 1H), 5.24 (ddd, *J* = 10.4, 2.3, 1.2 Hz, 1H), 4.67 (tt, *J* = 4.5, 1.3 Hz, 2H), 2.09 (s, 3H).

Synthesis of sulfone **9c**

The mesylate resulting from hydroxy ester, **9b**, was synthesized via an adaptation from the previously referenced work.⁷ The mesylate is then allowed to react in the same manner described for synthesis of sulfone **8h**, with the major exception of the procedure for isolation and purification of the mesylate prior to substitution (Silica chromatography 8:2 Hexanes:Ethyl acetate). The substitution reaction was carried out at room temperature, to afford sulfide ester in 96%. The oxidation was also carried out with *m*CPBA using standard conditions described for synthesis of sulfone **8h**. The enantioenrichment was determined to be 97% ee via chiral HPLC (Chiracel OD-H, 90:1 Hexanes/isopropanol, 1 mL/min, 210 nm, major t_r = 8.2 min., minor t_r = 15.2 min.). ¹H NMR (400 MHz, CDCl₃) δ 7.89 (d, *J* = 7.4 Hz, 2H, *o*-ArCH'sSO₂R), 7.68 (t, *J* = 7.5 Hz, 1H, *p*-ArCHSO₂R), 7.56 (t, *J* = 7.7 Hz, 2H, *m*-ArCH'sSO₂R), 7.24 – 7.18 (m, *J* = 7.1 Hz, 3H, ArCH'sCH₂R), 7.07 (d, *J* = 7.6 Hz, 2H, ArCH'sCH₂R), 4.09 (q, *J* = 7.1 Hz, 2H, ROCH₂R), 3.63 (d, *J* = 12.9 Hz, 1H, diastereotopic RCHHR), 3.05 (d, *J* = 13.0 Hz, 1H, RCHHR), 1.47 (s, 3H, qCCH₃), 1.15 (t, *J* = 7.1 Hz, 3H, RCH₂CH₃). ¹³C NMR (101 MHz, CDCl₃) δ 167.93 (RCO₂R), 136.00 (ArC), 134.48 (ArC), 134.42 (ArC), 130.83 (ArC's), 130.53 (ArC's), 128.96 (ArC's), 128.70 (ArC's), 127.59 (ArC), 73.97 (qC), 62.55 (ROCH₂R), 38.82 (PhCH₂R), 16.29 (qCCH₃), 14.02 (RCH₂CH₃).

Synthesis of sulfone **10c**

The precursor 1,2-diol was synthesized according to a literature prep.¹⁶ The diol was then oxidized to the hydroxy acid in the same manner as outlined in reference.¹³ The resulting hydroxy acid was esterified as outlined for **5h**. Mesylation, substitution and oxidation of the

resulting hydroxy ester followed the same procedure as that out-lined for sulfone **9c**. ¹H NMR (500 MHz, CDCl₃) δ 7.84 (dd, *J* = 8.3, 1.1 Hz, 2H, *o*-ArCH'sSO₂R), 7.66 (t, *J* = 7.5 Hz, 1H, *p*-ArCHSO₂R), 7.53 (t, *J* = 7.9 Hz, 2H, *m*-ArCH'sSO₂R), 6.80 – 6.67 (m, 4H, ROArCH'sOR), 5.79 (ddd, *J* = 16.3, 11.0, 5.8 Hz, 1H, RCHCH₂), 5.29 (dd, *J* = 17.2, 1.4 Hz, 1H, RCHCHH), 5.20 (d, *J* = 10.5 Hz, 1H, RCHCHH), 4.61 – 4.50 (m, 2H, ROCH₂CHCH₂), 4.07 – 3.91 (m, 2H, ArOCH₂R), 3.72 (s, 3H, ArOCH₃), 2.79 (ddd, *J* = 14.5, 8.8, 6.1 Hz, 1H, diastereotopic ArOCH₂CH₂qC), 2.30 (dt, *J* = 14.0, 4.9 Hz, 1H, diastereotopic ArOCH₂CH₂qC), 1.67 (s, 3H, qCCH₃). ¹³C NMR (126 MHz, CDCl₃) δ 167.91 (RCO₂R), 154.34 (ROqArCOR), 152.56 (RO-qArCOR), 135.70 (ArC), 134.42 (ArC), 131.24 (RCHCH₂), 130.81 (ArCHs), 128.95 (ArCH's), 119.35 (RCHCH₂), 115.58 (ROqArCH'sOR), 114.84 (ROqArCH'sOR), 71.81 (qCCH₃), 66.97 (ROCH₂CHR), 64.34 (ROCH₂CH₂R), 55.89 (ROCH₃), 32.58 (qCCH₂CH₂R), 16.33 (qCCH₃).

Acknowledgments

We thank the National Institute of General Medical Sciences (1R01GM079644) for support of this work.

References

1. Weaver JD, Tunge JA. *Org Lett* 2008;10:4657. [PubMed: 18785744]
2. Ratni H, Blum-Kaelin D, Dehmlow H, Hartman P, Jablonski P, Masciadri R, Maugeais C, Patiny-Adam A, Panday N, Wright M. *Bioorg Med Chem Lett* 2009;19:1654. [PubMed: 19231176]
3. Ohata K, Terashima S. *Tetrahedron Lett* 2006;47:2787.
4. (a) Cram DJ, Wingrove AS. *J Am Chem Soc* 1963;85:1100. (b) Corey EJ, Koenig H, Lowry TH. *Tetrahedron Lett* 1962:515.
5. Ikegai K, Pluemanupat W, Mukaiyama T. *Bull Chem Soc Jpn* 2006;79:780.
6. Corey EJ, Lowry TH. *Tetrahedron Lett* 1965:793.
7. Green JE, Bender DM, Jackson S, O'Donnell MJ, McCarthy JR. *Org Lett* 2009;11:807. [PubMed: 19159255]
8. (a) Li H, Wang B, Deng L. *J Am Chem Soc* 2006;128:732. [PubMed: 16417358] (b) Smith ND, Goodman M. *Org Lett* 2003;5:1035. [PubMed: 12659567] (c) Avenoza A, Busto JH, Jimenez-Oses G, Peregrina JM. *Org Lett* 2006;8:2855. [PubMed: 16774274]
9. Ikegai K, Pluemanupat W, Mukaiyama T. *Chem Lett* 2005;34:638.
10. (a) Blundell P, Ganguly AK, Girijavallabhan VM. *Synlett* 1994:263. (b) Fristrup P, Dideriksen BB, Tanner D, Norrby PO. *J Am Chem Soc* 2005;127:13672. [PubMed: 16190733] (c) Griesbach RC, Hamon DPG, Kennedy RJ. *Tetrahedron: Asymmetry* 1997;8:507. (d) Ishibashi H, Maeki M, Yagi J, Ohba M, Kanai T. *Tetrahedron* 1999;55:6075. (e) Lee D, Lee H, Kim S, Yeom CE, Kim BM. *Adv Synth Catal* 2006;348:1021. (f) Li A, Yue G, Li Y, Pan X, Yang TK. *Tetrahedron: Asymmetry* 2003;14:75. (g) Aladro FJ, Guerra FM, Moreno-Dorado FJ, Bustamante JM, Jorge ZD, Massanet GM. *Tetrahedron Lett* 2000;41:3209.
11. Benanni YL, Vanhessche KPM, Sharpless KB. *Tetrahedron: Asymmetry* 1994;5:1473.
12. Kolb HC, VanNieuwenhze MS, Sharpless KB. *Chem Rev* 1994;94:2483.
13. Ainge D, Ennis D, Gidlund M, Stefinovic M, Vaz LM. *Org Process Res Dev* 2003;7:198.
14. Prisinzano T, Hsin LW, Folk JE, Flippen-Anderson JL, George C, Jacobson AE, Rice KC. *Tetrahedron: Asymmetry* 2003;14:3285.
15. Brown DJ, Ford PW. *J Chem Soc Perkin 1* 1969;19:2720. [PubMed: 5391236]
16. Corey EJ, Guzman-Perez A, Noe MC. *J Am Chem Soc* 1995;117:10805.

Scheme 1.

Scheme 2.

Scheme 3.

Scheme 4.

Scheme 5.

Scheme 6.

a) **MsCl**, **DMAP**, **pyr**, -5 – 0°C ; b) **NaSPh**, **EtOH**, 0°C – rt ; (c) **mCPBA**, **DCM**, rt .

Scheme 7.

a) AD-mix- α *t*BuOH/H₂O, 0 °C, 61%; b) TFAA c) H₂, Pd/C EtOH, rt, 99% over 2 steps, >94% ee; d) MsCl, DMAP, pyr, rt, 78%; e) NaSPh, EtOH, 96%; f) mCPBA, DCM, rt, 99%; g) ee determined by chiral HPLC

Scheme 8.

a) AD-mix- α , *t*BuOH/H₂O, 0 °C, 90%; b) Pt/C, NaHCO₃ Air, H₂O, 70°C, 93%; c) K₂CO₃, Allyl bromide, Acetone rt, 99%; d) MsCl, DMAP, pyr rt, 81%; e) NaSPh, EtOH, 96%; f) mCPBA, DCM rt, 84%, 94% ee; g) ee determined by chiral HPLC PMP = *p*-methoxyphenyl.

Table 1

Ar	AD-mix	product	yield 3 (%)	product	yield 4 (%)
<i>p</i> -MeO ₂ CC ₆ H ₄	α	(<i>S</i>)-3a	79	(<i>S</i>)-4a	<10
<i>p</i> -BrC ₆ H ₄	β	(<i>R</i>)-3b	78	(<i>R</i>)-4b	81
<i>p</i> -ClC ₆ H ₄	α	(<i>S</i>)-3c	99	(<i>S</i>)-4c	85
<i>p</i> -MeC ₆ H ₄	α	(<i>S</i>)-3d	99	(<i>S</i>)-4d	64
<i>p</i> -FC ₆ H ₄	β	(<i>R</i>)-3e	98	(<i>R</i>)-4e	82

Table 2

Conditions for esterification of hydroxy acids

Entry	Acid	R	Allyl bromide		Yield	Product
			4b-e	5f		
1	4e	F			93%	5f
2	4d	Me			92%	5g
3	4b	Br			90%	5h
4	4b	Br			97%	5i
5	4c	Cl			99%	5j
6	4e	F			99%	5k

Allyl bromide	
Entry	Acid	R	Yield	Product	
7	4c	Cl	99%		5l
8	4d	Me	97%		5m

Table 3

entry	substrate	yield 6 (%)	ratio 6:7 ^a	yield 8 (%)	% ee 8 ^d
1	5f	29 %	2.5:1	77 %	46 %
2	5g	10 %	1:1	--	--
3	5h	41 %	19:1	38 %	97 %
4	5i	39 % ^c	19:1	48 %	80 % ^c
5	5j	23 %	ND ^b	38 %	89 %
6	5k	38 %	8:1	59 %	73 %
7	5l	50 %	5:1	72 %	61 %
8	5m	15 %	1:1	--	--

^aDetermined by ¹H NMR.^bRatio not determined before oxidation.^cMesylation inadvertently performed at rt.^dee determined by chiral HPLC.