Published in final edited form as:

Org Lett. 2010 June 4; 12(11): 2450–2452. doi:10.1021/ol1001686.

Efficient Diacetoxylation of Alkenes *via* Pd(II)/Pd(IV) Process with Peracetic Acid and Acetic Anhydride

Chan Pil Park, Joo Ho Lee, Kyung Soo Yoo, and Kyung Woon Jung*

Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California, Los Angeles, California 90089-1661

Abstract

A palladium-catalyzed diacetoxylation of alkenes in the presence of peracetic acid and acetic anhydride was developed to produce diacetates efficiently and diastereoselectively. Due to its mild conditions, this method was suitable for a broad range of substrates encompassing conjugated and non-conjugated olefins.

The Pd-catalyzed dioxygenation of alkenes is an important difunctionalization method in organic synthesis due to its efficiency, safety, and economy,¹ although osmium catalysts have been widely used for dioxygenation reactions.² Sigman proposed a Pd(II) catalyzed aerobic dialkoxylation of styrene derivatives requiring an *o*-phenol to facilitate the formation of a quinone methide intermediate.^{1b} On the other hand, we envisioned an alternative Pd(II)/Pd(IV) mechanism for dioxygenation, and embarked on the development of efficient processes for alkene diacetoxylation *via* peracetic acid as an oxidant. During the course of our research, Dong^{3a} and Jiang^{3b} reported olefin diacetoxylation catalyzed by palladium complexes in the presence of either PhI(OAc)₂ or molecular oxygen as an oxidant (eqs 1 and 2).

(1)

kwjung@usc.edu.

Supporting Information Available Experimental details for the preparation of compounds and spectral data of these compounds. This material is available free of charge via the internet at <http://pubs.acs.org>

(2)

These olefin dioxygenations would presumably occur by a distinct Pd(II) to Pd(IV) process, which is significantly broad in its application toward various alkenes.⁴ However, the use of $\text{PhI}(\text{OAc})_2$ as a stoichiometric oxidant produces a large amount of by-product while the use of molecular oxygen as an oxidant requires high pressure, temperature, and long reaction time. Therefore, our newly developed Pd(II)/Pd(IV) processes using peracetic acid can provide practical improvements over the previously reported reaction conditions.

Our initial studies included the identification of appropriate conditions and the understanding of the possible mechanism for the oxygenation using a palladium catalyst. As representatively shown in Scheme 1, diacetoxylation was facile in the presence of a combination of peracetic acid and acetic anhydride with palladium acetate, converting styrene (**1**) to diacetate **2** at room temperature efficiently. To validate the possibility of Pd(II)/Pd(IV) process, we compared similar conditions without peracetic acid or acetic anhydride. When the reaction was carried out with only acetic acid, styrene underwent acetoxylation and β -hydride elimination to furnish the vinyl acetate **3** presumably *via* Pd(0)/Pd(II) process. However, when the reaction was performed in the presence of peracetic acid as the oxidant at an ambient temperature, hydroxyacetoxylated products (**4**) were produced, implying a Pd(II)/Pd(IV) process instead of Pd(0)/Pd(II). In addition, when acetic anhydride was added to the mixture of **4** under similar conditions, conversion of **4** to **2** was not observed by ^1H NMR spectra. Therefore, diacetoxylation would require both peracetic acid and acetic anhydride, and further investigation was carried out.

As shown in Table 1, we examined the correlation between product selectivity and the ratio of peracetic acid and acetic anhydride. In the presence of only peracetic acid as an oxidant (entries 1 and 2), dioxygenation of styrene proceeded to give hydroxyacetoxylation product (**4**) preferably along with small amounts of diacetate **2**. Upon the addition of acetic anhydride, the chemoselectivity for the diacetoxylation product (**2**) improved significantly (entries 3 ~ 6), and its yield was proportional to the amount of acetic anhydride. Optimal results were obtained when styrene was added to the premixed solution of AcO_2H and Ac_2O at 0°C and the reaction mixture was slowly warmed to room temperature (see Supporting Information).

Based upon these results, it can be suggested that $\text{Pd}(\text{OAc})_2$ could be oxidized to $\text{Pd}(\text{IV})(\text{OH})(\text{OAc})_3$ without acetic anhydride or $\text{Pd}(\text{IV})(\text{OAc})_4$ with acetic anhydride. Keeping in mind this possibility, we propose a mechanism for the diacetoxylation of an alkene (Figure 1). The Pd(IV) species **II** (palladium tetraacetate), generated from $\text{Pd}(\text{OAc})_2$ and peracetic acid/acetic anhydride, is suggested to be the starting point of alkene acetoxylaion.⁵ The highly reactive Pd(IV) species would then undergo *syn*-addition onto the alkene to generate palladium intermediate **IV**, while maintaining the Pd(IV) oxidation state. This is followed quickly by reductive elimination to afford the diacetoxylation product from Pd(IV) center and regenerate $\text{Pd}(\text{OAc})_2$. The most notable feature of the proposed catalysis is its ability to facilitate diacetoxylation under mild conditions such as ambient temperature and a short reaction time.

Next, we screened various olefins using the optimal conditions. First, we examined 3-butenic acid as a representative non-conjugated olefin to afford diacetate product **6**, which was anticipated to cyclize to the valuable lactone **7**.⁶ In the absence of palladium catalysts, only a

trace amount of lactone compound was observed and most of unreacted 3-butenic acid was recovered (entry 1). When PdCl₂ effected the oxidation to a small degree (entry 2), Pd (OAc)₂ facilitated the oxidation and cyclization to offer lactone **7** in a good yield (entry 3). Other oxidants such as PhI(OAc)₂ were also tested, but none of them afforded significant efficiency (entry 4).⁷ By varying solvents, we sought optimal conditions. Wet acetic acid (entry 3) and organic solvents (entry 5) were not comparable to the use of anhydrous acetic acid (entry 6), which converted 3-butenic acid completely to the desired lactone **7** in an excellent yield.

Using the developed reaction conditions, we extended the diacetoxylation protocol to a number of olefins to verify the generality of this method. As shown in Table 3, the desired reaction products were synthesized generally in high yields. Terminal aliphatic olefins including 3-butenitrile, 3-methyl-3-butenyl acetate, and allylbenzene afforded their corresponding diacetoxylation products **8**, **9** and **10**, respectively, in 93%, 90%, and 86% (entries 1 – 3). In the case of the trisubstituted cyclic alkene, 1-methyl-1-cyclohexene underwent diacetoxylation smoothly to give the desired diacetate product **11** in 89% yield along with high diastereoselectivity (*syn/anti* = 15/1) (entry 4). Disubstituted aromatic alkenes also provided desired vicinal oxygenation products (entries 5 ~ 7), and high *syn*-addition of acetoxy group was confirmed again (entries 6 and 7) (*syn/anti* = 10/1, 12/1, respectively).^{3a} Furthermore, the reaction with 1-phenyl-1-cyclohexene as a trisubstituted conjugated olefin took place efficiently to provide the desired compound (**15**) in 87% yield along with high diastereoselectivity (*syn/anti* = 15/1) (entry 8).

In conclusion, we have successfully developed a highly reactive palladium-catalyzed diacetoxylation for a broad range of alkenes, including conjugated and non-conjugated, under ambient temperature and pressure. Mono- to trisubstituted alkenes were compatible with the developed method. Additionally, high stereoselectivity as well as functional group tolerance is worthy of note. Overall, our processes are believed to provide a convenient and practical method.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We acknowledge generous financial supports from the National Institute of Health (RO1 GM 71495 & S10 RR025432).

References

1. (a) Chevrin C, Le Bras J, Henin F, Muzart J. *Synthesis* 2005;2615–28618. (b) Schultz MJ, Sigman MS. *J. Am. Chem. Soc* 2006;128:1460–1463. [PubMed: 16448111] (c) Thiery E, Chevrin C, Le Bras J, Harakat D, Muzart J. *J. Org. Chem* 2007;72:1859–1862. [PubMed: 17274660] (d) Zhang Y, Sigman MS. *J. Am. Chem. Soc* 2007;129:3076–3077. [PubMed: 17298071] (e) Jensen KH, Sigman MS. *Org. Biomol. Chem* 2008;6:4083–4088. [PubMed: 18972034]
2. (a) Finn MG, Sharpless KB. *J. Am. Chem. Soc* 1991;113:113–126. (b) Crispino GA, Ho PT, Sharpless KB. *Science* 1993;259:64–66. [PubMed: 8418495] (c) Kolb HC, VanNieuwenhze MS, Sharpless KB. *Chem. Rev* 1994;94:2483–2547. (d) Kobayashi S, Endo M, Nagayama S. *J. Am. Chem. Soc* 1999;121:11229–11230. (e) Donohoe TJ, Blades K, Winter JJG, Stemp G. *Tetrahedron Lett* 2000;41:4701–4704. (f) Donohoe TJ. *Synlett* 2002;8:1223–1232. (g) Donohoe TJ, Harris RM, Butterworth S, Burrows JN, Cowley A, Parker JS. *J. Org. Chem* 2006;71:4481–4489. [PubMed: 16749777] (h) Christie SDR, Warrington AD. *Synthesis* 2008:1325–1341.
3. (a) Li Y, Song D, Dong VM. *J. Am. Chem. Soc* 2008;130:2962–2964. [PubMed: 18281992] (b) Wang A, Jiang H, Chen H. *J. Am. Chem. Soc* 2009;131:3846–3847. [PubMed: 19292485]

4. (a) Dick AR, Hull KL, Sanford MS. *J. Am. Chem. Soc* 2004;126:2300–2301. [PubMed: 14982422]
(b) Desai LV, Sanford MS. *Angew. Chem. Int. Ed* 2007;46:5737–5740.
5. (a) Nagata R, Saito I. *Synlett* 1990:291–301. (b) Chen MJY, Kochi JK. *J. Chem. Soc., Chem. Commun* 1977:204–205. (c) Muzart J. *J. Mol. Cat. A: Chemical* 2007;276:62–72.
6. (a) Ahmed MM, Berry BP, Hunters TJ, Tomcik DJ, O'Doherty GA. *Org. Lett* 2005;7:745–748. [PubMed: 15704940] (b) Wu Y, Guan M. *Ind. J. Chem., Sec. B: Org. Chem. Med. Chem* 1998;37B: 844–846. (c) Brzezinski LJ, Rafel S, Leahy JW. *J. Am. Chem. Soc* 1997;119:4317–4318. (d) Tiecco M, Testaferri L, Tingoli M. *Tetrahedron* 1993;49:5351–5358. (e) Uchiyama H, Kobayashi Y, Sato F. *Chem. Lett* 1985:467–470.
7. In Song's report(reference 3(a)), use of excess $\text{PhI}(\text{OAc})_2$ improved the yield and reduced β -hydride elimination

Figure 1.
Plausible Pd(II)/Pd(IV) pathway.

via Pd(0)/Pd(II)

via Pd(II)/Pd(IV)

Scheme 1.

Table 1

Diacetoxylation of Styrene with $\text{Pd}(\text{OAc})_2^a$

entry	time (h)	AcO_2H^b (mmol)	Ac_2O (mmol)	conversion (%)	
				1	2
1	4	0.13	0	41	53
2	4	0.26	0	12	78
3	4	0.26	1.3	4	79
4	4	0.26	2.6	0	62
5	4	0.26	5.2	0	21
6	4	0.26	7.8	0	6

^a All reactions were carried out with styrene (0.26 mmol) and $\text{Pd}(\text{OAc})_2$ (0.026 mmol) in acetic acid (500 μL).

^b 35.5 wt % in AcOH.

^c The conversions were determined by ^1H NMR in the presence of an internal standard.

Table 2

The Pd-Catalyzed Diacetoxylolation of 3-Butenoic acid.^a

entry	catalyst	oxidant	solvent ^b	time (h)	yield (%) ^c
1	none	AcO ₂ H	AcOH	12	trace
2	PdCl ₂	AcO ₂ H	AcOH	8	10
3	Pd(OAc) ₂	AcO ₂ H	AcOH	4	72
4	Pd(OAc) ₂	PhI(OAc) ₂	AcOH	12	trace
5 ^d	Pd(OAc) ₂	AcO ₂ H	AcOH/MeCN	4	62
6 ^e	Pd(OAc) ₂	AcO ₂ H	AcOH	4	92

^a 1.0 mmol scale (0.1 M solution), 5 mol % Pd catalyst, and 1.2 equiv oxidant, yield by ¹H NMR data spectra using an internal standard.

^b In entries 1–5, H₂O was evaporated to afford the cyclized product after reaction.

^c Isolated yields.

^d AcOH:MeCN = 1:1 (wt %).

^e Anhydrous acetic acid.

Table 3

Pd-Catalyzed Diacetoxylation of Alkenes^a

<div><div><div>$\text{R}-\text{CH}=\text{CH}-\text{R}_1 \xrightarrow[\text{anhydrous AcOH, 60 min, rt}]{\text{Pd(OAc)}_2 (2 \text{ mol}), \text{AcOOH, Ac}_2\text{O}}$</div></div></div>	
entry	substrate
1	
2	
3	
4

entry

substrate

5

entry

substrate

^{6d}

entry

substrate

7

entry

substrate

8

^a 1.0 mmol scale (0.1 M solution), 2 mol % Pd(OAc)₂, and 1.2 equiv CH₃CO₃H.

^b Isolated yields.

^c The diastereomeric ratios were determined by ¹H NMR analysis.

^d Reaction temperature was 45 °C.