Published in final edited form as:

J Agric Food Chem. 2009 March 11; 57(5): 1821–1826. doi:10.1021/jf8033732.

Herbicidal Activity of Glucosinolate Degradation Products in Fermented Meadowfoam (*Limnanthes alba*) Seed Meal

JAN F. STEVENS^{†,*}, RALPH L. REED[†], SUSAN ALBER[‡], LARRY PRITCHETT[§], and STEPHEN MACHADO[§]

[†]Department of Pharmaceutical Sciences, Oregon State University, Corvallis, Oregon 97331

[‡]Department of Statistics, Oregon State University, Corvallis, Oregon 97331

[§]Columbia Basin Agricultural Research Center, Oregon State University, Corvallis, Oregon 97331

Abstract

Meadowfoam (*Limnanthes alba*) is an oilseed crop grown in western Oregon. After extraction of the oil from the seeds, the remaining seed meal contains 2-4% of the glucosinolate, glucolimnanthin. We investigated the effect of fermentation of seed meal on its chemical composition and the effect of the altered composition on downy brome (*Bromus tectorum*) coleoptile emergence. Incubation of enzyme-inactive seed meal with enzyme-active seeds (1% by weight) resulted in complete degradation of glucolimnanthin and formation of 3-methoxybenzyl isothiocyanate in 28% yield. Fermentation in the presence of an aqueous solution of FeSO₄ (10 mM) resulted in the formation of 3-methoxyphenylacetone nitrile and 2-(3-methoxyphenyl)ethanethioamide, a novel natural product. The formation of the isothiocyanate, the nitrile and the thioamide, as a total, correlated with an increase of herbicidal potency of seed meal ($r^2 = 0.96$). The results of this study open new possibilities for the refinement of glucosinolate-containing seed meals for use as bioherbicides.

Keywords

meadowfoam; seed meal; *Limnanthes alba*; glucosinolate; glucolimnanthin; herbicide; isothiocyanate; nitrile; thioamide

INTRODUCTION

The continuing growth of organic farming practices calls for increased use of pesticides derived from naturally occurring materials. Many plants produce toxic phytochemicals when attacked by herbivores (phytoalexins) or deposit pro-toxic glycosides in leaves that prevent seed germination of other, neighboring plants when the aglycones are released from the shed leaves (allelochemicals). Glucosinolates form a group of allelochemicals produced by many species of the Brassicales, including *Brassica* and *Sinapis* spp. (mustards), *Lepidium* and *Nasturtium* spp. (cresses), and *Limnanthes* spp. (meadowfoams) (1). The interest in *Brassica* glucosinolates is primarily due to their presence in oilseeds which are retained in the marc after oil extraction (meal). Glucosinolate-containing seed meals are intensively investigated as biofumigants for weed control (2,3).

White meadowfoam (*Limnanthes alba* Hartw. ex Benth., Limnathaceae) is native to southern Oregon and northern California (4,5). Several cultivars have emerged from a meadowfoam

*To whom correspondence should be addressed. Phone: 541-737-9534. Fax: 541-737-3999. E-mail: fred.stevens@oregonstate.edu.

breeding program at Oregon State University (6). The species is cultivated in the Willamette valley of western Oregon for the seed oil which is rich in unusual 20:1 and 22:1 fatty acids (7). The oil has commercial value as an ingredient of skin care products. The spent seed material (meal) can be used as a bioherbicide due to the presence of allelochemicals. The meal contains the glucosinolate, glucolimnanthin **1** (Figure 1)(8), and 3-methoxyphenylacetone nitrile (also referred to as 3-methoxybenzyl cyanide, **2**), a known allelochemical (9) formed by heat-induced degradation of **1** during the oil extraction process. Because water is added at an early stage of the industrial oil extraction process to facilitate seed crushing, conversion of **1** into 3-methoxybenzyl isothiocyanate (**3**), catalyzed by myrosinase in the seed, would occur if the enzyme were not inactivated by application of heat. Early heat treatment is necessary to inactivate myrosinase and prevent contamination of the oil with non-polar breakdown products of **1** while late-stage heat treatment follows extraction of seed oil with an organic solvent with the purpose to remove residual solvent.

When seed meal is applied to soil for weed control, a disadvantage of the lack of myrosinase activity in seed meal is that breakdown of **1** depends on the soil micro-environment which may lead to variable results regarding degradation kinetics, profile of degradation products, and weed suppression. In this study, the conversion of **1** into the allelochemicals **2**, **3** (9) and 2-(3-methoxyphenyl)ethanethioamide (**4**) in enzyme-inactive meal was investigated by making use of active enzymes present in meadowfoam seeds and a co-factor, FeSO₄. We demonstrate that enzyme-treated meal products contain greater amounts of **2**, **3** or **4**, and show greater inhibitory activity in a soil-based seed germination assay, compared to untreated meadowfoam seed meal. The results of this study open new possibilities for the refinement of glucosinolate-containing seed meals for use as bioherbicides in (organic) farming and horticulture.

MATERIALS AND METHODS

General

NMR experiments were performed on a Bruker DPX400 instrument. High-resolution ESI-ToF MS measurements were performed on a Waters Micromass LCT Classic ToF instrument (Waters, Milford, MA) equipped with an electrospray ion source, interfaced to a Waters CapLC HPLC solvent delivery system.

Chemicals

HPLC water was produced from reversed-osmosis water by a Milli-Q water purification system. HPLC-grade acetonitrile and methanol were purchased from EMD Chemicals, San Diego, CA. 3-Methoxyphenyl acetone nitrile (**2**) was purchased from Sigma Aldrich, St. Louis, MO. 3-Methoxybenzyl isothiocyanate (**3**) was obtained from Oakwood Products, West Columbia, SC. 2-(3-Methoxyphenyl)ethanethioamide (**4**) was purchased from Endeavour Specialty Chemicals, Daventry, Northamptonshire, U.K., and 2-(3-methoxyphenyl) acetamide (**5**) was from Maybridge Trevillet, Tintagel, Cornwall, U.K.

HPLC

The HPLC equipment consisted of a Waters Delta 600 solvent delivery system, a Waters 717 plus Autosampler, a Waters 2996 photodiode array detector, a Waters 600 controller and a data acquisition/processing computer with Empower™ software (Waters, Milford, MA). In HPLC system 1, separations were achieved on a reverse-phase Lichrosphere 5 µm C18 column (4 × 250 mm, Phenomenex, Torrance, CA). The HPLC solvents were 0.1% aqueous trifluoroacetic acid (solvent A) and MeCN (solvent B). A linear solvent gradient was employed starting from 5% solvent B in solvent A to 100% B over 30 min at a flow rate of 1.0 mL/min. After returning to the starting conditions in 1 min, the column was equilibrated for 10 min before the next injection. The injection volume was 10 µL. On-line UV spectra were recorded in the range

210–500 nm and the λ 274 nm trace was used for calculation of peak areas for all compounds, except for the thioamide **4** which was recorded at λ 270 nm. Analyte concentrations were determined from calibration curves constructed for each analyte.

In HPLC system 2, used for monitoring fractions from a Sephadex LH-20 column, the HPLC column was an Agilent Zorbax 5 μ m SB-C18 column (2.1 $\times$ 50 mm). The column was eluted with solvent A for 2 min, then the proportion of solvent B was increased to 100% over 5 min, held at 100% for 0.5 min and decreased to 0% solvent B (100% solvent A) in 0.5 min. The column was equilibrated at 100% solvent A for 2.4 min before the next injection. The flow rate was 0.3 mL/min and the injection volume was 1.0 μ L.

Mass spectrometry

The LC-MS/MS instrument consisted of a 4000 QTrap hybrid linear ion trap–triple quadrupole mass spectrometer equipped with a pneumatically-assisted electrospray (Turbo V) source operated at 550 $^{\circ}$ C (Applied Biosystems/MDS Sciex, Concord, Ontario, Canada). Liquid nitrogen was the source for heating and nebulizing gas, curtain gas, and collision gas. The spray needle was kept at +5.5 kV (positive ion mode). Product ion (MS/MS) spectra were recorded by scanning in the range 50–200 a.m.u. at a cycle time of 0.6 s and at unit resolution for both Q1 and Q3. The collision energy was set at 27 eV and the collision gas was set at ‘medium’. Samples in MeCN-H₂O (1:1, v/v) were introduced into the mass spectrometer by loop injections (1 μ L) in a flow of MeCN-H₂O-HCOOH (50:50:0.1 by volume) at 0.2 mL/min. An HPLC system consisting of two Shimadzu Prominence LC-20AD pumps and a Shimadzu SIL-HTC autoinjector (Shimadzu Scientific Instruments, Columbia, MD) provided the solvent flow and performed the injections.

Isolation of glucolimnanthin (1) from meadowfoam seed meal

Factory-grade meal (250 g; Natural Plant Products Inc., Salem, Oregon) was soaked in 500 mL MeOH-H₂O (1:1, v/v) for 18 hrs and the slurry transferred to a 2 L percolator fitted at the bottom with a cotton plug and cotton cloth. MeOH-H₂O (1:1, v/v; 2.4 L) was passed through the column of seed meal at a flow rate of 0.2 L/hr. Three fractions (0–0.8 L, 0.8–1.6 L and 1.6–2.4 L) were collected and analyzed by HPLC using system 1. Fractions 1 and 2 were combined and taken to dryness by rotary evaporation (careful: foaming) and lyophilization. The residue was dissolved in 50 mL of water and diluted with 300 mL of MeOH. The resulting precipitate (carbohydrates and proteins) was removed by centrifugation, and the supernatant was concentrated *in vacuo*. The residue was taken up in 60 mL of MeOH and four 15 mL portions were fractionated by column chromatography on Sephadex LH-20 using MeOH as the eluting solvent at a flow rate of 1.6 mL/min. Fractions (10 mL) were collected, monitored by HPLC (system 2) for the presence of glucolimnanthin (**1**), and the fractions containing **1** were combined and taken to dryness by rotary evaporation. Crude **1** obtained from two column runs (2.6 g) was redissolved in 6 mL of MeOH and purified on the same Sephadex LH-20 column using the same chromatographic conditions, yielding 1.8 g of >95% pure **1** by NMR analysis. ¹H NMR (400 MHz, MeOH-*d*₄): δ H 7.26 (1H, *t*, *J* = 8 Hz, H-5'), 7.01 (1H, *s*, H-2'), 7.00 (1H, *d*, *J* = 8 Hz, H-6'), 6.83 (1H, *d*, *J* = 8 Hz, H-4'), 4.55 (1H, *d*, *J* = 9 Hz, H-1''), 4.25 (1H, *d*, *J* = 16 Hz, H-2), 4.05 (1H, *d*, *J* = 16 Hz, H-2), 3.87 (1H, *d*, *J* = 12 Hz, H-6''), 3.81 (3H, *s*, CH₃), 3.63 (1H, *dd*, *J* = 5, 12 Hz, H-6''), 3.37–3.25 (2H, *m*, H-3'' and H-4''), 3.21–3.12 (2H, *m*, H-2'' and H-5''). ¹³C NMR (100 MHz, MeOH-*d*₄): δ C 160.2 (C-3'), 159.5 (C-1), 137.6 (C-1'), 129.5 (C-5'), 120.2 (C-6'), 113.1 (C-2'), 112.6 (C-4'), 81.5 (C-1''), 80.9 (C-2''), 78.0 (C-3''), 72.8 (C-5''), 69.8 (C-4''), 61.4 (C-6''), 54.3 (CH₃), 38.3 (C-2). Assignment of these resonances was confirmed by ¹H-¹H COSY, ¹H-¹³C HSQC and HMBC experiments. Compound **1** was obtained as the K⁺-salt as determined by ESI-ToF MS: *m/z* 478.0269, calculated for C₁₅H₂₁NO₁₀S₂K: 478.0244 (5.2 ppm).

Isolation of compound 4

Two grams of meadowfoam seed meal was incubated overnight at room temperature with ground meadowfoam seeds (1% by wt) and 6 mL of an aqueous solution of FeSO₄ (10 mM). The incubation mixture was mixed with 6 mL of MeOH, sonicated for one minute, and allowed to stand overnight. The mixture was centrifuged, and the resulting supernatant was chromatographed by HPLC on a 10 × 250 mm Phenomenex Ultrex 5 μm C18 column using a solvent gradient from 5 to 95 % MeCN in H₂O containing 0.1 % HCOOH over 30 minutes at a flow rate of 5 mL/min. The peak eluting at 17 minutes (UV_{max} 270 nm) was collected manually and analyzed by electrospray mass spectrometry.

Synthesis of 3-methoxybenzyl thiocyanate (6)

The thiocyanate **6** was synthesized from 3-methoxybenzyl chloride and potassium thiocyanate using a general procedure for the conversion of alkyl halides into alkyl thiocyanates (10). Briefly, 3 g (17.2 mmol) of 1-*n*-butyl-3-methylimidazolium chloride [bmim]Cl (TCI America, Portland, OR) was dissolved in acetone (25 mL). After addition of KSCN (34.4 mmol), the mixture was stirred for 48 hrs at room temp. The resulting suspension was filtered and the filtrate was concentrated *in vacuo*. The residue was dissolved in CH₂Cl₂ and filtered again. After evaporation of the solvent, [bmim]SCN was obtained as a reddish liquid. A mixture of 395 mg of [bmim]SCN (2 mmol) and 157 mg (1 mmol) of 3-methoxybenzyl chloride (TCI America, Portland, OR) was stirred for 3 hrs at room temp and then kept overnight without stirring. The mixture was extracted with Et₂O (3 × 3 mL) by vigorous shaking. The combined Et₂O layers were dried with anhydrous Na₂SO₄ and concentrated *in vacuo*, yielding 3-methoxybenzyl thiocyanate (**6**) as a colorless oil (158 mg, 88% yield). ¹H NMR (400 MHz, CDCl₃): δ_H 7.28-7.35 (1H, *m*, H-5), 6.91-6.99 (3H, *m*, H-2, H-4 and H-6), 4.16 (2H, *s*, -CH₂-), 3.85 (3H, *s*, OCH₃). ¹³C NMR (100 MHz, CDCl₃): δ_C 160.0 (C-3), 135.8 (C-1), 130.2 (C-5), 121.2 (C-6), 114.54 (C-2), 114.46 (C-4), 112.0 (-C≡N), 55.3 (-OCH₃), 38.4 (-CH₂-). HR-EIMS (70 eV): C₉H₉NOS⁺ observed *m/z* 179.0409 (44% rel. int.), calculated 179.0405 (2.2 ppm), *m/z* 121 (100% rel. int., CH₃O-Ph-CH₂⁺).

Preparation of fermented meal products (treated seed meals, T1-T5)

Fermented meal was prepared by grinding 9.9 g meal together with 0.1 g of untreated meadowfoam seed (*Limnanthes alba* ssp. *alba* Benth., cultivar Ross) in a coffee grinder (model E160B, Proctor Silex, Washington, NC) for one minute. Ground batches were pooled and mixed with de-ionized water (3 mL/g meal), sonicated for five minutes, allowed to incubate for 18 hours at room temperature, freeze-dried, and re-ground for 30 seconds (**T1**). The freeze-drying step was considered necessary in order to be able to perform the germination assay. Iron-augmented meal was produced by the same procedure except 10 mM FeSO₄ was substituted for de-ionized water (**T2**). Control incubations, all without seeds, included unaltered meal (ground but not incubated with water and not freeze-dried, **T5**), sham-augmented meal (meal plus water alone, **T4**), and iron only (meal plus 10 mM FeSO₄, **T3**).

For analysis by HPLC-UV, 1.0 g aliquots of the fermented and freeze-dried meal products were mixed with 6 mL of 50 % methanol in screw-capped glass centrifuge tubes, vortexed for 30 seconds, and sonicated for 60 seconds. The glass tube contents were allowed to stand overnight in the dark at room temperature, vortexed and sonicated again, and centrifuged for 5 minutes on a clinical centrifuge. Supernatants were further centrifuged for ten minutes at 13,000 rpm using a micro-centrifuge, diluted 1:9 with 50 % methanol, and then analyzed directly using HPLC system 1. Samples were prepared in triplicate.

Assay for herbicidal activity

About 45 g of Walla Walla silt loam soil (coarse, silty, mixed, mesic Typic Haploxeroll, pH 6.1) was weighed into 10-cm diameter Petri dishes. The soil was taken from a grassland site to which no agrochemicals had been applied for 15 years, cleaned of straw and roots, and sieved through a 2.362 mm screen (#8). For germination testing of individual compounds (**1-5**), 15.0 mL of test solution was added to each dish. Test solutions of glucolimanthin **1** were made up in water and added to the dishes to give doses of 0 (control, $n = 12$), 0.09 ($n = 6$), 0.18 ($n = 12$), 0.28 ($n = 6$), 0.36 ($n = 12$), 0.47 ($n = 6$), 0.53 ($n = 6$), 0.71 ($n = 6$) and 0.89 ($n = 6$) mg per gram soil. Test solutions of compounds **2**, **3** and **5** were made up in ethanol and added to the dishes to give doses of 0 (control), 0.06, 0.11, 0.22, 0.44, and 0.89 mg per gram soil. Test solutions of compound **4** were made up in ethanol and added to the dishes to give doses of 0 (control), 0.18, 0.36, 0.53, 0.71 and 0.89 mg per gram soil. The ethanol added to soil was allowed to evaporate overnight in a hood, and then 15 mL of water was added to the dishes. Freeze-dried meal products were mixed with soil, followed by the addition of 15 mL of water. Fifteen seeds of *Bromus tectorum* were placed in concentric circles within each dish. For each dose, the number of replicates was 6 or 12 for compound **1** and 4 for compounds **2**, **3** and **5**, 6 for compound **4**, and $n = 4$ for **T1-T5**. Petri dishes with lids were placed in an incubator at 20 °C during day time (8 hrs) and at 15 °C at night time (16 hrs) for 7 days. Coleoptile emergence was recorded at the end of the incubation period. Seeds with emerged coleoptiles were considered to be germinated.

Statistical analysis

For compounds **1-5** and treated seed meals **T1-T5**, dose response relationships were estimated using logistic regression (linear in dose) allowing (via quasilielihood) for extrabinomial variation among the replicates at each dose (SAS 9.2 Genmod procedure, dscale option). For the compounds, the zero dose is the vehicle control (water for compound **1** and ethanol for compounds **2-5**). Therefore, compounds **2-5** have the same intercept in the model. For the treatments **T1-T5**, the zero dose is the same for all five groups. Therefore, in the model all 5 meal treatments have the same intercept. For compounds $i = 1, \dots, 5$, the logistic regression equation is $\ln[p_{ij}/(1-p_{ij})] = [\ln(\text{OR})]_i x_{ij} + b_i$, where $\ln$ is the natural logarithm, OR is the odds ratio, x_{ij} is the dose and p_{ij} is the probability of germination for dose j of compound i , and $[\ln(\text{OR})]_i$ is the slope and b_i is the intercept for compound i . The OR for compound i is the odds of germination at dose $x_{ij} = k + 1$ divided by the odds of germination at dose $x_{ij} = k$, where the odds of germination is $p_{ij}/(1-p_{ij})$. Since compounds **2-5** have a common intercept, $b_2 = b_3 = b_4 = b_5$. The same logistic regression equation was used for the seed meals, except that all seed meals have the same intercept. From the equation, the ED_{50} is $-b_i/[\ln(\text{OR})]_i$. Because the inverse of the ED_{50} is often used as an estimator of potency (11) and because the $\text{ED}_{50} = -b/\ln(\text{OR})$ and b is the same for all seed meals, we expressed the herbicidal potency of the treated seed meals as the negative logarithm of the odds ratio, $-\ln(\text{OR})$.

RESULTS AND DISCUSSION

Composition of meadowfoam seed meal

Analysis of meadowfoam seed meal by HPLC-UV shows glucolimanthin **1** (75 $\mu\text{mol/g}$) and its degradation product **2** (23 $\mu\text{mol/g}$) as the main constituents containing the 3-methoxybenzyl moiety ($\text{UV}_{\text{max}} 274 \text{ nm}$, Figure 2A and Table 1). The presence of substantial amounts of nitrile **2** in the meal is attributed to heat-induced degradation of glucolimanthin **1** during the industrial oil extraction process, because untreated meadowfoam seeds contain primarily **1** and only very small amounts of **2** (data not shown). This finding is consistent with continuous thermal formation of benzyl cyanide from benzylglucosinolate in seeds of the garden cress (*Lepidium sativum*) after heat-inactivation of myrosinase (12). The retention of **2** in meadowfoam seed meal after hexane extraction and **2** having sufficient solubility in hexane

suggest that the thermal conversion of **1** took place after oil extraction when the meal undergoes steaming to remove residual extraction solvent. The acetamide **5** was detected at trace levels in the meal, indicating that hydrolysis of **2** during meal steaming is negligible.

Degradation of glucolimnanthin in seed meal mediated by enzyme-active seeds

Fermentation of seed meal by inoculation with ground seeds resulted in complete degradation of glucosinolate **1** and formation of isothiocyanate **3** in 28% yield, while the content of nitrile **2** remained unchanged during fermentation (Table 1 and Figure 2B). This is in agreement with earlier work by Vaughn and co-workers (2) who reported 25% recovery of isothiocyanate **3** from **1** in a mixture of meadowfoam seed meal and soil. The moderate yield of the isothiocyanate **3** can be explained by a combination of incomplete extraction and covalent binding of the isothiocyanate moiety to proteins and amino acids in the seed meal and ground seeds. It is also possible that isothiocyanate **3** releases SCN^- which is known to be released from its analog, 4-hydroxybenzyl isothiocyanate (13), and has phytotoxic properties (14). However, 3-methoxybenzyl isothiocyanate (**3**) is expected to be more stable than 4-hydroxybenzyl isothiocyanate, because **3** cannot dissociate into quinone methide and SCN^- , the presumed mechanism for spontaneous degradation of 4-hydroxybenzyl isothiocyanate (13).

Degradation of glucolimnanthin in seed meal mediated by enzyme-active seeds in the presence of Fe^{2+}

Incubation of seed meal with ground seeds and 10 mM FeSO_4 resulted in complete degradation of glucosinolate **1**, formation of isothiocyanate in 13% yield and a doubling of the nitrile **2** content (Table 1 and Figure 2C). Incubation of seed meal with 10 mM FeSO_4 showed a 26% decrease of glucosinolate **1** which was not accounted for by an increase of nitrile **2** or isothiocyanate **3**. The origin of the apparent loss of **1** is unknown.

The enhanced formation of nitrile **2** in **T2** (active enzyme added) compared to **T3** (enzymes inactive) suggests a catalytic role for Fe^{2+} in conjunction with myrosinase or other enzymes to form nitrile **2**. Fe^{2+} has been identified by other researchers as a cofactor of nitrile-forming proteins that alter myrosinase-mediated hydrolysis of glucosinolates but lack hydrolytic activity by themselves (15,16). One form of nitrile-forming protein, epispecifier protein (ESP) from *Arabidopsis thaliana*, was shown by Burow *et al.* (15) to act in conjunction with myrosinase in the conversion of glucosinolates into the corresponding nitriles. They observed a three-fold increase in ESP activity when Fe^{2+} was added at 0.01 mM or 0.5 mM. Similar results were obtained by Matusheski *et al.* (17), who furthermore observed that the myrosinase-mediated conversion of glucoraphanin into sulphoraphane nitrile required higher concentrations of iron (0.1-1 mM Fe^{2+}) when ESP was absent. We observed very little conversion of glucosinolate **1** into nitrile **2** at FeSO_4 concentrations at 1 mM or less, which suggests that meadowfoam seeds have low nitrile-forming protein activity. On the other hand, meadowfoam seeds may contain normal levels of nitrile-forming protein compared to other species of Brassicales, but that the added iron is not sufficiently available due to binding to proteins or other cellular macromolecules in the seeds or seed meal. In support of ESP-independent Fe^{2+} catalysis, Bellostas *et al.* (18) recently hypothesized that Fe^{2+} coordinates with the sulphur atom of the thioglucoside moiety, thereby preventing myrosinase-directed Loessen rearrangement and so directing glucosinolate degradation to the nitrile at the expense of isothiocyanate formation.

In the seed meal incubations with myrosinase-active seeds in the presence of 10 mM FeSO_4 , we detected the formation of compound **4**, not present or present at very low levels in seed meal or in seed meal without addition of iron and seeds (Figure 2C). When the experiments were repeated with the addition of pure glucolimnanthin **1** (60 mg per gram meal), a four-fold

increase in the formation of compound **4** was observed compared to incubations without addition of **1**, indicating that **4** is another degradation product of **1**. Compound **4** was collected upon elution from a semi-preparative HPLC column. The peak fraction revealed a molecular ion with m/z 282 by ESI-MS and a fragment ion with m/z 121 (MeO-Ph-CH_2^+) (Figure 3). This suggested that compound **4** was identical with 2-(3-methoxyphenyl)ethanethioamide, which was confirmed by HPLC-UV and MS comparison of **4** with an authentic standard (Figure 3). Although Fe^{2+} -mediated hydrolysis and reduction of glucosinolates into the corresponding thioamides has previously been recognized by other researchers (18, 19), thioamide **4** has not previously been reported as a natural product nor has its herbicidal properties been examined. Our results indicate the involvement of an enzyme in the Fe^{2+} -mediated formation of thioamide **4** because **4** was not formed in the absence of enzyme-active seeds when iron was added (Table 1). Regarding the mechanism of thioamide formation, Bellostas *et al.* (18) proposed that Fe^{2+} coordinates with the imine nitrogen, a sulfate oxygen and a hydroxyl group at C-2 in the side chain of the aglycone. The formation of thioamide **4** in our study suggests that there is no requirement for a C-2 hydroxyl to form a favorable glucosinolate- Fe^{2+} transition complex, because glucolimnanthin does not have such a hydroxyl group. An alternative explanation for the role of iron in the enzyme-mediated formation is that iron prevents Loessen rearrangement, possibly by coordination with the thioglucoside sulphur (18), resulting in an intermediate hydrolysis product, *N*-hydroxy-2-(3-methoxyphenyl)thioacetamide, that is subsequently reduced by two equivalents of Fe^{2+} to form the thioamide **4**.

Thiocyanate (**6**) formation

Thiocyanates are often reported as degradation products of glucosinolates (1,2,20) and therefore we investigated the presence of thiocyanate **6** in the fermented seed meals. Synthetic 3-methoxybenzyl thiocyanate (**6**) showed a retention time of 22.2 min in HPLC system 1 and a UV maximum at 280.5 nm, thus eluting between nitrile **2** and isothiocyanate **3**. Thiocyanate **6** was not detected in any of the (fermented) seed meals (Figure 2).

Herbicidal activity of glucolimnanthin (**1**) and degradation products (**2-5**)

For compounds **1-5**, the estimated log odds ratios are given in Table 2. The predictions from the logistic regression model were back-transformed onto the percent germination scale to yield dose-response curves are shown in Figure 4. As can be seen in Figure 4 which show the predicted dose-response curve together with the observed values, the regression model gives a good fit to the data. The absolute value of the log odds ratio, and therefore also the herbicidal potency of the compounds, decreases in the order: thioamide **4** > nitrile **2** > acetamide **5** > glucolimnanthin **1** > isothiocyanate **3**. There was no significant difference between **1** and **3** or between **1** and **5** (both $P > 0.2$). All other the pairs of compounds were significantly different (all $P < 0.001$). The fact that **3** and **5** are significantly different, but **1** and **5** are not, is largely due to the smaller standard error of the estimated difference between **3** and **5** as compared to **1** and **5**.

Herbicidal activity of fermented meadowfoam seed meal products

For seed meals **T1-T5**, the estimated log odds ratios are given in Table 2 and the predicted dose-response curves are shown in Figure 5. As can be seen in Figure 5 the regression model gives a good fit to the data. Seed meals **T3**, **T4**, and **T5** were not significantly different from each other ($P = 0.43$). Treatment of seed meal with enzyme active seeds significantly increased the herbicidal potency compared to the water treatment ($P < 0.001$) and compared to untreated meal ($P < 0.001$). Addition of 10 mM FeSO_4 to the incubation mixture of seed meal with enzyme active seeds further enhanced the herbicidal potency as compared to **T1** ($P = 0.012$). This treatment (**T2**) was also significantly different from **T3** ($P < 0.0001$).

To understand the increases in meal potency as a result of glucosinolate degradation, we examined the relationships between the content of the glucolimnanthin breakdown products and herbicidal potency. When the $-\ln(\text{OR})$ was plotted separately against the content of the compounds **1**, **2**, **3**, and **4** in the seed meals, **T1-T5** (Table 1), only **1** showed a regression curve with slope that was significantly different from zero ($r^2 = 0.94$, $P = 0.0068$) (curve not shown). The negative sign of the slope indicated that the decrease of **1** in the meals correlated with an increase of meal potency. When the $-\ln(\text{OR})$ was plotted against the total content of the compounds **2**, **3**, and **4** in the seed meals, a regression curve was obtained with a positive slope that was significantly different from zero ($r^2 = 0.96$, $P = 0.0033$) (Figure 6). These analyses demonstrate that the formation of **2**, **3**, and **4** from **1** strongly correlates with an increase in herbicidal potency of the treated seed meals. Diverting the degradation pathway away from **3** toward the formation of **2** and **4** results in further increases of meal potency, consistent with the potency order: thioamide **4** > nitrile **2** > isothiocyanate **3** (Figure 4 and Table 2).

We have explored a simple and inexpensive approach to convert the glucosinolate **1** into more potent herbicidal degradation products. The fermented seed meal products have greater herbicidal potency than factory-grade seed meal, whose herbicidal activity largely depends on its constitutive content of nitrile **2** and soil factors for *in situ* degradation of **1**. *Brassica* manure is a well-documented source of myrosinase in the soil (21), but seed meals lack active myrosinase. *Aspergillus* fungi from soil have been reported to possess myrosinase activity (22,23). However, the soil microflora is variable and may not completely convert glucosinolate, resulting in under-utilization of the seed meal's herbicidal potential. Therefore, fermented glucosinolate-containing seed meals may offer more effective bioherbicides with more predictable activity for use in (organic) farming.

Acknowledgments

This study was supported by the U.S. Department of Agriculture (grant # USDA/CREES 2005-34407-15670), the National Institutes of Health (grants # S10RR022589 and P30ES000210) and Natural Plant Products, Inc., Salem, Oregon.

References

1. Fahey JW, Zalcmann AT, Talalay P. The chemical diversity and distribution of glucosinolates and isothiocyanates among plants. *Phytochemistry* 2001;56:5–51. [PubMed: 11198818]
2. Vaughn SF, Palmquist DE, Duval SM, Berhow MA. Herbicidal activity of glucosinolate-containing seedmeals. *Weed Science* 2006;54:743–748.
3. Brown PD, Morra MJ. Glucosinolate-containing plant tissues as bioherbicides. *J Agric Food Chem* 1995;43:3070–3074.
4. Mason, CT. A systematic study of the genus *Limnanthes*. University of California; Berkeley: 1952. p. 455-512.
5. Jain, SK. Domestication of *Limnanthes* (Meadowfoam) as a new oil crop; Plant domestication induced mutation: Proceedings of an advisory group meeting on the possible use of mutation breeding for rapid domestication of new crop plants; Vienna, Austria. 1986; p. 121-134.
6. Knapp, SJ.; Crane, JM. Breeding advances and germplasm resources in meadowfoam: a very long chain oilseed. In: Janick, J., editor. Perspectives on new crops and new uses. ASHS Press; Alexandria, VA: 1999. p. 225-233.
7. Miller RW, Daxenbichler ME, Earle FR. Search for new industrial oils, VIII. The genus *Limnanthes*. *J Am Oil Chem Soc* 1964;41:167–196.
8. Ettlinger MG, Lundeen AJ. The mustard oil of *Limnanthes douglasii* seed, m-methoxybenzylisothiocyanate. *J Am Chem Soc* 1956;78:1952–1954.
9. Vaughn SF, Boydston RA, Mallory-Smith CA. Isolation and identification of (3-methoxyphenyl) acetonitrile as a phytotoxin from meadowfoam (*Limnanthes alba*) seedmeal. *Journal of Chemical Ecology* 1996;22:1939–1949.

10. Kamal A, Chouhan G. A task-specific ionic liquid [bmim]SCN for the conversion of alkyl halides to alkyl thiocyanates at room temperature. *Tetrahedron Letters* 2005;46:1489–1491.
11. Piegorsch, WW.; Bailer, AJ. *Statistics for Environmental Biology and Toxicology: Interdisciplinary Statistics*. Chapman and Hall; London: 1997. p. 579
12. Hasapis X, MacLeod AJ. Benzylglucosinolate degradation in heat-treated *Lepidium sativum* seeds and detection of a thiocyanate-forming factor. *Phytochemistry* 1982;21:1009–1013.
13. Borek V, Morra MJ. Ionic thiocyanate (SCN[−]) production from 4-hydroxybenzyl glucosinolate contained in *Sinapis alba* seed meal. *J Agric Food Chem* 2005;53:8650–4. [PubMed: 16248567]
14. Hansson D, Morra MJ, Borek V, Snyder AJ, Johnson-Maynard JL, Thill DC. Ionic thiocyanate (SCN[−]) production, fate, and phytotoxicity in soil amended with Brassicaceae seed meals. *J Agric Food Chem* 2008;56:3912–7. [PubMed: 18442242]
15. Burow M, Markert J, Gershenzon J, Wittstock U. Comparative biochemical characterization of nitrile-forming proteins from plants and insects that alter myrosinase-catalysed hydrolysis of glucosinolates. *FEBS J* 2006;273:2432–46. [PubMed: 16704417]
16. Lambrix V, Reichelt M, Mitchell-Olds T, Kliebenstein DJ, Gershenzon J. The Arabidopsis epithiospecifier protein promotes the hydrolysis of glucosinolates to nitriles and influences *Trichoplusia ni* herbivory. *Plant Cell* 2001;13:2793–807. [PubMed: 11752388]
17. Matusheski NV, Swarup R, Juvik JA, Mithen R, Bennett M, Jeffery EH. Epithiospecifier protein from broccoli (*Brassica oleracea* L. ssp. *italica*) inhibits formation of the anticancer agent sulforaphane. *J Agric Food Chem* 2006;54:2069–76. [PubMed: 16536577]
18. Bellostas N, Sorensen AD, Sorensen JC, Sorensen H. Fe²⁺-catalyzed formation of nitriles and thionamides from intact glucosinolates. *J Nat Prod* 2008;71:76–80. [PubMed: 18163583]
19. Austin FL, Gent CA, Wolff IA. *J Agric Food Chem* 1968;16:752–755.
20. Heany, RK.; Fenwick, GR. *Methods for Glucosinolate Analysis*. In: Waterman, PG., editor. *Methods in Plant Biochemistry*. Academic Press; London: 1993. Alkaloids and Sulphur Compounds. 8:531–550.
21. Borek V, Morra MJ, M JP. Myrosinase activity in soil extracts. *Soil Science Society of America Journal* 1996;60:1792–1797.
22. Ohtsuru M, Hata T. Fungus myrosinase. V. General characteristics of the intracellular myrosinase from *Aspergillus niger*. *Agricultural and Biological Chemistry* 1973;37:2543–2548.
23. Butrindr B, Niamsup H, Shank L, Rakariyatham N. Myrosinase overproducing mutants of *Aspergillus* sp. NR463. *Annals of Microbiology* 2004;54:493–501.

Figure 1. Enzymatic degradation of the glucosinolate, glucolimnanthin (1), forming the nitrile 2, the isothiocyanate 3, and the thioamide 4. The nitrile 2 may hydrolyze to form the acetamide 4.

Figure 2. HPLC analysis of untreated meal (panel A), meal treated with 1% myrosinase-active meadowfoam seeds (panel B), and meal incubated with a 10 mM solution of FeSO_4 in the presence of 1% myrosinase-active meadowfoam seeds (panel C). The UV trace was recorded at 274 nm.

Figure 3. Identification of thioamide **4** in seed meal by electrospray MS/MS (panel A) and HPLC-UV (panel B).

Figure 4.

Herbicidal activity of compounds 1-5. Symbols represent averages of measurements: up-pointing triangles ($n = 12$ at dose = zero and $n = 6$ or 12 for other doses), compound 1; filled circles ($n = 4$), compound 2; filled squares ($n = 4$), compound 3; down-pointing triangles ($n = 6$), compound 4; diamonds ($n = 4$), compound 5.

Figure 5. Herbicidal activity of seed meal treatments **T1-T5**. Symbols represent averages of measurements ($n = 4$): circles, **T1**; down-pointing triangles, **T2**; filled squares, **T3**; up-pointing triangles, **T4**; diamonds, **T5**.

Figure 6.
Correlation between herbicidal potency expressed as $-\ln(OR)$ and total content of compounds 2-4 in seed meals T1-T5.

Table 1

Composition of treated and untreated meadowfoam seed meal

	Average $\mu\text{mol/g}$ meal $\pm$ SD (n=3)				
	Glucosinolate 1	Nitrile 2	Isothiocyanate 3	Thioamide 4	Total of 1-4
Meal + 1% seed (T1)	n.d. ^a	22.1 $\pm$ 0.3	20.8 $\pm$ 0.1	n.d.	42.9
Meal + 1% seed + FeSO ₄ (T2)	n.d.	49.4 $\pm$ 0.8	10.0 $\pm$ 0.2	0.82 $\pm$ 0.01	60.2
Meal + FeSO ₄ (T3)	55.8 $\pm$ 0.1	25.9 $\pm$ 0.1	0.76 $\pm$ 0.02	n.d.	82.5
Meal + water (T4)	71.8 $\pm$ 0.3	18.3 $\pm$ 0.3	0.19 $\pm$ 0.01	n.d.	90.3
Untreated meal (T5)	75.2 $\pm$ 0.8	23.2 $\pm$ 0.3	n.d.	n.d.	98.4

^a n.d., not detected.

Table 2Estimation of herbicidal potency for test compounds and treated meals^a

Compounds	$-\ln(\text{OR})^b$ (std error)	$-\ln(\text{OR})$ Difference ($P < 0.05$) ^c
Glucolimnanthin 1	2.86 (0.602)	a, d
Nitrile 2	6.26 (0.592)	b
Isothiocyanate 3	2.15 (0.429)	a
Thioamide 4	10.47 (0.703)	c
Acetamide 5	3.77 (0.406)	d
Treated meals		
Meal + 1% seed (T1)	0.152 (0.0139)	f
Meal + 1% seed + FeSO ₄ (T2)	0.1913 (0.0156)	g
Meal + FeSO ₄ (T3)	0.0438 (0.0148)	h
Meal + water (T4)	0.0223 (0.0162)	h
Untreated meal (T5)	0.0412 (0.0149)	h

^a Calculations were performed on mg/g soil basis.^b OR, odds ratio (see under 'Statistical analyses'); $-\ln(\text{OR})$ is the slope of the regression curve, a measure of herbicidal potency.^c Within compound and seed meal groups, shared letters indicate lack of difference at the 95% confidence level.