Published in final edited form as:

Org Lett. 2010 June 4; 12(11): 2517–2519. doi:10.1021/ol100772u.

Copper-Catalyzed Cyanation of Heterocycle Carbon-Hydrogen Bonds

Do Hien-Quang and Olafs Daugulis *

Department of Chemistry, University of Houston, Houston, TX 77204-5003

Abstract

A method for regioselective cyanation of heterocycles has been developed. A number of aromatic heterocycles as well as azulene can be cyanated in reasonable to good yields by using a copper cyanide catalyst and an iodine oxidant.

The nitrile functional group is found in many pharmaceuticals and agrochemicals.^{1,2} Sandmeyer and Rosenmund-von Braun reactions are classic methods for the synthesis of aromatic nitriles (scheme 1A).³ However, both procedures employ a stoichiometric copper(I) cyanide reagent and prefunctionalized starting materials. Cyanation of aryl halides by employing catalytic copper or palladium is also well-known.⁴ In contrast, direct cyanation through C-H bond functionalization is more attractive due to use of readily available reactants. Direct cyanation can be achieved by employing an appropriate oxidant in combination with a cyanide source (scheme 1B).⁶ Methods for direct cyanation of pyridines,^{6b} thiophenes,^{6c,e} substituted indoles,^{6c,e,h} pyrroles,^{6c,e} and 2-phenylpyridines^{6d,f,g} have been reported in the literature. However, direct transitionmetal catalyzed cyanation of acidic heterocycles such as azoles, imidazoles, and triazoles has not been described yet.^{6j,k} We report here a method for regioselective, direct cyanation of aromatic heterocycles as well as azulene by employing copper catalysis and an iodine oxidant.

Several key issues need to be addressed for achieving a general cyanation procedure. First, high regioselectivity is required for the procedure to be synthetically relevant. High regioselectivity observed in direct arylation⁷ and halogenation reactions⁸ (scheme 2A) is imparted by the initial deprotonation step. This protocol rules out the use of positive cyanide sources due to their incompatibility with strong bases. Consequently, simple cyanide salts such as NaCN or KCN must be employed. Second, choice of base is important. As described earlier,^{7,8} *t*BuOLi allows for functionalization of a wide range of substrates possessing DMSO *pK_a*'s below 35–37. Third, the oxidant has to be compatible with all other reagents. It should not react with the organolithium intermediate producing byproducts that can not be converted to the

*olafs@uh.edu.

Supporting Information Available: Experimental details, data and spectra for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

desired cyanation product. Oxygen reactivity with some arylmetals complicates the catalytic system by requiring the use of zinc or magnesium amide bases.⁹ Iodine is compatible with copper catalysts and aryl iodides formed in the reaction of ArLi with I₂ can be converted to aryl cyanides.^{4,8} Consequently, we decided to employ iodine as the oxidant in conjunction with *t*BuOLi base and a copper catalyst (scheme 2B). Benzothiazole was chosen as model substrate and copper(I) cyanide/phenanthroline catalyst was used to screen for suitable reaction conditions. DMF solvent was chosen as a starting point for optimization because it has been successfully used in base-mediated iodination.⁸ Several cyanating reagents such as NaCN, KCN, Zn(CN)₂, and K₃[Fe(CN)₆] were screened. However, none of these reactions afforded any cyanation product. Solvent screening revealed that 1,4-dioxane yields 42% conversion to the desired product if NaCN is employed as the cyanating reagent (Table 1). The reaction mixture was monitored to gain insights into the reaction pathway. The results indicate that the cyanation is a stepwise reaction. Benzothiazole is iodinated to afford 2-iodobenzothiazole followed by copper-catalyzed cyanation yielding the desired product **2**. Consequently, reaction conditions should be optimized with respect to both iodination and cyanation. Among tested cyanide sources, sodium cyanide provides highest reactivity due to its solubility in dioxane. However, it has been well established that high concentration of cyanide will deactivate copper catalyst due to strong complexation.⁴ Careful tuning of the solvent polarity led to significant improvement of conversion to the product (Table 1).

Ligand optimization is shown in Table 2. The cyanation did not proceed in absence of phenanthroline. In addition, considerable amount of benzothiazole dimer byproduct was observed if less than 20 mol % of phenanthroline was used. Bipyridine ligand was shown to be almost as efficient as phenanthroline (entry 4). Copper(I) chloride and copper(I) iodide can also be employed as catalysts. The optimized conditions involve CuCN catalyst, phenanthroline ligand, sodium cyanide, iodine oxidant, and mixed dioxane/*m*-xylene solvent. These conditions were applied to cyanation of a variety of heterocycles (Table 3).

Heterocycles such as benzoxazole, benzothiazole, benzimidazole, caffeine, and triazoles are cyanated in reasonable to good yields (entries 1-7). Electron-deficient 2-phenylpyridine oxide is also cyanated in a good yield (entry 8). Pyridine derivatives with electron-withdrawing (fluorine) or electron-donating (methoxy) substituents are reactive under the general conditions affording products in acceptable yields (entries 9-10). In both cases, the cyano group was introduced at 2-position. Similar selectivity has been reported by Katritzky in the direct cyanation of pyridines by addition/elimination approach.^{6b}

Indoles constitute an important motif of aromatic compounds encountered in many natural products and pharmaceuticals. Hence, selective functionalization of indoles has attracted much attention in recent years.¹⁰ Wang and coworkers have recently developed a method for palladium-catalyzed cyanation of 1- and 2-substituted indoles by employing 3 equiv of Cu(OAc)₂ reoxidant.^{6h}

Minor modifications of our general conditions enabled regioselective cyanation of *N*-methylindole and azulene. The following changes in procedure were required for achieving high yield of the product. First, pyridine additive prevents the solidification of reaction mixture. Second, sodium phosphate base was found to be more efficient than *t*BuOLi in the iodination step. Third, copper catalyst was added to the reaction mixture after the iodination step was complete.

Consequently, the combination of iodine and sodium phosphate in dioxane provided effective iodination conditions that are compatible with the subsequent copper-catalyzed cyanation. Copper catalyst was added after iodination step in entries 11 and 12 to prevent the oxidation of copper(I) catalyst.^{7d} This procedure selectively afforded 3-cyano-1-methylindole in

excellent yield (entry 11). Azulene was cyanated in moderate yield (36%) due to low reactivity of iodoazulene intermediate in the cyanation step.

In conclusion, we have developed a method for copper-catalyzed cyanation of aromatic heterocycles and azulene. The use of mild iodine oxidant allows for regioselective sequential iodination/cyanation of heterocycles with acidic C-H bonds, methyindole, and azulene.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the Welch Foundation (Grant No. E-1571), the NIGMS (Grant No. R01GM077635), A. P. Sloan Foundation, Norman Hackerman Advanced Research Program, and Camille and Henry Dreyfus Foundation for supporting this research.

References

- (1) (a). Murdoch D, Keam SJ. *Drugs* 2005;65:2379. [PubMed: 16266205] (b) Njar VCO, Brodie AMH. *Drugs* 1999;58:233. [PubMed: 10473018]
- (2). Ozoe Y, Ishikawa S, Tomiyama S, Ozoe F, Kozaki T, Scott JG. *ACS Symp. Ser* 2007;948:39.
- (3)(a). Sandmeyer T. *Ber. Dtsch. Chem. Ges* 1884;17:1633. (b) Galli C. *Chem. Rev* 1988;88:765. (c) Rosenmund KW, Struck E. *Chem. Ber* 1919;52:1749. (d) von Braun J, Manz G. *Liebigs Ann. Chem* 1931;488:111. (e) Connor JA, Leeming SW, Price RJ. *Chem. Soc., Perkin Trans. 1* 1990:1127. (f) Ellis GP, Romney-Alexander TM. *Chem. Rev* 1987;87:779.
- (4) (a). Zanon J, Klapars A, Buchwald SL. *J. Am. Chem. Soc* 2003;125:2890. [PubMed: 12617652] (b) Cristau H-J, Ouali A, Spindler J-F, Taillefer M. *Chem. Eur. J* 2005;11:2483. (c) Schareina T, Zapf A, Magerlein W, Muller N, Beller M. *Chem. Eur. J* 2007;13:6249. (d) Sundermeier M, Zapf A, Beller M. *Eur. J. Inorg. Chem* 2003:3513. (e) Sundermeier M, Zapf A, Beller M. *Angew. Chem., Int. Ed* 2003;42:1661.
- (5). Selected reviews: (a) Lewis JC, Bergman RG, Ellman JA. *Acc. Chem. Res* 2008;41:1013. [PubMed: 18616300] (b) Campeau L-C, Fagnou K. *Chem. Soc. Rev* 2007;36:1058. [PubMed: 17576474] (c) Ackermann L. *Synlett* 2007:507. (d) Chen X, Engle KM, Wang D-H, Yu J-Q. *Angew. Chem., Int. Ed* 2009;48:5094. (e) Seregin IV, Gevorgyan V. *Chem. Soc. Rev* 2007;36:1173. [PubMed: 17576484] (f) Lyons TW, Sanford MS. *Chem. Rev* 2010;110:1147. [PubMed: 20078038] (g) Daugulis O, Do H-Q, Shabashov D. *Acc. Chem. Res* 2009;42:1074. [PubMed: 19552413] (h) Alberico D, Scott ME, Lautens M. *Chem. Rev* 2007;107:174. [PubMed: 17212475]
- (6) (a). So Y-H, Miller LL. *J. Am. Chem. Soc* 1980;102:7119. (b) Katritzky AR, Scriven EFV, Majumder S, Tu H, Vakulenko AV, Akhmedov NG, Murugan R. *Synthesis* 2005:993. (c) Dohi T, Morimoto K, Kiyono Y, Tohma H, Kita Y. *Org. Lett* 2005;7:537. [PubMed: 15704888] (d) Chen X, Hao X-S, Goodhue CE. *J. Am. Chem. Soc* 2006;128:6790. [PubMed: 16719450] (e) Dohi T, Morimoto K, Takenaga N, Goto A, Maruyama A, Kiyono Y, Tohma H, Kita Y. *J. Org. Chem* 2007;72:109. [PubMed: 17194088] (f) Jia X, Yang D, Zhang S, Cheng J. *Org. Lett* 2009;11:4716. [PubMed: 19764735] (g) Jia X, Yang D, Wang W, Luo F, Cheng J. *J. Org. Chem* 2009;74:9470. [PubMed: 19938811] (h) Yan G, Kuang C, Zhang Y, Wang J. *Org. Lett* 2010;12:1052. [PubMed: 20141185] (i) Mariampillai B, Alliot J, Li M, Lautens M. *J. Am. Chem. Soc* 2007;129:15372. [PubMed: 18020336] 2-Cyanoimidazoles can be prepared by the reaction of the corresponding imidazole with cyanogen bromide: (j) Whitten JP, McCarthy JR, Matthews DP. *Synthesis* 1988:470. Electrochemical oxidation of caffeine in the presence of cyanide has been shown to afford 8-cyanocaffeine: (k) Sono M, Toyoda N, Shizuri Y, Tori M. *Tetrahedron Lett* 1994;35:9237.
- (7) (a). Yotphan S, Bergman RG, Ellman JA. *Org. Lett* 2009;11:1511. [PubMed: 19260648] (b) Do H-Q, Daugulis O. *J. Am. Chem. Soc* 2008;130:1128. [PubMed: 18181627] (c) Do H-Q, Khan RMK, Daugulis O. *J. Am. Chem. Soc* 2008;130:15185. [PubMed: 18855471] (d) Do H-Q, Daugulis O. *Chem. Commun* 2009:6433.
- (8). Popov I, Do H-Q, Daugulis O. *J. Org. Chem* 2009;74:8309. [PubMed: 19827765]

- (9) (a). King AE, Brunold TC, Stahl SS. J. Am. Chem Soc 2009;131:5044. [PubMed: 19309072] (b) Do H-Q, Daugulis O. J. Am. Chem. Soc 2009;131:17052. [PubMed: 19899771] (c) Lam PYS, Bonne D, Vincent G, Clark CG, Combs AP. Tetrahedron Lett 2003;44:1691. (d) Demir AS, Reis ö. Emruallahoglu M. J. Org. Chem 2003;68:10130. [PubMed: 14682710]
- (10). For recent reviews, see (a) Humphrey GR, Kuethe JT. Chem. Rev 2006;106:2875. [PubMed: 16836303] (b) Cacchi S, Fabrizi G. Chem. Rev 2005;105:2873. [PubMed: 16011327] (c) Bandini M, Eichholzer A. Angew. Chem., Int. Ed 2009;48:9608. (d) Joucla L, Djakovitch L. Adv. Synth. Catal 2009;351:673.

A. Sandmeyer and Rosenmund - von Braun Reactions

$\text{X} = \text{N}_2^+$ Sandmeyer

$\text{X} = \text{Cl}, \text{Br}, \text{I}$ Rosenmund-von Braun

B. Direct Cyanation

Scheme 1.
Cyanation Methods

A. Deprotonation/Functionalization Approach

B. This work

Scheme 2.
Method Development

Table 1

Optimization of Cyanation Conditions

entry	solvent	MCN	2 ^a
1	DMF	NaCN, KCN, Zn(CN) ₂ , or K ₃ [Fe(CN) ₆]	0%
2	Dioxane	NaCN	42%
3	Dioxane	KCN, Zn(CN) ₂ , K ₃ [Fe(CN) ₆]	> 2%
4	Xylene	NaCN	0%
5	Dioxane/Xylene (9/1)	NaCN	54%
6	Dioxane/Xylene (4/1)	NaCN	72%
7	Dioxane/Xylene (7/3)	NaCN	90%
8	Dioxane/Xylene (3/2)	NaCN	78%
9	Dioxane/Xylene (1/1)	NaCN	59%

^aConversion by GC reported

Table 2

Optimization of Copper Catalyst

entry	catalyst/ligand	2 ^a	3 ^a
1	CuCN/no ligand	<2%	<2%
2	CuCN/Phenanthroline (1/1)	60%	16%
3	CuCN/Phenanthroline (1/2)	90%	< 2%
4	CuCN/Bipyridine (1/2)	88%	< 3%
5	CuCl/Phenanthroline (1/2)	87%	< 2%
6	CuI/Phenanthroline (1/2)	90%	< 2%

^aConversion by GC reported. Dioxane/m-xylene solvent (7/3).

Table 3

Cyanation Scope^a

entry	arene
1
2
3

entry

arene

entry

arene

5

entry

arene

6

entry

arene

entry

arene

8

entry

arene

9^b

10

entry

arene

11^c 12^c

^a Substrate (1 equiv), tBuOLi (2-2.13 equiv), iodine (1.35-1.5 equiv), NaCN (1.3 equiv), CuCN (0.1 equiv), and phenanthroline (0.2 equiv). Yields are isolated yields.

^b Cyanide as the limiting reagent.

^c Substrate (1.5-2 equiv), pyridine (0.76 equiv), Na₃PO₄ (2 equiv), iodine (1-1.35 equiv), and dioxane solvent. Copper(I) catalyst (0.2 equiv), phenanthroline (0.3 equiv), and sodium cyanide (1-1.3 equiv) were added after iodination step. See Supporting Information for details.