Published in final edited form as:

J Steroid Biochem Mol Biol. 2010 April ; 119(3-5): 135–140. doi:10.1016/j.jsbmb.2010.02.005.

Identification and Characterization of an Androgen-responsive Kap Promoter Enhancer Located in the Intron II Region of human Angiotensinogen Gene

Li-qiang Fan^{1,2,*}, Dianne O. Hardy^{2,*}, James F Catterall², Jian Zhao¹, and Su-xia Li¹

¹ The State Key Laboratory of Bioreactor Engineering, East China University of Science and Technology, Shanghai 200237, P. R. China

² Centre for Biomedical Research, Population Council, New York 10021 U.S.A.

Abstract

Transgenic expression of the human angiotensinogen (HAGT) gene directed by the mouse kidney androgen-regulated protein (*Kap*) gene promoter is proximal tubule cell-specific and androgen-regulated *in vivo*. The same *Kap* promoter fragment did not support similar regulation of other genes, but a transgene based on the original chimeric KAP-hAGT construct successfully directed NHE3 to kidney, suggesting that sequences within the HAGT gene fragment of the construct contributed to the regulation of its expression *in vivo*. In the present study, androgen-responsive regulatory sequences in the HAGT gene portions of the transgene were examined in transfected renal cells. A 1.4 kb enhancer between exons 2 and 3 was identified that increased the basal expression of *Kap* promoter 1.5 to 2-fold, its induction by dihydrotestosterone (DHT) 2 to 3-fold and its induction by dexamethasone (Dex) 4 to 5-fold. Sequence analysis revealed two potential hormone responsive elements. Mutational assays and electrophoretic mobility shift assay showed one of these elements was androgen-specific. These findings may influence future strategies for the design of inducible, cell-specific transgenes.

Keywords

androgen; glucocorticoid; enhancer; gene regulation; hAGT; *Kap* promoter

1 Introduction

Steroid hormone receptors (SR) are intracellular receptors that act upon binding of their ligand to regulate gene transcription. Upon activation by their cognate hormone, the SRs form homodimers that are able to bind to an imperfect palindromic DNA sequence, called the hormone response element (HRE). Investigation as to how SR regulates hormone-responsive promoters and enhancers, and mechanisms of steroid receptor specificity are ongoing in several experimental model systems [1-5]. HREs are found in various genes as hexameric palindromes

*Corresponding authors: Li-qiang Fan, Ph.D. PO Box 365, The State Key Laboratory of Bioreactor Engineering, East China University of Science and Technology, Shanghai, P. R China. 200237; Tel: 86-21-64252257; Fax: 86-21-64252257; fanglq@ecust.edu.cn; Dianne O Hardy, Ph.D. Centre for Biomedical Research, Population Council, 1230 York Avenue, New York, NY 10021 USA; Tel: 1-212-327-8754; Fax: 1-212-327-7678; Dianne@popcbr.rockefeller.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

or hexameric direct repeats (DR) with 3 to 5 bp spacers [6-9]. *In vitro*, when androgen receptor (AR) bound with random sequences preceding TGTCT, AR selected a partial overlapping direct repeat of three hexamers [9]. Less is known about the selectivity for AR at HREs *in vivo*.

The kidney androgen-regulated protein gene (*Kap*) is tissue-specific and regulated by androgen [10]. The *Kap* promoter has been for the study of androgen regulated gene expression [10] and also as an inducible gene-targeting tool [11-13]. The expression of a transgene consisting of the *Kap* promoter and the human Angiotensinogen (HAGT) gene is proximal tubule cell-specific and androgen regulated *in vivo* [14]. The degree of androgen induction of HAGT mRNA in these transgenic mice was 10-15 times higher than for endogenously expressed *Kap* [15]. As enhancers within introns or the 3'-untranslated region have been shown to increase the transcriptional efficiency and expression of chimeric transgenes in transgenic mice [11,15] and many attempts to use the *Kap* promoter to direct expression of intronless DNA have failed, the question arose of whether the HAGT gene contained an androgen-inducible enhancer. Enhancers in HAGT have been reported to direct cell type-specific expression of hAGT [16-18], however none have been associated with androgen response.

The aim of the current study was to locate portions of the hAGT gene that may act as enhancers in androgen-regulated expression in kidney cells and then identify its HREs *in vitro*. Experimental strategies used included overlap analysis to find the location of downstream enhancers, site-directed mutagenesis and DNA binding assay to evaluate putative response elements.

2. Materials and methods

2.1. Materials

OK (opossum kidney epithelial), was obtained from American Type Culture Collection (ATCC, Manassas, VA, USA). DMEM, and fetal bovine serum were obtained from Gibco. All oligonucleotides were synthesized at Shanghai Sangon (P.R. China). Transfection reagent Fugene 6 was purchased from Roche (USA). Luciferase reporter plasmids and luciferase substrate were obtained from Promega (USA). Dihydrotestosterone (DHT) and dexamethasone (Dex) were obtained from Sigma. All enzymes used for molecular cloning were from New England Biolabs (USA).

2.2. Plasmid Construction

First, pGL3-Kap1542 was constructed by inserting *Kap* promoter fragments -1542/-1 into Sac I and *Hind* III sites of pGL3-Basic as previously described[24]. Sequence analysis showed single restriction site in hAGT for *Bgl* II, *Bam* H I and *Xho* I site and double restriction sites for *Sal* I and *Kpn* I. A diagram of the restriction fragments from hAGT that were inserted as enhancer fragments is shown in Figure 1A. Because *Bgl* II leaves ends compatible with those left by *Bam* H I and *Xho* I leaves ends compatible with those left by *Sal* I, subclones *Bgl*-*Sal* (7.9kb), *Bgl*-*Bam*(2.7kb), *Bam*-*Sal*(5.2 kb), *Sal*-*Sal* (1.7kb), *Bam*-*Xho* and *Xho*-*Sal* were constructed by inserting corresponding fragments into a *Bam* H I and(or) *Sal* I cloning site following the poly A region of pGL3-Kap, that was inserted upstream of the promoter fragments. Subclone *Bgl*-*Sal*-*Sal*(7.9kb) containing the whole hAGT genomic region in *Kap*-HAGT transgenic construct was constructed by inserting 1.7 kb *Sal* I fragment into the *Sal* I site of subclone *Bgl*-*Sal* and selecting clones with correct orientation of the *Sal* I fragment after sequence analysis. The 1.4 kb *Kpn* I fragment was inserted into the *Kpn* I site of the polylinker in pGL3-Kap1542 to make subclone *Kpn*/*Kap*1542.

The positive control vector for glucocorticoid and androgen response was constructed by insertion of the luciferase reporter (excised from pGL3-Basic with *Nhe* I and *Xba* I) into the *Nhe* I site of pMSG-CAT (Pharmacia, Piscataway, NJ, USA). pET-rARDBD1 was constructed by insertion of a polymerase chain reaction DNA fragment with primer ARDBD1 (GGGGATCCCCCATCGACTATTACTTC) and ARDBD2 (GCGAATTCTTCTCCTTCTCCTGTAG), encoding the DNA binding domain of rat AR, from amino acids 530-627, into the *Bam*HI & *Eco*R I site of and pET28a (Novagen). The pET-ARDBD1 construct generated a 97-amino acid protein with a His tag on the amino terminus.

2.3 Site-directed Mutagenesis

The QuikChange® II kit (Stratagene, La Jolla, CA, USA) was used for site-directed mutagenesis. For the alteration of potential AREs in the *Kpn* I fragment of HAGT, the primers 5'-GTCTGCCAGTGGTTGCTATGTTCTGAGCATGGG, 5'-GGGACAAGAGTTATATGGGAAGTTTCTCGATTCTGCTCGATTTTGCTGTGAA, 5'-ACAAGAGTTATATGGGAAGTTTCTCTGTTTCTGCTCGATTTTGCTGTGAAC and the exact complements of each were used as pairs. All constructions were confirmed by DNA sequence analysis.

2.4 Cell culture and transient transfection assay

Opossum kidney epithelial cells (OK cells) were grown in DMEM (Life Technologies, Inc.) supplemented with 10% fetal bovine serum (FBS), L-glutamine, and antibiotics (penicillin/streptomycin). Transfection assays on OK cells were done using Fugene® 6 (Roche Molecular Biochemicals, Indianapolis, IN, USA) under the manufacturers directions. In detail, cells were seeded in 48-well tissue culture plates at a density of 2.5×10^4 cells in phenol red-free DMEM supplemented with 10% charcoal/dextran-treated FBS. 24h later, when cells reached about 50–60% confluence, FuGene was resuspended (5 min at room temperature) in serum-free medium and allowed to complex with DNA constructs at the ratio of 6:1 (reagent in microliters to DNA in micrograms) for 15 min at room temperature. The plasmid/FuGene mixture was then overlaid on the cells in a final volume of 0.4 ml of fresh charcoal-stripped complete medium. Each assay well contained 5 ng *Renilla* luciferase vector, phRL-TK (Promega) as the internal control, and 25ng pSG5rAR as the source of androgen receptor(AR) or 25ng pSG5hGR as a source of glucocorticoid receptor(GR). Each experiment included pGL3-Basic as the negative control, and 50 ng/well pMSG-Luc as a positive control for androgen and glucocorticoid induction. The experimental promoter constructs were transfected in amounts equimolar to each other and to the pGL3-Basic control. After incubation for 24h at 37°C, medium was replaced and cells were treated with/without 15nM dihydrotestosterone, 15nM dexamethasone or ethanol vehicle for 24-30 hours. Each transfection assay included six wells for each DNA constructs (three with and three without steroid) and was carried out at least three times on separate days.

2.5 Luciferase Reporter Assay

Cells were harvested for assay 48 hours after transfection. The medium was removed, the wells were washed once with PBS, and 50 ul of passive lysis buffer (Promega) was added to each well. The lysates were incubated at room temperature for 2 hours. Aliquots of 20ul cleared lysate were assayed for Firefly and *Renilla* luciferase activities in a luminometer(Luminoskan Ascent, Thermo Electron Corporation, Helsinki, Finland) using the reagents and directions provided in the Dual-Luciferase Reporter assay System(Promega). The results shown in the figures are a single experiment that is representative of at least three experiments, and plotted using Microsoft Excel software. The inductions given in the text of the results section are the average of three experiments.

2.6 Expression and purification of ARDBD

The *Escherichia coli* strain BL21(DE3) was transformed with pET-rARDBD1. 100ml cells were grown to OD₆₀₀=0.4-0.6 and induced with 0.5mM isopropyl-1-thio-β-D-galactopyranoside(IPTG) for 3h. After harvesting, cells were resuspended in 5ml of Buffer A (20mM Tris-HCl, pH7.9, 20% glycerol, 0.5mM phenylmethylsulfonyl fluoride) and lysed by sonication. The supernatant was incubated with 1.5ml of Ni-NTA-agarose (Qiagen) resin in batch, rocked gently for 30min at 4°C, and then loaded on to a small Bio-Rad Econo-column. After successively washing the resin with 15ml of Buffer A and 15ml of Buffer A containing 20mM imidazole, ARDBD was eluted with Buffer A containing 150mM imidazole. The fractions were analyzed by SDS-polyacrylamide gel electrophoresis and Coomassie Blue staining. Peak fractions were combined. The final preparation contained 100μg/ml ARDBD.

2.7 Electrophoretic mobility shift assay (EMSA)

Mobility shift experiments were performed with 5 μg of purified ARDBD/GRDBD and 5' end DIG-labeled double-stranded oligonucleotides containing the desired HRE sequences. Typically, the binding reactions were carried out in 20 μl containing 10 mM Tris-HCl (pH 7.5), 65 mM NaCl, 5 mM dithiothreitol, 5 mM MgCl₂, 0.05% Nonidet P-40, 10% glycerol, 1 mg/ml bovine serum albumin, and 25 μg/ml poly(dI-dC) as a carrier for 30 min at 4 °C. In competition experiments, a 100-fold molar excess of the cold competitor was preincubated with purified ARDBD/GRDBD for 30 min at 4 °C before the labeled DNA fragment were added. The complexes were resolved on nondenaturing 6% polyacrylamide gels in 0.5 ×TBE buffer for 1 h at 14 V/cm, and viewed following ECL detection.

3. Results

3.1 Androgen Response and enhancer effect of hAGT gene

In vivo data from transgenic mice suggested that a 1542 bp *Kap* promoter fragment (*Kap*1542) alone was sufficient to drive the PTC-specific, androgen-regulated hAGT expression, but not several other genes, whereas the chimeric *Kap*1542-HAGT could successfully drive the PTC-specific, androgen-regulated human renin and rat NHE3 expression. Therefore *Kap* promoter sequences seemed necessary but not sufficient for such expression and additional regulatory elements within the hAGT genomic sequence might contribute to tissue-specific expression. Therefore, the relationship between the *Kap* promoter and hAGT reporter gene was studied.

To analyze 1542bp *Kap* promoter activity *in vitro*, the fragment (nt -1542 to +1 for *Kap*) was cloned into a luciferase reporter plasmid (see *Materials and Methods*). The reporter activity was assayed by transient transfection in OK cells, Cos-1 cells, NIH3T3 cells and LNCaP cells. The AR was co-transfected into all lines except for LNCaP, which has an endogenous AR. Luciferase activity from the *Kap* promoter construct was induced 1.2 fold in the presence of 10⁻⁶ M 5α-DHT only in OK cells. Under the same conditions, the mouse mammary tumor virus (MMTV)-Luc construct (positive control) showed 8- to 10-fold steroid-induced transcription in all cell lines, and pGL3-basic (negative control), showed no induction in the presence of androgens (data not shown). Therefore it seemed that OK cells, but not other cell lines, provided factors necessary for *Kap* promoter activity, so OK cells were chosen for additional experiments.

The hAGT reporter used in the transgenic mouse construct included exons 2 through 5, intronic regions, and some 3'-flanking sequence. This 9.7 kb sequence was inserted into the enhancer site of the *Kap*1542, and referred to as *Bgl-Sal-Sal*(Figure 1a). The chimeric construct was tested in OK cells. *Bgl-Sal-Sal* had a 1.5 to 3 fold higher luciferase expression than that of *Kap*1542 without the HAGT and was induced an additional 2-3 fold by DHT, whereas *Kap*1542 was not or slightly induced (Figure 1b). From this result, we inferred that the HAGT

sequence included androgen responsive sequences, although the level of induction in transfection assay was much lower than that for the transgene *in vivo*.

3.2 Identification of an androgen-responsive enhancer region in the HAGT reporter

Two liver cell-type dependent enhancer elements have been reported in the downstream region of the HAGT gene [17,18]. Therefore, chimeric constructs consisting of the mouse *Kap* promoter and hAGT restriction fragments that either included or excluded known enhancers were subcloned and assayed by transient transfection in OK cells (Figure 1b). The cleavage of the HAGT gene by *Sal I* yielded the subclones *Bgl-Sal*(7.9kb) and *Sal-Sal*(1.7kb)(Figure 1a). Luciferase reporter activity from the *Sal-Sal*, which contained the previously identified core enhancer d61, was about two-fold higher than the basal activity of *Kap1542*, but was not induced by DHT. However, induction by DHT of luciferase activity of the *Bgl-Sal* was a slightly higher than that of *Bgl-Sal-Sal*. The subclone *Xho-Sal* included the previously identified core enhancer B2-5 but was not induced by androgen and did not increase the basic promoter activity of *Kap1542*. The subclone *Bam-Xho* had a similar androgen response to *Bam-Sal*(Figure 1b). We concluded that neither B2-5 nor d61 were androgen-responsive enhancers of *Kap1542*.

A smaller *Kpn I* fragment (+2987 to +4171), located within *Bam-Xho* (Figure 1a), increased the basal activity and androgen induced activity of the *Kap147* promoter by 2-3 fold (Figure 1b). This induction was higher than for the other constructs containing portions of HAGT, therefore the *Kpn I* fragment was most likely to contain an androgen response element.

3.3 Steroid specificity of the hAGT *Kpn I* fragment

We tested the responses of the construct containing hAGT *Kpn I* fragment to both androgen and glucorticoid in OK cells. To measure androgen-response, the luciferase construct was co-transfected with pSG5-mAR and treated with 5 α -DHT(Figure 2a). To measure glucocorticoid-response, the hAGT construct was co-transfected with pSG5-hGR and treated with dexamethasone (Figure 2b). The positive control in each case was pMSG-luc, which is inducible by either androgen or glucorticoid. As shown in Figure 2, expression of *KpnKap1542* was induced 2.4-fold by DHT and expression of *KpnKap 1542* was induced 4.5-fold by DEX. The inductions by DEX were greater than by DHT. Therefore the hAGT *Kpn I* fragment did not appear to be selective for androgen.

3.4 Analysis of Sequences Involved in the Steroid Response in HAGT *Kpn I*

Initially, we looked for potential AREs/GREs consisting of closely spaced hexamers resembling the core motif TGTTCT forming indirect or direct repeats. Finding none, the criteria were lowered to include partial direct repeats including any hexamer following the pattern A/TGTnCA/T. We found two possible AREs/GREs: a DR3, a direct repeat of hexamers separated by three nts close to the 5' *Kpn I* site, with the sequence AGTGCTtgcTGTGCT (designated A); and another DR3 nearer to the 3' *Kpn I* site with the sequence GGAActtctGTACT (designated B)(Table 1).

To assess the function of the putative ARE/GREs, we used site-directed mutagenesis to eliminate guanine or cytosine contact sites. The alteration of site A, to the sequence AGTGGTtgcTATGTT (C19) (Table 1), eliminated any occurrence of nGnnCn. This alteration had little effect on androgen induction but no effect on glucocorticoid induction (Figure 3). Site B was altered in either of two ways: C1292 with sequence GGAAGTtctTCGATT resulted in alteration of both hexamer half sites, and G1303 with sequence GGAActtctTCTGTT resulted in alteration of the right hexamer only (Table 1). Inductions of C1292 and G1303 by androgen were decreased from 3.6-fold to 2.2- and 1.5-fold respectively (Figure 3A).

Inductions of C1292 by glucocorticoid were one-tenth those of the unaltered sequence (Figure 3B). These data showed that site B was involved in both AR and GR response.

3.5 Electrophoretic mobility shift assay (EMSA) of ARE/GRE in hAGT *Kpn* I

To study sequence-specific binding of AR and GR to site A and site B of hAGT *Kpn* fragment, we used 24-mer oligonucleotide probes corresponding to site A (HAGTA) and site B (HAGTB), in EMSA with AR-DBD and GR-DBD proteins. As shown in Figure 4, HAGTA bound only to the AR-DBD but not GR-DBD, while HAGTB bound to both AR-DBD and GR-DBD. This was consistent with the result from site-directed mutagenesis that site A was a potential androgen-specific HRE and site B was not an androgen-specific HRE.

We used a set of unlabeled competitor oligonucleotides representing standard HAGTA, HAGTB, and variants in which the A/TGTnCA/T motif had been mutated (C19g, C1292). Sequence-specific complexes between HAGTA and purified ARDBD were completely suppressed by an excess of unlabeled competitor HAGTA, but not by mutant competitor C19g (Figure 4a). Complex formations between HAGTB and purified AR-DBD or GR-DBD were completely suppressed by an excess of unlabeled competitor HAGTB, and partly by mutant competitor C1292 (Figure 4b).

4 Discussion

In vitro, the receptors for glucocorticoids, mineralocorticoids, progestins, and androgens recognize the same DNA consensus sequence, which consists of an inverted repeat of two 6-bp half-sites, separated by a three-nucleotide (nt) spacer 5'-GGTACAnnnTGTCT-3'. The requirements for androgen receptor-specific recognition *in vivo* are less clear cut. The palindromic pattern sometimes applies to androgen-response elements found *in vivo*, such as in the prostatein gene [6], but direct repeats of hexamers form the androgen-response elements in the genes for probasin [5], sex-limited protein [7], and secretory component [8]. The spacing between hexamers can be variable, as in the Pem homeobox gene, the ARE of which is a direct repeat of hexamers separated by five nt [19].

In our previous study, a human gene fragment, hAGT, conferred androgen inducibility to a tissue-specific promoter, *Kap*, in transgenic mice [14] and in the current study, the same fragment conferred androgen-inducibility to a *Kap*-luciferase reporter construct *in vitro*. Therefore, we used sequence homology algorithms to search for potential AR and GR contact sites in the hAGT *Kpn* I fragment. When looking for potential AREs/GREs with TGTTCT hexamers that formed indirect or direct repeats, we found none. When the criteria were lowered to the hexamer A/TGTnCA/T and two AREs/GREs were located. Site-specific mutagenesis and EMSA in the DR3DR3 motif showed that site A was selective for androgen receptor, but either androgen receptor or glucocorticoid receptor could interact with site B.

The hAGT *Kpn* I fragment did not overlap with other portions of *HAGT* shown to have enhancer activity for liver-specific expression. The enhancer elements identified elsewhere in *HAGT* are known as B2-5 and d61 [17,18]. When *HAGT* fragments including either B2-5 or d61 were tested in luciferase reporter constructs under the same conditions as the *Kpn*I fragment, neither conferred androgen response to the *Kap* promoter.

The *HAGT* promoter directed expression of its own mRNA in liver, kidney, heart, adrenal gland, ovary, brain and adipose tissue of HuAoGen transgenic mice, furthermore, the *HAGT* expression in the kidney and adrenals of HuAoGen mice was regulated by androgen [21]. Substitution of the *Kap* promoter for the *HAGT* promoter restricted *HAGT* transgene expression to the proximal tubules and to the epididymis [14], another site of endogenous *Kap* expression. The expression of the *Kap*-*HAGT* transgene is normally undetectable in the

kidney of female mice, can be dramatically induced in response to testosterone treatment [14]. We identified androgen response element in the *Kap* promoter [22], and in the current study showed that sequences from the HAGT contributed to hormonal response. This is the first report to suggest a basis for the androgen or glucocorticoid response in the downstream portion of this gene.

In summary, we found that an androgen-regulated downstream enhancer with two HREs is localized to a 1.4kb sequence within intron 2 of the HAGT gene, one of which is specific to androgen. Future experiments will address the question of whether the hybrid HAGT Kpn I enhancer-Kap promoter can be used to specifically target other genes to the proximal tubule cells and how is it regulated in vivo.

Acknowledgments

The androgen responsive promoter pPB(-285/+32)-luc, the androgen receptor expression vector pSG5-rAR and glucocorticoid receptor expression vector pSG5-hGR were kindly provided by Jorma Palvimo, University of Helsinki, Finland. Supported by Science and Technology Commission of Shanghai, Shanghai Pujiang Program 06PJ14024; NIH grant DK069312.

References

1. Claessens F, Verrijdt G, Schoenmaker E, Haelens A, Peeters B, Verhoeven G, Rombauts W. Selective DNA binding by the androgen receptor as a mechanism for hormone-specific gene regulation. *J Steroid Biochem Mol Biol* 2001;76:23–30. [PubMed: 11384860]
2. Hsu CL, Chen YL, Ting HJ, Lin WJ, Yang Z, Zhang Y, Wang L, Wu CT, Chang HC, Yeh S, Pimplikar SW, Chang C. Androgen receptor (AR) NH2- and COOH-terminal interactions result in the differential influences on the AR-mediated transactivation and cell growth. *Mol Endocrinol* 2005;19:350–61. [PubMed: 15514032]
3. Nelson CC, Hendy SC, Shukin RJ, Cheng H, Bruchovsky N, Koop BF, Rennie PS. Determinants of DNA sequence specificity of the androgen, progesterone, and glucocorticoid receptors: Evidence for differential steroid receptor response elements. *Mol Endocrinol* 1999;13:2090–2107. [PubMed: 10598584]
4. Robins DM. Androgen receptor and molecular mechanisms of male-specific gene expression. *Novartis Foundation Symposium* 2005;268:42–52. [PubMed: 16206874]
5. Zhang J, Gao N, Kasper S, Reid K, Nelson C, Matusik RJ. An androgen-dependent upstream enhancer is essential for high levels of probasin gene expression. *Endocrinology* 2004;145:134–148. [PubMed: 14500578]
6. Tan J, Marschke KB, Ho KC, Perry ST, Wilson EM, French FS. Response elements of the androgen-regulated C3 gene. *J Biol Chem* 1992;267:4456–4466. [PubMed: 1537831]
7. Loreni F, Stavenhagen J, Kalff M, Robins DM. A complex androgen responsive enhancer resides 2 kilobases upstream of the mouse *Slp* gene. *Mol Cell Biol* 1988;8:2350–2360. [PubMed: 3165490]
8. Haelens A, Verrijdt G, Callewaert L, Peeters B, Rombauts W, Claessens F. Androgen-receptor-specific DNA binding to an element in the first exon of the human secretory component gene. *Biochem J* 2001;353:611–620. [PubMed: 11171058]
9. Zhou Z, Corden JL, Brown TR. Identification and characterization of a novel androgen response element composed of a direct repeat. *J Biol Chem* 1997;272:8227–8235. [PubMed: 9079641]
10. Soler M, Tornavaca O, Solé E, Menoyo A, Hardy D, Catterall JF, Vandewalle A, Meseguer A. Hormone-specific regulation of the kidney androgen-regulated gene promoter in cultured mouse renal proximal tubule cells. *Biochem J* 2002;366:757–766. [PubMed: 12030848]
11. Malstrom SE, Tornavaca O, Meseguer A, Purchio AF, West DB. The Characterization and Hormonal Regulation of Kidney Androgen-Regulated Protein (*Kap*)- Luciferase Transgenic Mice. *Toxicol Sci* 2004;79:266–277. [PubMed: 15056803]
12. Lavoie JL, Lake-Bruse KD, Sigmund CD. Increased blood pressure in transgenic mice expressing both human renin and Angiotensinogen in the renal proximal tubule. *Am J Physiol Renal Physiol* 2004;286:F965–F971. [PubMed: 15075192]

13. Bianco, Robert A.; Keen, Henry L.; Lavoie, Julie L.; Sigmund, Curt D. Untraditional methods for targeting the kidney in transgenic mice. *Am J Physiol Renal Physiol* 2003;285:F1027–1033. [PubMed: 14600026]
14. Ding Y, Davisson RL, Hardy DO, Zhu LJ, Merrill DC, Catterall JF, Sigmund CD. The kidney androgen-regulated protein promoter confers renal proximal tubule cell-specific and highly androgen-responsive expression on the human angiotensinogen gene in transgenic mice. *J Biol Chem* 1997;272:28142–28148. [PubMed: 9346970]
15. Hardy DO, Niu EM, Catterall JF. *Kap* promoter analysis *in vivo*: A regulatory role for a truncated L1 repeat. *Mol Cell Endocrinol* 2001;181:57–67. [PubMed: 11476941]
16. Brinster RL, Allen JM, Behringer RR, Gelinas RE, Palmiter RD. Introns. *Proc Natl Acad Sci USA* 1988;85:836–840. [PubMed: 3422466]
17. Takabashi S, Fukamizu A, Hasegawa T, Yokoyama M, Nomura T, Katsuki M, Murakami K. *Biochem Biophys Res Commun* 1991;180:1103–1109. [PubMed: 1659395]
18. Nibu Y, Takahashi S, Tanimoto K, Murakami K, Fukamizu A. Identification of cell type-dependent enhancer core element located in the 3'-downstream region of the human angiotensinogen gene. *J Biol Chem* 1994;269:28598–28605. [PubMed: 7961807]
19. Nibu Y, Tanimoto K, Takahashi S, Ono H, Murakami K, Fukamizu A. A cell type-dependent enhancer core element is located in exon 5 of the human angiotensinogen gene. *Biochem Biophys Res Comm* 1994;205:1102–1108. [PubMed: 7802638]
20. Barbeslescu K, Geserik C, Schüttke I, Schluning W, Haendler B. New androgen response elements in the murine *pem* promoter mediate selective transactivation. *Mol Endocrin* 2001;15:1803–1816.
21. Yang G, Merrill DC, Thompson M, Robillard JE, Sigmund CD. Functional expression of the human angiotensinogen gene in transgenic mice. *J Biol Chem* 1994;269:32497–502. [PubMed: 7798251]
22. Fan L, Hardy DO, Catterall JF, Zhao J, Li S. Identification and characterization of the minimal androgen-regulated kidney-specific androgen-regulated protein gene promoter. *Acta Biochim Biophys Sin* 2008;40:979–988. [PubMed: 19089304]

A

B

C

Figure 1. Transient transfection assay of the androgen-responsible enhancer effect of hAGT gene
A Map of overlap constructs. The HAGT reporter includes exons II, III, IV and V, as well as intervening sequences between the included exons and some 3' untranslated sequence. The overlapping fragments were made using available restriction sites, and the lines indicate the portions of the structural gene represented. Kap promoter fragments were inserted into the multiple cloning sites preceding the luciferase reporter (LUC). The HAGT fragments were inserted into a *Bam*H I and *Sal* I cloning site following the poly A region, except for the *Kpn* I fragment, that was inserted upstream of the promoter fragments. **B Transfection.** Kap1542 plasmids containing various fragments of the HAGT gene were co-transfected with androgen receptor expression vector into OK cells in equimolar amount to the negative control, pGL3-Basic. pPB-Luc was used as androgen inducible positive control. pRL-TK was used as the internal control. Each construct was represented by six wells of a 48-well assay plate, three of which were exposed to 15 nM DHT. Cells were harvested 48 hours after transfection. Activities of firefly and Renilla luciferases were measured sequentially from the same aliquots using the Dual-Luciferase Reporter Assay System (Promega). A single experiment that is representative of at least three experiments was shown.

The inductions by DHT were: PB, 20-fold, Kap1542, 1.2-fold, Bgl-Sal-Sal, 2.4-fold, Bgl-Bam, 2.5-fold, Bam-Sal, 1.7-fold, Bam-Xho, 1.6 fold, Xho-Sal, 1.1 fold, Kpn, 3.2-fold, and Sal, 1.2-fold. **C Average induction fold by DHT.** In each experiment, the induction fold of every sample was divided by that of the negative control, pGL3-Basic. Average data from five independent transfection experiments were shown ($\pm$ S.E.).

A

B

Figure 2. Steroid specific of hAGT *Kpn I* fragment.

Constructs were co-transfected with pSG5-hGR into OK cells and assayed in the absence and presence of 15 nM Dexamethasone (DEX) or 15nM DHT. (A) The inductions by DHT were, Kap1542, 1-fold, KpnKap1542, 2.4-fold. (B) The inductions by DEX were, Kap1542, 2-fold, KpnKap1542, 4.5-fold.

A.

B.

Figure 3.

Mutational analysis of ARE-like/GRE-like sequences in the HAGT Kpn I enhancer fragment. There were 2 possible androgen-responsive elements in the HAGT Kpn fragment (A and B) and three kinds of mutation (C19, C1292, C1302)

(A) Induction by DHT. The unaltered construct was induced 3.6-fold by DHT. C19 contained a point mutation of a DR3 near the upstream *Kpn* I site, this was induced 3.1-fold. C1292 and C1302 contain point mutations in a DR3DR3 near the downstream *Kpn* I site, these mutations resulted in lowered induction in response to DHT, 2.2-fold and 1.53 fold.

(B) Induction by DEX. The unaltered construct was induced 27.9-fold by DEX. C19 contained a point mutations of a DR3 near the upstream *Kpn* I site, this was induced 31.3-fold. C1292

contain point mutations in a DR3DR3 near the downstream Kpn I site, this mutation resulted in lowered induction in response to DEX, 3.96-fold.

HAGTA	+	+	+	+	+	+
Comp.	-	-	-	HAGTA	HAGTA	C19G
ARDED	+	-	-	-	+	+
GRDED	-	-	+	+	-	-

HAGTB	+	+	+	+	+	+	+
AR	+	+	+	-	-	-	-
GR	-	-	-	-	+	+	+
Comp.	C1292	HAGTB	-	-	-	HAGTB	C1292

Electrophoretic mobility shift assay (EMSA) for interactions of the hAGT Kpn fragments with purified recombinant ARDBD and GRDBD. Specific complexes are suppressed by 50ng of unlabeled hAGT Kpn HRE(s) (standard HAGTA, HAGTB) and unlabeled mutant hAGT Kpn HRE(s) (C19g, C1292) competitor. **(A) HAGTA (B) HAGTB**

Table 1Original and mutant DNA sequences of two possible Steroid Response elements in HAGT *Kpn* I

Original	HAGT(<i>Kpn</i>) 20-34 A	AGTGCTtgcTGTGCT	DR3
Altered	C19g	AGTGGTtgcTATGTT	
Original	HAGT(<i>Kpn</i>)1291-1317 B	GGAACtTtcTGTACTtTcTGCTCGA	DR3DR3
Altered	C1292	GGAAGTtTcTCTATTtTcTGCTCGA	
Altered	C1302	GGAACtTtcTCTGTTtTcTGCTCGA	

Column 1 states whether this is the original sequence or the altered sequence. Column 2 refers to the location of the original sequence or the designation of the plasmid containing the altered sequence. Column 3 shows the putative androgen-response element, with the hexamer in upper case and the spacer in lower case. Column 4 refers to the kind of repeat - direct repeat (DR) or inverted repeat (IR) - and number of bases in the spacer.