Published in final edited form as:

J Chem Eng Data. 2010 May 13; 55(5): 1837–1841. doi:10.1021/je9008377.

Local semi-empirical formulae for the contact values of the singlet distribution functions of a double layer

Whashington Silvestre-Alcantara,

Department of Physics and Laboratory for Theoretical Physics, University of Puerto Rico, San Juan, Puerto Rico 00931-3343

Lutful Bari Bhuiyan, and

Department of Physics and Laboratory for Theoretical Physics, University of Puerto Rico, San Juan, Puerto Rico 00931-3343

Douglas Henderson

Department of Chemistry and Biochemistry, Brigham Young University, Provo UT 84602

Abstract

The electrochemical double layer is an important practical and theoretical problem. Generally speaking, experiment gives valuable information about quantities, such as potential differences, that involve integrals of density and charge profiles but does not provide direct information about the profiles themselves. Computer simulations have given numerical information about these profiles. However, explicit expressions are useful in understanding these data. For some years an exact expression has been known for the contact value of total density profile of the ions in the double layer but, until recently, an expression for the contact value of the more important charge profile has been lacking. A few years ago, a semi-empirical local result for the charge profile, valid at low electrode charge, was proposed and, very recently, extended to higher electrode charge. This expression contains a parameter; the effect of variations in this parameter is explored in this paper and the result is compared with a large set of simulation data for the contact values of various profiles that we have accumulated in the past few years. The agreement of the semi-empirical expression with our simulation results is excellent. The best values for this parameter are fairly close to the value suggested by theory.

Introduction

Josef Barthel has made extensive contributions to theory and experiment for electrolytes, including the correlation and systematization of experimental data. We dedicate this paper to him and hope that our own modest efforts towards the systematization of data about the density and charge profiles of inhomogeneous electrolytes will be of interest to him and our colleagues.

As is well known, the double layer is the interfacial region formed by an electrolyte near an electrode. The name *double layer* comes from the intuitive idea that the charge of the electrode is one charged layer and the charge of the interfacial electrolyte is a second layer that is equal in magnitude, but opposite in sign, to that of the electrode charge. Although the charge in the interfacial electrolyte is, in total, equal and opposite to the electrode charge (known as *local electroneutrality* in the double layer literature), the density and profile of the interfacial

Correspondence to: Lutful Bari Bhuiyan.

Supporting Information **Available:** A fuller set of figures containing a comparison of the results of our simulations and fit for the double layer correlations. This material is available free of charge via Internet at <http://pubs.acs.org>.

electrolyte can be nonmonotonic and the charge profile of the interfacial electrolyte can even change sign. Actually, the charged layer in the electrode can also be nonmonotonic but this is rarely considered. Even so, the name *double layer* persists.

An understanding of the double layer is helpful for studies of energy storage, corrosion, colloidal suspensions and, if the ‘electrode’ is a membrane or protein, physiology. Electrochemical experiments provide information about the voltage difference across the interfacial region as a function of the charge of the electrode but provide only limited information about the local distribution (the profiles) of charge within the double layer whereas an understanding of corrosion, colloidal forces, and membrane function really requires such local information.

The ‘standard’ theory of the double layer is the venerable classical theory due to Gouy, Chapman, and Stern^{1–3} (GCS). It is based on what is usually called the *primitive model* (PM) of the electrolyte where the solvent is replaced by a dielectric continuum whose dielectric constant, ϵ_r , is equal to that of the solvent and the ions are mimicked by rigid spheres with charges at their centres. For simplicity, the electrode next to the electrolyte is assumed to be planar with all the charge of the electrode confined at the surface (zero skin depth). If all the ions have the same diameter, the PM is called the *restricted primitive model* (RPM) with the resultant double layer being called the RPM planar double layer. The GCS theory assumes that the ions are point ions, that their diameters may be neglected, and that an ion interacts with the electrode and the other ions only through a mean field raised by all the ions. Thus, the theory neglects interionic correlations beyond this mean field interaction. To avoid an electrostatic singularity, in the GCS theory the ions are given a distance of closest approach to the electrode. This is inconsistent but better than a singularity. The GCS theory provides information about both the local profiles and the nonlocal voltage difference across the double layer. This is exemplified by the GCS result for the density profiles at contact. Assuming further for simplicity, that the ions are symmetric in charge and size, the GCS contact values are given by⁴

$$g_{\pm}^{GCS}(d/2) = 1 + \frac{b^2}{2} \pm b \sqrt{1 + \frac{b^2}{4}}, \quad (1)$$

where d is the common ion diameter, z and e are the magnitudes of the common valence and elementary charge, and $d/2$ the distance of closest approach. The distance of closest approach need not be the ion radius but this is the simplest choice and the one most consistent with the primitive model. The parameter $b = (\beta z e \sigma) / (\epsilon_0 \epsilon_r)$ is a dimensionless measure of electrode surface charge density, σ . The quantity $\beta = 1/k_B T$, where T is the temperature and k_B is the Boltzmann constant. The parameter κ is the Debye inverse screening length, whose square is given by

$$\kappa^2 = \frac{\beta e^2}{\epsilon_0 \epsilon_r} \sum z_s^2 \rho_s = \frac{\beta e^2 z^2 \rho}{\epsilon_0 \epsilon_r}, \quad (2)$$

where $\rho = \sum \rho_s$ is the density of all of the ions (counterions and coions). We assume without loss of generality, that the electrode is negatively charged, so that the counterions will have positive sign, while the coions will have negative sign. Thus, $g_{ctr}(d/2) = g_+(d/2)$ is the contact value of the counterions and $g_{co}(d/2) = g_-(d/2)$ is the contact value of the coions. The total density and charge profile results are proportional to the sum and difference of $g_{ctr}(x)$ and

$g_{co}(x)$, respectively, with x being the perpendicular distance from the electrode surface into the electrolyte. Thus, $g_{sum}(x) = [g_{ctr}(x) + g_{co}(x)]/2$ and $g_{diff}(x) = [g_{ctr}(x) - g_{co}(x)]/2$. Note that $g_{sum}^{GCS}(d/2)$ and is equal to the first two terms in eq 1 and $g_{sum}^{GCS}(d/2)$ is equal to the last term in eq 1. Another profile of interest is the product profile, $g_{ctr}(x)g_{co}(x)$. It is seen from eq 1 that the GCS product profile at contact is one. Within the GCS theory, this is actually true for all x . The GCS theory assumes that $g_{co}(x)$ is exactly the reciprocal of $g_{ctr}(x)$. There is no reason to suppose that this is true in general. Thus, the value of the product profile and its departure from one is a useful test of the accuracy of the GCS theory.

Contact value theorems

The reason for our interest in the contact values is that they provide insight into the accuracy of a theory. Any decent theory will satisfy local electroneutrality, that is, will have the correct value for the integral of the charge profile and, thus, the correct value for the integral of $g_{diff}(x)$. If a function has both the correct contact value and the correct integral, there is reason to hope that the function is fairly reliable.

Henderson and Blum⁵ and Henderson *et al.*⁶ have obtained an exact sum rule for the contact value of $g_{sum}(x)$. For a symmetric valence RPM double layer, their result is

$$g_{sum}(d/2) = a + \frac{b^2}{2}, \quad (3)$$

where $a = p/\rho kT$ and p is the bulk (osmotic) pressure. Note that the GCS expression satisfies eq 3 only if $a = 1$. This is a result of the neglect of ion diameters and the ion-ion interaction, which leads to an ideal gas approximation for a . If a is near unity, the GCS theory might be expected to be a fairly reasonable theory for $g_{sum}(x)$, but not necessarily for $g_{diff}(x)$. However, a is often not close to unity. Under such circumstances, the GCS theory will likely fail even for $g_{sum}(x)$.

A contact value sum rule for $g_{diff}(d/2)$ would be of more value since it relates directly to the electrical properties of the double layer. Until recently, such a result was not available. However, independently two groups have recently proposed sum rules for $g_{diff}(d/2)$. The result of Holovko *et al.*⁷⁻⁹ is exact and general but is nonlocal and difficult to use. Their expression is

$$g_{diff}(d/2) = -\beta z e \int_{d/2}^{\infty} \frac{\partial \phi(t)}{\partial t} g_{sum}(t) dt. \quad (4)$$

The function $\phi(x)$ is the potential difference between x and ∞ and is obtained from

$$\phi(x) = -\frac{e}{\epsilon_0 \epsilon_r} \sum_i z_i \rho_i \int_x^{\infty} (t - x) g_i(t) dt, \quad (5)$$

which leads to

$$\frac{\partial \phi(x)}{\partial x} = -\frac{ez\rho}{\epsilon_0\epsilon_r} \int_x^\infty g_{diff}(t)dt, \quad (6)$$

for a symmetric valence $z:z$ electrolyte. On substituting eq 6 in eq 4, it is seen that the latter now involves two integrations, is highly nonlocal, and is difficult to use. The result of Henderson and Boda¹⁰ on the other hand,

$$g_{diff}(d/2) = ab + O(b^3), \quad (7)$$

is local and is easy to use but is valid only for small b . The parameter a can be determined from simulations of $g_{sum}(d/2)$ at small b or from the mean spherical approximation that we have found to be quite accurate.¹¹ Equation 7 is semi-empirical and was obtained by inference from simulation results of Henderson and Boda.¹⁰ If there is a method of obtaining eq 7 from eq 4, it is not obvious. The possibility that eq 7 might fail in some application beyond those covered by the Henderson-Boda simulations cannot be discounted but is not likely as eq 7 has since been tested using a large set of simulations^{12–16} covering a broad range of states. Using eqs 3 and 7, a formula for the contact value of the product profile that is applicable for small b is

$$g_{ctr}(d/2)g_{co}(d/2) = a^2 + (a - a^2)b^2 + O(b^4). \quad (8)$$

This result correctly predicts the initial slope of $g_{ctr}(d/2)g_{co}(d/2)$ as a function of b .

The extension of eqs 7 and 8 to large b has been difficult. Because of the presence of the term containing $(a - a^2)$ in the right hand side of eq 8, the possibility of unphysical negative values of the product $g_{ctr}(d/2)g_{co}(d/2)$ cannot be excluded. In a recent paper, henceforth to be denoted by I, Henderson and Bhuiyan¹⁷ have suggested an empirical relation for $g_{ctr}(d/2)g_{co}(d/2)$, viz.,

$$g_{ctr}(d/2)g_{co}(d/2) = (a^2 + ab^2) \exp\left(-b^2 \frac{1 + \rho^* b/2}{1 + \alpha b^2}\right), \quad (9)$$

where $\rho^* = \rho d^3$ and α is a parameter.

As in I, the above equation may be combined with the exact contact condition, eq 3 to give

$$g_{diff}(d/2) = ([g_{sum}(d/2)]^2 - g_{ctr}(d/2)g_{co}(d/2))^{1/2}, \quad (10)$$

$$g_{ctr}(d/2) = a + \frac{b^2}{2} + g_{diff}(d/2), \quad (11)$$

and

$$g_{co}(d/2) = a + \frac{b^2}{2} - g_{diff}(d/2). \quad (12)$$

Equation (9) does not lead to negative values for the product of the contact values and was inspired by an expression of Lou and Lee¹⁸ who developed a theory that is applicable at large b . Henderson and Bhuiyan took their result at large b together with eq 8 at small b with the eq 9 being an interpolation between these two results. The parameter α does not appear explicitly in the theory of Lou and Lee. Their expression arises from what they call a modified Poisson-Boltzmann theory and is obtained from eq 9 in the limit of large b with a factor of 1/2 in place of α . A physical significance of the parameter α may be had by noting that α controls the rate at which the contact value of the product profile tends to zero at large b . The particular value of $\alpha = 1/2$ is well founded in the Lou-Lee theory. This value was used in I and gave good results. However, the results were obtained at one temperature only. In this paper we will explore eq 9 with α as a parameter for more states covering a broader temperature range and will adjust α to give best fit to the simulations.^{13–17} A variable α is likely to offer insight as to the asymptotic behaviour of the product function $g_{ctr}(d/2)g_{co}(d/2)$ at large b under different physical conditions. We have also supplemented the simulation data by performing fresh simulations for more states. Although the ‘best’ values of α differ from 1/2, they are never far from 1/2.

Monte Carlo simulations

The MC simulations were done in the canonical ensemble using the standard Metropolis algorithm. The central MC box was a rectangular parallelepiped with one face being the planar charged surface at $x = 0$ and the other face at $x = l_x$ being uncharged. The parallel *charged sheets* method of Torrie and Valleau¹⁹ was adopted to account for the long range Coulomb interactions together with the usual minimum image plus periodic boundary conditions along the remaining y and z axes. This technique has been used in recent years by other groups.^{20, 21} Although Torrie and Valleau used the grand canonical ensemble, in the canonical ensemble the required bulk concentration can be achieved by adjusting the MC cell length l_x . The typical number of configurations was around 10^8 with the first 10^7 being employed for equilibration of the system. The uncertainty in reproducing the bulk concentration was never more than $\pm 2\%$.

Results and Discussion

Results are presented in Figures 1–5 for a spectrum of states covering a range of electrolyte temperature and density. These are only a representative subset of 20 that the authors have examined. For the purposes of this discussion it is convenient to use universal reduced (non-dimensional) parameters, viz., the reduced temperature $T^* = (4\pi\epsilon_0\epsilon_r k_B T d) / (z^2 e^2)$ and the reduced density $\rho^* = \rho d^3$. Note that $T^* = 1/\Gamma_p$, where Γ_p is the plasma coupling constant. The physical states studied ranged in T^* from 0.15 to 0.6 encompassing both 1:1 and 2:2 valence electrolytes, while generally both low (0.00925) and high values (0.0925) of ρ^* were chosen to sample dilute and concentrated solutions. We have compared eq 9 to a large set of earlier simulation values^{12–16} and new simulation values for $g_{ctr}(d/2)$ and $g_{co}(d/2)$.

For each physical state we have adjusted α by trial and error to optimize the fit of eq 9 to the corresponding simulation value of the product $g_{ctr}(d/2)g_{co}(d/2)$. This optimum α was then used to evaluate $g_{diff}(d/2)$, $g_{ctr}(d/2)$, and $g_{co}(d/2)$ from eqs 10, 11, and 12, respectively. In the figures, results for $\alpha = 1$ are also given for comparison purposes. Additionally, in panel (a) of Figure 1 we have included the result for $\alpha = 1/2$ from I to make contact with the earlier work.

It is clear from this figure that the optimum α (≈ 0.6) leads to a closer agreement of eq 9 with the simulations than does the earlier result with $\alpha = 1/2$. Indeed in the course of the current calculations we have tested all the situations presented in I, and in every case the results with a variable (optimum) α were superior to those with $\alpha = 1/2$ vis-a-vis the simulations. These results are not shown here for reasons of clarity and brevity. A fuller set of our figures is accessible electronically as Supporting Information (SI) in the World Wide Web edition of this paper.

The presence of the exponential factor in the right hand side of eq 9 ensures that the product $g_{ctr}(d/2)g_{co}(d/2)$ goes to zero at large b . But the manner in which this occurs is determined by the parameter α . For instance, because of the negative sign of the exponent, the exponential factor is relatively larger when α is large (≥ 1) compared to when α is small (< 1). It is evident from the figures that with an optimum α , eq 9 generally reproduces the simulation data very well. We note here that the optimum value of α in all cases is fairly close to the value $1/2$ suggested by the Lou-Lee theory.¹⁸ Values of α that differ substantially from the value of $1/2$ can lead to quite different results. For example, the value of 1 for α can lead to very large values of $g_{ctr}(d/2)g_{co}(d/2)$ and a pronounced maximum. Such a maximum in this function was seen earlier in the MC simulation data of Bhuiyan *et al.*¹², which is smaller and has a quite different origin. In the present case, it is apparent from eq 8 that when $\alpha < 1$ that the initial slope of this function will be positive. Since eq 9 guarantees that this function tends to zero at large b , under these circumstances there must be a maximum in $g_{ctr}(d/2)g_{co}(d/2)$. However, since in the simulations this results from the initial slope of this function and the value of α , it is initially flat for $\alpha \approx 1$ or sloping downwards for $\alpha > 1$.

Of the four functions that are presented in the figures, the product function, $g_{ctr}(d/2)g_{co}(d/2)$, gives the best test of our contact value expression. The counterion contact value, $g_{ctr}(d/2)$ is the least sensitive. This is not surprising since the coion contact value, $g_{co}(d/2)$ is small at large b and, as a result, at large b $g_{ctr}(d/2)$ will resemble $g_{sum}(d/2)$, which is given by eq 3. This demonstrates the utility of the product function and eq 9.

We are aware that a convincing, rigorous justification of eq 9 is lacking. Even though it is empirical, its accuracy in reproducing *exact* machine simulated data, given an optimum α , is unmistakable. This coupled with the fact that the equations derived from it, viz., eqs 10-12 are all accurate further enhances arguments in favour of eq 9. From an experimental perspective, knowledge of the various contact properties can be useful, for example, in electrochemistry experiments that pertain to the electrode-electrolyte interface. It ought to be emphasized that obtaining an expression of the product function that goes to very small values at large b without passing through unphysical negative values has been a nontrivial exercise that has occupied our attention in varying degrees for several years. It is not correct to say that with an adjustable parameter one can fit any set of data points. The essential point of our paper is eq 9. The adjustment of the parameter α is secondary. With eq 9 we are able to fit a wide range of data with a single parameter that, as our results show, never varies appreciably from the value suggested by theory.¹⁸ Indeed, as is seen in our study, a large variation of α from $1/2$, for example, $\alpha = 1$, gives markedly poorer results.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

LBB acknowledges an institutional grant through Fondos Institucionales Para la Investigacion (FIPI), University of Puerto Rico, while WSA is thankful for a studentship from FIPI. DH acknowledges the support of the National Institutes of Health through Grant No 3R01GM076013-04S1.

Literature Cited

1. Gouy M. Sur la constitution de la charge electrique a la surface d'un electrolyte. *J Physique* 1910;9:457–468.
2. Chapman DL. A contribution to the theory of electrocapillarity. *Phil Mag* 1913;25:475–481.
3. Stern O. Zur theorie der electrolytischen doppelschicht. *Z Elektrochem* 1924;30:508–516.
4. Henderson D, Boda D. Insights from theory and simulation on the electrical double layer. *Phys Chem Chem Phys* 2009;11:3822–3830. [PubMed: 19440608]
5. Henderson D, Blum L. Some exact results and the application of the mean spherical approximation to charged hard spheres near a charged hard wall. *J Chem Phys* 1978;69:5441–5449.
6. Henderson D, Blum L, Lebowitz JL. An exact formulae for the contact value of the density profile of a system of charged hard spheres near a charged wall. *J Electroanal Chem* 1979;102:315–319.
7. Holovko M, Badiali JP, di Caprio D. Contact conditions for the charge in the theory of the electrical double layer. *J Chem Phys* 2005;123:234705-1–234705-5. [PubMed: 16392942]
8. Holovko M, Badiali JP, di Caprio D. New results from the contact theorem for the charge profile for symmetric electrolytes. *J Chem Phys* 2007;127:014106-1–014106-3. [PubMed: 17627336]
9. Holovko M, di-Caprio D. On the contact conditions for the charge profile in the theory of the electrical double layer for nonsymmetrical electrolytes. *J Chem Phys* 2008;128:174702-1–174702-5. [PubMed: 18465931]
10. Henderson D, Boda D. On a conjecture of Fawcett. *J Electroanal Chem* 2005;582:16–20.
11. Henderson D, Lamperski S, Outhwaite CW, Bhuiyan LB. A mean spherical approximation study of the capacitance of an electric double layer formed by a high density electrolyte. *Coll Czech Chem Comm.* in the press.
12. Bhuiyan LB, Outhwaite CW, Henderson D. Some simulation and modified Poisson-Boltzmann theory results for the contact values of an electrolyte near a charged electrode. *J Electroanal Chem* 2007;607:54–60.
13. Henderson D, Bhuiyan LB. Evidence for a second contact value formula for the electric double layer. *Molec Sim* 2005;33:953–957.
14. Bhuiyan LB, Outhwaite CW, Henderson D, Alawneh M. A further Monte Carlo and modified Poisson-Boltzmann analysis of two recent results in the electric double layer theory. *Bangladesh J Phys* 2007;4:93–102.
15. Henderson D, Bhuiyan LB. Contact value formulae for the density profiles of the electroc double layer. *Collect Czech Chem Comm* 2008;73:558–574.
16. Silvestre-Alcantara W, Bhuiyan LB, Outhwaite CW, Henderson D. A modified Poisson-Boltzmann study of the electrode-ion singlet distribution functions at contact in the planar electric double layer. *Collect Czech Chem Comm.* preprint.
17. Henderson D, Bhuiyan LB. Simple expressions for contact values of density profiles in a planar double layer. *J Chem Theory Comput* 2009;5:1985–1989.
18. Lou P, Lee J. Exact analytic result of the contact value for the density in a modified Poisson-Boltzmann theory of an electrical double layer. *J Chem Theory Comput* 2009;5:1079–1083.
19. Torrie GM, Valleau JP. Electric double layers. I. Monte Carlo study of a uniformly charged surface. *J Chem Phys* 1980;73:5807–5816.
20. Boda D, Henderson D, Chan KY. Monte Carlo study of the capacitance of the double layer in a model molten salt. *J Chem Phys* 1999;110:5346–5350.
21. Lamperski S, Bhuiyan LB. Counterion layering at high surface charge in an electric double layer. Effect of local concentration approximation. *J Electroanal Chem* 2003;540:79–87.

Figure 1.

The product $g_{ctr}(d/2)g_{co}(d/2)$, $g_{diff}(d/2, b)$, $g_{co}(d/2)$, and $g_{ctr}(d/2)$ as functions of b in a RPM planar double layer at the reduced temperature $T^* = 0.15$, and reduced electrolyte density $\rho^* = 0.3$. The symbols are the MC simulation data, while the solid, dash, and dash-dotted lines are the results from eq 9 with $\alpha = 0.6, 1$, and $1/2$, respectively.

Figure 2.

The product $g_+(d/2)g_-(d/2)$, $g_{diff}(d/2, b)$, $g_{co}(d/2)$, and $g_{ctr}(d/2)$ as functions of b in a RPM planar double layer at the reduced reduced temperature $T^* = 0.20$, and reduced electrolyte density $\rho^* = 0.0925$. The symbols are the MC simulation data, while the solid and the dashed lines are the results from eq 9 with $\alpha = 0.5$ and 1, respectively.

Figure 3.

The same as Figure 2 but at the reduced reduced temperature $T^* = 0.30$, and reduced electrolyte density $\rho^* = 0.00925$. The solid and the dashed lines are the results from eq. 9 with $\alpha = 0.335$ and 1, respectively.

Figure 4.

The same as Figure 2 but at the reduced reduced temperature $T^* = 0.40$, and reduced electrolyte density $\rho^* = 0.0925$. The solid and the dashed lines are the results from eq 9 with $\alpha = 0.4$ and 1, respectively.

Figure 5.

The same as Figure 2 but at the reduced temperature $T^* = 0.60$, and reduced electrolyte density $\rho^* = 0.00925$. The solid and the dashed lines are the results from eq 9 with $\alpha = 0.3$ and 1, respectively.