CORRESPONDENCE

Yeasts in the Gut: From Commensals to Infectious Agents

by PD Dr. rer. nat. habil. Jürgen Schulze, Dr. rer. nat. Ulrich Sonnenborn
in volume 51–52/2009

Immunostimulating Treatment

I can agree with Schulze, and Sonnenborn with regard to the allergenic potential of *Candida* and other yeast infections as depicted by Truss.

If non-evidence based reports of experience gathered in successful practices are still regarded as useful then I wish to point out a 80% cure rate in more than 2500 chronically allergic patients with mixed infection whose medical histories date back up to 30 years. However, for practical reasons I have stopped using the methods mentioned in the article since comparative studies were published that included 650 patients in whom serological diagnostic evaluation as well as diagnostic tests of stool specimens and results from electroacupuncture diagnosis after Voll (1) were compared. Because of a lack of long term effectiveness, I do not treat my patients with antibiotic or antimycotic agents but with immunostimulants; increasingly, I am using live lactobacilli and bifidus preparations (92). Removing glucose and saccharose from patients' diets for 4–11 weeks is part of the dual strategic approach. The effectiveness of this approach has become so well known among patients that my practice has to give out appointments up to 6 months in advance. This approach has also been used in 720 patients with rheumatic disorders, according to a recent study report (3).

In severely immunosuppressed patients, however, this approach will remain unsuccessful.

DOI: 10.3238/arztebl.2010.0368a

REFERENCES

1. Frick G: Behandlung von Mykosen mit Sanum-Mitteln, Mora-Therapie und UVB. Erfolgsbeurteilung mit dem OET nach Linke, Stuhlkontrollen und Antikörper-Verlauf. Sanum-Post 2001; 54: 24–7.
2. Frick G: Thema Darmgesundheit. Komplement integr Med 2008; 49: 12–5.
3. Frick G: Die andere Rheumatherapie. Limburg: Multiprint Verlags GmbH 2009.
4. Schulze J, Sonnenborn U: Yeasts in the gut: from commensals to infectious agents [Pilze im Darm – von kommensalen Untermietern zu Infektionserregern]. Dtsch Arztebl Int 2009; 106: 837–42.

Dr. med. habil. Gerhard Frick
Amtsstr. 11 B
14469 Potsdam, Germany

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Heroic Self Experiment

Perhaps it is of interests that the doyen of German mycology, Professor Dr Hans-Heinrich Rieth (1914–1994), showed as early as in the 1950s to 1960s in a heroic self experiment that *Candida albicans* solution, if imbibed by a healthy subject, is entirely innocuous. However, the scenario is different in patients who have intestinal damage or who are immunosuppressed; in such patients the so called thrush-sepsis cycle (1) may affect the mucosa saprogenically as well as hematogenically. This has been shown for generalized *Candida* mycosis in panmyelophthisis (2).

DOI: 10.3238/arztebl.2010.0368b

REFERENCES

1. Ruprecht KW: Zum Pathomechanismus der haematogenen *Candida*-Enteritis. Mykosen 1968; 11: 843–6.
2. Ruprecht KW: Generalisierte *Candida*-Mykose bei Panmyelophthise. Mykosen 1969; 12: 1–16.
3. Schulze J, Sonnenborn U: Yeasts in the gut: From commensals to infectious agents [Pilze im Darm – von kommensalen Untermietern zu Infektionserregern]. Dtsch Arztebl Int 2009; 106: 837–42.

Prof. (em.) Dr. med. Klaus W. Ruprecht
Steigackerweg 6
87637 Seeg, Germany

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Contaminated Food

I was rather surprised that yeasts in the gut are the subject of serious discussion in the medical literature.

As Professor Ottenjann (text book author, joint founding father of flexible endoscopy in Germany), with whom I exchanged letters and had discussions regarding possible intestinal yeast infections in the early 1980s, explained to me at the time: “Yeasts require squamous epithelium; this ends at the Z line and re-surfaces at the dentate line. Only in that location are you likely to find yeasts on the mucosa, with corresponding inflammatory reactions.”

Since I was doubtful he asked me to send him an image and a case report if I ever came upon such a finding on endoscopy of the stomach or gut.

In spite of undertaking more than 25 000 endoscopies, with 3–5 times the number of associated biopsies, I have remained unsuccessful. The only scenario where I found yeast colonization distal to the esophagus was in necrotically disintegrating tumors of the stomach and intestine (cylinder epithelium)—and in these cases, the yeast was the least of the patients' problems.

Only for completeness's sake, I wish to mention that orally ingested fungal spores are not digested, and any laboratory should be able to grow yeasts from a stool specimen. Spores are ubiquitous and constantly contaminate foodstuffs.

DOI: 10.3238/arztebl.2010.0368c

REFERENCES

1. Schulze J, Sonnenborn U: Yeasts in the gut: from commensals to infectious agents [Pilze im Darm – von kommensalen Untermietern zu Infektionserregern]. Dtsch Arztebl Int 2009; 106: 837–42.

Dr. med. Gerd Helmecke

Königstr. 4
53773 Hennef, Germany
Dok-Helmecke@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Eradication Is Impossible

“Yeasts in the gut” are a long running topic of interest—a review article in the *Deutsches Ärzteblatt* could form a solid foundation of information. However, the article by Schulze et al does not do justice to the topic. Why might this be so? There are few technical mistakes: data about glucan exposure are cited unilaterally, expression of pathogenicity factors does not enable yeasts to form biofilms. However, a closer look shows that it is suggestive phrases that may result in misunderstandings. Instead of “Candida colonization” the authors refer to “Candida load” [note: in the German original, not the English translation]. Irritable bowel syndrome is described as the indication with the closest—if unconfirmed—similarity to Candida colonization. Rather, it would be correct to state that thus far no clue exists for any such association (1)—and on which basis this is described as a “possible connection” is not clear. The conclusion, “The eradication of intestinal yeasts is advised only for certain clearly defined indications,” is not helpful. However, it still implies that “eradication” may be advised under certain (not precisely defined) circumstances and ignores the fact that this is confirmed not to be possible (1). No new knowledge can be gained from this. The authors’ remaining “open questions” are not articulated; the “additional facts” that have been “clarified” since 2004 remain obscure.

The authors’ activities for Ardeypharm GmbH are correctly reported as a conflict of interests—the committee at the Robert Koch-Institute cites one of the authors (US) as a “protagonist of the Candida hypothesis” (1). This in itself is not reprehensible—the question it raises is that of appropriate and functioning peer review. How could such an article—which is as medically unenlightening as it is highly suggestive—be published in a renowned journal such as *Deutsches Ärzteblatt*?

DOI: 10.3238/arztebl.2010.0369a

REFERENCES

1. Kommission „Methoden und Qualitätssicherung in der Umweltmedizin“: Pathogenetische Bedeutung der intestinalen Candidabesiedlung. Bundesgesundheitsbl Gesundheitsforsch Gesundheitsschutz 2004; 47: 587–600.
2. Schulze J, Sonnenborn U: Yeasts in the gut: from commensals to infectious agents [Pilze im Darm – von kommensalen Untermietern zu Infektionserregern]. Dtsch Arztebl Int 2009; 106: 837–42.

Prof. Dr. med. Oliver Kurzai

Septomics Research Centre
Friedrich-Schiller-Universität Jena
Leibniz-Institut für Naturstoff-Forschung und Infektionsbiologie –
Hans-Knöll-Institut
Beutenbergstr. 11 a
07745 Jena, Germany
oliver.kurzai@hki-jena.de

Conflict of interest statement

The author has received research grants from SIRS-Lab and travel expenses and honoraria from Abbott, Gilead, and Pfizer.

In Reply:

We thank all our readers (including those who communicated their feedback to the authors directly) for contributing to the discussion of our article (1), which was controversial and thus underlines the continuing interest in the topic.

Hans Rieth (1914–1994) vitally contributed to developments in medical mycology in the German speaking area, together with a small number of clinical mycologists in both parts of Germany, in the 1960s to 1980s, by conducting elegant microbiological and experimental clinical studies (including courageous self experiments). However, at this time, the so called “Omics” technologies of molecular biology were in their infancy, at best. Sometimes, the deducted practical conclusions were exaggerated—one reason why colleagues from other specialties, especially gastroenterologists, opposed the “mycophobic” tendencies that started to develop among patients. Rudolf Ottenjann (1925–2005), a gifted endoscopist, opposed these trends vehemently and declared that only the treatment of Candida induced thrush on the squamous epithelium, which could be (macro)endoscopically confirmed, was necessary (2). New, high resolution techniques, such as confocal endoscopy (3), may open up opportunities to detect microorganisms during endoscopy on the cylinder epithelium without requiring a biopsy (see image in our article).

Candida yeasts—which multiply only by means of budding and, in contrast to molds, do not form spores—are able to overcome the host’s defenses by a reversible change between yeast and hyphal stages. These changes of cell morphology are accompanied by a reduction in the immunogenic β -glucan content of the yeast cell wall (4). The phenotypic switching that is associated with biofilm formation heads in the same direction (immuno-evasion) (5). The ability to form biofilms in itself is not a marker of pathogenicity, but it

protects the yeasts from being attacked by the immune system.

The pathogenesis of irritable bowel syndrome remains unclear to this day. Current discussions are focused on changed reactions of the enteric nervous system, visceral sensitivity and mobility, and persistent underlying inflammatory conditions. Preclinical studies have shown the influence of intestinal microbes on the functions and reactions of the intestinal tract. This is especially true for the irritable bowel syndrome that develops subsequent to an infection (“postinfectious irritable bowel syndrome”) (6). The association between *Candida* and irritable bowel syndrome—not proved so far—has assumed relevance for *Candida* problems by way of the current discussion of the pathogenetic background of the irritable bowel syndrome.

Review articles relate facts; their purpose is not to relate pre-formed opinions, nor semantically unclear phrases. We therefore follow in this context the request to substitute “*Candida* load”—used once in the German manuscript—with the term “*Candida* colonization”—used more than 10 times in the German manuscript.

The situation is different for formulating conclusions and key messages—these require distilling the topic at hand into central key statements without any accompanying explanations. The authors may assume that the entire article has been read in order to understand these.

It was not planned in a review article (1) to comment on individual case reports and studies that were conducted in a manner not consistent with GCP (Good Clinical

Practice); as a result, many reported results, especially from the naturopathic field, were not included. It is therefore not possible to assess the entire multitude of therapeutic strategies, which would exceed the size of this authors’ reply.

DOI: 10.3238/arztebl.2010.0369b

REFERENCES

1. Schulze J, Sonnenborn U: Yeasts in the gut: from commensals to infectious agents [Pilze im Darm – von kommensalen Unternehmern zu Infektionserregern]. *Dtsch Arztebl Int* 2009; 106: 837–42.
2. Anonymus: Streitthema Darmsanierung – Kontra: Prof. Dr. med. Rudolf Ottenjann. *Naturamed* 1994; (12): 13–4.
3. Hoffmann A, Kiesslich R: Gastrointestinale Endoskopie – erweiterte Bildgebung. *Gastroenterologie up2date* 2009; 5: 145–63.
4. Netea MG, Brown GD, Kullberg BJ, Gow NAR: An integrated model of the recognition of *Candida albicans* by the innate immune system. *Nat Rev Immunol* 2008; 6: 67–78.
5. Ramirez-Zavala B, Reuß O, Park YN, Ohlsen K, Morschhäuser J: Environmental induction of white-opaque switching in *Candida albicans*. *PLoS Pathog* 2008; 4: e1000089.
6. McKeown ES, Parry SD, Stansfield R, Barton JR, Welfare MR: Postinfectious irritable bowel syndrome may occur after non-gastrointestinal and intestinal infection. *Neurogastroenterol Motil* 2006; 18: 839–43.

PD Dr. rer. nat. habil. Jürgen Schulze

Alice-Bloch-Str. 7
14558 Nuthetal, Germany
JuR.Schulze@t-online.de

Conflict of interest statement

Until 2005, Dr. Schulze was an employee of the company Ardeypharm GmbH. Since January 2006, he has not maintained any financial or non-material relations with the company.