Frog Virus 3 Replication: Induction and Intracellular Distribution of Polypeptides in Infected Cells

R. M. ELLIOTT† AND D. C. KELLY*

*Natural Environment Research Council, Unit of Invertebrate Virology, and Department of Forestry,
University of Oxford, Oxford, OX1 3UB, United Kingdom*

The synthesis of the polypeptides induced in frog virus 3-infected cells was analyzed by high-resolution sodium dodecyl sulfate-polyacrylamide gel electrophoresis of radiolabeled cell extracts. Purified frog virus 3 contained 22 polypeptides, with molecular weights in the range 9×10^3 to 114×10^3 . All of the structural and an additional seven nonstructural polypeptides were detected in infected cell lysates. The following three classes of induced polypeptides (under temporal control) were observed in BHK cells: at 2 h, four α polypeptides; at 4 h, 13 β polypeptides; and at 6 h, the remaining 12 γ polypeptides. The total molecular weight of the infected cell-specific polypeptides (ICPs) was $\approx 1.5 \times 10^6$, which accounts for about 30% of the coding capacity of the viral genome. At least 10 of the induced polypeptides were phosphorylated, but none was glycosylated or sulfated. No evidence for posttranslation cleavage of polypeptides in pulse-chase and inhibition experiments was obtained. The synthesis of γ polypeptides was not detected in the presence of the viral DNA replication inhibitors cytosine arabinoside and hydroxyurea, but halogenated nucleotides apparently had no effect. These results suggest that α and β polypeptides are "early" events and that detectable γ polypeptide synthesis is dependent on the production of progeny viral DNA. The regulation of frog virus 3-induced polypeptide synthesis in infected BHK cells was examined by using inhibitors of protein and RNA synthesis and amino acid analogs. These experiments confirmed the existence of three sequentially synthesized, coordinately regulated classes of polypeptides, designated α , β , and γ . The requirements for the synthesis of each class were as follows: (i) α polypeptides did not require previous cell protein synthesis; (ii) β polypeptides required a prescribed period of α polypeptide synthesis and new mRNA synthesis; and (iii) γ polypeptides required prior synthesis of functional β polypeptides and new mRNA synthesis. α polypeptide synthesis was controlled by β and γ polypeptides, and α and β polypeptides were involved in the suppression of host cell polypeptide synthesis. Indirect evidence was obtained for the temporal regulation of frog virus 3 transcription. The intracellular distribution of virus-induced polypeptides in cells infected with frog virus 3 was investigated by using standard cell fractionation techniques. Most of the 29 induced polypeptides were bound to structures within the nucleus, and only two ICPs were not associated with purified nuclei. When isolated nuclei were incubated in an infected cell cytoplasm preparation, all of the nuclear ICPs were incorporated in vitro. All of the ICPs were associated with ribosomal and rough endoplasmic reticulum fractions of infected cells, and a number of ICPs were found on smooth intracellular membranes. Most of the ICPs were also associated with purified plasma membranes of infected cells, and one polypeptide (ICP 58) was highly enriched in the plasma membrane compared with whole cell extracts or purified frog virus 3.

Icosahedral cytoplasmic deoxyriboviruses are a group of viruses isolated from a diverse range of hosts, and this group includes iridescent viruses of insects, lymphocystis viruses of fish,

African swine fever viruses and, amphibian polyhedral viruses (44, 55). They are characterized as having an icosahedral virion greater than 100 nm in diameter and a linear double-stranded DNA genome, and their replication is apparently restricted to the cytoplasm. The most extensively studied icosahedral cytoplasmic deoxyri-

† Present address: Department of Microbiology, Mount Sinai School of Medicine, City University of New York, New York, NY 10029.

bovirus is frog virus 3 (FV3), which was isolated from the leopard frog, *Rana pipiens*, and has subsequently been shown to replicate in cells of amphibian, piscine, avian, and mammalian origin (30). The optimum temperature for viral replication is about 28°C, and replication is inhibited at 33°C, although several viral functions are expressed at the nonpermissive temperature (49).

The virion of FV3 is composed of at least 17 polypeptides (75), and a variety of enzyme activities have been associated with purified virus particles, including nucleases (39), nucleotide phosphohydrolase (78), protein phosphatase (69), protein kinase (69–72) and DNA-dependent RNA polymerase (24). The genome of FV3 is a double-stranded RNA having a molecular weight of 100×10^6 (43).

The synthesis of polypeptides in FV3-infected cells has been studied by Granoff and colleagues, who reported that 20 virus-specified polypeptides were synthesized within 2 h of infection and that the induction of these polypeptides was independent of viral DNA replication (27). However, these investigations were performed by using the technique of slicing and counting polyacrylamide tube gels, which has a low power of resolution (36). More recently, these workers claim to have resolved 35 polypeptides on polyacrylamide gradient slab gels (82).

We have made a detailed study of the induction, regulation, and distribution of the polypeptides synthesized in FV3-infected cells by using high-resolution polyacrylamide gel electrophoresis (PAGE). This paper describes the identification of virus-specified polypeptides in infected cells, the minimal processing of the induced polypeptides by cleavage and glycosylation, the phosphorylation of FV3 polypeptides, and the dependence of certain polypeptides on viral DNA replication.

FV3 has a number of similarities with both herpesviruses and poxviruses, which led McAuslan and Armentrout (54) to postulate that FV3 may be a "missing link" between these groups. The size and morphology of the virion and size of the genome are herpesvirus-like, whereas the cytoplasmic site of replication and a number of virion-associated enzyme activities are poxvirus-like (24, 44, 54). The existence of a postreplicative class of induced polypeptides is also a shared feature with poxviruses (56; see below). The polypeptides induced in herpes simplex virus-infected cells form at least three sequentially synthesized, coordinately regulated groups, designated α , β , and γ . The main features of this "cascade" regulation are as follows: (i) synthesis of α polypeptides does not require previous infected cell protein synthesis; (ii) the

initiation of β polypeptide synthesis requires the synthesis of functional α polypeptides as well as new RNA synthesis; and (iii) γ polypeptide synthesis has a similar requirement for one or more β polypeptides. Inhibition of viral DNA synthesis does not prevent the synthesis of α , β , or γ polypeptides, but does reduce the amounts of γ polypeptides produced (37). Since there was a similarity in the sequential induction of polypeptides in FV3- and herpesvirus-infected cells, we designed experiments to determine whether similar regulatory mechanisms existed. Specifically, we wished to answer the following questions. (i) Was the synthesis of one class of polypeptides dependent on the synthesis of another? (ii) Was new mRNA synthesis required for each class of polypeptide? And (iii), did the synthesis of a later class of polypeptides exert a controlling influence over an earlier polypeptide class? We employed inhibitors of protein and RNA syntheses and amino acid analogs added at various times during the viral cycle to answer these questions and to demonstrate a cascade regulation of FV3-induced polypeptide synthesis. Other inhibitors of RNA synthesis were used to examine further the expression of the viral genome.

There is considerable evidence for the interaction of FV3 with the nucleus and with various membranes in infected cells. Electron microscope observations of FV3-infected cells have shown that the virus apparently replicates in the cytoplasm (6, 15, 41), although late in infection mature virus particles have been found in some nuclei (the number of such nuclei varied with the cell line) and the nuclear membrane was apparently intact (42). Nuclear involvement of FV3 is also implicated from other data; the virus selectively inhibits RNA polymerase II (nuclear RNA polymerase) (10, 13), and it is claimed that FV3 prevents the replication of some nuclear RNA viruses (4). The presence of fibrillar bodies in hepatocyte nuclei of mice during the early stages of the toxic degenerative hepatitis produced by FV3 has also been observed (5). The requirement of a functional cell nucleus for FV3 replication to occur has also been demonstrated (29), and a fraction of FV3 DNA is synthesized in the nucleus (28, 55). The virus may leave the cell by budding through the plasma membrane, and modifications of the membrane occur at localized areas (16, 42, 77). The accumulation of virus-specific antigens at the plasma membrane has been detected by immunological and cytochemical methods involving electron microscopy (76).

The association of infected cell-specific polypeptides (ICPs) with the nucleus, the plasma membrane, or the cytoplasmic membranes in

FV3-infected cells has not been investigated. We extended our studies of FV3-induced polypeptide synthesis to examine the intracellular distribution of polypeptides by using standard cell fractionation techniques. We report here the association of virus-specific polypeptides within the nucleus and with plasma and cytoplasmic membranes.

MATERIALS AND METHODS

Virus and cells. A clonal isolate of FV3 was kindly supplied by A. Granoff, St. Jude Children's Research Hospital, Memphis, Tenn. Procedures for the maintenance of baby hamster kidney (BHK 21) and fathead minnow (FHM) cells in Glasgow minimal essential medium and for the titration of FV3 have been described elsewhere (16).

Chemicals and isotopes. Reagents for PAGE, fluorescein mercuric acetate, and sodium deoxycholate were purchased from British Drug Houses Ltd., Poole, England. Cytosine arabinoside (ara-C) was obtained from Koch Light, Colnbrook, England. Tween 40, hydroxyurea, 5'-bromodeoxyuridine, 5'-fluorodeoxyuridine, *N*- α -tosyl-L-lysine chloromethyl ketone (TLCK), L-1-tosylamide-2-phenyl-ethylchloromethyl ketone (TPCK), actinomycin D, L-azetidine 2-carboxylic acid, L-canavanine, cordycepin (3'-deoxyadenosine), and DL-*p*-fluorophenylalanine (FPA) were obtained from Sigma London Chemical Co. Ltd., London, England. Cycloheximide (Calbiochem) was a gift from M. L. Fenwick, Sir William Dunn School of Pathology, University of Oxford. Camptothecin was a gift from the Drug Development Branch, National Cancer Institute, Bethesda, Md. Rifampicin was obtained from Mann Research Laboratories, Becton, Dickinson & Co. Ltd., London, England.

The following radioisotopes were supplied by the Radiochemical Centre, Amersham, England: L-[³⁵S]methionine (840 to 1,120 Ci/mmol); [³⁵S]sulfate (carrier-free); D-[1-¹⁴C]glucosamine hydrochloride (60 mCi/mmol); U-¹⁴C-protein hydrolysate (56 mCi/milatom of carbon); and ³²P_i (30 to 100 Ci/mg).

Growth, isotopic labeling, and purification of FV3. FHM cells were grown to confluent monolayers in 32-ounce (960-ml) glass bottles (Flow Laboratories Ltd., Irvine, Scotland) and infected with 0.1 to 1 PFU/cell. After adsorption for 1 h at room temperature, growth medium was added and incubation was continued at 28°C.

To grow [³⁵S]methionine-labeled virus, the medium was replaced at 3 h postinfection with 5 ml of Earle balanced salt solution (EBSS) containing 2% dialyzed fetal calf serum and 40 μ Ci of [³⁵S]methionine per ml, and incubation was continued for 2 h at 28°C; growth medium was then added to bring the final concentration of isotope to 4 μ Ci/ml.

After 72 h of incubation, infected cells and released viruses were collected by centrifugation (2 h at 50,000 $\times g$), and the pellet was resuspended in reticulocyte standard buffer (RSB; 10 mM Tris-hydrochloride, pH 7.2, 10 mM KCl, 1.5 mM MgCl₂). Cells were swollen for at least 1 h at 4°C and then disrupted with about 25 strokes of a Dounce homogenizer. Cell debris was

removed by centrifugation (1,000 $\times g$ for 10 min), and the virus in the supernatant was pelleted at 95,000 $\times g$ for 30 min. The pellet was resuspended in 0.01 M Tris-hydrochloride buffer (pH 8.0), and FV3 was purified by rate gradient centrifugation on a 25 to 50% (wt/vol) sucrose gradient (1 h at 42,000 $\times g$), followed by equilibrium centrifugation for 16 h at 42,000 $\times g$ in a 40 to 80% (wt/vol) sucrose gradient (all sucrose solutions were prepared in the Tris-hydrochloride buffer). Purified virus was pelleted at 95,000 $\times g$ for 30 min and stored in Tris-hydrochloride buffer at a concentration of 1 to 2 mg/ml at -20°C.

Isotopic labeling of cells for analysis of polypeptides. Cells were grown in 30-mm Nunc plastic petri dishes (GIBCO BioCult Ltd., Paisley, Scotland) containing 3×10^5 cells and were infected at 25 to 30 PFU/cell for 1 h at room temperature. The monolayers were then washed with medium (at zero time postinfection), fresh medium was added, and incubation was continued at 28°C. At various times after infection, the medium was removed and replaced with 1 ml of labeling medium containing 2% dialyzed fetal calf serum and one of the following radioactive precursors: [³⁵S]methionine (20 μ Ci/ml) in EBSS; [¹⁴C]-labeled amino acids (10 μ Ci/ml) in EBSS; [¹⁴C]glucosamine (8 μ Ci/ml) in EBSS-glucose; ³²P_i (100 μ Ci/ml) in EBSS with 25% phosphate; or [³⁵S]sulfate (100 μ Ci/ml) in EBSS with Mg(SO₄)₂ replaced by MgCl₂. To ensure efficient incorporation of ¹⁴C-labeled compounds, the cells were pretreated for 15 min with labeling medium without isotope.

After labeling for 1 h at 28°C, the label was removed, and the cells were washed with cold phosphate-buffered saline (PBS) and scraped into 200 μ l of sample buffer (1 M Tris-hydrochloride, pH 6.8, containing 10% [wt/vol] sodium dodecyl sulfate, 10% [vol/vol] 2-mercaptoethanol, 15% [vol/vol] glycerol, and 0.002% bromophenol blue). The samples were boiled for 2 min and stored at room temperature.

PAGE. Analysis of polypeptides was performed on 16-cm sodium dodecyl sulfate-polyacrylamide slab gels as previously described (19, 52). Unless otherwise stated, 12% gels with a ratio of acrylamide to bisacrylamide of 75:1 were used. The samples were boiled again immediately before electrophoresis; usually 40- μ l portions were loaded onto the gels, and electrophoresis was performed for 16 h at 60 to 70 V.

After electrophoresis, gels were fixed in 45% methanol-5% acetic acid for 1 h and either stained with Coomassie brilliant blue (19) and dried directly (³²P-labeled samples) or prepared for fluorography (7) before drying. Dried gels were exposed to Fuji RX medical X-ray film at -70°C. Molecular weights of Coomassie brilliant blue-stained polypeptides were calculated with respect to standard markers electrophoresed on the same gel, as previously described (19).

Use of inhibitors. Stock solutions of inhibitors were made in growth medium (minimal essential medium) at concentrations of 1 mg/ml. Cycloheximide at 50 μ g/ml was used to inhibit protein synthesis. The amino acid analogs azetidine, canavanine, and FPA were used at 500 μ g/ml. Cordycepin was used at 50 μ g/ml, and camptothecin was used at 100 μ g/ml. Actinomycin D was used at 10 μ g/ml.

After treatment with the protein synthesis inhibitors and amino acid analogs, the drug-containing medium was removed, and the cell monolayers were rinsed five times with 2-ml portions of minimal essential medium maintained at 28°C. This procedure was accomplished within 5 min and had no deleterious effect on the cells. After the monolayers were rinsed, fresh minimal essential medium was added and incubation was continued at 28°C.

Cell fractionation procedures. (i) Nucleus-cytoplasm fractionation. BHK cells were fractionated by using methods based on those described by Penman (62) and Fenwick et al. (22). Cells (10^6) were grown in 25-cm² flasks and infected for 1 h with 25 to 30 PFU of FV3. At various times after infection, the cells were labeled for 1 h with 100 μ Ci of [³⁵S]methionine in 2 ml of EBSS containing 2% dialyzed fetal bovine serum. After labeling, the cells were washed with PBS and scraped into 1 ml of RSB. A 50- μ l portion was removed and dissociated with 50 μ l of PAGE sample buffer for use as a whole cell extract. The cells were swollen for 15 to 20 min on ice and homogenized with 20 strokes of a tight-fitting Dounce homogenizer, and 0.2 volume of 60% (wt/vol) sucrose in RSB was added. Nuclei were pelleted by centrifugation for 5 min at $2,000 \times g$. The supernatant was removed, the nuclei were resuspended in 1 ml of 10% (wt/vol) sucrose in RSB, and 150 μ l of detergent mix (2 parts of 10% Tween 40 plus 1 part of 10% sodium deoxycholate) (32) was added. The nuclei were shaken on a Vortex mixer for 4 s and then pelleted by centrifugation. The supernatant was collected and added to the first supernatant; this represented the cytoplasmic fraction. The proteins were precipitated by adding 5 volumes of acetone and leaving at -20°C overnight. The precipitate was collected by centrifugation for 2 h at $1,500 \times g$. The pellet was air dried and resuspended in 100 μ l of PAGE sample buffer. The pellet of purified nuclei was also resuspended in 100 μ l of PAGE sample buffer.

(ii) Preparation of cytoplasmic membranes. Cytoplasmic membranes from BHK cells were prepared by using a microscale version of the technique described by Caligiuri and Tamm (9). Cells (8×10^6) were grown in 175-cm² plastic flasks and either infected with 25 to 30 PFU of FV3 per cell for 1 h or left uninfected as a control. The cells were labeled with 300 μ Ci of [³⁵S]methionine in 10 ml of EBSS. After labeling, the cells were washed twice with cold PBS, scraped into 10 ml of PBS, and pelleted at $1,000 \times g$ for 10 min. The cells were resuspended in 3 ml of RSB, swollen on ice for 20 min, and homogenized with about 30 strokes in an all glass Dounce homogenizer with a tight pestle. Nuclei and cell debris were removed by centrifugation ($900 \times g$ for 10 min), and the membrane-containing supernatant was made up to 30% in sucrose by adding an equal volume of 60% (wt/vol) sucrose in RSB. The membrane suspension was included in a discontinuous sucrose gradient made in a 25-ml centrifuge tube and composed of the following sucrose-RSB solutions (weight/weight): 1.5 ml of 60% sucrose, 3.5 ml of 45% sucrose, 3.5 ml of 40% sucrose, 5 ml of 30% sucrose containing the cytoplasmic extract, 3.5 ml of 25% sucrose, and 1.5 ml of RSB. The gradient was centrifuged at $86,000 \times g$ for 16 h. Fractions were

collected as described by Caligiuri and Tamm (9) and Compans (11), diluted in RSB, and pelleted for 2 h at $100,000 \times g$. The fractions were dissociated in 200 μ l of PAGE sample buffer.

(iii) Preparation of plasma membranes. The preparation of BHK plasma membranes was based on methodology described by Warren and Glick (80) and employed fluorescein mercuric acetate to stabilize the membranes. Cells (3×10^6) were grown in 80-cm² plastic flasks and either infected for 1 h with 25 to 30 PFU of FV3 or left uninfected. Cells were labeled with 100 μ Ci of [³²S]methionine in 6 ml of EBSS. After labeling, the cells were washed twice with PBS and removed from the flask by incubation for 5 min at room temperature in 3 ml of 3 mM EDTA. The cells were pelleted at $1,000 \times g$ for 5 min and washed twice with 0.16 M NaCl, and the final cell pellet was resuspended in 0.5 ml of 0.16 M NaCl. The suspension was transferred to a 3.0-ml all glass Dounce homogenizer with a tight pestle. A 1.5-ml amount of saturated fluorescein mercuric acetate (0.0022 M fluorescein mercuric acetate in 0.02 M Tris-hydrochloride, pH 8.1, freshly made and filtered before use) was added, and the suspension was incubated for 15 min at room temperature, followed by 15 min on ice. The cells were disrupted with about 25 slow strokes of the pestle, and the homogenization procedure was monitored by phase-contrast microscopy. A 2-ml amount of 60% (wt/wt) sucrose in water was added, and the mixture was layered onto 10 ml of 45% (wt/wt) sucrose in a 50-ml centrifuge tube. The tube was centrifuged for 1 h at $150 \times g$ and 4°C, and the upper phase was removed. The meniscus region was collected separately, diluted with 0.1 volume of water, and centrifuged on 10 ml of 45% (wt/wt) sucrose for 30 min at $850 \times g$. The upper phase was collected, combined with that of the first centrifugation, diluted with 0.1 volume of water, and centrifuged through 10 ml of 35% (wt/wt) sucrose for 1 h at $1,800 \times g$. The pellet of purified plasma membranes was dissociated with PAGE sample buffer, and the proteins were analyzed by PAGE.

Electron microscopy. Electron microscopy was performed as previously described (16, 41, 42).

RESULTS

Structural polypeptides of FV3. Figure 1 shows the polypeptides of purified FV3 analyzed on a 12% sodium dodecyl sulfate-polyacrylamide gel compared with standard molecular weight markers. A total of 22 structural polypeptides were resolved, with molecular weights ranging from 9,000 to 114,000; the major structural polypeptide had a molecular weight of 55,000. The total molecular weight of the structural polypeptides was 1.196×10^6 (Table 1).

The polypeptide profile of purified [³⁵S]methionine-labeled FV3 was similar to that of Coomassie brilliant blue-stained virus (Fig. 1b).

Identification of ICPs. A comparison of [³⁵S]methionine-labeled FV3-infected BHK cell extracts and purified labeled FV3 is shown in Fig. 1b. We were able to resolve 29 polypeptides

FIG. 1. Analysis of FV3 polypeptides on a 12% polyacrylamide slab gel. (a) Purified FV3 compared with standard molecular weight markers stained with Coomassie brilliant blue. The markers were: a, phosphorylase A (molecular weight, 108,000); b, transferrin (90,000); c, bovine serum albumin (69,000); d, ovalbumin (45,000); e, lactic dehydrogenase (35,000); f, carbonic anhydrase (31,000); g, α -chymotrypsinogen (25,000); h, myoglobin (16,900); i, cytochrome c (12,400). VP = virus structural polypeptide molecular weight ($\times 10^{-3}$). (b) Comparison of the polypeptides synthesized in uninfected and FV3-infected BHK cells and the polypeptides in purified FV3. In this and other figures, the numbers above each track indicate the time (hours p.i.) of the start of a 1-h radioactive pulse. U, Uninfected cells; VP, virus structural polypeptide; NS, nonstructural polypeptide. ICPs are identified by their molecular weights ($\times 10^{-3}$).

in the infected cells by 9 h postinfection (p.i.), which either had no counterparts in the uninfected cells or which increased in intensity after infection, and we considered these to be ICPs. All 22 structural polypeptides were detected on the basis that their electrophoretic mobilities were similar to those of the polypeptides of purified labeled FV3 electrophoresed in an adjacent well on the same slab gel. An additional seven polypeptides not present in purified FV3 were observed, and these have been classed as nonstructural ICPs (Table 1).

The total molecular weight of the ICPs is 1.469×10^6 , which is considerably less than the maximum theoretical coding capacity for a genome of 100×10^6 daltons (43). To determine whether more polypeptides could be resolved on other one-dimensional PAGE systems, we analyzed the infected cell extracts on a 6% acryl-

amide gel, a 15 to 25% acrylamide linear gradient gel, a 12% acrylamide gel cross-linked with *N,N'*-diallyltartardiamide (26), and 7 to 20% acrylamide-*N,N'*-diallyltartardiamide gradient gels (66). We were unable to resolve any more polypeptides above 9,000 molecular weight in the cell extracts with these different systems, but a polypeptide of 7,500 molecular weight, which runs with the front on our standard gel systems, was resolved on the 15 to 25% gradient gel. The higher resolving power of two-dimensional gel electrophoresis (60) to analyze FV3-induced polypeptides is considered in an accompanying paper (18).

Suitability of [35 S]methionine as the radioactive precursor. To ensure that [35 S]methionine was a suitable precursor to label all of the ICPs, we compared cell extracts that had been labeled with either [35 S]methionine or a

TABLE 1. *Catalog of ICPs in FV3-infected cells detected by one-dimensional gel electrophoresis*

ICP ^a	Classification of ICP ^b	Class ^c		Labeled with ³² PO ₄
114	VP	γ	]	+
104	VP	γ		+
98	VP	β		
93	VP	β		
88	NS	β		
78	VP	γ		
76	VP	γ		
69	VP	β		+
66	NS	β		
61	VP	α		+
58	VP	β		
55	VP	γ		+
52	VP	β		
50	VP	β		
46	VP	β		
44	VP	α		+
40	NS	γ		
37	VP	γ		
35	VP	γ		+
34	VP	β		
29	VP	β	]	+
27	VP	γ		+
23	NS	β		+
22	NS	γ	]	+
18	NS	γ		+
16	NS	α		+
15	VP	α	]	+
12	VP	γ		+
9	VP	β		

^a ICPs are identified by their molecular weights ($\times 10^3$).

^b VP indicates a counterpart of similar electrophoretic (virion polypeptide) mobility in purified FV3; NS (nonstructural) indicates no counterpart in FV3.

^c Class of ICP is based on time of appearance and dependence on replication of viral DNA.

¹⁴C-labeled amino acid mixture. The resulting fluorogram showed that no additional bands were detected by labeling with the amino acid mixture (data not shown). Also, when the intensities of the bands on the fluorogram were compared, there were no relatively methionine-rich or methionine-deficient polypeptides. Thus, [³⁵S]methionine was considered to be the precursor of choice.

Sequential appearance of polypeptides in FV3-infected BHK cells. From the experiment shown in Fig. 1b, it was obvious that not all of the ICPs were synthesized at the same time. To investigate this aspect further, a series of infected BHK monolayers were labeled at hourly intervals, and the cell extracts were analyzed by PAGE. There appeared to be three phases of polypeptide induction temporally spaced, which we have designated α , β , and γ . The α polypep-

tides were first detected at 2 h p.i. (e.g., ICP 44 and ICP 15); β polypeptides were first seen at 4 h p.i. (e.g., ICP 98 and ICP 34); and γ polypeptides were first detected at 6 h (e.g., ICP 55 and ICP 12). A complete catalog of ICPs is given in Table 1. All 29 ICPs were being synthesized by 6 h, and their polypeptide profiles remained constant to 21 h p.i. The exact time of induction of each phase of polypeptide synthesis was found to vary (± 1 h) from one experiment to another, but the polypeptides appearing within each phase were always consistent.

Effect of temperature on FV3-induced polypeptides. To study the effect of temperature on FV3-induced polypeptide synthesis, BHK monolayers were infected at room temperature for 1 h with FV3, the monolayers were washed, and incubation was continued at one of the following temperatures: 21°C (suboptimum), 28°C (optimum), 34°C (the cut-off temperature for virus replication [49]), and 37°C. The cells were labeled at various times after infection and analyzed by PAGE (Fig. 2). The synthesis of FV3-induced polypeptides at 21°C was considerably slower than that at 28°C, but by 24 h p.i. all ICPs were being synthesized. At the nonpermissive temperatures (34 and 37°C) only the synthesis of α polypeptides (e.g., ICPs 44, 16, and 15) was detected, and the suppression of host cell polypeptides was less effective than at 28°C.

Effect of heat-inactivated FV3 on induced polypeptides. Heat-inactivated FV3 (56°C, 30 min) (Δ FV3) has been reported to shut off host cell DNA, RNA, and polypeptide syntheses, while allowing subsequent infection with active FV3 to proceed apparently normally (27). We heat inactivated (56°C, 30 min) a purified preparation of FV3 suspended in PBS and observed that heat treatment reduced the titer of the virus by 10^5 -fold or more. Monolayers of BHK cells were treated with Δ FV3 for 1 h at room temperature at an equivalent of 50 PFU/cell, and the monolayers were washed and then incubated at 28°C for 3 h; this procedure is similar to that described by Goorha and Granoff (27). Control monolayers were treated with PBS. The cultures were infected with active FV3 as usual, and polypeptide synthesis was examined at various times after infection (Fig. 3). Prior treatment with Δ FV3 had no effect on the induction of polypeptides, which had a temporal appearance similar to that in the PBS-treated samples. No induced polypeptides were observed in the culture treated with Δ FV3 alone. However, the use of Δ FV3 did reduce the overall incorporation of radioactive precursor into the cells (Fig. 3), but Δ FV3 had little effect on the rate of disappear-

FIG. 2. Effect of incubation temperature on FV3-induced polypeptides in infected BHK cells. Cells were infected for 1 h at room temperature, and then incubation was continued at 21, 28, 34, or 37°C. All ICPs were synthesized at 21°C but at slower rates than at 28°C. Only the four α ICPs were synthesized at the nonpermissive temperatures (34 and 37°C).

FIG. 3. Effect of heat-inactivated Δ FV3 on the polypeptides synthesized in BHK cells. The cells were treated with 50 PFU equivalents of Δ FV3 (+ Δ FV3) per cell or with PBS (+PBS) for 1 h at room temperature. The cells were incubated for 3 h at 28°C after removal of Δ FV3 or PBS and then infected with active FV3. The cells were pulse-labeled at various times after infection. Prior treatment with Δ FV3 did not affect the sequential induction of FV3 polypeptides, but did reduce the synthesis of host polypeptides in uninfected cells.

ance of host polypeptides compared with the PBS-treated cells.

Processing of FV3-induced polypeptides.

(i) **Glycosylation.** We were unable to demonstrate any glycosylation of ICPs when [^{14}C]glucosamine was used as the precursor in medium devoid of glucose.

(ii) **Sulfation.** No evidence for sulfation of FV3-induced polypeptides was obtained by labeling infected cells with $^{35}\text{SO}_4$, although sulfation of a number of host cell proteins was observed.

(iii) **Phosphorylation.** Figure 4 shows a comparison of FV3-infected BHK cell lysates which had been labeled with either [^{35}S]methionine or $^{32}\text{P}_i$. At least 10 ICPs were labeled with $^{32}\text{PO}_4$, but because phosphate-labeled polypeptides produced more diffuse bands on the autoradiogram, some bands may represent two or three polypeptides, as indicated on Fig. 4. The major phosphorylated polypeptides were ICPs 114/104, 44, 35, and 12; ICPs 15/16 (α polypeptides) were phosphorylated some time after they were first detected by [^{35}S]methionine labeling. The major structural polypeptide, ICP 55, was only slightly phosphorylated.

To confirm that the bands detected on the

autoradiogram represented polypeptides and not RNA molecules, replicate cell extracts were treated with either RNase or protease (66) before dissociation and electrophoresis. RNase treatment had no effect on the polypeptide profiles, but no bands were detected by PAGE after protease treatment. This indicated that the bands observed were proteins.

(iv) **Cleavage.** We examined the possibility of both rapid posttranslational cleavage and slow cleavage of FV3-induced polypeptides. The use of inhibitors of trypsin and chymotrypsin (TLCK and TPCK, respectively) prevents the cleavage of the primary transcription product of poliovirus (48, 74) and these compounds have been used to examine cleavage in herpes simplex virus-infected cells (36). We labeled FV3-infected BHK cells in the presence of either 10^{-4} M TPCK or 10^{-4} M TLCK at various times after infection, but we could not detect any high-molecular-weight polypeptides that are not found under usual labeling conditions.

Pulse-chase experiments were performed to determine whether any gradual degradation of induced polypeptides occurred. A series of infected BHK monolayers were labeled for 1 h at either 3 or 7 h p.i., and then the medium was

FIG. 4. Phosphorylation of FV3-induced polypeptides. Infected BHK cells were labeled with either [^{35}S]methionine (^{35}S met) or $^{32}\text{P}_i$ for 1 h at various times after infection. Certain phosphorylated ICPs were not resolved cleanly and are bracketed together.

replaced with minimal essential medium containing unlabeled methionine for chase periods of up to 3 h. Again no evidence for cleavage was observed, because the polypeptide profiles of the pulsed and pulse-chased samples were similar. The only polypeptides that declined in synthesis during the chase period were of host origin.

Dependence of γ polypeptide synthesis on viral DNA replication. FV3 DNA replication occurs around 5 to 8 h p.i. in BHK cells, as determined by [3 H]thymidine incorporation or CsCl density gradient centrifugation of labeled cell lysates (17; unpublished data). Since the onset of γ polypeptide synthesis occurred at the same time as viral DNA replication, we examined whether this class of polypeptides was dependent on the production of progeny DNA.

Figure 5 shows the effect of 30 μ g of ara-C per ml (added after adsorption of the virus and maintained through the experiment) on FV3-induced polypeptides. The drug had no effect on the production of α polypeptides or on the onset of β polypeptide synthesis, with the exception of ICP 69 (whose synthesis declined in the presence of ara-C) and ICP 46 (which was not synthesized). The production of γ polypeptides, e.g. ICPs 76, 55, and 12, was not detected in the presence of ara-C. The suppression of host cell polypeptides was unaffected by the drug, although there was an overall diminution in the uptake of [35 S]methionine. The fact that there was a class of FV3-induced polypeptides which were dependent on viral DNA replication was contrary to published data (27). Therefore, we investigated the effect of ara-C on FV3-induced polypeptides in FHM cells and obtained a similar result, namely that the synthesis of γ ICPs was not detected.

The effects of other inhibitors of DNA replication were also examined. Hydroxyurea at a concentration of 50 μ g/ml inhibited the production of γ polypeptides in a manner similar to ara-C (Fig. 6a), but later in infection (24 h p.i.) some γ polypeptide synthesis occurred (e.g., ICPs 55 and 12). Reversal of the block effect of hydroxyurea by extensive washing (five times with warm medium) resulted in the initiation of detectable γ polypeptide synthesis within 2 h at a normal rate (Fig. 6b). The polypeptide patterns of 24-h untreated infected cells and infected cells labeled 18 h after reversal of hydroxyurea were similar, although the synthesis of ICPs 66 and 37 were depressed in the drug-treated samples.

5'-Fluorodeoxyuridine (50 μ g/ml) and 5'-bromodeoxyuridine (at concentrations up to 500 μ g/ml) had no effect on the synthesis of induced polypeptides.

FIG. 5. Effect of ara-C on FV3-induced polypeptides. Infected BHK cells were incubated in the presence of 30 μ g of ara-C per ml or left untreated. Cells treated with ara-C were labeled in the presence of the drug (+ara-C). The syntheses of α and β ICPs were unaffected by ara-C, but no γ ICPs were detected in the presence of the drug.

Effect of inhibition of protein synthesis in FV3-infected cells on subsequent synthesis of ICPs. The experiments described here and below were designed to determine whether the synthesis of one class of ICPs was dependent on the prior synthesis of one or more ICPs. We performed experiments in which protein synthesis was inhibited by the addition of cycloheximide (50 μ g/ml) at the end of the adsorption period. Infected cells were incubated for various times in the presence of the drug, and the cells were washed to remove the drug and pulse-labeled at various times thereafter.

Figure 7 shows the effect of prolonged treatment with cycloheximide on polypeptide synthesis in infected cells. Cultures were incubated with the inhibitor for 5 or 9 h and then washed and pulse-labeled. The polypeptides synthesized immediately after removal of cycloheximide at 5 h p.i. were only a subset of the ICPs detected at 5 h in untreated cells. These polypeptides ICPs 61, 44, 16, and 15) were the only ICPs detected on removal of cycloheximide at 9 h and corresponded to the α polypeptides. The syntheses of ICPs 61 and 44 were amplified upon removal of the inhibitor. Three other polypep-

FIG. 6. Effect of hydroxyurea on FV3-induced polypeptides. (a) Infected cells were incubated and labeled in the presence of 50 μ g of hydroxyurea per ml or were left untreated. The synthesis of γ ICPs was inhibited by hydroxyurea treatment, but some leak through of γ ICPs was detected at 24 h p.i. (b) Infected BHK cells were maintained in the presence of 50 μ g of hydroxyurea per ml for 6 h and then washed five times with medium to reverse the action of the drug. Cells were labeled immediately or at various times after reversal, and the polypeptides were analyzed by PAGE. An untreated, 24-h p.i. cell extract was co-electrophoresed in an adjacent track. γ ICPs were detected 2 h after reversal of the drug. The 24-h p.i. polypeptide profiles of treated and untreated cell extracts were similar, but the syntheses of ICPs 66 and 37 were depressed in drug-treated cells.

tides which had electrophoretic mobilities similar to those of both host polypeptides and ICPs 93, 88, and 58 could also be detected after the inhibitor was removed, but because their syntheses were not amplified, we did not consider them to be in the same class as the α ICPs. The syntheses of ICPs 16 and 15 were amplified after the drug treatment but not as extensively as those of ICPs 61 and 64 after the drug was removed; however, as these polypeptides did not have counterparts in the uninfected cells, we considered them to be α ICPs. The interval before β ICP synthesis was detected and the

following interval before γ ICPs were synthesized were both 2 h, again similar to temporal regulation of α , β , and γ polypeptide syntheses in untreated cells. These results indicated that a prescribed period of α ICP synthesis was required for the production of β polypeptides and that the synthesis of γ ICPs had a similar requirement for a period of β ICP synthesis. These requirements were always the same and were independent of the length of inhibition by cycloheximide.

Effect of time of addition of cycloheximide. These experiments were designed to ex-

FIG. 7. Effect of prolonged treatment with cycloheximide on the polypeptides synthesized in FV3-infected cells. Infected cells were incubated in the presence of the drug for 0 to 5 or 0 to 9 h and then pulse-labeled at various times after reversal of cycloheximide block. Just the four α ICPs were synthesized immediately after removal of the drug (ICPs 61, 44, 16, and 15). There was a delay of 2 h before β ICPs were synthesized and a further delay before γ ICPs were synthesized.

amine the effect on ICP synthesis when cycloheximide was added after the synthesis of α or β ICPs had been initiated. Infected cells were treated with inhibitor at 2 to 5 h (this allowed the synthesis of α but not β ICPs) or at 4 to 8 h (allowing α and β ICP synthesis but not γ ICP synthesis) (Fig. 8). The polypeptides synthesized immediately after treatment with cycloheximide at 2 to 5 hours were the same as those in untreated 5-h cells (i.e., all of the α and β ICPs), but the synthesis of γ ICPs was not detected until 3 to 4 h later. If addition of cycloheximide was delayed until 4 p.i., all FV3-induced polypeptides were synthesized after removal of the drug. These results indicated that a prescribed interval of infected cell polypeptide synthesis was required before the addition of cycloheximide for the production of certain ICPs after removal of the drug.

Requirement of functional induced proteins. The data presented so far indicated that β ICP synthesis required prior α ICP synthesis and that γ ICP synthesis had a similar requirement for β ICP synthesis. The following experiments were performed to determine whether

there was a functional involvement of any α or β polypeptides in the synthesis of the subsequent polypeptide class. If functional polypeptides are required, then the substitution of one or more amino acids by corresponding analogs could prevent the transition from one polypeptide class to the next.

Azetidine, canavanine, and FPA (analog of proline and hydroxyproline, arginine, and phenylalanine, respectively) were added singly to unmodified medium at 500 μ g/ml. BHK cells were infected with FV3, and after adsorption the cells were incubated for various periods in the presence of one of the analogs. The cells were labeled either immediately after removal of the analog or at various times after removal to determine whether further viral protein synthesis occurred.

Figure 9 shows the effect on ICP synthesis of adding azetidine at 0 to 5, 2 to 5, or 4 to 8 h. Treatment at these times allowed no ICP synthesis, only α ICP synthesis, or only α and β ICP synthesis, respectively, before addition of the analog. After treatment at 0 to 5 h with azetidine, both α and β ICPs were synthesized; γ ICPs were not synthesized until 4 h after removal of azetidine. When azetidine was added at 2 to 5 h, only α and β ICP syntheses were detected in the 5- to 6-h pulse. The amounts of α and β ICPs

FIG. 8. Polypeptides synthesized in FV3-infected cells after cycloheximide treatment at 2 to 5 or 4 to 8 h. The drug was reversed by washing, and the cells were pulse-labeled at intervals thereafter. Both α and β ICPs were synthesized after 2- to 5-h treatment, and all ICPs were detected after the 4- to 8-h treatment.

FIG. 9. Effect of azetidine on FV3-induced polypeptides. Infected cells were incubated in medium containing azetidine for 0 to 5, 2 to 5, or 4 to 8 h p.i., the analog was removed, and the cells were pulse-labeled at various times thereafter. Both α and β ICPs were synthesized on removal of the analog after the 0- to 5-h or the 2- to 5-h treatment; γ ICPs were delayed by 2 to 4 h after removal of the analog. All ICPs were synthesized immediately after removal of azetidine which had been present from 4 to 8 h.

synthesized were reduced compared with untreated infected cells. Some γ polypeptide synthesis could be detected 2 h after removal of azetidine (which was present from 2 to 5 h), and the amounts synthesized 4 h after removal of the analog were comparable to the 6-h untreated infected cells. Treatment with azetidine at 4 to 8 h did not inhibit the synthesis of any ICP when infected cells were labeled at 8 to 9 h.

Experiments with the arginine analog canavanine produced results similar to those with azetidine. Treatment at 0 to 5 h had no effect on the syntheses of α and β ICPs at 5 h, and some γ ICPs were also detected at this time. All γ ICPs were detected on removal of the analog after treatment at 2 to 5 h. Treatment at 4 to 8 h had no effect on the synthesis of any of the polypeptide classes.

Experiments with FPA (Fig. 10) gave results similar to those with azetidine, except that γ ICP synthesis was detected 2 h after removal of FPA, which was present at 0 to 5 h.

Treatment of infected cells with amino acid analogs did not affect the synthesis of β poly-

peptides. This could be because (i) functional α ICPs are not a requirement of β ICP synthesis or (ii) either proline, hydroxyproline, arginine, or phenylalanine is not at a functional site and thus the analogs of these amino acids did not produce nonfunctional α ICPs. Functional β polypeptides were involved in the synthesis of γ ICPs (Fig. 9 and 10), although it is unlikely that arginine is at a functional site because incorporation of its analog (canavanine) did not interfere with γ ICP synthesis.

Effect of actinomycin D on ICP synthesis.

The preceding experiments suggested that syntheses of β and γ ICPs were probably dependent on prior α ICP synthesis. We next performed experiments to determine whether α polypeptide synthesis was the only requirement to turn on β and γ syntheses or whether the synthesis of new RNA was also required. Inhibition of protein synthesis with cycloheximide allows transcription of mRNA to proceed; if all of the mRNA's are transcribed during the cycloheximide block, the immediate addition of actinomycin D after reversal of cycloheximide treat-

FIG. 10. Effect of FPA on FV3-induced polypeptides. The experiment was similar to that described in the legend to Fig. 4. After treatment with FPA at 0 to 5 or 2 to 5 h, α and β ICPs were synthesized at 5 h. γ ICP synthesis was delayed by 2 h. All ICPs were detected after the 4- to 8-h treatment with FPA.

ment should allow subsequent translation of all the mRNA. We therefore added actinomycin D (10 μ g/ml, a previously determined concentration which inhibited the production of all FV3-induced polypeptides) to infected cells immediately after reversal of cycloheximide treatment.

The results of these experiments are shown in Fig. 11. The polypeptides synthesized immediately after removal of cycloheximide were the same whether actinomycin D was present or not. Thus, in cells treated at 0 to 3 or 0 to 5 h only α ICPs were synthesized, and in cells treated at 2 to 5 h α and β ICPs were synthesized on removal of cycloheximide.

Some cycloheximide-treated cells were washed, actinomycin D was added, and the cells were labeled 1.5 and 3 h later. The polypeptide profiles during these subsequent periods remained consistent with the profiles obtained immediately after removal of the cycloheximide. Actinomycin D prevented the onset of the next class of ICP, which indicated that new mRNA synthesis was also a requirement for ICP synthesis.

When does the synthesis of mRNA for each class of ICP occur? These experiments were devised to determine whether all of the mRNA's for each ICP in any one class were synthesized at the same time and whether there

were any differences in the stabilities of these RNAs. We investigated the synthesis of the mRNA's indirectly by analyzing their translation products in cell extracts. Actinomycin D was added to groups of replicate infected BHK monolayers at hourly intervals. Treated monolayers from each group were pulse-labeled with [35 S]methionine at various intervals after the addition of the inhibitor.

The PAGE analysis of the cell extracts is shown in Fig. 12, and the times of appearance of the ICPs are summarized in Table 2. All of the α ICPs were detected at the same time, but the synthesis of β polypeptides apparently occurred in two phases. Similarly, the majority of γ ICPs were detected at 5 h, but some were not synthesized until 6 h.

The polypeptide profiles remained consistent within any one group of cultures, which suggested that no posttranscriptional modifications occurred in the presence of actinomycin D. The

FIG. 11. Analysis of polypeptides synthesized in FV3-infected cells in the presence of actinomycin D added immediately after cycloheximide treatment to examine the requirement for new mRNA for each ICP class. Infected cells were incubated in the presence of cycloheximide for the times shown (0 to 3 h, 0 to 5 h, and 2 to 5 h), and then immediately after reversal of cycloheximide, actinomycin D was added to prevent further transcription. Cells were pulse-labeled in the presence of actinomycin D immediately or 1.5 or 3 h later. No transition to the next class of ICP occurred in the absence of further transcription.

FIG. 12. Effect of actinomycin D added at various times after infection on the synthesis of FV3-induced polypeptides to determine when the synthesis of mRNA for each ICP occurred. Actinomycin D was added to a series of infected cells at each of the times indicated, and individual cultures were pulse-labeled (in the presence of actinomycin D) at various times after the addition of the drug. The polypeptide profiles remained constant within any one group of treated cultures. There was a sequential appearance of ICPs, which provides strong evidence for sequentially controlled transcription of the FV3 genome.

stabilities of the mRNA's appeared to be similar, as all ICPs appeared to decline at the same rate in the presence of actinomycin D; it was not possible to determine the half-lives of the mRNA's, but these must have been in excess of 6 h.

Addition of actinomycin D reduced the suppression of host cell polypeptides, and the earlier during infection the drug was added, the more the suppression was reduced. This suggested that α and β ICPs were responsible for the suppression of host cell polypeptides and that the more α and β ICPs synthesized before actinomycin D was added, the greater the suppression.

Effect of cordycepin on ICP synthesis. Cordycepin inhibits RNA synthesis either by competitive inhibition of the RNA polymerases (68) or as a chain terminator which is incorporated into the RNA (67). ICP synthesis in the presence of cordycepin (50 μ g/ml) delayed the onset of the production of each class of polypeptides; α ICPs were not produced in significant quantities until 4 h p.i. (2 h in untreated cells), β ICP synthesis was delayed until 6 h p.i. (4 h p.i. in untreated cells), and the synthesis of γ ICPs could not be detected until 9 h p.i. (instead of 6 h). All induced polypeptides were synthesized in the presence of cordycepin, however, and the drug had no effect on the suppression of host cell polypeptides compared with untreated infected cells.

TABLE 2. Times of appearance of ICPs in FV3-infected cells after addition of actinomycin D

Time of addition of actinomycin D (h)	ICPs first detected during this treatment	
	Class	ICPs
1		
2	α	61, 44, 16, 15
3	β	98, 93, 58, 52, 50(?), 34, 29
4	β	69, 46, 23
5	γ	104, 78, 76, 55, 40, 37, 35, 27, 22, 9
6	γ	18, 12

Effect of camptothecin on ICP synthesis. Camptothecin inhibits both RNA synthesis and DNA synthesis in mammalian cells by random nicking of template DNA (8, 46, 79). This drug was added to FV3-infected cells immediately after the absorption period, and the effects on the syntheses of induced polypeptides are shown in Fig. 13. The production of just α induced polypeptides was detected, and no transition to β or γ ICP synthesis occurred. The syntheses of ICPs 61 and 44 markedly declined in the presence of the drug, but the syntheses of ICPs 16 and 15 were not greatly affected. The suppression of host cell polypeptide synthesis was reduced by treatment with camptothecin, and many high-molecular-weight cell polypeptides were synthesized at 24 h p.i.

Distribution of ICPs between nucleus and cytoplasm. (i) Association of FV3-induced

FIG. 13. Effect of camptothecin on FV3-induced polypeptides. The drug was added after the adsorption period and maintained throughout the experiment. Only α ICPs were synthesized in the presence of camptothecin.

polypeptides in purified nuclei. To determine whether any of the induced polypeptides were associated with nuclei of infected cells, monolayers were pulse-labeled for 1 h with [35 S]methionine at 2.5 and 6 h p.i. and then fractionated to give nuclear and cytoplasmic preparations. The nuclei were examined by phase-contrast microscopy and were devoid of contaminating material. Figure 14 shows a comparison of the polypeptides found in the preparations together with those found in whole cell extracts and purified FV3. The majority of ICPs were found in the nuclei, although the relative amounts of these polypeptides were different from those in the cytoplasm or in whole cell extracts. The ICPs could be divided into the following categories: (i) predominantly cytoplasmic (ICPs 61 and 50); (ii) predominantly nuclear (ICPs 98, 78, 76, 34, and 23); and (iii) both nuclear and cytoplasmic (ICPs 114, 104, 93, 88, 69, 66, 58, 55, 52, 46, 44, 37, 35, 29, 27, 22, 16, and 15).

(ii) **Treatment of isolated nuclei.** We investigated the stability of the nuclear association of the ICPs by isolating nuclei from cells labeled 6 to 7 h p.i. and subjecting them to various treatments. The nuclear pellet was resuspended

in one of the following solutions: RSB, 0.5 M NaCl, or 0.1 M NaCl-0.01 M EDTA. The nuclei were then collected by centrifugation and lysed in PAGE sample buffer. Nuclei were also prepared without the detergent treatment and lysed for analysis by PAGE (Fig. 15). Treatment with 0.5 M NaCl reduced the amount of ICP 98 in the nuclei (Fig. 15a, tracks 1 and 2), but the relative amounts of the other nuclear ICPs were unaffected. Washing the nuclei with 0.1 M NaCl-0.01 M EDTA had no effect on the amounts of nuclear ICPs. Nuclei isolated without the Tween 40-deoxycholate treatment carried the same polypeptides as nuclei prepared with detergent treatment, with the exception that the amount of ICP 58 was greatly increased in non-detergent-treated nuclei. This suggests that this polypeptide may be associated with the outer nuclear membrane because the mixed detergent (Tween 40 plus deoxycholate) removes the outer nuclear membrane (35).

FIG. 14. Distribution between nucleus and cytoplasm of polypeptides in FV3-infected BHK cells. Infected cells were pulse-labeled with [35 S]methionine for 1 h at 2.5 and 6 h p.i., the cells were homogenized, and the nuclei (n) were separated from the cytoplasm (c). Polypeptides in the nuclear and cytoplasmic fractions, in whole cell extracts, and in purified FV3 were analyzed by PAGE. All ICPs except ICPs 61 and 50 were observed in nuclei. ICPs 78 and 23 were not present in the cytoplasmic preparations.

(iii) **Uptake of polypeptides by isolated nuclei.** The affinities of virus-induced polypeptides for nuclear structures were investigated further by incubating purified unlabeled uninfected nuclei in a radiolabeled cytoplasmic preparation derived from infected cells. Nuclei from 5×10^6 uninfected BHK cells were incubated in a cytoplasmic fraction prepared from 10^6 infected cells labeled at 6 to 7 h p.i. The detergent washing step was omitted in both of these preparations. After incubations for 1 h at room temperature, the nuclei were reisolated by centrifugation, washed with Tween 40-deoxycholate, and lysed in PAGE sample buffer. All of the polypeptides found in the nuclei of infected cells were taken up by nuclei *in vitro*. The uptake of ICPs 98 and 76 appeared less efficient *in vitro*, but the relative amounts of other ICPs were similar to those in nuclei *in vivo*. No polypeptide that was not present in the nuclei of infected cells was taken up by the isolated nuclei.

Distribution of ICPs on cytoplasmic membranes. (i) Cell fractionation. Fractionation of cytoplasmic extracts of BHK cells on discontinuous sucrose gradients resolved six visible bands. The top band (fraction 1) contained just lipid and was discarded. The other five bands (fractions 2 through 6) were collected, diluted with RSB, pelleted, and prepared for electron microscopy or PAGE. Transmission electron microscope examination of thin sections showed that the fractions contained structures similar to those observed with fractions derived from other cell systems using the same procedure (9, 11, 47). Fractions 2 and 3 contained smooth membranes, fraction 4 contained a mixture of smooth and rough membranes, fraction 5 contained rough membranes and mitochondria, and fraction 6 was composed predominantly of free ribosomes. A pellet at the bottom of the gradient designated fraction 7 (11) was not detected in this microscale procedure. It has been established previously that BHK plasma membranes are removed by the low-speed centrifugation step in this procedure (11, 57), and the identities of the membranes in each fraction have been confirmed by enzyme analyses (9, 57).

(ii) **Association of viral polypeptides with cytoplasmic membranes.** BHK cells infected with FV3 were labeled with [35 S]methionine and fractionated as described above, and the membrane fractions were analyzed by PAGE (Fig. 16). Cells labeled at 3 to 4 h p.i. contained α and β ICPs. The α ICPs (e.g., ICPs 44, 16, and 15) were found associated with all fractions but were present in larger amounts on the rough membranes (fractions 4 and 5) and in the ribosomal fraction (fraction 6). β ICPs (e.g., ICPs 98, 52,

FIG. 15. (a) Treatment of isolated nuclei derived from infected cells pulse-labeled with [35 S]methionine at 6 to 7 h p.i. Nuclei were resuspended in RSB (track 1), RSB-0.5 M NaCl (track 2), or RSB-0.1 M NaCl-0.01 M EDTA (track 3) and then collected by centrifugation and lysed for PAGE. Nuclei were also prepared without the mixed detergent treatment (track 4). (b) Uptake of polypeptides by isolated nuclei. Uninfected nuclei were incubated in [35 S]methionine-labeled infected cytoplasm for 1 h, and the nuclei (N) were reisolated by centrifugation and treated with mixed detergents. The nuclei *in vitro* incorporated all of the nuclear ICPs.

and 29) had a similar distribution but were relatively more abundant on fractions 4, 5, and 6 than on fractions 2 and 3 compared with the α ICPs.

The predominant polypeptide on the smooth membranes (fractions 2 and 3) had a molecular weight of approximately 35,000, and we have designated this protein 35a. ICP 35 is a γ polypeptide, but a number of other virus-induced polypeptides with molecular weights of about 35,000 have been detected by two-dimensional gel electrophoresis (18). We suggest that protein 35a is one or more of these polypeptides and

FIG. 16. Polypeptides associated with cytoplasmic membranes of FV3-infected BHK cells. Cells were pulse-labeled for 1 h with [^{35}S]methionine, and at 3–4 or 7–8 h after infection the cells were fractionated by the method of Caligiuri and Tamm (9). Fractions 2 through 6 were collected, diluted with RSB, pelleted by centrifugation, and lysed in PAGE sample buffer as described in the text. The majority of ICPs were associated with fractions containing ribosomes (fractions 4 through 6). Smooth membrane fraction 3 contained most of the ICPs, but few ICPs were associated with smooth membrane fraction 2.

that these polypeptides can be more easily detected on isolated membranes than in whole cell extracts.

Cells labeled at 7 to 8 h p.i. contained α , β , and γ virus-induced polypeptides. All ICPs were found associated with the ribosomal fraction (fraction 6) and with rough membranes (fractions 4 and 5). ICP 46 was found in greater amounts in fraction 6. The majority of ICPs were also detected on smooth membrane fraction 3, but the amounts of some of the polypeptides (e.g., ICPs 98, 46, and 34) were much less compared with the amounts on the rough membranes. Only ICPs 55, 52, and 16 were detected in fraction 2.

Association of induced polypeptides with the plasma membrane. The purity of BHK plasma membranes prepared by using fluorescein mercuric acetate has been discussed by Klenk and Chopin (47). The plasma membrane

preparations described here were examined by microscopy and resembled those described by these authors and on morphological grounds were similar to those of other workers (80, 81).

BHK cells infected with FV3 were labeled with [^{35}S]methionine at 4 to 5 or 12 to 13 h p.i. Figure 17 shows the polypeptide profiles of purified plasma membranes derived from these cells compared with the profiles of the corresponding whole cell extracts and purified FV3. The relative amounts of polypeptides in the plasma membrane were different from those in infected whole cell extracts and in purified virus. Some α ICPs (ICPs 44, 16, and 15) and β ICPs (ICPs 98 and 58) were found in plasma membranes derived from cells labeled at 4 to 5 h p.i. The amount of ICP 44 was enriched compared with ICP 16 and ICP 15. The predominant plasma membrane-associated polypeptides labeled 12 to 13 h p.i. were ICP 58 and ICP 55. ICP 58 was

FIG. 17. Association of polypeptides with the plasma membrane in FV3-infected cells. Plasma membranes were isolated from [35 S]methionine-labeled cells and prepared for analysis by PAGE. The predominant polypeptides were ICPs 58 and 55. ICP 58 was enriched relative to ICP 55 compared with either whole cell extracts or purified FV3.

considerably enriched relative to ICP 55 compared with whole cell extracts and with purified FV3. ICPs 98, 69, 16, and 15 were also detected in proportions similar to those found at 4 h p.i., but ICP 44 was reduced compared with the 4- to 5-h labeling period.

DISCUSSION

Analysis by PAGE of radiolabeled cell extracts showed that FV3 induces at least 29 polypeptides in infected BHK or FHM cells. The absolute criterion for the viral specificity of a polypeptide is its production in a cell-free protein-synthesizing system programmed with mRNA complementary to viral DNA (37). We used the following criteria as evidence of viral specificity of the 29 ICPs: (i) their presence in infected cells but not in uninfected cells; (ii) stimulation of their synthesis after infection; and (iii) similar infected cell polypeptide profiles in a number of cell lines. The identification of ICPs as structural polypeptides was based on the sim-

ilarity of the electrophoretic mobilities of these polypeptides to those of the polypeptides in purified FV3, and the seven ICPs which did not fulfill this criterion were assumed to be nonstructural. Ideally, one should demonstrate chemical identity between the viral structural polypeptides and their presumptive counterpart ICPs to differentiate the structural and nonstructural polypeptides (37).

The 29 ICPs have an aggregate molecular weight of 1.469×10^6 , which would account for about 29.4% of the coding capacity of the viral genome; of this, 5.5% (2.73×10^5) comprises the nonstructural polypeptides. Summation of the ICPs in each phase of synthesis shows that the α ICPs (molecular weight, 1.36×10^5) represent 2.7%, the β ICPs (molecular weight, 7.15×10^5) represent 14.3%, and the γ ICPs (molecular weight, 6.18×10^5) represent 12.4% of the coding capacity.

The remaining 70% of the genetic information, if expressed, is either synthesized in amounts too small to be detected or is masked by other polypeptides. Analysis by a variety of other one-dimensional PAGE systems failed to resolve additional ICPs, although more induced polypeptides have been resolved by two-dimensional gel electrophoresis and are described in an accompanying paper (18).

Posttranslational modifications of FV3-induced polypeptides. Modifications of polypeptides after translation include (i) rapid posttranslational cleavages, (ii) slow posttranslational cleavages occurring during transport of polypeptides within the cell or during assembly of virions, and (iii) addition of various prosthetic groups (e.g., phosphorylation, glycosylation). In this paper we have demonstrated that there is little posttranslational modification of FV3 ICPs. The use of protease inhibitors (TPCK and TLCK) and pulse-chase experiments showed that neither type of posttranslational cleavage occurred.

No glycosylation of ICPs was detected when radiolabeled glucosamine was used as the precursor. These results confirm unpublished observations cited by Willis et al. (82) but contrast with both herpesvirus and poxvirus replication, where slow posttranslation cleavages and glycosylation have been demonstrated to be important modifications of the induced polypeptides (36, 63, 72).

Pinter and Compans (65) have suggested that a general characteristic of enveloped viruses is the sulfation of certain virion glycoproteins, and studies on herpesviruses have shown that both structural and nonstructural glycoproteins are sulfated (20, 40). The sulfation of non-glycosyl-

ated polypeptides has not been demonstrated (40, 65) and is considered to be a specific event for glycoproteins (58). It was not unexpected, therefore, that we were unable to detect sulfated ICPs in FV3-infected cells.

Purified FV3 particles possess a protein kinase activity which has been extensively characterized (70, 75), and it has been shown in an *in vitro* assay that at least nine virion polypeptides (with apparent molecular weights of 111,000, 102,000, 53,000, 46,000, 29,000, 23,000, 16,000, 14,000, and 10,000) were phosphorylated (69). We have resolved at least 10 phosphorylated ICPs, and allowing for differences in molecular weight estimates in the different electrophoretic conditions, all of the above structural polypeptides were phosphorylated *in vivo*. ICPs 69 and 61 were phosphorylated *in vivo* but apparently not *in vitro*.

Temporal control of induced polypeptides. We have presented evidence for the temporal control of FV3-induced polypeptides in cells; similar results have been found in FV3-infected *Xenopus laevis* cells (16) and FHM cells (R. M. Elliott, D. Phil. thesis, University of Oxford, Oxford, England, 1979). Three coordinately regulated phases of synthesis occurred, which we have designated α , β , and γ . These results are in marked contrast to the data of Goorha and Granoff (27) and Willis et al. (82), who claimed that all induced polypeptides were synthesized at the same time (by 2 h p.i.), although the rates of synthesis differed as replication progressed. One reason for this may be that the experiments were performed in FHM cells (in which FV3 replicates quickly), and so the length of the radioactive pulse and the time of the pulse may have been unsuitable to reveal sequential polypeptide synthesis. Another factor we considered that could account for the difference was the use of Δ FV3 by this group to inhibit host cell macromolecular synthesis before infection with active FV3; however, only cells to be labeled early in infection (before 6 h p.i.) were given this treatment (27). We have not found it necessary to inhibit host cell functions to detect early viral expression, because experiments mimicking the work described by Goorha and Granoff (27) failed to show that prior treatment with Δ FV3 had an effect on the sequential synthesis of induced polypeptides.

Suppression of host cell polypeptide synthesis. Host cell DNA, RNA, and protein syntheses are suppressed after infection with FV3 at permissive or nonpermissive temperatures, with heat-inactivated or irradiated FV3, or with a detergent-solubilized viral preparation, and this suggests that a structural component of

the virus is responsible for the suppression (1, 2, 27, 32, 50, 53). We have demonstrated that the suppression of host cell polypeptide synthesis is not due to a γ ICP, because the suppression of host proteins occurs unabated in the presence of ara-C. Suppression of host polypeptide synthesis was found to be temperature dependent, the rate of suppression being less at the nonpermissive temperatures. We detected the synthesis of only α ICPs at nonpermissive temperatures, in contrast to the data of Costanzo et al. (12), who claimed that all induced polypeptides were produced at the nonpermissive temperature. The production of just α ICPs could be explained by functional α ICP being required for the synthesis of subsequent induced polypeptides and by the fact that one or more α ICPs become nonfunctional at the nonpermissive temperature.

Viral DNA replication is required for γ polypeptide synthesis. The polypeptides induced by vaccinia virus can be divided into the following two groups: "prereplicative" polypeptides, which are synthesized before viral DNA replication; and "postreplicative" polypeptides, which are synthesized after viral DNA replication (63). The synthesis of detectable γ ICPs in FV3-infected cells was prevented when viral DNA replication was inhibited by ara-C, hydroxyurea, or phosphonoacetic acid (17). The γ ICPs therefore represent postreplicative polypeptides, and in this respect there is a similarity between the polypeptides induced by FV3 and those induced by vaccinia virus. Goorha and Granoff (27) were unable to demonstrate a postreplicative class of FV3-induced polypeptides when they inhibited viral DNA replication with ara-C or BUdR. The situation in FV3-infected cells is different from that in herpesvirus-infected cells, where synthesis of late polypeptides (γ phase) is reduced by two-thirds when viral DNA replication is prevented (36; R. W. Honess, personal communication).

The thymidine analogs 5'-bromodeoxyuridine and 5'-fluorodeoxyuridine had no effect on the production of γ polypeptides, and this could reflect the relatively low adenine plus thymidine content (44%) (43) of the FV3 genome. However, it has been reported that 15 μ g of 5'-fluorodeoxyuridine per ml (less than $\frac{1}{2}$ the concentration we used) blocks FV3 viral DNA replication (27). In vaccinia virus-infected cells, 5'-bromodeoxyuridine did not inhibit the synthesis of postreplicative polypeptides, although it did interfere with the processing of some of them (63); however, a number of postreplicative vaccinia virus-induced polypeptides were not synthesized in the presence of 5'-fluorodeoxyuridine (21).

The synthesis of α ICPs did not require pre-

vious cell protein synthesis since α ICPs were produced immediately after removal of inhibitors of protein synthesis, which had been present from the beginning of infection. In the presence of cycloheximide, the α polypeptide mRNA's were synthesized and remained functionally stable; on removal of the drug, the synthesis of two α ICPs (ICPs 61 and 44) was amplified, suggesting either the accumulation of these mRNA's during cycloheximide treatment or the stimulation by cycloheximide of a factor which enhances the translation of the mRNA's when the drug was removed. The other α ICPs (ICPs 16 and 15) were synthesized at a lower rate than ICPs 61 and 44 after removal of cycloheximide, suggesting that α polypeptides may represent more than one class of polypeptides, although all four are coordinately induced temporally in the presence of camptothecin and at supraoptimal temperatures, and there is no other criterion as yet by which we may subdivide α polypeptides into classes.

The synthesis of β polypeptides was dependent on a specified period of infected cell protein synthesis, presumably the production of α ICPs, because β ICP synthesis was not detected until 2 h after α polypeptide production had begun. This interval of α polypeptide production was about 2 to 3 h long and was independent of the length of previous inhibition of protein synthesis. The experiments with cycloheximide and actinomycin D indicated that new mRNA synthesis in the presence of α ICPs was also required for the synthesis of β polypeptides. We suggest, therefore, that one or more α ICPs were involved in the transcription of β polypeptide mRNA's and that transcription of these RNAs occurred after the 2-h period of α ICP synthesis. We were unable to demonstrate the production of nonfunctional α polypeptides by using the amino acid analogs azetidine, canavanine, and FPA, although we had previously suggested that functional α ICPs were required for subsequent β polypeptide synthesis, because only α ICPs were produced at nonpermissive temperatures. The use of other amino acid analogs should be investigated to attempt to produce nonfunctional α ICPs.

The production of γ polypeptides also had a requirement for infected cell protein syntheses and required a specific interval (about 2 h) of β ICP production. This indicated that the synthesis of γ ICPs may involve one or more β polypeptides or β polypeptides plus α polypeptides. The synthesis of new mRNA was also required, as no γ ICP syntheses was detected in infected cells producing β polypeptides when actinomycin D was added. Nonfunctional β polypeptides

produced by incubating infected cells in the presence of amino acid analogs did not switch on γ ICP syntheses. We have shown previously that the syntheses of viral progeny DNA is required for the production of γ ICPs; we now extend this requirement to include the production of functional β ICP syntheses is inhibited but when viral DNA replication does occur (data not shown), i.e. during the 2- to 5-h block with cycloheximide.

In herpesvirus-infected cells, the majority of α ICPs are nonstructural polypeptides, β ICPs are nonstructural or minor structural components of the virion, and γ ICPs are typically major structural polypeptides. The induced polypeptides in FV3-infected cells are similar, although some α ICP minor structural components and some nonstructural polypeptides are synthesized in the γ phase. The functions of the induced polypeptides, other than the involvement of α and β ICPs in the syntheses of β and γ polypeptides, respectively, are not known, although one can speculate on the role of α ICPs from the results of the experiments with camptothecin. We suggest that α polypeptides may be concerned with uncoating of the viral DNA and that until α ICPs are synthesized, the DNA is protected from the action of camptothecin. Once α ICPs are produced (via the virion-associated RNA polymerase [24]), the DNA is uncoated and subjected to cleavages by camptothecin, and further transcription is prevented. We have described three alternative means for obtaining cells that are synthesizing predominately α ICPs, i.e. cycloheximide treatment, camptothecin treatment, and incubation at nonpermissive temperatures, and these could be exploited to explore the properties of α polypeptides.

The synthesis of α ICPs is prolonged and amplified in the absence of β and γ polypeptides. This suggests that the mechanism(s) for switching off α ICP synthesis is present in the β or γ phase. We could not directly ascribe the switch-off mechanism to a β polypeptide(s) because there was no rapid depression of α ICP synthesis when β polypeptide synthesis began. Significant quantities of α ICPs were still detected at 24 h p.i. in all experiments, so the switch off was never complete. This differs from the regulation of herpesvirus-induced polypeptides, where the synthesis of α ICPs declines rapidly in the presence of β polypeptides. The switch off of β and γ ICP syntheses was less marked, and the control mechanisms are not yet understood.

The suppression of host cell polypeptide synthesis is not caused by a γ polypeptide, as we have previously shown that the suppression of host proteins continues in the absence of γ ICPs.

The suppression was less marked under conditions where only α ICPs were synthesized than under conditions where both α and β ICPs were produced, and hence we suggest that suppression of host cell protein synthesis is a combined effect of one or more α and β polypeptides.

RNA synthesis in FV3-infected cells. Studies on RNA synthesis in FV3-infected cells have yielded conflicting data. Armentrout and McAuslan (1) claimed that there was no temporal regulation of FV3 transcription and that all RNA species were synthesized at the same relative rates throughout the infectious cycle. Subsequently, Willis et al. (82) reported that there was a temporally ordered control of FV3 transcription, and more recently Willis and Granoff have presented evidence for two classes of early mRNA by examining the RNAs produced after treating infected cells with cycloheximide or FPA (86). We have examined the synthesis of RNA in FV3-infected cells indirectly by analyzing the translation products, and the results obtained in the actinomycin D experiment provide strong evidence for temporal regulation of FV3 transcription, which correlates with temporal induction of polypeptides.

Willis and Granoff (86) were unable to detect any viral polypeptides after removal of cycloheximide and so were unable to identify the translation products of the "immediate" early mRNA; presumably these polypeptides should correlate with the α ICPs observed in this study. Willis et al. (82) examined the polypeptides synthesized after FPA treatment, and these appeared to be similar to the α and β polypeptides we observed. However, the mRNA for these polypeptides represented 36% of the genome, whereas we calculated that 17% of the coding capacity was required for the α and β polypeptides. Neither late mRNA nor viral DNA was produced in the presence of FPA, but Willis and Granoff (86) do not consider these events to be related. The late mRNA detected by Willis and Granoff (86) hybridized to a further 14% of the genome, which is in agreement with the 12.4% of the genome required to code for γ polypeptides.

Cordycepin is reported to have a greater effect on polyadenylic acid polymerase than on other polymerases in mammalian cells, although in vaccinia virus-infected cells mRNA synthesis was more severely inhibited than polyadenylic acid synthesis (59). Armentrout and McAuslan (1) reported the presence of polyadenylic acid sequences on FV3 mRNA, but Willis and Granoff (85) were unable to confirm this. Cordycepin delayed the synthesis of induced polypeptides, and this could be ascribed to inhibition of the polyadenylation of α ICP mRNA. Since α ICP mRNA represents a small part of the genome,

Willis and Granoff (85) could have failed to detect polyadenylic acid sequences at such low levels.

Intracellular distribution of ICPs. In FV3-infected cells, 25 virus-specific polypeptides are bound in or to the nuclei. Some ICPs are predominantly nuclear (e.g., ICPs 98, 78, 76, 34, and 23), and only ICPs 61 and 50 are not associated with nuclei. The other ICPs are found in both nuclear and cytoplasmic fractions but not necessarily in equivalent amounts. None of the treatments of isolated nuclei caused a general release of polypeptides from the nuclei, which suggests that the ICPs are not in solution within the nucleus. Treatment of nuclei with high salt (which would be expected to elute soluble proteins) only partially removed ICP 98. This indicates that the ICPs are bound to large intranuclear structures, e.g. the internal nuclear membrane, or to DNA. The attachment of the ICPs does not involve divalent cations because EDTA fails to remove these ICPs. ICP 58 is probably bound to the outer nuclear membrane since it is removed by detergent treatment. All of the nuclear ICPs were taken up by isolated uninfected nuclei *in vitro*, which suggests that these polypeptides are able to bind to normal nuclear material. The uptake is probably passive because the isolated nuclei are presumably not metabolically active.

The results presented here complement the evidence for nuclear involvement in FV3 replication and suggest that the nucleus may be an alternative site of virus replication and assembly. The failure to detect two minor structural polypeptides (ICPs 61 and 50) in the nucleus does not invalidate this suggestion, as the location of these polypeptides within the virion is unknown and they could be envelope proteins. On the other hand, since viral polypeptides were taken up by isolated nuclei *in vitro*, the nuclear association of viral polypeptides may be artifactual and may have occurred during or after breakage of the cells.

McAuslan and Armentrout (54) have suggested that FV3 is a missing link between herpesviruses and poxviruses. If FV3 multiplication can occur in the nucleus, it is likely that the strategy of FV3 replication more closely resembles that of herpesviruses. Certain poxvirus replicative events have been reported to occur in infected cell nuclei, including virus-specific DNA and RNA syntheses (25, 51). Hardy et al. (33) claimed that they detected a virus-induced polypeptide in the nuclei of cells infected with fowl poxvirus, but the detection of mature poxvirus or viral structural polypeptides in nuclei has not been reported. The biological significance of these observations is unclear because vaccinia

virus can synthesize viral DNA and proteins in enucleated cells, but maturation does not occur (64). FV3 is unable to replicate in cells lacking a functional nucleus (29), and Ortin and Vinuela (61) have reported that another icosahedral cytoplasmic deoxyribovirus, African swine fever virus, also requires a nucleus for replication. Whether this is a characteristic of all members of the iridovirus group is not yet known.

Most viral polypeptides synthesized at 3 to 4 h p.i. (i.e., α and β ICPs) were found in purified plasma membrane preparations. Later in infection when the virus buds from the cell, the majority of all ICPs (α , β , and γ) were associated with the plasma membrane. The predominant polypeptides were ICPs 58 and 55, and the amount of ICP 58 was greatly enriched with respect to ICP 55 (a major structural virion polypeptide) compared with both whole cell extracts and purified FV3. It is possible that ICP 58 may be involved in modifying the plasma membrane, resulting in the changes of this membrane described by Tripiet et al. (76). This may enable the virus to acquire its external envelope. Whether ICP 58 is a major constituent of the viral envelope is not known because it is not possible to purify enveloped virus adequately.

The polypeptides associated with cytoplasmic membranes of infected BHK cells prepared by the procedure of Caligiuri and Tamm (9) were analyzed by PAGE. There appeared to be no specific association of ICPs with particular membrane fractions. All ICPs were found in the ribosomal fraction (fraction 6) and on rough endoplasmic reticulum (fraction 5), and to some extent the ICPs were found on smooth membranes (fraction 3). Four polypeptides were detected in smooth membrane fraction 2. ICP 46 was mainly found in the ribosomal fraction and ICP 35a appeared predominantly on smooth membrane fractions 2 and 3.

The role of intracellular membranes in FV3 replication is unclear. None of the induced polypeptides has been demonstrated to be glycosylated, and so the addition of carbohydrate groups in a stepwise manner during migration from rough to smooth membranes, as in the case of the formation of the influenza virus hemagglutinin (11, 73), does not occur. FV3 probably contains an internal lipid bilayer (83) similar to the internal membrane of iridescent viruses (45). The composition of the FV3 membrane does not resemble that of the host cell membranes, and thus Willis and Granoff (83) concluded that the viral membrane was synthesized *de novo*. In this respect, FV3 resembles bacteriophage PM2 and vaccinia virus, in which lipoproteins are also synthesized *de novo* (14, 23, 34). Phage PM2 has a striking morphological resemblance to irido-

viruses, although it is smaller (60 nm in diameter). The assembly of PM2 is similar to that of iridoviruses in that the virus appears to be formed in the cytoplasm away from any cellular membrane structures (23). The membrane of PM2 has a further similarity to the iridovirus membrane because no glycoproteins are present (23). *De novo* synthesis of viral membrane components also occurs in poxvirus-infected cells, and the earliest recognizable virus-like structures seen in the cytoplasm are incomplete membranes; later in infection these envelope the viral DNA and proteins (14).

FV3 appears to be unique in having an internal lipid membrane that is synthesized *de novo* and an external envelope derived from the plasma membrane of the cell. The results presented in this paper provide a basis to elucidate the relationship between the assembly of the viral membrane and the intracellular membranes of the host.

ACKNOWLEDGMENTS

We thank R. W. Honess and M. L. Fenwick for helpful discussions of this work and J. S. Robertson for critically reading the manuscript.

R.M.E. received a postgraduate studentship from the Science Research Council.

LITERATURE CITED

1. Armentrout, R. W., and B. R. McAuslan. 1974. RNA synthesis in cells infected with an icosahedral cytoplasmic deoxyvirus (frog virus 3). *J. Virol.* 13:1083-1092.
2. Aubertin, A. M., C. Decker, and A. Kirn. 1970. Effect of γ -rays on infectivity and capacity for nuclear-associated and cytoplasmic DNA replication of FV3 in chick embryo fibroblasts. *Radiat. Res.* 44:178-186.
3. Aubertin, A. M., C. Hirth, C. Travo, H. Nonenmacher, and A. Kirn. 1973. Preparation and properties of an inhibitory extract from frog virus 3 particles. *J. Virol.* 11:694-701.
4. Barzilai, R., and B.-E. Lachmi. 1973. The effect of frog virus 3 on the biological activity of various RNA viruses. *J. Gen. Virol.* 21:201-204.
5. Bingen, A., and A. Kirn. 1975. Fibrillar bodies in the hepatocyte nuclei during the course of the toxic hepatitis produced by frog virus 3 in mice. *J. Ultrastruct. Res.* 50:167-173.
6. Bingen-Brendel, A., F. Tripiet, and A. Kirn. 1971. Etude morphologique sequentielle du developement du FV3 sur cellules BHK 21. *J. Microsc. (Oxford)* 11:249-258.
7. Bonner, W. M., and R. A. Laskey. 1974. A film detection method for tritium-labeled proteins and nucleic acids in polyacrylamide gels. *Eur. J. Biochem.* 46:83-88.
8. Bosmann, H. B. 1970. Camptothecin inhibits macromolecular synthesis in mammalian cells but not in isolated mitochondria or *E. coli*. *Biochem. Biophys. Res. Commun.* 41:1412-1417.
9. Caligiuri, L. A., and I. Tamm. 1970. The role of cytoplasmic membranes in poliovirus biosynthesis. *Virology* 42:100-111.
10. Campadelli-Fiume, G., F. Costanzo, A. Mannini-Palenzona, and M. LaPlaca. 1972. Impairment of host cell ribonucleic acid polymerase II after infection with frog virus 3. *J. Virol.* 9:698-700.
11. Compan, R. W. 1973. Influenza virus proteins. II. Asso-

- ciation with components of the cytoplasm. *Virology* 51: 56-70.
12. Costanzo, F., F. Campadelli-Fuime, L. Foa-Tomasi, and M. LaPlaca. 1976. Effect of non-permissive temperature on protein synthesis of frog virus 3 infected cells. *Boll. Ist. Sieroter. Milan* 55:267-271.
 13. Costanzo, F., M. LaPlaca, A. M. Palenzona, F. Novello, and F. Stirpe. 1970. Selective inhibition of a nuclear RNA polymerase in cells infected with frog virus. *Nature (London)* 227:294-295.
 14. Dales, S., and E. H. Mosbach. 1968. Vaccinia as a model of membrane biogenesis. *Virology* 35:564-583.
 15. Darlington, R. W., A. Granoff, and D. C. Breeze. 1966. Viruses and renal carcinoma of *Rana pipiens*. II. Ultrastructural studies and sequential development of virus isolated from normal and tumor tissue. *Virology* 29: 149-156.
 16. Elliott, R. M., M. K. Arnold, and D. C. Kelly. 1979. The replication of frog virus 3 in an amphibian cell line (XTC-2) derived from *Xenopus laevis*. *J. Gen. Virol.* 44:89-98.
 17. Elliott, R. M., A. Bateson, and D. C. Kelly. 1980. Phosphonacetic acid inhibition of frog virus 3 replication. *J. Virol.* 33:539-542.
 18. Elliott, R. M., R. Bravo, and D. C. Kelly. 1980. Frog virus 3 replication: analysis of structural and nonstructural polypeptides in infected BHK cells by acidic and basic two-dimensional gel electrophoresis. *J. Virol.* 33: 18-27.
 19. Elliott, R. M., T. Lescott, and D. C. Kelly. 1977. Serological relationships of an iridescent virus (type 25) recently isolated from *Tipula* sp. with two other iridescent viruses (types 2 and 22). *Virology* 81:309-316.
 20. Erickson, J. S., and A. S. Kaplan. 1973. Synthesis of proteins in cells infected with herpes virus. IX. Sulfated proteins. *Virology* 55:94-102.
 21. Esteban, M., and D. H. Metz. 1973. Early virus protein synthesis in vaccinia virus-infected cells. *J. Gen. Virol.* 19:201-216.
 22. Fenwick, M. L., M. J. Walker, and J. M. Petkevich. 1978. On the association of virus proteins with the nuclei of cells infected with herpes simplex virus. *J. Gen. Virol.* 39:519-530.
 23. Franklin, R. M. 1974. Structure and synthesis of bacteriophage PM2, with particular emphasis on the viral lipid bilayer. *Curr. Top. Microbiol. Immunol.* 68:107-159.
 24. Gaby, N. S., and L. S. Kucera. 1974. DNA-dependent RNA polymerase activity associated with subviral particles of polyhedral cytoplasmic deoxyribovirus. *J. Virol.* 14:231-238.
 25. Gafford, L. G., and C. C. Randall. 1976. Virus-specific RNA and DNA in the nuclei of cells infected with fowlpox virus. *Virology* 69:1-14.
 26. Gibson, W., and B. Roizman. 1974. Proteins specified by herpes simplex virus. X. Staining and radiolabeling properties of β capsid and virion proteins in polyacrylamide gels. *J. Virol.* 13:155-165.
 27. Goorha, R., and A. Granoff. 1974. Macromolecular synthesis in cells infected by frog virus 3. I. Virus-specific protein synthesis and its regulation. *Virology* 60:237-250.
 28. Goorha, R., M. Murti, A. Granoff, and R. Tirey. 1978. Macromolecular synthesis in cells infected by frog virus 3. VIII. The nucleus is a site of frog virus 3 DNA and RNA synthesis. *Virology* 84:32-50.
 29. Goorha, R., D. B. Willis, and A. Granoff. 1977. Macromolecular synthesis in cells infected by frog virus 3. Frog virus 3 replication is dependent on the cell nucleus. *J. Virol.* 21:802-805.
 30. Granoff, A. 1969. Viruses of amphibia. *Curr. Top. Microbiol. Immunol.* 50:107-137.
 31. Granoff, A., P. E. Came, and D. C. Breeze. 1966. Viruses and renal carcinoma of *Rana pipiens*. I. The isolation and properties of virus from normal and tumour tissue. *Virology* 29:133-148.
 32. Guir, J., J. Braunwald, and A. Kirn. 1971. Inhibition of host-specific DNA and RNA synthesis in KB cells following infection with frog virus 3. *J. Gen. Virol.* 12: 293-301.
 33. Hardy, C. L., C. C. Randall, and L. G. Gafford. 1978. A viral-induced protein in the nuclei of cells infected with fowlpox virus. *Virology* 84:236-241.
 34. Harrison, S. C., D. L. C. Caspar, R. D. Camerini-Otero, and R. M. Franklin. 1971. Lipid and protein arrangement in bacteriophage PM2. *Nature (London)* New Biol. 229:197-201.
 35. Holtzman, E., I. Smith, and S. Penman. 1966. Electron microscope studies of detergent treated HeLa cell nuclei. *J. Mol. Biol.* 17:131-135.
 36. Honess, R. W., and B. Roizman. 1973. Proteins specified by herpes simplex virus. XI. Identification and relative molar rates of synthesis of structural and nonstructural herpes virus polypeptides in the infected cell. *J. Virol.* 12:1347-1365.
 37. Honess, R. W., and B. Roizman. 1974. Regulation of herpes virus macromolecular synthesis. I. Cascade regulation of the synthesis of three groups of viral proteins. *J. Virol.* 14:8-19.
 38. Horwitz, M. S., and S. B. Horwitz. 1971. Intracellular degradation of HeLa and adenovirus type 2 DNA induced by camptothecin. *Biochem. Biophys. Res. Commun.* 45:723-727.
 39. Kang, H. S., and B. R. McAuslan. 1972. Virus-associated nucleases: location and properties of deoxyribonucleases and ribonucleases in purified frog virus 3. *J. Virol.* 10:202-210.
 40. Kaplan, A. S., and T. Ben Porat. 1976. Synthesis of proteins in cells infected with herpes virus. XI. Sulfated, structural glycoproteins. *Virology* 70:561-563.
 41. Kelly, D. C. 1975. Frog virus 3 replication: electron microscope observations on the sequence of infection in chick embryo fibroblasts. *J. Gen. Virol.* 26:71-86.
 42. Kelly, D. C., and M. A. Atkinson. 1975. Frog virus 3 replication: electron microscope observations on the terminal stages of infection in chronically infected cell cultures. *J. Gen. Virol.* 28:391-407.
 43. Kelly, D. C., and R. J. Avery. 1974. Frog virus 3 deoxyribonucleic acid. *J. Gen. Virol.* 24:339-348.
 44. Kelly, D. C., and J. S. Robertson. 1973. Icosahedral cytoplasmic deoxyriboviruses. *J. Gen. Virol.* 20:17-41.
 45. Kelly, D. C., and D. E. Vance. 1973. The lipid content of two iridescent viruses. *J. Gen. Virol.* 21:417-423.
 46. Kessel, D., and R. Dysard. 1973. Effects of camptothecin on RNA synthesis in L1210 cells. *Biochim. Biophys. Acta* 312:716-721.
 47. Klenk, H.-D., and P. W. Choppin. 1969. Lipids of plasma membranes of monkey and hamster kidney cells and of parainfluenza virions grown in these cells. *Virology* 38: 255-268.
 48. Korant, B. D. 1972. Cleavage of viral precursor proteins in vivo and in vitro. *J. Virol.* 10:751-759.
 49. Kucera, L. S. 1970. Effects of temperature on frog polyhedral cytoplasmic deoxyribovirus multiplication: thermosensitivity of initiation, replication and encapsidation of viral DNA. *Virology* 42:576-589.
 50. Kucera, L. S., and A. Granoff. 1968. Viruses and renal carcinoma of *Rana pipiens*. VI. Interrelationships of macromolecular synthesis and infectious virus production in frog virus 3 infected BHK 21/13 cells. *Virology* 34:240-249.
 51. LaColla, P., and A. Weissbach. 1975. Vaccinia virus infection of HeLa cells. I. Synthesis of vaccinia DNA in host cell nuclei. *J. Virol.* 15:305-315.

52. Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* **227**:680-685.
53. Maes, R., and A. Granoff. 1967. Viruses and renal carcinoma of *Rana pipiens*. IV. Nucleic acid synthesis in frog virus 3 infected BHK 21/13 cells. *Virology* **33**:491-502.
54. McAuslan, B. R., and R. W. Armentrout. 1974. The biochemistry of icosahedral cytoplasmic deoxyviruses. *Curr. Top. Microbiol. Immunol.* **68**:77-105.
55. McAuslan, B. R., and W. R. Smith. 1968. Deoxyribonucleic acid synthesis in FV3-infected mammalian cells. *J. Virol.* **2**:1006-1015.
56. Moss, B. 1974. Reproduction of pox viruses, p. 405-474. In H. Fraenkel-Conrat and R. R. Wagner (ed.), *Comprehensive virology*, vol. 3. Plenum Press, New York.
57. Mosser, A. G., L. A. Caligiuri, A. S. Scheid, and I. Tamm. 1972. Chemical and enzymatic characteristics of cytoplasmic membranes of poliovirus-infected HeLa cells. *Virology* **47**:30-38.
58. Nakamura, K., and R. W. Compans. 1978. Effects of glucosamine, 2-deoxyglucose and tunicamycin on glycosylation, sulfation and assembly of influenza viral proteins. *Virology* **84**:303-319.
59. Nevins, J. R., and W. K. Joklik. 1975. Poly(A) sequences of vaccinia virus messenger RNA: nature, mode of addition and function during translation in vitro and in vivo. *Virology* **63**:1-14.
60. O'Farrell, P. H. 1975. High resolution two-dimensional electrophoresis of proteins. *J. Biol. Chem.* **250**:4007-4021.
61. Ortin, J., and E. Vinuela. 1977. Requirement of cell nucleus for African swine fever virus replication in Vero cells. *J. Virol.* **21**:902-905.
62. Penman, S. 1969. Preparation of purified nuclei and nucleoli from mammalian cells, p. 35-48. In K. Habel and N. P. Salzman (ed.), *Fundamental techniques in virology*. Academic Press Inc., New York.
63. Pennington, T. H. 1974. Vaccinia virus polypeptide synthesis: sequential appearance and stability of pre- and post-replicative polypeptides. *J. Gen. Virol.* **25**:433-444.
64. Pennington, T. H., and E. A. C. Follett. 1974. Vaccinia virus replication in enucleate BSC-1 cells: particle production and synthesis of viral DNA and proteins. *J. Virol.* **13**:488-493.
65. Pinter, A., and R. W. Compans. 1975. Sulfated components of enveloped viruses. *J. Virol.* **16**:859-866.
66. Russell, W. C., and G. E. Blair. 1977. Polypeptide phosphorylation in adenovirus-infected cells. *J. Gen. Virol.* **34**:19-35.
67. Sawicki, S. G., W. Jelinek, and J. E. Darnell. 1977. 3' Terminal addition to HeLa cell nuclear and cytoplasmic poly(A). *J. Mol. Biol.* **113**:219-235.
68. Siev, M., R. Weinberg, and S. Penman. 1969. The selective interruption of nucleolar RNA synthesis in HeLa cells by cordycepin. *J. Cell Biol.* **41**:510-520.
69. Silberstein, H., and J. T. August. 1973. Phosphorylation of animal virus proteins by a virion protein kinase. *J. Virol.* **12**:511-522.
70. Silberstein, H., and J. T. August. 1976. Purification and properties of a virion protein kinase. *J. Biol. Chem.* **251**:3176-3184.
71. Silberstein, H., and J. T. August. 1976. Characterization of a virion protein kinase as a virus-specified enzyme. *J. Biol. Chem.* **251**:3185-3190.
72. Spear, P. G., J. M. Keller, and B. Roizman. 1970. Proteins specified by herpes simplex virus. II. Viral glycoproteins associated with cellular membranes. *J. Virol.* **5**:121-131.
73. Stanley, P., S. S. Gandhi, and D. O. White. 1973. The polypeptides of influenza virus. VII. Synthesis of the haemagglutinin. *Virology* **53**:92-106.
74. Summers, D. F., E. N. Shaw, M. L. Stewart, and J. V. Maizel. 1972. Inhibition of cleavage of large poliovirus-specific precursor proteins in infected HeLa cells by inhibitors of proteolytic enzymes. *J. Virol.* **10**:880-884.
75. Tan, K. B., and B. R. McAuslan. 1971. Proteins of polyhedral cytoplasmic deoxyviruses. I. The structural polypeptides of FV3. *Virology* **45**:200-207.
76. Tripier, F., J. Braunwald, L. Markovic, and A. Kirm. 1974. Budding of frog virus 3 studied by immunological and cytochemical methods in electron microscopy. *Intervirology* **3**:305-318.
77. Tripier, F., L. Markovic, J. Brawnwald, and A. Kirm. 1975. Donnees nouvelles sur la structure et la bourgeonnement du FV3 (frog virus 3). *Ann. Inst. Pasteur Paris* **126B**:447-460.
78. Vilagines, R., and B. R. McAuslan. 1971. Proteins of polyhedral cytoplasmic deoxyvirus. II. Nucleotide phosphohydrolase activity associated with frog virus 3. *J. Virol.* **7**:619-624.
79. Wall, M. E., M. C. Wani, C. E. Cook, K. H. Palmer, A. T. McPhail, and G. A. Sim. 1966. Plant antitumor agents. I. The isolation and structure of camptothecin, a novel alkaloidal leukemia and tumor inhibitor from *Camptotheca acuminata*. *J. Am. Chem. Soc.* **88**:3888-3890.
80. Warren, L., and M. C. Glick. 1969. Isolation of surface membranes of tissue culture cells, p. 66-77. In K. Habel and N. P. Salzman (ed.), *Fundamental techniques in virology*. Academic Press Inc., New York.
81. Warren, L., M. C. Glick, and M. K. Nass. 1966. Membranes of animal cells. I. Methods of isolation of the surface membrane. *J. Cell. Physiol.* **68**:269-288.
82. Willis, D. B., R. Goorha, M. Miles, and A. Granoff. 1977. Macromolecular synthesis in cells infected by frog virus 3. VII. Transcriptional and posttranscriptional regulation of virus gene expression. *J. Virol.* **24**:326-342.
83. Willis, D., and A. Granoff. 1974. Lipid composition of frog virus 3. *Virology* **61**:256-269.
84. Willis, D. B., and A. Granoff. 1976. Macromolecular synthesis in cells infected by frog virus 3. IV. Regulation of virus-specific RNA synthesis. *Virology* **70**:399-410.
85. Willis, D. B., and A. Granoff. 1976. Macromolecular synthesis in cells infected with frog virus 3. V. The absence of polyadenylic acid in the majority of virus-specific RNA species. *Virology* **73**:543-547.
86. Willis, D. B., and A. Granoff. 1978. Macromolecular synthesis in cells infected by frog virus 3. IX. Two temporal classes of early viral RNA. *Virology* **86**:443-453.