Published in final edited form as:

J Nat Prod. 2010 March 26; 73(3): 485–488. doi:10.1021/np900728x.

Paltolides A–C, Anabaenopeptin-type Peptides from the Palau Sponge *Theonella Swinhoei*[†]

Alberto Plaza[‡], Jessica L. Keffer[‡], John R. Lloyd[‡], Patrick L. Colin[§], and Carole A. Bewley^{*,‡}

Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland 20892-0820 Coral Reef Research Foundation, Republic of Palau

Abstract

Three new anabaenopeptin-like peptides, named paltolides A–C, were isolated from a deep-water specimen of the marine sponge *Theonella swinhoei* from Palau. Paltolides belong to a rare subgroup of sponge-derived anabaenopeptins that have in common a C-terminal tryptophan residue linked to the ϵ -amine of a lysine bearing a D configuration. The structures of paltolides A–C were determined by NMR and tandem MS techniques. Paltolide A is the first anabaenopeptin structure where a non-*N*-methylated amino acid precedes the C-terminal residue.

Sponges belonging to the genus *Theonella* have proven to be one of the most prolific sources of bioactive natural products, some of which are almost certainly biosynthesized by symbiotic bacteria or cyanobacteria.¹ Among this group, the sponge *Theonella swinhoei* (order Lithistida, family Theonellidae) in particular has yielded a number of unusual peptides with unique structures including the antifungals cyclolithistide A,^{2a} theonegramides^{2b} and theopalauamide^{2c}, the HIV-inhibitors papuamides³, and the cytotoxic polytheonamides^{4a}. In our continuing studies of lithistid demosponges, we identified a deep water Palauan specimen of *T. swinhoei* with a red-purple ectosome and cream endosome that showed activity in HIV-1 neutralization assays. Fractionation of its methanol extract led to the isolation of the known glycopeptide theopalauamide^{2c} and six anabaenopeptin-like peptides (1–6). Paltolides A–C (1–3) are new peptides while compounds 4–6 have been previously isolated from the Australian sponge *Melophlus* sp.⁵ Their structures were elucidated by extensive spectroscopic methods including 1D (¹H and ¹³C) and 2D (DQF-COSY, HOHAHA, HMBC, HSQC, and ROESY) NMR experiments as well as ESI-MSⁿ analysis. The absolute configurations of the paltolides were established by LC-MS analysis of their advanced Marfey's derivatives.⁶

[†]Dedicated to the late Dr. John W. Daly of NIDDK, NIH, Bethesda, Maryland and to the late Dr. Richard E. Moore of the University of Hawaii at Manoa for their pioneering work on bioactive natural products.

^{*}To whom correspondence should be addressed. Tel.: (301) 594-5187. Fax (301) 402-4182. caroleb@mail.nih.gov.

[‡]NIDDK

[§]Coral Reef Research Foundation

Supporting Information Available: 1D and 2D NMR spectra for compounds 1–3, and NMR spectroscopic data for compound 4. This material is available free of charge via the Internet at <http://pubs.acs.org>.

1	X = H	Y = H	R ¹ = H
2	X = H	Y = OH	R ¹ = Me
3	X = Br	Y = H	R ¹ = Me
4	X = Cl	Y = OH	R ¹ = Me
5	X = Cl	Y = H	R ¹ = Me
6	X = H	Y = H	R ¹ = Me

Following lyophilization of the frozen sponge, extracts were prepared through sequential extraction of the freeze dried sponge with hexanes, CHCl₃, and MeOH. The MeOH extract was partitioned with *n*-BuOH–H₂O and the organic layer was fractionated on Sephadex LH-20. C12 reversed-phase HPLC of the fractions containing peptides yielded compounds **1**–**6**. The most polar compound among this group was paltolide A (**1**). Its HR-ESI-MS showed a major ion peak at *m/z* 812.4775 [M+H]⁺ corresponding to a molecular formula of C₃₉H₆₁N₁₁O₈. An IR band at ν 1649 cm⁻¹ and HSQC correlations between six α -proton signals at δ 3.90–4.70 and six carbon resonances at δ 51.0–56.5 indicated that **1** was a peptide containing six amino acid residues. Detailed analysis of the 2D NMR data obtained from HSQC, HMBC, 2D-HOHAHA, and DQF-COSY experiments revealed the presence of arginine, lysine, alanine,

and two leucine residues (see Table 1). Moreover the downfield region of the ^1H NMR spectrum contained a singlet signal at δ 7.11 (1H, s), along with proton signals characteristic of a 1,2-disubstituted benzene at δ 7.57 (1H, br d, $J = 8.1$ Hz), 7.31 (1H, br d, $J = 8.1$ Hz), 7.07 (1H, t, $J = 7.3$ Hz), and 7.00 (1H, t, $J = 7.3$ Hz). These data together with diagnostic HMBC correlations (see Table 1) from the proton signals at δ 7.11 (H-2'_{Trp}) and 7.57 (H-4'_{Trp}) to the carbon resonances at δ 111.5 (C-3'_{Trp}), 128.6 (C-9'_{Trp}), and 138.0 (C-8'_{Trp}) indicated the presence of an indole ring. Subsequently, long-range correlations from the β -methylene protons at δ 3.48 and 3.20 to the carbon resonances at δ 111.5, 124.5, 128.6, and 173.7 (C-1_{Trp}) allowed us to assign the sixth residue as tryptophan.

The amino acid sequence of **1** could be established from interpretation of a single HMBC experiment where long-range correlations between α -protons and carbonyl carbons belonging to adjacent amino acids provided the partial sequence Lys-Ala-Leu1-Leu2-Trp (Table 1). Cyclization between the C-terminal tryptophan and the lysine ω -amine was apparent from HMBC correlations between the ϵ -methylene protons of lysine (δ_{H} 3.51, 3.11, δ_{C} 40.1) and the carbonyl resonance at δ 173.7 (C-1_{Trp}). An unassigned carbon resonance at δ 160.1 remained, and HMBC correlations from α -protons at δ 4.32 (H-2_{Arg}) and 4.08 (H-2_{Lys}) to this carbon were indicative of the presence of a ureido moiety connecting the arginine and N-terminal lysine residues establishing the structure of paltolide A as depicted in **1**.

Additional evidence confirming this peptide sequence was obtained from ESI-MSⁿ experiments. Fragmentation of the major ion peak at m/z 812 $[\text{M}+\text{H}]^+$ displayed an intense ion at m/z 612 $[\text{M}+\text{H}-\text{Arg}-\text{CO}]^+$, and two fragments of small intensity at m/z 484 $[\text{M}+\text{H}-\text{Arg}-\text{CO}-\text{Lys}]^+$ and m/z 413 $[\text{M}+\text{H}-\text{Arg}-\text{CO}-\text{Lys}-\text{Ala}]^+$. Further MS³ fragmentation of the ion daughter at m/z 612 yielded ion fragments at m/z 499 $[\text{M}+\text{H}-\text{Arg}-\text{CO}-\text{Leu}]^+$, m/z 386 $[\text{M}+\text{H}-\text{Arg}-\text{CO}-\text{Leu}-\text{Leu}]^+$, and m/z 200 $[\text{M}+\text{H}-\text{Arg}-\text{CO}-\text{Leu}-\text{Leu}-\text{Trp}]^+$. Therefore the MSⁿ fragmentation patterns were in total agreement with the structure proposed for **1** by NMR.

Acid hydrolysis of paltolide A followed by derivatization with L- and D-FDLA (1-fluoro-2,4-dinitrophenyl-5-L/D-leucinamide) and subsequent LC-MS analysis⁶ in comparison to respective standards allowed us to establish the absolute configurations of Arg, Ala, Trp, and the two Leu residues as L, while the configuration of Lys was established as D.

The molecular formula of paltolide B (**2**) was determined to be $\text{C}_{40}\text{H}_{63}\text{N}_{11}\text{O}_9$ by HR-ESI-MS (m/z 842.4880 $[\text{M}+\text{H}]^+$, calcd for $\text{C}_{40}\text{H}_{64}\text{N}_{11}\text{O}_9$, 842.4888). Its ^1H NMR spectrum showed signals characteristic of a peptide including exchangeable NH signals at δ 7.70–10.00 and a signal at δ 1.89 corresponding to an upfield-shifted methylamide. Furthermore, HSQC correlations and the coupling patterns of its aromatic signals at δ 6.63–7.12 suggested the presence of a 2',5'-disubstituted indole ring (see Table 1). Analysis of the 2D NMR data established the presence of Lys, Ala, Leu, NMeLeu, 5'-hydroxytryptophan (5'-OHTrp), Arg and an urea moiety. Additionally, the HMBC spectrum indicated paltolide B has a very similar amino acid sequence with respect to that of the previously reported compound **4** (see Table S1 for NMR data of **4**). In fact comparison between their NMR data showed that the only difference between these two compounds was the absence of chlorine at the 6' position of the tryptophan residue in paltolide B thereby establishing **2** as the des-chloro analogue of **4**. The presence of bromine in paltolide C (**3**) was apparent from the isotopic pattern of 1:1 observed for the major ion peaks at 904 $[\text{M}+\text{H}]^+$ and 906 $[\text{M}+2\text{H}]^+$, and its molecular formula was assigned as $\text{C}_{40}\text{H}_{62}\text{BrN}_{11}\text{O}_8$ by HR-ESI-MS (m/z 904.4030 $[\text{M}+\text{H}]^+$). Comparison of the 2D NMR spectra of paltolide C with the spectra of paltolide B showed signals assigned to 5'-hydroxytryptophan (5'-OHTrp) to be replaced with signals belonging to 6'-bromotryptophan (6'-BrTrp). LC-MS analysis of the L/D-FDLA-derivatized hydrolysates of **2** and **3** revealed L configurations for NMeLeu, Leu, Arg, and Ala, and D configuration for Lys. Due to insufficient quantities of peptide and poor yields of the substituted Trp residues following acid hydrolysis and

derivatization, we were unable to determine the absolute configuration of the substituted tryptophan residues by LC-MS. However, the NMR data and chemical composition of **2** and **3** are highly similar to those of the known compound **4** whose absolute configuration was determined by X-ray crystallography to include the D configuration of Lys and L configuration for all other amino acids, including the substituted tryptophan.⁵ On the basis of these data and the co-existence of paltolides A–C with the known compound **4** in this *T. swinhoei* sample, we suggest that the 5'-OHTrp residue in paltolide B (**2**) and 6'-BrTrp residue in paltolide C (**3**) also possess an L configuration.

Paltolide B and compound **4** did not inhibit HIV-1 entry or show cytotoxicity toward HCT-116 or a control mammalian cell line at concentrations up to 100 µg/mL. Instead, the HIV-1 neutralizing activity of the extract was traced to the known glycopeptide theopalauamide.

Anabaenopeptins, which have been isolated from cyanobacteria and sponges, are a well-described family of ureido-containing hexapeptides characterized by a 5-residue lactam ring formed by cyclization between the C-terminal acid and the ε-amine of an invariant N-terminal lysine residue. Exocyclic to the lactam, a sixth amino acid is linked to the α-amine of Lys through a urea bridge. On the basis of their structural features, paltolides A–C (**1–3**) can be placed within a rare subgroup of anabaenopeptins that have in common a tryptophan residue at the C-terminus linked to the ε-amine of the N-terminal lysine. The remaining members of this subgroup include six carboxypeptidase U inhibitors from *Melophlus* sp. described in a patent by researchers in Australia;⁵ and konbamide,⁷ a calmodulin antagonist isolated from an Okinawan collection of *Theonella* sp. Another signature of all anabaenopeptins regardless of source is the presence of an *N*-methylated amino acid appearing adjacent to and before the C-terminal residue.^{8–10} Surprisingly, paltolide A (**1**) possesses a standard leucine residue at this position presenting the first example where the amino acid at this position lacks an *N*-methyl group. Paltolides A–C also represent the first example of anabaenopeptin-type peptides coming from *T. swinhoei*. Other sponge-derived anabaenopeptins have come from collections of *Theonella* spp. (konbamide,⁷ keramamides A11a and L11b and mozabamides A and B12), *Psammocinia* aff. *bulbosa* (psymbamide A8), and *Melophlus* sp.⁵

It is also interesting to note that the lysine residue in each of the paltolides has a D configuration and the remaining amino acids an L configuration. All twenty-nine cyanobacterial-derived anabaenopeptins contain a D-Lys and the remaining amino acids have L configurations, while anabaenopeptins isolated from sponges typically show L configurations for all their amino acids including Lys.^{10,13} Similar to the configuration seen in paltolides A–C, the sponge-derived compounds psymbamide A and anabaenopeptins from *Melophlus* sp. also contain a D-lysine with L configurations for the other five amino acids. This distribution is remarkable given that several of the L-Lys-containing peptides were isolated from sponges of the genus *Theonella*, and that to date anabaenopeptins have not been found in numerous other Palauan *T. swinhoei* samples collected at shallower depths (less than 100 ft). These findings extend the diversity of chemistry associated with *Theonella swinhoei*, and probably indicate that cyanobacteria or other prokaryotes are involved in the synthesis of these peptides. Indeed, Piel, Crews and their co-workers recently showed a prokaryotic origin for psymberrin, an antitumor natural product found in some *P. aff. bulbosa* chemotypes such as those that yield psymbamide A.¹⁴

Although paltolides A–C did not show inhibition of HIV-1 entry nor cytotoxicity toward HCT-116, their structural features warrant further biological evaluation. As mentioned above, paltolides A–C are closely related to a group of anabaenopeptins that are submicromolar inhibitors of carboxypeptidase U with greater than 50-fold selectivity over other carboxypeptidases.⁵

Experimental Section

General Experimental Procedures

Optical rotations were measured with a Jasco P-2000 polarimeter, UV spectra were recorded on an Agilent 8453 spectrophotometer, and IR spectra were recorded on a Perkin Elmer FT-IR Spectrum One spectrometer. NMR spectra were recorded in CD₃OD on Bruker spectrometers operating at 600 and 500 (cryoprobe) MHz for ¹H, and 150 and 125 MHz for ¹³C NMR. DQF-COSY, 2D-HOHAHA, HSQC, HMBC, and ROESY experiments were recorded using standard pulse programs. HSQC experiments were optimized for ¹J_{C-H} = 145 Hz, and HMBC spectra were optimized for ^{2,3}J_{C-H} = 5, 6 and 8 Hz. The accurate mass electrospray ionization (ESI) mass spectra were measured on a Waters LCT Premier time-of-flight (TOF) mass spectrometer. The instrument was operated in ω -mode at a nominal resolution of 10,000. The electrospray capillary voltage was set at 2KV and the sample cone voltage at 60 volts. The desolvation temperature was set to 275 °C and nitrogen was used as the desolvation gas with a flow rate of 300 L/h. Accurate masses were obtained using the internal reference standard method. ESI-MS/MS data were obtained using a Thermo-Scientific (San Jose, CA, USA) LTQ ion Trap mass spectrometer. Sample was infused into the mass spectrometer using an Advion BioSciences (Ithaca, NY, USA) Triversa chip based nano-electrospray ionization system. The nitrogen gas pressure was 0.25 PSI and the electrospray tip voltage was 1.4 KV. The CID MS/MS collision energy was 35 V and the parent ion isolation width was 3 Daltons. The maximum injection time for parent ions was 700 ms and 500 ms for daughter ions. The maximum AGC ion target setting was 1 × 10⁵ for parent ions and 5 × 10⁴ for daughter ions. QTOF-MS/MS data were obtained using a Waters (Milford, MA USA) Q-TOF-2 mass spectrometer operated in positive ion mode. The ESI capillary voltage was 3.5 KV, the desolvation gas was nitrogen with a flow of 300 L/h, and the desolvation temperature was 250 °C. The collision gas was argon and the collision energy was 45 V. The parent fragment ion was generated by in-source fragmentation with a cone voltage of 40 V.

Sponge Material

The marine sponge *Theonella swinhoei* (lithistid Demospongiae: family Theonellidae) was collected by hand using SCUBA on Uchelbeluu reef in Palau at a depth of 330 feet in June 2008 and frozen within 2 h. Samples were flown to Maryland frozen and stored at −80 °C until freeze-drying prior to extraction.

Isolation

The lyophilized sponge (30 g) was sequentially extracted with hexanes, CHCl₃, and MeOH. The MeOH extract (4 g) was partitioned between *n*-BuOH-H₂O (1:1) and the organic layer (1 g) was fractionated on Sephadex LH-20. Fractions containing peptides (95 mg) were purified by reversed-phase HPLC (Jupiter Proteo C12, 250 × 10 mm, 4 μ , DAD at 220 and 280 nm) eluting with a linear gradient of 50–80% MeOH in 0.05% TFA in 50 min to afford compounds **1** (0.5 mg, *t*_R = 19.0 min), **2** (1.5 mg, *t*_R = 20.7 min), **3** (0.6 mg, *t*_R = 36.5 min), **4** (6.8 mg, *t*_R = 25.2 min), **5** (0.3 mg, *t*_R = 35.3 min), **6** (24.8 mg, *t*_R = 26.3 min), and theopalauamide (0.9 mg, *t*_R = 39.8 min).

Paltolide A (1): colorless amorphous powder; [α]_D²³ −22 (*c* 0.05, MeOH); UV (MeOH) $\lambda_{\max}$ (log ϵ) 226 (4.29), 262 (3.95) nm; IR (film) $\nu_{\max}$ 3318, 2956, 1649, 1619, 1550, 1458, 1203, 1139, 1086, 1026, 838, 721 cm^{−1}; ¹H and ¹³C NMR data, see Table 1; HR-ESI-MS *m/z* 812.4775 [M+H]⁺ (calcd for C₃₉H₆₂N₁₁O₈, 812.4783)

Paltolide B (2): colorless amorphous powder; [α]_D²³ −71 (*c* 0.2, MeOH); UV (MeOH) $\lambda_{\max}$ (log ϵ) 226 (4.88), 260 (4.58) nm; IR (film) $\nu_{\max}$ 3321, 2960, 1649, 1641, 1553, 1468, 1203,

1183, 1139 cm^{-1} ; ^1H and ^{13}C NMR data, see Table 1; HR-ESI-MS m/z 842.4880 $[\text{M}+\text{H}]^+$ (calcd for $\text{C}_{40}\text{H}_{64}\text{N}_{11}\text{O}_9$, 842.4888).

Paltolide C (3): colorless amorphous powder; $[\alpha]_D^{23} -19$ (c 0.07, MeOH); UV (MeOH) λ_{max} (log ϵ) 226 (4.39), 260 (4.08) nm; IR (film) ν_{max} 3318, 2965, 1669, 1637, 1548, 1458, 1204, 1135 cm^{-1} ; ^1H and ^{13}C NMR data for Lys, Ala, Leu, *N*MeLeu, urea, and Arg are identical to those reported for **2** in Table 1; ^1H NMR (CD_3OD , 500 MHz) 6'-BrTrp: δ 4.72 (1H, m, H-2d), 3.50 (1H, dd, J = 14.4, 3.6 Hz H-3a), 2.96 (1H, overlapped, H-3b), 6.77 (1H, s, H-2'), 7.54 (1H, d, J = 8.4 Hz, H-4'), 7.11 (1H, dd, J = 8.2, 1.3 Hz, H-5'), 7.46 (1H, d, J = 1.3 Hz, H-7'), 9.01 (1H, d, J = 8.5 Hz, NH); ^{13}C NMR (CD_3OD , 125 MHz) 6'-BrTrp: δ 174.7 (C-1), 29.4 (C-2), 125.5 (C-2'), 112.0 (C-3'), 120.5 (C-4'), 122.9 (C-5'), 115.5 (C-6'), 115.0 (C-7'), 138.4 (C-8'), 127.4 (C-9'); HR-ESI-MS m/z 904.4030 $[\text{M}+\text{H}]^+$ (calcd for $\text{C}_{40}\text{H}_{63}\text{BrN}_{11}\text{O}_8$, 904.4044).

LC/MS Analysis of L/D-FDLA Derivatives

Approximately 0.4 mg of compounds **1–3** were separately hydrolyzed with 5 N HCl (LabChem Inc, traceable to NIST) (0.8 mL) in an Ace high pressure tube for 16 h at 90 °C, dried, and dissolved in H_2O (100 μL). To a 50 μL aliquot of each was added 1 N NaHCO_3 (20 μL) and 1% 1-fluoro-2,4-dinitrophenyl-5-L-leucinamide (L -FDLA or D-FDLA solution in acetone, 100 μL), and the mixtures were heated to 40 °C for 40 min, allowed to cool to rt, neutralized with 2 N HCl (20 μL), and evaporated to dryness. Residues were dissolved in CH_3CN and analyzed by LC-MS. Analyses of the L- and L/D-FDLA (mixture of D- and L-FDLA) derivatives were performed using a Phenomenex Jupiter Proteo C12 column (4 μm , 150 $\times$ 4.6 mm) aqueous CH_3CN containing 0.01 % TFA was used as a mobile phase eluting with a linear gradient of 25–70% CH_3CN in 45 min at a flow rate of 0.5 mL/min. An Agilent Series 1100 MSD mass spectrometer was used for detection in negative ESI mode. The fragmentor and capillary voltage were kept at 70 and 1000 V, respectively, and the ion source at 350 °C. A mass range of m/z 100–1000 was scanned in 0.1 min. Retention times (t_R , min) of the FDLA-derivatized amino acids for compounds **1–3**: L-Lys 38.3, D-Lys 38.8 m/z 735 $[\text{M}+\text{H}]^-$ (bis derivative); L-Trp 30.0, D-Trp 32.8 m/z 497 $[\text{M}-\text{H}]^-$; L-*N*MeLeu 32.8, D-*N*MeLeu 35.5 m/z 438 $[\text{M}-\text{H}]^-$; L-Ala 24.5, D-Ala 27.0 m/z 382 $[\text{M}-\text{H}]^-$; L-Leu 30.6, D-Leu 36.8 m/z 424 $[\text{M}-\text{H}]^-$; L-Arg 15.9, D-Arg 13.9 m/z 467 $[\text{M}-\text{H}]^-$.

Biological assays

Cytotoxicity assays were carried out using an MTT cell proliferation assay kit (American Type Culture Collection) according to the instructions provided. Briefly, HCT-116, BSC-1, or TZM-BL cells were seeded in 96-well tissue culture plates at a density of 2×10^4 cells/well in 50 μL of growth media and allowed to adhere for 18 h. Attached cells were incubated with inhibitors for 24 h (as controls for the neutralization assay), after which time the media was either replaced or diluted 3-fold with fresh growth media. Following an additional 48 h incubation period, cell viability was assessed upon treatment with MTT (A_{570} , Molecular Devices 96-well absorbance plate reader). Single round HIV-1 neutralization assays were performed with viruses pseudotyped with SF162 Envelope using published conditions.¹⁵ At least two separate experiments were performed for each assay, and all assays were performed in duplicate.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank H. Baker for HIV neutralization assays and the AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH, for reagents and cell lines used therein; and the Republic of Palau for collecting permits. This

work was supported in part by the NIH Intramural Research Program (NIDDK) and the Intramural AIDS Targeted Antiviral Program, Office of the Director, NIH (C.A.B.).

References and Notes

1. (a) Piel J, Hui D, Wen G, Butzke D, Platzer M, Fusetani N, Matsunaga S. *Proc Natl Acad Sci USA* 2004;101:16222–16227. [PubMed: 15520376] (b) Matsunaga S, Fusetani N. *Curr Org Chem* 2003;7:945–966. (c) Bewley CA, Faulkner DJ. *Angew Chem Int Ed Engl* 1998;37:2162–2178.
2. (a) Clark DP, Carroll J, Naylor S, Crews P. *J Org Chem* 1998;63:8757–8764. (b) Bewley CA, Faulkner DJ. *J Org Chem* 1994;59:4849–4852. (c) Schmidt EW, Bewley CA, Faulkner DJ. *J Org Chem* 1998;63:1254–1258.
3. Ford PW, Gustafson KR, McKee TC, Shigematsu N, Maurizi LK, Pannell LK, Williams DE, Dillip de Silva E, Lassota P, Allen TM, Van Soest R, Andersen RJ, Boyd MR. *J Am Chem Soc* 1999;121:5899–5909.
4. Hamada T, Matsunaga S, Yano G, Fusetani N. *J Am Chem Soc* 2005;127:110–118. [PubMed: 15631460]
5. Bjoerquist P, Buchanan M, Campitelli M, Carroll A, Hyde E, Neve J, Polla M, Quinn R. *PCT Int Appl* 2005 WO 2005039617.
6. (a) Fujii K, Ikai Y, Oka H, Suzuki M, Harada K-I. *Anal Chem* 1997;69:5146–5151. (b) Fujii K, Ikai Y, Mayumi T, Oka H, Suzuki M, Harada K-I. *Anal Chem* 1997;69:3346–3352.
7. (a) Kobayashi J, Sato M, Murayama T, Ishibashi M, Walchi MR, Kanai M, Shoji J, Ohizumi Y. *J Chem Soc, Chem Commun* 1991:1050–1052. (b) Shibashi M, Li Y, Sato M, Kobayashi J. *Nat Prod Lett* 1994;4:293–296.
8. Robinson SJ, Tenney K, Yee DF, Martinez L, Media JE, Valeriote FA, van Soest RWM, Crews P. *J Nat Prod* 2007;70:1002–1009. [PubMed: 17559267]
9. Reshef V, Carmeli S. *J Nat Prod* 2002;65:1187–1189. [PubMed: 12193029]
10. Grach-Pogrebinsky O, Carmeli S. *Tetrahedron* 2008;64:10233–10238.
11. (a) Kobayashi J, Sato M, Ishibashi M, Shigemori H, Nakamura T, Ohizumi Y. *J Chem Soc, Perkin Trans 1* 1991:2609–2611. (b) Uemoto H, Yahiro Y, Shigemori H, Tsuda M, Takao T, Shimonishi Y, Kobayashi J. *Tetrahedron* 1998;54:6719–6724.
12. Schmidt EW, Harper MK, Faulkner DJ. *J Nat Prod* 1997;60:779–782.
13. Gresner-Apter S, Carmeli S. *J Nat Prod* 2009;72:1429–1436. [PubMed: 19650639]
14. Fisch KM, Gurgui C, Heycke N, van der Sar SA, Anderson SA, Webb VL, Taudien S, Platzer M, Rubio BK, Robinson SJ, Crews P, Piel J. *Nat Chem Biol* 2009;5:494–500. [PubMed: 19448639]
15. Gustchina E, Bewley CA, Clore GM. *J Virol* 2008;82:10032–10041. [PubMed: 18667502]

Table 1

NMR Spectroscopic Data for Paltolides A (1) and B (2) (CD₃OD).

		1		2		
position	δ_C^a	δ_H^b (J in Hz)	HMBC ^c	δ_C^a	δ_H^b (J in Hz)	HMBC ^c
Lys						
1	175.8			175.8		
2	56.0	4.08, t (4.7)	1, 3, 4, 1 _{Urea}	56.1	4.13, br t (4.8)	1, 3, 4, 1 _{Urea}
3	32.4	1.84, 1.73, m	2	32.5	1.92, 1.68, m	2
4	22.5	1.64, 1.41, m		21.5	1.58, 1.32, m	
5	30.0	1.56, 1.50, m		28.8	1.58, m	
6	40.1	3.51, 3.11, m	4, 5, 1 _{Trp}	39.6	3.79, 2.93, m	4, 5, 1 _{5'-OHTrp}
NH-6					7.95, br d (7.6)	1 _{5'-OHTrp}
Ala						
1	176.0			175.4		
2	51.4	4.20, q (6.8)	1, 3, 1 _{Lys}	50.7	4.26 ^d	1, 3, 1 _{Lys}
3	17.3	1.39, d (6.8)	1, 2	17.1	1.48, d (6.8)	1, 2
Leu1						
1	176.0			175.4		
2	54.5	4.12, dd (8.5, 6.5)	1, 3, 4, 1 _{Ala}	49.6	4.92, dd (11.9, 2.8)	3, 4
3	41.7	1.54, m	1, 2, 5, 6	41.4	1.80, 1.32, m	1, 2, 4, 5, 6
4	25.9	1.69, m	3, 5, 6	26.3	1.94, m	3, 5, 6
5	23.3	0.99, d (6.6)	3, 4, 6	23.8	1.03, d (6.8)	3, 4, 6
6	22.2	0.96, d (6.6)	3, 4, 5	22.4	1.01, d (6.8)	3, 4, 5
NH					8.71, br s	1 _{Ala}
Leu2						
1	174.4			172.1		
2	55.7	3.87, dd (10.4, 4.7)	1, 3, 4, 1 _{Leu1}	60.0	4.95, t (7.3)	1, 3, 4, NMe, 1 _{Leu1}
3	39.5	1.75, 1.39, m	1, 2, 5, 6	38.3	1.77, 1.30, m	1, 2, 4, 5, 6
4	26.0	1.49, m	3, 5, 6	25.8	1.47, m	3, 5, 6
5	23.4	0.86, d (6.5)	3, 4, 6	23.7	0.95, d (6.4)	3, 4, 6

1			2			
position	δ_C^a	δ_H^b (<i>J</i> in Hz)	HMBC ^c	δ_C^a	δ_H^b (<i>J</i> in Hz)	HMBC ^c
6	21.5	0.82, d (6.5)	3, 4, 5	23.7	0.93, d (6.4)	3, 4, 5
NMe				29.0	1.89, s	2, 1 _{Leu}
		Trp			5'-OHTrp	
1	173.7			174.7		
2	56.1	4.57, dd (9.8, 4.0)	1, 3, 5, 1 _{Leu2}	55.5	4.67, m	1, 3, 3', 9', 1 _{N_{MeLeu}}
3a	27.8	3.48, dd (14.6, 4.0)	1, 2, 2', 3', 9'	30.1	3.44, dd (15.0, 2.8)	1, 2, 2', 3', 9'
3b		3.20 ^d	1, 2, 2', 3', 9'		2.92 ^d	1, 2, 2', 3', 9'
NH-1'					9.96, s	
2'	124.5	7.11, s	3, 3', 8', 9'	125.6	6.70, s	2, 3, 3', 4', 8', 9'
3'	111.5			110.4		
4'	119.1	7.57, br d (8.1)		103.5	6.98, d (1.9)	2', 5', 7', 8'
5'	119.7	7.00, t (7.3)	3', 6', 7', 8', 9'	150.9		
6'	122.3	7.07, t (7.3)	6', 7', 9'	112.6	6.63, dd (8.8, 1.9)	4', 5', 8'
7'	112.3	7.31, br d (8.1)	4', 8'	112.7	7.12, d (8.8)	4', 5', 9'
8'	138.0		4', 5', 9'	132.6		
9'	128.6			128.5		
NH					8.97, d (8.5)	2, 3, 1 _{N_{MeLeu}}
		Urea			Urea	
1	160.1			160.1		
		Arg			Arg	
1	176.0			176.3		
2	53.6	4.32, m	1, 3, 4, 1 _{Urea}	54.3	4.25, m	1, 3, 4, 1 _{Urea}
3	30.7	1.94, 1.70, m	2, 4	30.4	1.94, 1.70, m	2, 4
4	26.2	1.72, 1.65, m	5	26.0	1.72, m	5
5	41.1	3.25, 3.21, m	3, 4, 6	41.9	3.23, m	3, 4, 6
6	158.3			158.5		

^aRecorded at 125 MHz; referenced to residual CD₃OD at δ 49.1 ppm.

^bRecorded at 500 MHz; referenced to residual CD₃OD at δ 3.30 ppm.

^cProton showing HMBc correlation to indicated carbon.

^dOverlaped signal.