Published in final edited form as:

Neuroreport. 2010 July 14; 21(10): 675–679. doi:10.1097/WNR.0b013e32833add6e.

Illusory force perception following a voluntary limb movement

Carl P.T. Jackson^{1,2,*} and R. Chris Miall²

¹Centre for Neuroscience Studies, Queen's University

²Behavioural Brain Sciences, School of Psychology, University of Birmingham

Abstract

We present a novel illusion in which participants report constant forces on their hand as steadily increasing. Participants made discrete reaching movements perturbed by a lateral force that increased with distance moved; when stationary at the end of the movement, a true constant force was perceived to increase. We tested perceived subjective equality by increasing or decreasing the force. The illusion was significantly stronger when the perturbation was applied during active movement. We conclude that the unusual context of moving against lateral spring-forces results in participants failing to predict steady lateral forces at the end of their movement, and causes an illusion of increasing forces even after movement termination. This result further emphasizes the role of action-prediction in sensory perception.

Keywords

human; motor control; arm; illusion; force; psychophysics; sensorimotor

Introduction

When we make a reaching movement, it is important that our hand gets to the right destination. To accomplish this, the motor system uses a combination of feedforward prediction and visual/proprioceptive feedback [1,2,3]. Movement planning takes account of the context of the movement [4,5,6] to specify the initial feedforward motor commands to the limb [2]. Visual and proprioceptive feedback from the ongoing action is used to correct for perturbations in limb trajectory [7,8].

It is unclear how the motor system behaves in a novel context, when it has no experience of a particular set of sensory consequences given a certain set of motor commands. Many studies have been done on reaching in both stable and unstable force fields [9,10,11,12] – but these reaching studies generally involve adapting to contexts that the motor system is used to handling. A *completely* novel context is rare, and as such it is assumed that the motor system would be unable to accurately predict the sensory consequences of an action in this case.

Here we report investigations into a newly discovered sensory illusion that takes place in such a novel context. Participants made a movement in a position-dependent field in which

*Corresponding author Centre for Neuroscience Studies Botterell Hall, Rm. 234 Queen's University Kingston, ON K7L 3N6 CANADA carl@biomed.queensu.ca Tel: +1(613) 533-6000 ext. 74924 Fax: +1(613) 533-6840.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

an increasing force was applied to their arm perpendicular to their movement direction. At the end of the movement they had to stabilise their arm against a maintained constant force level. Under these conditions we found a surprising perceptual illusion: the force on the limb at the end of the movement appeared to increase over the course of a few seconds, but crucially the illusion only seemed to appear after a brisk reaching movement.

To quantify this illusion using the principle of perceptual constancy, we imposed a force ramp (gradient) at the end of each reaching movement that increased or decreased over time. From this we constructed a psychophysical function using a two-alternative forced choice paradigm to find force ramps that would cancel the participants' impression of an increasing force. We hypothesized from our pilot data that we would see a bias toward participants reporting static forces as 'increasing', thus shifting the psychophysical function. We also included another condition designed to test whether it was the active movement that was important or just the impact of the force on a stationary arm. We also hypothesized that the illusion was at least partly due to the unfamiliar context of a lateral force impacting on a forward reaching movement and thus we conducted a follow-up experiment that aimed to explore the effect of force and target direction on its presentation.

Materials and Methods

Participants

10 participants took part in the experimental condition (3 male, 7 female, age range 19 to 37 years, median age 25.1 years, one left-handed) and 10 different participants took part in the control condition (all female, age range 18 to 31 years, median age 22.7 years, two left-handed). 13 further participants took part in the follow-up experiment (8 male, 5 female, age range 19 to 50 years, median age 29 years, four left-handed). All gave their informed consent. None of the participants had any known motor or sensory abnormalities, and all were normal healthy adults with normal or corrected-to-normal vision. The experiment was approved by the local ethics committee at the University of Birmingham.

Apparatus and materials

Participants made reaching movements using their right hand on a vBOT robotic manipulandum [13]. They viewed a 0.5 cm red circular cursor showing the position of their hand using a projection-mirror system so that the cursor appeared to be in the same plane as their hand. Responses were gathered using the right foot on a pair of foot pedals.

Procedure

In the movement condition (Figure 1A), participants moved the red cursor to a white starting circle at the bottom of the screen at the start of each trial. After 500 ms a blue target appeared 16 cm in front of the white starting circle and participants were instructed to move to the target within 500 ms. If they did so the target turned yellow, otherwise it turned red (movement too quick) or green (movement too slow). Along the way to the target, participants received a position-dependent force that pushed their arm to the right with force constant 0.72 N/cm measured along the axis from start to target, peaking at the target position at $f_{\text{Max}} = 11.52 \text{ N}$. Participants were required to counteract this force and stabilise their hand at the target. As soon as the target changed colour, participants experienced one of eleven force ramps over 1000 ms ranging from -50% (decreasing from 11.52 to 5.76N) to +50% (increasing from 11.52 to 17.28N) and including a 0% constant force. Force ramps were randomly interleaved from trial to trial. After 1000 ms, the target returned to its original blue colour and the ramp continued while participants had a further 500 ms to respond with the foot pedal as to whether the force was 'increasing' (toe) or 'decreasing' (heel). After this further 500 ms, if participants had not made a response, the ramp was

turned off and the trial was reshuffled into the remaining trials. 220 trials were performed in total.

In the static condition (Figure 1B), participants moved to the white starting circle as in the movement condition. Another white circle then appeared 16 cm in front of the first one and the participants were instructed to move to this circle. There were no imposed forces during their movement. When they had reached it, the target turned blue and a rightward force that increased over 500 ms with a minimum-jerk profile [14] was imposed on their static arm, again peaking at 11.52 N. Participants were again required to stabilise their hand at the target and resist this force. Immediately after the 500ms force profile, the target turned yellow and the trial proceeded in the same way as in the movement condition with increasing or decreasing force ramps and a response by the participant.

In the follow-up experiment, the same protocol as the movement condition was followed except that the target could now be in one of four directions (forward, backward, left and right with respect to the start position; left and right flipped during analysis for left-handed participants) and the force could also act in one of these four directions. Thus sometimes the force was assistive, sometimes resistive and sometimes orthogonal to the movement direction. Participants performed 25 trials in each combination of target/force directions in a single block; as before, the gradients were randomized across trials. Blocks were randomised and participants were informed before each block what the target and force direction would be. Each participant received 400 trials in total.

Results

We hypothesized from pilot data that if participants perceived an increasing force at the end of their movement even when the force was constant then a) the psychometric function should be left shifted to reflect this bias and b) the number of 'increasing' responses at constant force should be significantly greater than chance. Figure 2 (solid line) shows the psychometric function formed from 'increasing' responses vs. gradient, and this function is indeed left shifted. Furthermore, the mean percentage of 'increasing' responses at constant force was 74.0% (SD: 13.9%), which was significantly greater than chance [$t(9) = 5.46$, $p < 0.001$]. We have used t-tests rather than d' values throughout, because at constant force the gradient was neither increasing nor decreasing so there was no correct response.

We wondered whether the illusion demonstrated in the movement condition was simply caused by an unexpected force on the hand rather than a consequence of the predictive movement system. To control for this possibility, we performed a static condition in which participants made similar judgements as before, but this time forces were only applied while their hand was still. Figure 2 (dashed line) shows the psychometric function plotting 'increasing' responses vs. gradient in the static condition; the function is still left shifted, but less than in the movement condition. Furthermore, the mean percentage of 'increasing' responses at constant force was only 60.0% (SD: 14.5%), which was not significantly greater than chance [$t(9) = 2.18$, $p = 0.06$], although there is a trend towards significance. However, a two-tailed, two-sample t-test on the percentage of 'increasing' responses at constant force across the two groups showed that the movement group responded 'increasing' significantly more than the static group [$t(18) = 2.20$, $p = 0.04$, Cohen's $d = 1.04$] and thus it seems clear that the illusion is stronger when moving through a novel movement-dependent force field than when an external force is applied to the static hand.

We were concerned that participants might have had an underlying bias toward saying 'increasing' rather than 'decreasing', since in the movement condition participants correctly identified the largest positive gradient as 'increasing' 90% of the time (91% in the static

condition) but reported the largest negative gradient as 'decreasing' only 78.5% (77.5%) of the time. The fact that the force increased as participants moved may have caused this disparity. To correct for it, we shifted the entire curve down until the percentages of correct responses at the extremes were equal. After this correction, the mean percentage of 'increasing' responses at constant force in the movement condition was reduced from 74.0 to 68.3% (SD: 17.8%), but was still significantly greater than chance [$t(9) = 3.24, p = 0.01$]. In the static condition, the correction reduced the mean percentage of 'increasing' responses at constant to 53.0% (SD: 9.1%), which was still not significantly greater than chance [$t(9) = 1.04, p = 0.33$]. A two-tailed, two-sample t-test between the two groups still showed a difference [$t(18) = 2.41, p = 0.02$, Cohen's $d = 1.14$].

Discussion

In the experiments presented, we found that participants experienced a perceptual illusion of increasing force when they moved their arm rapidly to a target in the presence of a laterally perturbing force field. The illusion was absent or much reduced when the force was presented following the movement rather than concurrent with it.

What is the source of these kinds of perceptual illusions? While visual illusions are well known (see for example [15]) it is less common to find illusions in other sensory systems – perhaps because vision is our dominant sense. One recent example of an illusory force that is similar to our illusion comes from Diedrichsen and colleagues [16]. In their study, participants supported an object with one hand and lifted it with the other. When the participants were presented with a force on their supporting hand, ostensibly from the object but that persisted after lifting, they reported that the force was perceived to be increasing when in reality it was constant. The authors demonstrated that prediction errors and visual information both contributed to the maintenance of the illusion. We cannot entirely rule out an effect of vision in our experiment. Opposing the lateral forces in order to maintain the cursor on-target implies that participants used visual guidance even after movement completion. However preliminary data, not reported here, show that the illusion is also present in the absence of vision.

As in their study it seems likely that requiring the participants to oppose a novel, movement-related force was the primary cause of the effect. When we move our arms we are usually able to predict the sensory consequences of our action extremely well, since we have had lifelong experience in this particular arena. However, when these predictions are erroneous (in this case when the external force on the arm is applied in an unusual direction, atypical of normal object interactions) this mismatch between prediction and feedback might cause an illusory sensation. Note however that our illusion differs from the one described by Diedrichsen and colleagues [16] in that it has its effect in a single hand only.

Furthermore our results from the static condition suggest that an important factor causing the illusion is reaching in the unusual force field. When participants made an unperturbed movement, stopped, and then had the force impact on their hand the illusion was much reduced – it only appeared consistently when the force impacted *during* the movement. This important difference is further evidence for the role of action prediction in the illusion, specifically the prediction of the consequences of the action from the efference copy of the motor command [3,17]. Presumably the reduction of the illusion in the static condition as the prediction that the arm will come to a halt with zero external force is fulfilled. When the force impacts on the stationary arm, the participant is now performing a postural stability task and the force is perceived to be an external event unrelated to the previous action. Another possible explanation is that sensory attenuation, known to occur during active

movement (e.g. [18]) is gradually reduced following the reach, leading to the false percept of increasing force.

One interesting point is the atypical relationship between movement direction and imposed force, which tends to deviate the arm from its intended trajectory. We predicted that the illusion would therefore be greatest when forces were orthogonal to movement direction. Preliminary results from a follow up study that tests the presence of the illusion in different movement orientations and for different force directions indicate that the illusion is strongest when the force on the limb pushes it to the left or right with respect to the trunk, rather than forward or backward. When we reach in an unstable environment, the motor system must selectively control the impedance geometry – i.e. the stiffness of the limb – to stabilise the arm [19]. While participants tend to increase their stiffness depending on the direction of instability in the environment [20], there is an inherent variation in stiffness in the neuromuscular system across different arm orientations. Our result, shown in Table 1, supports the idea that the illusion is affected by the inherent stability of the limb, perhaps because we are able to deal with perturbations in the forward/backward axis more easily than the left/right direction.

Conclusion

In conclusion, we have demonstrated the existence of a novel force illusion that presents as the apparent increase in perceived force at the end of a movement when in fact the force on the limb is constant. Our identification of a perceptual sensory illusion operating within a single limb, and that is apparently not dependent on visuo-motor discrepancies, is therefore novel. The illusion does not appear as strongly when the force impacts on the stationary limb, and may be due to the mismatch between the predicted and actual outcome of voluntary movement, or the reduction of sensory attenuation following a movement. Our hypothesis that the illusion is strongest when the force perturbation is applied orthogonal to the movement direction (which is unusual in everyday experience) was only partly supported and instead there is evidence – albeit weak – that the illusion may be modulated by the anisotropic stiffness of the arm in different reaching movements.

Acknowledgments

This work was supported by the Wellcome Trust [GR069439].

Reference List

1. Kawato M. Internal models for motor control and trajectory planning. *Curr Opin Neurobiol.* 1999; 9:718–727. [PubMed: 10607637]
2. Miall RC, Wolpert DM. Forward models for physiological motor control. *Neural Netw.* 1996; 9:1265–1279. [PubMed: 12662535]
3. Wolpert DM, Ghahramani Z. Computational principles of movement neuroscience. *Nat Neurosci.* 2000; 3:1212–1217. [PubMed: 11127840]
4. Vetter P, Wolpert DM. Context estimation for sensorimotor control. *J Neurophysiol.* 2000; 84:1026–1034. [PubMed: 10938325]
5. Blakemore SJ, Goodbody SJ, Wolpert DM. Predicting the consequences of our own actions: The role of sensorimotor context estimation. *J Neurosci.* 1998; 18:7511–7518. [PubMed: 9736669]
6. Haruno M, Wolpert DM, Kawato M. MOSAIC model for sensorimotor learning and control. *Neural Comput.* 2001; 13:2201–2220. [PubMed: 11570996]
7. Bagesteiro LB, Sarlegna FR, Sainburg RL. Differential influence of vision and proprioception on control of movement distance. *Exp Brain Res.* 2006; 171:358–370. [PubMed: 16307242]

8. Scheidt RA, Conditt MA, Secco EL, Mussa-Ivaldi FA. Interaction of visual and proprioceptive feedback during adaptation of human reaching movements. *J Neurophysiol.* 2005; 93:3200–3213. [PubMed: 15659526]
9. Franklin DW, Osu R, Burdet E, Kawato M, Milner TE. Adaptation to stable and unstable dynamics achieved by combined impedance control and inverse dynamics model. *J Neurophysiol.* 2003; 90:3270–3282. [PubMed: 14615432]
10. Milner TE, Franklin DW. Impedance control and internal model use during the initial stage of adaptation to novel dynamics in humans. *J Physiol.* 2005; 567:651–664. [PubMed: 15961421]
11. Lackner JR, DiZio P. Rapid adaptations of coriolis force perturbations of arm trajectory. *J Neurophysiol.* 1994; 72:299–313. [PubMed: 7965013]
12. Shadmehr R, Mussa-Ivaldi FA. Adaptive representation of dynamics during learning of a motor task. *J Neurosci.* 1994; 14:3208–3224. [PubMed: 8182467]
13. Howard IS, Ingram JN, Wolpert DM. A modular planar robotic manipulandum with end-point torque control. *J Neurosci Methods.* 2009; 181:199–211. [PubMed: 19450621]
14. Flash T, Hogan N. The coordination of arm movements - an experimentally confirmed mathematical-model. *J Neurosci.* 1985; 5:1688–1703. [PubMed: 4020415]
15. Changizi MA, Hsieh A, Nijhawan R, Kanai R, Shimojo S. Perceiving-the-present and a systematization of illusions. *Cogn Sci.* 2008; 32:459–503. [PubMed: 21635343]
16. Diedrichsen J, Verstynen T, Hon A, Zhang Y, Ivry RB. Illusions of force perception: the role of sensori-motor predictions, visual information, and motor errors. *J Neurophysiol.* 2007; 97:3305–3313. [PubMed: 17344369]
17. Wolpert DM, Flanagan JR. Motor prediction. *Curr Biol.* 2001; 11:R729–732. [PubMed: 11566114]
18. Shergill SS, Bays PM, Frith CD, Wolpert DM. Two eyes for an eye: the neuroscience of force escalation. *Science.* 2003; 301:187. [PubMed: 12855800]
19. Burdet E, Osu R, Franklin DW, Milner TE, Kawato M. The central nervous system stabilizes unstable dynamics by learning optimal impedance. *Nature.* 2001; 414:446–449. [PubMed: 11719805]
20. Franklin DW, Liaw G, Milner TE, Osu R, Burdet E, Kawato M. Endpoint stiffness of the arm is directionally tuned to instability in the environment. *J Neurosci.* 2007; 27:7705–7716. [PubMed: 17634365]

Fig. 1.

Experimental procedure. **a** Schematic of procedure for movement condition showing the bell-shaped velocity profile of a typical forward reaching movement, and the position dependent lateral force imposed on the arm. At the end of the reaching movement one of 11 increasing or decreasing force ramps was applied. **b** Similar schematic for static condition but showing that the force onset was after movement completion.

Fig. 2. Psychophysical response functions from movement (solid lines) and static (dashed lines) conditions showing the percentage of 'increasing' responses against each force gradient. Note the significant difference between movement and static at constant force. Each point in the group mean ± 1 S.E.M.

Table 1

Percentage of 'increasing' responses at constant force during movements and force perturbations applied in 4 directions. The illusion was apparent in all target directions and when the force was applied in the left and right directions, but not forwards and backwards. The 4×4 ANOVA of force by target direction showed a main effect of force direction [$F(3,192) = 2.85$, $p = 0.039$] but no effect of target direction and no interaction. Statistics in the table are based on uncorrected t-tests between the percentage of 'increasing' responses for each force or target direction vs. chance (50%).

	Forward	Backward	Left	Right
Force direction	54.2%	56.9%	65.0% **	66.2% ***
Target direction	61.2% **	59.6% *	58.1% *	63.5% **

* $p < 0.05$

** $p < 0.01$

*** $p < 0.001$