Published in final edited form as:

AIDS. 2010 June 1; 24(9): 1281–1290. doi:10.1097/QAD.0b013e328339e228.

Microbial translocation induces persistent macrophage activation unrelated to HIV-1 levels or T cell activation following therapy

Mark A. Wallet,

Department of Pathology, Immunology and Laboratory Medicine, University of Florida, Gainesville, FL, USA

Carina A. Rodriguez,

Department of Pediatrics, Division of Allergy Immunology and Rheumatology, University of South Florida and All Children's Hospital, St. Petersburg, FL, USA

Li Yin,

Department of Pathology, Immunology and Laboratory Medicine, University of Florida, Gainesville, FL, USA

Sara Saporta,

Department of Pediatrics, Division of Allergy Immunology and Rheumatology, University of South Florida and All Children's Hospital, St. Petersburg, FL, USA

Sasawan Chinratanapisit,

Department of Pediatrics, Division of Allergy Immunology and Rheumatology, University of South Florida and All Children's Hospital, St. Petersburg, FL, USA

Wei Hou,

Division of Biostatistics, Department of Epidemiology and Health Policy Research, University of Florida, Gainesville, FL, USA

John W. Sleasman, and

Department of Pediatrics, Division of Allergy Immunology and Rheumatology, University of South Florida and All Children's Hospital, St. Petersburg, FL, USA

Corresponding author and reprint requests: Maureen M. Goodenow, PhD, Department of Pathology, 2033 Mowry Road, Room 280, Gainesville, FL 32610, goodenow@ufl.edu.

J.W.S. and M.M.G. contributed equally to this work

Conflicts of interest: None

Author Contributions: **Mark A. Wallet** conceived and designed the study, performed most experiments, collected and analyzed data and wrote the manuscript.

Carina A. Rodriguez enrolled patients and performed flow cytometry to assess T cell activation and contributed to writing and revising the manuscript.

Li Yin performed TREC assays and contributed to writing and revising the manuscript.

Sara Saporta performed EndoCAB ELISA.

Sasawan Chinratanapisit performed sCD14 ELISA.

Wei Hou performed statistical analyses and contributed to revising the manuscript.

John W. Sleasman enrolled patients, oversaw the conception and study design and contributed to writing and revising the manuscript.

Maureen M. Goodenow oversaw the entire project including conception and study design, patient enrolment, specimen archiving, clinical database management and contributed to writing and revising the manuscript.

List of Supplemental Digital Content

Supplemental Digital Content 1. doc

Supplemental Digital Content 2. doc

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Maureen M. Goodenow

Department of Pathology, Immunology and Laboratory Medicine, University of Florida, Gainesville, FL, USA

Abstract

Objective—HIV-1 replication and microbial translocation occur concomitant with systemic immune activation. This study delineates mechanisms of immune activation and CD4 T cell decline in pediatric HIV-1 infection.

Design—Cross-sectional and longitudinal cellular and soluble plasma markers for inflammation were evaluated in 14 healthy and 33 perinatally HIV-1-infected pediatric subjects prior to and over 96 weeks of protease-inhibitor-containing combination antiretroviral treatment [ART]. All HIV-1-infected subjects reconstituted CD4 T cells either with suppression of viremia or rebound of drug-resistant virus.

Methods—Systemic immune activation was determined by polychromatic flow cytometry of blood lymphocytes and ELISA for plasma soluble CD27 [sCD27], soluble CD14 [sCD14], and tumor necrosis factor [TNF]. Microbial translocation was evaluated by limulus amebocyte lysate assay to detect bacterial lipopolysaccharide [LPS] and ELISA for anti-endotoxin core antigen IgM antibodies. Immune activation markers were compared to viral load, CD4% and LPS by regression models. Comparisons between healthy and HIV-1 infected or between different viral outcome groups were performed by non-parametric rank sum.

Results—Microbial translocation was detected in healthy infants but resolved with age ($P<0.05$). LPS and sCD14 levels were elevated in all HIV-1 infected subjects ($P<0.05$ and $P<0.0001$, respectively) and persisted even if CD4 T cells were fully reconstituted, virus optimally suppressed, and lymphocyte activation resolved by ART. Children with CD4 T cell reconstitution but viral rebound following ART continued to display high levels of sCD27.

Conclusions—Microbial translocation in pediatric HIV-1-infection is associated with persistent monocyte/macrophage activation independent of viral replication or T cell activation.

Keywords

HIV-1; microbial translocation; monocyte/macrophage; systemic immune activation; children and adolescents; lipopolysaccharide; inflammation

Introduction

HIV-1 infection develops with acute viremia and rapid depletion of CD4 T cells within mucosal-associated lymphoid tissues [MALT], particularly in gut lymphoid compartments [1-3]. HIV-1-induced disruption of MALT results in intestinal fibrosis and translocation of microbial products across the intestinal mucosa into the peripheral circulation producing high plasma levels of lipopolysaccharide [LPS] and bacterial DNA[4-6] that persist throughout chronic HIV-1 infection[7]. Microbial translocation appears related to systemic leukocyte activation and elevated plasma levels of inflammatory proteins[5,8,9], although whether or not microbial translocation and consequent inflammation are direct causes of progression to AIDS or merely sequelae of HIV-1 infection seems to differ among cohorts[5,7-9].

Chronic HIV-1 infection is characterized by progressive activation of CD4 and CD8 T cells, as well as CD14 monocyte/macrophages[5,10-14], and gradual depletion of peripheral naïve and memory CD4 T cells. Microbial translocation is implicated in systemic T cell activation associated with HIV-1 infection[8] and idiopathic CD4 lymphocytopenia[15], providing evidence that circulating LPS or other microbial products may contribute to acquired and/or

primary immunodeficiency. The premise that microbial translocation is a causative factor for systemic T cell activation, CD4 T cell decline, and disease progression is supported by studies of natural SIV infection of sooty mangabeys that develop high levels of viral replication in the absence of systemic T cell activation, CD4 T cell decline, or microbial translocation[16-18].

In HIV-1 infection, systemic immune activation drives viral replication generating a vicious cycle of viremia, immune activation, and CD4 T cell attrition that ultimately results in AIDS [14]. Effective combination antiretroviral treatment [ART] rapidly suppresses viremia, restores peripheral CD4 T cells, reduces lymphocyte activation, and restores most immune functions [19]. However, ART fails to fully restore memory T cells in MALT[20,21] or completely reverse microbial translocation even following complete suppression of viremia[5]. A unique group of individuals receiving ART reconstitute T cell immunity but fail to control viral replication due to the development of drug resistant virus[22]. This discordant outcome is not uncommon among HIV-1-infected children who develop improved T cell function despite viral rebound[23,24]. Persistent viral replication following ART in conjunction with immune reconstitution in HIV-1-infected children provides an opportunity to study the correlates of pathogenesis, including systemic immune activation and microbial translocation, in the presence of high viral replication in a human host that might be modeled by SIV infection in sooty mangabeys[22-25].

Effects of microbial translocation in HIV-1-infected infants and children may be unique due to inherent differences in gut flora and LPS responses during early development[26-28]. Robust thymic output in younger HIV-1-infected individuals[29] may compensate for CD4 T cell activation and attrition induced by microbial translocation, although systemic activation of monocytes, macrophages, microglia or other cell types potentially contribute to HIV-1-associated illnesses such as neurological disorders, dementia, cardiovascular events or cancer [13,30-32]. Determining the causes and consequences of inflammatory HIV-1 complications is particularly critical for infants and children who may survive, due to effective antiretroviral therapies, for several decades with chronic inflammation.

Methods

Study Design

Prospectively stored plasma and cryopreserved peripheral blood mononuclear cells were obtained from a cohort of HIV-infected infants and children enrolled in a clinical trial of protease inhibitors plus two nucleotide reverse transcriptase inhibitors as previously described. [23,24,33]. Briefly, all HIV-1-infected subjects were infected perinatally and viral and immune outcomes were known. The current study included 33 of the 40 subjects who displayed T cell immune reconstitution and completed 96 weeks on the study. The protocol was approved by University of Florida and University of South Florida/All Children's Hospital Institutional Review Boards. Control subjects were volunteer healthy infants and children with no underlying medical conditions, recruited over the same time frame according to a separate protocol approved by both institutions. Subjects with gastrointestinal conditions were excluded. Whole blood samples were collected, using phlebotomy, in sterile Vacutainer™ (Becton Dickinson) acid citrate dextrose tubes, and processed within 12 hours[34-36]. PBMC and plasma samples were stored at -180° C in liquid nitrogen or at -80° C, respectively, in non-pyrogenic polypropylene cryovials (Nunc Cryotubes™).

Healthy and HIV-1 infected subjects ranged in age from 0.4 to 15 years (median = 1.4; 25th / 75th quartiles = 0.4 / 13.0) and 0.2 to 17.7 years (median = 6.6; 25th / 75th quartiles = 4.5 / 11.6), respectively (P>0.05, Mann-Whitney U test). Viral load, CD4% and CD4 CD45RO% were analyzed at 0, 4, 12, 48 and 96 weeks post-ART. Prior to ART, HIV-1 infected subjects had plasma viral loads ranging from 2.9 to 6.1 log₁₀ HIV-1 RNA copies/ml (median = 4.6

log₁₀). By 24 weeks of ART, all subjects experienced increased CD4% either with viral suppression [<400 viral RNA copies/ml plasma] that was sustained for 96 weeks or viral rebound [>400 viral RNA copies/ml plasma] (24 week median = 3.9 log₁₀ copies/ml; 96 week median = 4.0 log₁₀ copies/ml)[23,24,33].

Measurement of CD4 and CD8 T cell activation

Polychromatic flow cytometry analysis was performed with fluorescence-tagged monoclonal antibodies described previously[29,37] and a multiparameter LSR2 flow cytometer incorporating the cytometer setup and tracking program and Diva software (BD Biosciences, San Jose, CA, USA). Activated effector/memory CD4 T cells were defined as CD3 CD4 cells expressing CD45RO. [38]. Activated CD3 CD8 T cells were defined as CD45RA⁺ CD27⁻ CD11a⁺⁺ (bright), a phenotype associated with HIV-1 disease progression [29,39-41]. The frequency of recent thymic emigrant T cells was determined by performing T cell receptor excision circle [TREC] analysis as previously described[29].

Measurement of soluble markers for immune activation and microbial translocation

Plasma samples were diluted 1:4 in 0.15 M NaCl, heat inactivated at 65° C for 30 minutes, then used to measure soluble CD27 [sCD27] by PeliKine human soluble CD27 ELISA kit (RDI Fitzgerald Industries, Acton, MA, USA); soluble CD14 [sCD14] by Human sCD14 Immunoassay, (R&D Systems, Minneapolis, MN, USA); LPS levels by limulus amebocyte assay [LAL] Chromogenic Endpoint Assay (Lonza Group, Ltd, Allendale, NJ, USA); TNF (TNF superfamily, member 2; TNF α) levels by human TNF ELISA (BD Biosciences); and anti-endotoxin core antigen antibodies [EndoCAb] IgM levels by EndoCAb ELISA (Cell Sciences Inc., Canton, MA, USA). Specimens were handled in an endotoxin-free environment and controls were performed with all LAL assays to rule out endotoxin contamination.

Statistical analysis

Pearson's correlation and simple linear regression analyses were performed to determine the relationships between two variables. Cross-sectional comparisons of median clinical values between different groups of subjects were performed by Mann-Whitney U test. Longitudinal comparison between pre-ART and post-ART values was performed by Wilcoxon matched pairs test. Power analysis was performed based on a one-way ANOVA for three group comparisons. Sample size has an 80% power to identify a minimum detectable effect size of 0.95 S.D. in the group differences. Effects of continuous predictors and their interactions on CD4 percent were measured and tested for significance by using a multiple regression model. Analyses were performed with GraphPad Prism 5.0 software (La Jolla, CA) or SAS software (Cary, NC) and considered significant when P value was less than 0.05.

Results

Pre-therapy CD4 CD45RO T cell frequency and viremia differentially correlate with CD4 T cell decline

Increased frequency of effector/memory CD4 CD45RO T cells, reduced frequency of CD4 CD45RA naïve T cells and activation of CD8 T cells are associated with CD4 T cell decline and progression to AIDS in HIV-1 infection[29,42]. As expected, prior to ART, HIV-1-infected subjects displayed significant inverse correlations between overall CD4 T cell percent and frequency of CD4 CD45RO T cells ($P<0.0001$), frequency of CD8 CD45RA T cells with an activated CD27-CD11a bright phenotype ($P<0.05$), or plasma viral load ($P<0.01$) (Figure 1A, 1C, and 1E). Pre-ART measures of T cell frequency, T cell activation, or viral load were similar between groups who ultimately suppressed [VS] or failed to suppress [VF] viral burden following ART (Figure 1B, 1D, and 1F).

To determine the effects of length of infection [LOI], which was identical to age as all subjects were infected perinatally, a multiple regression model was applied. CD4 decline was related inversely to the frequency of CD4 CD45RO cells, a correlation more pronounced in infants than children and adolescents (Figure 1G, $P < 0.01$). Conversely, a trend in the relationship between viral load and CD4 T cell attrition was more pronounced among older rather than younger children (Figure 1H, $P = 0.097$). These analyses indicated that age/LOI is a determining factor in the relationship between viral load, CD4 T cell activation and CD4 T cell decline.

Microbial translocation develops concomitant with multilineage leukocyte activation in HIV-1 infection

Microbial translocation develops coincident to systemic inflammation in some, but not all HIV-1-infected subjects[5,7,8]. Microbial translocation (LPS) was evident among all therapy naïve HIV-1 infected subjects, as well as a subset of healthy children (Figure 2A, $P < 0.05$). Among healthy subjects, microbial translocation was more pronounced in infants (0-2 years), than in children or adolescents (> 2 -18 years) (Figure 2B, $P < 0.05$). Levels of LPS in HIV-1 infected infants were comparable to healthy infants, while infected children had higher levels than healthy children ($P < 0.01$). The monocyte/macrophage soluble activation marker sCD14 was detected in the plasma of all healthy and HIV-1-infected pediatric subjects. In our study sCD14 levels (median sCD14 = 20.4 ng/ml) were markedly lower in healthy pediatric subjects, including infants who had microbial translocation, than sCD14 levels reported for healthy adults[5,7] (Figure 2B and 2C). In contrast, sCD14 levels were significantly elevated (median sCD14 = 931.8 ng/ml) in HIV-1 infected pediatric subjects compared to healthy subjects (Figure 2C, $P < 0.0001$). EndoCAb levels were significantly reduced in HIV-1-infected subjects (Figure 2D, $P < 0.05$), although unrelated to LPS levels in HIV-1-infected or healthy children (Figure 2E or data not shown). Levels of sCD27, reflecting generalized lymphocyte activation, were elevated in HIV-1-infected subjects compared to healthy controls (Figure 2F, $P < 0.0001$).

The cytokine TNF, a key mediator of systemic inflammation, was undetectable among healthy subjects, but detected in 8 of 33 HIV-1-infected subjects prior to ART (Figure 3A). No difference in LPS levels between TNF(-) and TNF(+) subjects was identified (Figure 3B). Microbial translocation displayed a trend toward a positive correlation with activation of monocyte/macrophages (Figure 3C, $P = 0.055$). In contrast, LPS levels showed no relationship to general lymphocyte activation measured by sCD27 levels or frequency of CD4 CD45RO T cells or activated CD8 T cells (Figure 3D-3F). Together, these results indicate that microbial translocation was independent of lymphocyte activation.

Microbial translocation or systemic inflammation may contribute to CD4 T cell attrition in HIV-1 infection. Prior to ART, plasma levels of LPS were unrelated to CD4 T cell levels (Figure 4A). Likewise, neither sCD14 nor sCD27 levels was associated with CD4% (Figure 4B and 4C). Overall, pre-ART levels of markers for microbial translocation or multilineage leukocyte activation were unrelated to CD4 T cell decline and similar between subjects who developed viral success or failure following ART (Figure 4D, 4E and 4F).

Microbial translocation and monocyte/macrophage activation both persist independent of ART-related CD4 T cell reconstitution

ART results in significant immune reconstitution and improved immune function in individuals who optimally control viral replication, as well as in subjects who develop viral rebound with drug resistant virus[24]. In longitudinal analyses, viral success subjects suppressed viremia below limits of detection by 24 weeks of ART (Figure 5A, $P < 0.0001$) with concomitant improvement in CD4 T cell frequency (Figure 5B, $P < 0.01$). In contrast, viral failure subjects experienced transiently reduced viral loads at 4 weeks post-ART that rebounded by 24 weeks

to an average $>4.0 \log_{10}$ copies per ml (Figure 5A), even though CD4 T cell frequencies improved significantly and persisted at levels $>25\%$ (Figure 5B, $P<0.001$).

Frequency of CD4 CD45RO T cells declined by 24 weeks of ART in both outcome groups to levels approaching those in healthy subjects (Figure 5C). Decline in CD4 CD45RO T cell frequency was significantly associated with increasing T cell receptor excision circles [TREC] (Figure 5D, $P<0.0001$). Plasma sCD27 levels remained unchanged by ART in viral failure subjects, but were reduced in viral success subjects (Figure 5E, $P<0.05$).

Pretherapy levels of sCD14 were unchanged at 24 weeks of ART, but declined by 96 weeks to significantly lower levels in both viral success ($P<0.001$) and viral failure ($P<0.05$) groups. Nonetheless, elevated sCD14 persisted among HIV-1 infected individuals for two years after initiation of ART at levels about 20-fold higher than healthy controls (Figure 5F). Likewise, microbial translocation was unchanged over 96 weeks of ART in viral success or viral failure subjects and remained well above levels detected in healthy controls (Figure 5G). No subject with undetectable TNF pre-ART developed measurable levels post-ART, while pretherapy plasma TNF, detected among 8 subjects, persisted for 96 weeks of ART (Figure 5H). Elevated TNF had no association with CD4%, microbial translocation or monocyte/macrophage activation and was independent of viral success or viral failure outcome (data not shown).

Longitudinal changes in viremia, immune activation and microbial translocation were compared over the course of treatment by linear regression to contemporaneous changes in CD4%. No significant associations between changes in CD4% and viral load, sCD27, sCD14 or LPS following 24 or 96 weeks of ART were identified. Decreasing frequency of CD4 CD45RO T cells was the only parameter related to sustained improvement in CD4 T cell frequency during ART in viral success and viral failure subjects (24 weeks: $P<0.05$; 96 weeks: $P<0.001$). (See Table S2, Supplemental Digital Content).

Discussion

HIV-1 infection results in systemic immune activation, which depends on virus replication and/or microbial translocation, and is resolved to an extent by antiretroviral therapy[5,9,14]. Systemic immune activation drives viral replication creating a cycle of T cell attrition. When combined with viral effects on thymic T cell output, dissecting the contribution of each disease variable to immune activation poses significant challenges. Our studies show that during the natural history of pediatric HIV-1 infection, T-cell activation and CD4 T-cell attrition are related to levels of viral replication depending on age/length of infection. In contrast, macrophage activation is related to both microbial translocation and effects of HIV-1 infection that persist even when viremia is suppressed below limits of detection. Sustained immune reconstitution by ART, independent of viral outcomes, induced concordant attenuation of CD4 and CD8 T cell activation[29], even though microbial translocation or macrophage activation persisted indicating that systemic immune activation in T cell versus monocyte/macrophage compartments can be dissociated based on distinct mechanisms.

While viral replication is the principle factor driving lymphocyte activation, microbial translocation is also implicated in systemic T cell activation in some, but not all, natural history studies of HIV-1 infection[5,7,8]. In contrast our study of perinatally-infected children found no association between microbial translocation and T cell activation or immune deficiency pre-ART, highlighting the complex relationships between different causes and consequences of immune activation. Following ART, reconstitution of CD4 T cells and increased TREC occurred in our cohort despite persistent microbial translocation indicating that thymic output, a major contributor to post-ART lymphocyte reconstitution[29,43,44], is independent of LPS-driven inflammation. Failure of ART to attenuate microbial translocation and LPS levels in

children, unlike ART treatment of HIV-1 infected adults[5], may reflect irreversible intestinal fibrosis induced by early viremia during infancy[45]. HIV-1-infection during infancy, compared to infection during adolescence or adulthood may have different effects on MALT due to normal processes of gut remodeling that occur during growth[46] when intestinal permeability is increased and adaptive immunity is still developing. Indeed, transient microbial translocation found in our study of healthy and HIV-1-infected infants resolved after 2 years of age only in healthy children, indicating that HIV-1 disrupted normal gut development. It would be interesting to further investigate the nature of microbial translocation in early infancy versus older HIV-1-infected children to determine if disruption of gut epithelial and endothelial tissues differs by age and illness. Microbial translocation, as a consequence of HIV-1 pathogenesis in adults, is inversely correlated with EndoCAb levels[5,47]. In our study, EndoCAb levels were reduced among HIV-1 infected children, but no relationship between EndoCAb and LPS levels was apparent. Developmental regulation of natural antibody production could impair the ability of perinatally-infected children to neutralize and clear circulating LPS during early infection.

The LPS detected in plasma of HIV-1 infected subjects is biologically active and inflammatory *in vivo*, reflected by the link with monocyte/macrophage activation, although healthy infants with similarly elevated LPS levels displayed no concomitant increase in monocyte/macrophage activation. HIV-1 might function as a co-factor for LPS-induced inflammation in infants. The overall extent of monocyte/macrophage activation found with HIV-1 infection could reflect direct activating effects of HIV-1 on monocytes[6,48,49] or priming by HIV-1, which sensitizes macrophages to subsequent activation with toll-like receptor [TLR] ligands including LPS[50]. Whether sCD14 detected in plasma is derived from monocytes or differentiated macrophages is unclear, but both scenarios are plausible. *In vivo*, macrophages are efficient targets for productive HIV-1 infection whereas monocytes are rarely infected [51,52], although binding of HIV-1 envelope proteins to CCR5 on monocytes induces cell signaling that enhances monocyte survival[53] and may induce activation and shedding of sCD14.

Lack of monocyte/macrophage responsiveness to LPS in healthy infants is characteristic of normal immune development[26,27]. For example, in atopic allergy, an immune hypersensitivity disorder, sCD14 levels are lower in neonates versus older children, even though neonates have increased frequency of CD14⁺ cells[54]. Breast-feeding is another variable that regulates inflammatory versus tolerogenic responses to microbial translocation. Secretory IgA in breast milk can have significant effects on gut microbial flora through alterations in colonization and prevention of trans-mucosal sepsis, and other soluble milk components can mediate anti-inflammatory effects[28]. None of the neonates in our study was breast fed, which may itself contribute to immune activation. Clearly, systemic monocyte/macrophage activation is less pronounced during childhood and HIV-1 infection significantly disrupts normal immune homeostasis. Although ART for HIV-1-infected subjects in this study improved health with significant increases in height and weight and decreased prevalence of AIDS-defining illnesses[24], inability of therapy to resolve either microbial translocation or monocyte/macrophage activation may contribute to accelerated immune senescence with increased long-term morbidity and mortality from non-AIDS related inflammatory conditions [55].

CD27, a member of the tumor necrosis factor receptor family and its ligand, CD70, are predominantly expressed by lymphocytes[56]. Lymphocyte activation in both the T cell and B cell compartments results in high levels of the soluble protein in the plasma and can be used to measure systemic immune activation in HIV-1 infection[56]. The divergence in sCD27 levels in post-ART viral success versus viral failure subjects may be linked to qualitative differences in HIV-1 phenotype and co-receptor utilization. HIV-1 isolated from subjects who

suppress viremia primarily use only CCR5 co-receptor for target cell entry, while HIV-1 from viral failure subjects use both CCR5 and CXCR4 co-receptors before and after ART[33]. Persistently elevated sCD27 demonstrates that replication of drug-resistant HIV-1 induces some component of lymphocyte activation in spite of improved CD4 T cells counts and normal immune function[24], indicating that systemic immune activation and its consequences are complex and multifactorial.

Discordant outcomes following ART represent an important paradigm in HIV-1 pathogenesis, which may be modeled by high levels of SIV replication in its natural host[18]. However, our study demonstrates fundamental differences between discordant therapy outcome in HIV-1 infection and natural SIV infection of sooty mangabeys in that CD4 T cell pathogenesis is uncoupled from microbial translocation in HIV-1 infection. In sooty mangabeys, viremia fails to induce intestinal fibrosis or microbial translocation which is related to the overall lack of CD4 T cell pathogenesis or immune deficiency in this model[5,17]. In contrast, HIV-1-infected children experience increased naïve CD4 T cell counts and TREC levels after ART regardless of viral outcome, suggesting that persistently elevated LPS two years into therapy does not adversely impact thymic output[29,57,58]. Post therapy viruses do not induce the same degree of T cell activation as wild type virus and are associated with higher TREC levels. In the discordant response group, post therapy drug resistant viruses are less fit for replication in the thymus resulting in a high proportion of naïve T cells in the peripheral blood[59,60]. Thus, improved CD4 T cell frequency in post-ART discordant response subjects is more likely a consequence of attenuated viral pathogenesis than changes in gut pathogenesis or microbial translocation. Similar to adults who continue failing ART regimens[22], discordant response in children and adolescents was associated with evidence of CD4 T cell declines by 96 weeks, although the proportion of CD45RO CD4 T cells failed to increase, an indication that robust thymic output persisted.

Our finding that microbial translocation is not a major factor in HIV-1 immunodeficiency in infants, children and adolescents is similar to a study of HIV-1 subtype A and D infections in African adults, who progressed to AIDS in the absence of HIV-1-specific microbial translocation and chronic HIV-1-associated inflammation[7]. Nonetheless microbial translocation is clearly an important component of innate immune inflammation in some HIV-1-infected subjects. In our study, both outcome groups developed post-ART declines in sCD14 levels, but monocyte/macrophage activation failed to normalize in either group, even following two years of ART, indicating that microbial translocation contributes to sustained macrophage activation that can be independent of high level viral replication. Prolonged systemic monocyte/macrophage activation may contribute to HIV-1-associated inflammatory conditions including dementia, increased risk for cardiovascular events and thrombophilia or enhanced angiogenesis in malignancy[13,30-32]. The extent of monocyte/macrophage activation and resulting inflammatory conditions may differ in perinatally-infected children compared to infected adults due to greater infectivity and HIV-1 replication in neonatal or cord blood-derived monocyte/macrophages[61,62]. Considering that perinatally-infected neonates will likely survive several decades with ART, the implications of persistent microbial translocation and monocyte/macrophage activation should be a major focus of investigation in coming years.

Acknowledgments

The authors wish to thank the study volunteers for participating, the clinical coordinator Carla Duff, R.N., C.C.R.P. for enrollment and maintenance of the cohort and Brent Gardener and Amanda Lowe for managing the clinical specimen repository and clinical database. We also wish to thank Dr. Grace Aldrovandi for critical reading of this manuscript.

Sources of financial support: *M.A.W.* – NIH T32 training grant fellowship AR007603 and the Laura McClamma Fellowship for Immune Deficiency.

C.A.R., L.Y. and W.H. – Florida Center for AIDS Research

S.S. –Eleanor Dana Charitable Trust

J.W.S. – NIH grant AI047723

M.M.G. – NIH grants HD032259 and AI065265; Center for Research in Pediatric Immune Deficiency; and Stephany W. Holloway University Chair for AIDS Research

References

1. Brenchley J, Schacker T, Ruff L, Price D, Taylor J, Beilman G, et al. CD4+ T cell depletion during all stages of HIV disease occurs predominantly in the gastrointestinal tract. *J Exp Med* 2004;200:749–759. [PubMed: 15365096]
2. Mehndru S, Poles M, Tenner-Racz K, Horowitz A, Hurley A, Hogan C, et al. Primary HIV-1 infection is associated with preferential depletion of CD4+ T lymphocytes from effector sites in the gastrointestinal tract. *J Exp Med* 2004;200:761–770. [PubMed: 15365095]
3. Arthos J, Cicala C, Martinelli E, Macleod K, Van Ryk D, Wei D, et al. HIV-1 envelope protein binds to and signals through integrin $\alpha 4 \beta 7$, the gut mucosal homing receptor for peripheral T cells. *Nat Immunol* 2008;9:301–309. [PubMed: 18264102]
4. Stein T, Koerner B, Schluter M, Leskiw M, Gprindachvilli T, Richards E, et al. Weight loss, the gut and the inflammatory response in aids patients. *Cytokine* 1997;9:143–147. [PubMed: 9071566]
5. Brenchley J, Price D, Schacker T, Asher T, Silvestri G, Rao S, et al. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med* 2006;12:1365–1371. [PubMed: 17115046]
6. Jiang W, Lederman M, Hunt P, Sieg S, Haley K, Rodriguez B, et al. Plasma Levels of Bacterial DNA Correlate with Immune Activation and the Magnitude of Immune Restoration in Persons with Antiretroviral-Treated HIV Infection. *J Infect Dis*. 2009
7. Redd A, Dabito D, Bream J, Charvat B, Laeyendecker O, Kiwanuka N, et al. Microbial translocation, the innate cytokine response, and HIV-1 disease progression in Africa. *Proc Natl Acad Sci U S A* 2009;106:6718–6723. [PubMed: 19357303]
8. Marchetti G, Bellistri G, Borghi E, Tincati C, Ferramosca S, La Francesca M, et al. Microbial translocation is associated with sustained failure in CD4+ T-cell reconstitution in HIV-infected patients on long-term highly active antiretroviral therapy. *AIDS* 2008;22:2035–2038. [PubMed: 18784466]
9. Anselmi A, Vendrame D, Rampon O, Giaquinto C, Zanchetta M, De Rossi A. Immune reconstitution in human immunodeficiency virus type 1-infected children with different virological responses to anti-retroviral therapy. *Clin Exp Immunol* 2007;150:442–450. [PubMed: 17956580]
10. Lien E, Aukrust P, Sundan A, Müller F, Frøland S, Espevik T. Elevated levels of serum-soluble CD14 in human immunodeficiency virus type 1 (HIV-1) infection: correlation to disease progression and clinical events. *Blood* 1998;92:2084–2092. [PubMed: 9731066]
11. Cadogan M, Dalglish A. HIV immunopathogenesis and strategies for intervention. *Lancet Infect Dis* 2008;8:675–684. [PubMed: 18818125]
12. Appay V, Sauce D. Immune activation and inflammation in HIV-1 infection: causes and consequences. *J Pathol* 2008;214:231–241. [PubMed: 18161758]
13. Ancuta P, Kamat A, Kunstman K, Kim E, Autissier P, Wurcel A, et al. Microbial translocation is associated with increased monocyte activation and dementia in AIDS patients. *PLoS ONE* 2008;3:e2516. [PubMed: 18575590]
14. Grossman Z, Meier-Schellersheim M, Paul W, Picker L. Pathogenesis of HIV infection: what the virus spares is as important as what it destroys. *Nat Med* 2006;12:289–295. [PubMed: 16520776]
15. Lee P, Ciccone E, Read S, Asher A, Pitts R, Douek D, et al. Evidence for translocation of microbial products in patients with idiopathic CD4 lymphocytopenia. *J Infect Dis* 2009;199:1664–1670. [PubMed: 19432548]

16. Gordon S, Klatt N, Bosinger S, Brechley J, Milush J, Engram J, et al. Severe depletion of mucosal CD4+ T cells in AIDS-free simian immunodeficiency virus-infected sooty mangabeys. *J Immunol* 2007;179:3026–3034. [PubMed: 17709517]
17. Silvestri G, Paiardini M, Pandrea I, Lederman M, Sodora D. Understanding the benign nature of SIV infection in natural hosts. *J Clin Invest* 2007;117:3148–3154. [PubMed: 17975656]
18. Kaur A, Grant R, Means R, McClure H, Feinberg M, Johnson R. Diverse host responses and outcomes following simian immunodeficiency virus SIVmac239 infection in sooty mangabeys and rhesus macaques. *J Virol* 1998;72:9597–9611. [PubMed: 9811693]
19. Hunt P, Brechley J, Sinclair E, McCune J, Roland M, Page-Shafer K, et al. Relationship between T cell activation and CD4+ T cell count in HIV-seropositive individuals with undetectable plasma HIV RNA levels in the absence of therapy. *J Infect Dis* 2008;197:126–133. [PubMed: 18171295]
20. Guadalupe M, Sankaran S, George M, Reay E, Verhoeven D, Shacklett B, et al. Viral suppression and immune restoration in the gastrointestinal mucosa of human immunodeficiency virus type 1-infected patients initiating therapy during primary or chronic infection. *J Virol* 2006;80:8236–8247. [PubMed: 16873279]
21. Guadalupe M, Reay E, Sankaran S, Prindiville T, Flamm J, McNeil A, et al. Severe CD4+ T-cell depletion in gut lymphoid tissue during primary human immunodeficiency virus type 1 infection and substantial delay in restoration following highly active antiretroviral therapy. *J Virol* 2003;77:11708–11717. [PubMed: 14557656]
22. Deeks S, Barbour J, Martin J, Swanson M, Grant R. Sustained CD4+ T cell response after virologic failure of protease inhibitor-based regimens in patients with human immunodeficiency virus infection. *J Infect Dis* 2000;181:946–953. [PubMed: 10720517]
23. Perez E, Rose S, Peyser B, Lamers S, Burkhardt B, Dunn B, et al. Human immunodeficiency virus type 1 protease genotype predicts immune and viral responses to combination therapy with protease inhibitors (PIs) in PI-naïve patients. *J Infect Dis* 2001;183:579–588. [PubMed: 11170983]
24. Ghaffari G, Passalacqua D, Caicedo J, Goodenow M, Sleasman J. Two-year clinical and immune outcomes in human immunodeficiency virus-infected children who reconstitute CD4 T cells without control of viral replication after combination antiretroviral therapy. *Pediatrics* 2004;114:e604–611. [PubMed: 15492356]
25. Deeks S, Hoh R, Grant R, Wrin T, Barbour J, Narvaez A, et al. CD4+ T cell kinetics and activation in human immunodeficiency virus-infected patients who remain viremic despite long-term treatment with protease inhibitor-based therapy. *J Infect Dis* 2002;185:315–323. [PubMed: 11807713]
26. Qing G, Howlett S, Bortolussi R. Lipopolysaccharide binding proteins on polymorphonuclear leukocytes: comparison of adult and neonatal cells. *Infect Immun* 1996;64:4638–4642. [PubMed: 8890218]
27. Cohen L, Haziot A, Shen D, Lin X, Sia C, Harper R, et al. CD14-independent responses to LPS require a serum factor that is absent from neonates. *J Immunol* 1995;155:5337–5342. [PubMed: 7594548]
28. Wold A, Adlerberth I. Breast feeding and the intestinal microflora of the infant--implications for protection against infectious diseases. *Adv Exp Med Biol* 2000;478:77–93. [PubMed: 11065062]
29. Yin L, Rodriguez C, Hou W, Potter O, Caplan M, Goodenow M, et al. Antiretroviral therapy corrects HIV-1-induced expansion of CD8+ CD45RA+ CD2-) CD11a(bright) activated T cells. *J Allergy Clin Immunol* 2008;122:166–172. e161-162. [PubMed: 18538835]
30. Sun J, Zheng J, Zhao M, Lee S, Goldstein H. Increased in vivo activation of microglia and astrocytes in the brains of mice transgenic for an infectious R5 human immunodeficiency virus type 1 provirus and for CD4-specific expression of human cyclin T1 in response to stimulation by lipopolysaccharides. *J Virol* 2008;82:5562–5572. [PubMed: 18353948]
31. Funderburg N, Mayne E, Sieg S, Asaad R, Jiang W, Kalinowska M, et al. Increased tissue factor expression on circulating monocytes in chronic HIV infection: relationship to in vivo coagulation and immune activation. *Blood*. 2009
32. Mullerat J, Perrett C, Deroide F, Winslet M, Bofill M, Poulter L. The role of macrophages in angiogenesis. Comparison between HIV+ and HIV- populations with anal dysplasia and anal cancer. *Anticancer Res* 25:693–699. [PubMed: 15868898]

33. Ho S, Perez E, Rose S, Coman R, Lowe A, Hou W, et al. Genetic determinants in HIV-1 Gag and Env V3 are related to viral response to combination antiretroviral therapy with a protease inhibitor. *AIDS* 2009;23:1631–1640. [PubMed: 19625947]
34. Weinberg A, Zhang L, Brown D, Erice A, Polsky B, Hirsch M, et al. Viability and functional activity of cryopreserved mononuclear cells. *Clin Diagn Lab Immunol* 2000;7:714–716. [PubMed: 10882680]
35. Weinberg A, Song L, Wilkening C, Sevin A, Blais B, Louzao R, et al. Optimization and limitations of use of cryopreserved peripheral blood mononuclear cells for functional and phenotypic T-cell characterization. *Clin Vaccine Immunol* 2009;16:1176–1186. [PubMed: 19515870]
36. HIV/AIDS Network. ACTG/IMPAACT Lab Manual. 2009. In: HIV/AIDS Network Coordination; www.hanc.info/labs/Pages/ACTGIMPAACTLabManual.aspx
37. Tuttle D, Coberley C, Xie X, Kou Z, Sleasman J, Goodenow M. Effects of human immunodeficiency virus type 1 infection on CCR5 and CXCR4 coreceptor expression on CD4 T lymphocyte subsets in infants and adolescents. *AIDS Res Hum Retroviruses* 2004;20:305–313. [PubMed: 15117454]
38. Yin L, Kou Z, Rodriguez C, Hou W, Goodenow M, Sleasman J. Antiretroviral therapy restores diversity in the T-cell receptor Vbeta repertoire of CD4 T-cell subpopulations among human immunodeficiency virus type 1-infected children and adolescents. *Clin Vaccine Immunol* 2009;16:1293–1301. [PubMed: 19605599]
39. Rabin R, Roederer M, Maldonado Y, Petru A, Herzenberg L. Altered representation of naive and memory CD8 T cell subsets in HIV-infected children. *J Clin Invest* 1995;95:2054–2060. [PubMed: 7738172]
40. Stent G, Cameron P, Crowe S. Expression of CD11/CD18 and ICAM-1 on monocytes and lymphocytes of HIV-1-infected individuals. *J Leukoc Biol* 1994;56:304–309. [PubMed: 7916028]
41. Scala E, Ansotegui I, Bellioni B, Guerra E, Aiuti F, Paganelli R. Expansion of CD11abright cells in CD8+CD45RA+ from HIV-infected patients: a new early marker for disease progression? *AIDS Res Hum Retroviruses* 1995;11:1327–1333. [PubMed: 8573389]
42. Froebel K, Doherty K, Whitelaw J, Hague R, Mok J, Bird A. Increased expression of the CD45RO (memory) antigen on T cells in HIV-infected children. *AIDS* 1991;5:97–99. [PubMed: 1676273]
43. Douek D, Koup R, McFarland R, Sullivan J, Luzuriaga K. Effect of HIV on thymic function before and after antiretroviral therapy in children. *J Infect Dis* 2000;181:1479–1482. [PubMed: 10762580]
44. Kou Z, Puh J, Wu S, Goodenow M, Sleasman J. Combination antiretroviral therapy results in a rapid increase in T cell receptor variable region beta repertoire diversity within CD45RA CD8 T cells in human immunodeficiency virus-infected children. *J Infect Dis* 2003;187:385–397. [PubMed: 12552422]
45. Schacker T, Nguyen P, Beilman G, Wolinsky S, Larson M, Reilly C, et al. Collagen deposition in HIV-1 infected lymphatic tissues and T cell homeostasis. *J Clin Invest* 2002;110:1133–1139. [PubMed: 12393849]
46. Neu J. Gastrointestinal maturation and implications for infant feeding. *Early Hum Dev* 2007;83:767–775. [PubMed: 17913404]
47. Balagopal A, Philp F, Astemborski J, Block T, Mehta A, Long R, et al. Human immunodeficiency virus-related microbial translocation and progression of hepatitis C. *Gastroenterology* 2008;135:226–233. [PubMed: 18457674]
48. Lee A, Sharp E, O'Mahony A, Rosenberg M, Israelski D, Nolan G, et al. Single-cell, phosphopeptide-specific analysis demonstrates cell type- and pathway-specific dysregulation of Jak/STAT and MAPK signaling associated with in vivo human immunodeficiency virus type 1 infection. *J Virol* 2008;82:3702–3712. [PubMed: 18216116]
49. Lester R, Yao X, Ball T, McKinnon L, Kaul R, Wachihi C, et al. Toll-like receptor expression and responsiveness are increased in viraemic HIV-1 infection. *AIDS* 2008;22:685–694. [PubMed: 18356597]
50. Brown J, Kohler J, Coberley C, Sleasman J, Goodenow M. HIV-1 activates macrophages independent of Toll-like receptors. *PLoS One* 2008;3:e3664. [PubMed: 19048100]
51. McElrath M, Pruett J, Cohn Z. Mononuclear phagocytes of blood and bone marrow: comparative roles as viral reservoirs in human immunodeficiency virus type 1 infections. *Proc Natl Acad Sci U S A* 1989;86:675–679. [PubMed: 2492110]

52. McElrath M, Steinman R, Cohn Z. Latent HIV-1 infection in enriched populations of blood monocytes and T cells from seropositive patients. *J Clin Invest* 1991;87:27–30. [PubMed: 1985101]
53. Giri M, Nebozyhn M, Raymond A, Gekonge B, Hancock A, Creer S, et al. Circulating monocytes in HIV-1-infected viremic subjects exhibit an antiapoptosis gene signature and virus- and host-mediated apoptosis resistance. *J Immunol* 2009;182:4459–4470. [PubMed: 19299747]
54. Holmlund U, Höglind A, Larsson A, Nilsson C, Sverremark Ekström E. CD14 and development of atopic disease at 2 years of age in children with atopic or non-atopic mothers. *Clin Exp Allergy* 2003;33:455–463. [PubMed: 12680860]
55. Deeks S. Immune dysfunction, inflammation, and accelerated aging in patients on antiretroviral therapy. *Top HIV Med* 2009;17:118–123. [PubMed: 19890183]
56. Lens S, Tesselaar K, van Oers M, van Lier R. Control of lymphocyte function through CD27-CD70 interactions. *Semin Immunol* 1998;10:491–499. [PubMed: 9826582]
57. Rodriguez C, Koch S, Goodenow M, Sleasman J. Clinical implications of discordant viral and immune outcomes following protease inhibitor containing antiretroviral therapy for HIV-infected children. *Immunol Res* 2008;40:271–286. [PubMed: 17952387]
58. Chavan S, Bennuri B, Kharbanda M, Chandrasekaran A, Bakshi S, Pahwa S. Evaluation of T cell receptor gene rearrangement excision circles after antiretroviral therapy in children infected with human immunodeficiency virus. *J Infect Dis* 2001;183:1445–1454. [PubMed: 11329124]
59. Stoddart C, Liegler T, Mammano F, Linquist-Stepps V, Hayden M, Deeks S, et al. Impaired replication of protease inhibitor-resistant HIV-1 in human thymus. *Nat Med* 2001;7:712–718. [PubMed: 11385509]
60. Sufka S, Ferrari G, Gryszowka V, Wrin T, Fiscus S, Tomaras G, et al. Prolonged CD4+ cell/virus load discordance during treatment with protease inhibitor-based highly active antiretroviral therapy: immune response and viral control. *J Infect Dis* 2003;187:1027–1037. [PubMed: 12660916]
61. Sundaravaradan V, Saxena S, Ramakrishnan R, Yedavalli V, Harris D, Ahmad N. Differential HIV-1 replication in neonatal and adult blood mononuclear cells is influenced at the level of HIV-1 gene expression. *Proc Natl Acad Sci U S A* 2006;103:11701–11706. [PubMed: 16868088]
62. Sperduto A, Bryson Y, Chen I. Increased susceptibility of neonatal monocyte/macrophages to HIV-1 infection. *AIDS Res Hum Retroviruses* 1993;9:1277–1285. [PubMed: 8142145]

Figure 1. T cell activation and uncontrolled viremia mediate age-dependent pathogenesis in pre-therapy HIV-1 infection

The frequency of (A) CD4 CD45RO T cells ($n = 31$) or (C) activated CD8 CD45RA CD27⁻ CD11a⁺⁺ T cells ($n = 17$) were compared to the overall frequency of CD4 T cells by linear regression analysis. (B) Comparison of CD4% and frequency of CD4 CD45RO T cells between viral success [VS] versus viral failure [VF] subjects. (D) Comparison of CD8% and frequency of activated CD8 T cells between VS and VF subjects. (E) The relationship between plasma viral load [VL] and CD4% was determined by linear regression analysis ($n=33$). (F) Comparison of viral load between VS and VF subjects. (For panels A, C and E, * $P < 0.05$, ** $P < 0.01$, and § $P < 0.0001$; Pearson correlation.) A multiple regression model was used to

determine age/LOI effects on correlations between CD4% and CD45RO frequency (**G**) or viral load (**H**). Solid line represents correlations at the median age of the cohort (6.6 yrs) and dashed lines represent correlations at birth (0 yrs) or maturity (18 yrs). (* $P < 0.05$ and $\phi P = 0.097$ represent interaction between age/LOI and either CD45RO or viral load by a multiple regression model.) The specific values for interactions between age/LOI and either CD4 CD45RO T cell frequency or viral load used to generate Figures 1G and 1H are presented in Table S1 (See Supplemental Digital Content).

Figure 2. Microbial translocation and multi-lineage leukocyte activation develop simultaneously in HIV-1 infected subjects

(A) Plasma levels of LPS were determined by LAL assay in healthy controls and HIV-1-infected subjects and (B) LPS data were divided into groups based upon age (Infants, 0-2 yrs; Children, >2-18 yrs). Plasma levels of (C) sCD14 and (D) EndoCAB IgM levels were determined by ELISA and (E) linear regression analysis was performed to compare EndoCAB to LPS in HIV-1-infected subjects. (F) Levels of sCD27 were determined by ELISA in healthy controls and HIV-1-infected subjects. (*P<0.05, **P<0.01 and §P<0.0001; Mann-Whitney U test).

Figure 3. Microbial translocation is associated with monocyte/macrophage activation independent of lymphocyte activation

(A) Levels of TNF in plasma of healthy controls and HIV-1-infected subjects were compared (dotted line represents limit of detection 300 pg/ml). (B) HIV-1-infected subjects were delineated into TNF (-) and TNF (+) groups and the levels of plasma LPS in each group was compared. The relationships between levels of plasma LPS and (C) monocyte/macrophage activation or (D) systemic lymphocyte activation were determined by linear regression analyses. Association of plasma LPS with frequency of (E) effector/memory CD4 T cells or (F) activated CD8 T cells were determined by linear regression analyses.

Figure 4. Microbial translocation and multilineage leukocyte activation develop independent of CD4 T cell attrition

The relationships between CD4 T cell frequency and plasma levels of (A) LPS, (B) sCD14 or (C) sCD27 were determined by linear regression analyses. The levels of (D) LPS, (E) sCD14 and (F) sCD27 were compared among viral success and viral failure subjects by Mann-Whitney U-test.

Figure 5. Longitudinal analyses of response to ART among viral success and viral failure subjects (A) Plasma viral load (B) CD4% and (C) the frequency of effector/memory CD4 T cells were determined just prior to ART (0 weeks) and after 4, 12, 24, 48 and 96 weeks of ART in viral success (black circles) and viral failure (shaded circles) subjects (Panel B, dotted line represents limit of detection 2.6 log₁₀ copies / ml). (D) The changes in CD4 CD45RO T cell frequency and log₁₀ TREC following 24-48 weeks of ART were compared by linear regression. Plasma levels of (E) sCD27, (F) sCD14 and (G) LPS were determined prior to ART and after 24 and 96 weeks of ART in viral success and viral failure subjects. (Panels C, E, F and G, dashed lines represent mean values among healthy control subjects) (H) Longitudinal analysis of 8 subjects who had detectable plasma TNF. (Panel H, dotted line represents limit of detection 300 pg/ml TNF; Heavy dashed line with boxes represents mean value for all 8 subjects) (*P<0.05, **P<0.01, †P<0.001 and §P<0.0001, Wilcoxon matched pairs test for 0 wk vs. 24 wk or 0 wk vs. 96 wk, or Mann-Whitney U test for viral success versus viral failure).