Published in final edited form as:

Dev Psychopathol. 2010 ; 22(1): 23–33. doi:10.1017/S095457940999023X.

Parent Conflict Predicts Infants' Vagal Regulation in Social Interaction

Ginger A. Moore

The Pennsylvania State University

Abstract

Parent conflict during infancy may affect rapidly developing physiological regulation. To examine the association between parent conflict and infants' vagal tone functioning, mothers ($N = 48$) reported levels of parent conflict and their 6-month-old male and female infants' respiratory sinus arrhythmia (RSA) was measured in the Still-Face Paradigm (SFP). Higher parent conflict was related to lower RSA at baseline and each episode of the SFP. Infants in higher conflict families showed diminished RSA reactivity and unexpected RSA withdrawal when interacting with mothers. Findings suggest atypical RSA regulation and reliance on self-regulation for infants in families with moderate levels of parent conflict. Implications for later development and future research are discussed.

Keywords

marital conflict; infant development; emotion regulation; physiological arousal

Pathways to risk or resilience for children in the context of parent conflict are likely to involve multiple interactions among factors inherent within children and their environments. An extensive body of research has found that conflict between parents is associated with higher rates of child behavior and emotional problems (e.g., Cox, Paley, & Harter, 2001; Cummings & Davies, 1994; Katz & Gottman, 1993) and with greater physiological arousal and slower rates of recovery from arousal (Gottman & Katz, 1989, 1993; Katz, 2001), but that effective physiological regulation, specifically regulation of vagal tone, may buffer children from negative behavioral consequences of parent conflict (El-Sheikh, Harger, & Whitson, 2001; Katz & Gottman, 1995, 1997). Together this research suggests that the relation between parent conflict and behavior disorders may be moderated by children's vagal regulation of stress.

Understanding how parent conflict is related to the development of children's vagal regulation is a critical area of research, particularly during infancy, as early experiences in emotionally challenging environments can have a significant impact on neurological and physiological systems when they are undergoing rapid development (Panksepp, 2001; Pollack, 2005; Schore, 2000; Siegel, 2001), signifying the possibility of a sensitive period for the development of physiological regulatory systems.

To date, research that has examined the relation between parent conflict and physiological regulation during early childhood is rare, even though very young children are more likely than older children to be exposed to parent conflict (Fantuzzo, Boruch, Beriama, & Atkins, 1997) and the highest levels of parent conflict occur during infancy and preschool years (Belsky & Rovine, 1990). As an initial step, given the paucity of research examining parent conflict during

infancy, the current study aimed to assess relations between parent conflict and infants' regulation of vagal tone during an experimental procedure designed to present a mild social stressor.

Cardiac vagal tone, typically indexed by respiratory sinus arrhythmia (RSA), is thought to play a central role in the organization of behavior, emotion, and attention. In infancy, RSA regulation (including measures of baseline RSA levels and RSA reactivity) has been found to be related to soothability, attentional control, and emotion regulation (Calkins, 1997; Huffman, Bryan, Del Carmen, Pedersen, Doussard-Roosevelt, & Porges, 1998; Porges, Doussard-Roosevelt, Portales, & Suess, 1994; Stifter & Corey, 2001; Suess, Porges, & Plude, 1994), which are precursors of later behavioral and emotional adjustment. In children and adolescents, less effective RSA regulation, indicated by low RSA and atypical RSA reactivity to challenging situations, has been linked consistently to behavioral and emotional problems (see review by Beauchaine, Gatzke-Kopp, & Mead, 2007).

There are various mechanisms to explain how parent conflict could be related to the development of RSA regulation: genetic factors shared between parents and children may be distally implicated in the association; parent conflict may directly increase demands on infants to regulate arousal, and higher conflict between parents may diminish parents' abilities to respond sensitively to infants' needs for support in regulation. Because vagal tone is theorized to have evolved to mediate social affiliation behaviors (Porges, 2007) and because the development of RSA is sensitive to socialization experiences across 3 to 12 months of age (e.g., Field et al., 1995; Propper et al., 2008), the current research was guided by the theory that parent conflict is associated with RSA regulation as a function of the spillover that parent conflict may have on parenting. Although parenting was not directly measured in the study, infants' RSA regulation was examined during a social interaction that typically functions as a social stressor for infants and that experimentally manipulated parents' availability to support infants' RSA regulation, the Still-Face Paradigm (SFP; Tronick, Als, Adamson, Wise, & Brazelton, 1978).

Parent-Infant Interaction and Infants' Developing RSA Regulation

Infants' self-regulation mechanisms are rudimentary at first and the development of effective RSA regulation may be a function of experiences in social interaction with parents during early infancy (Feldman, Greenbaum, & Yirmiya, 1999; Propper & Moore, 2006). For example, although baseline levels of RSA typically increase across the first years of life (e.g., Bar-Haim, Marshall, & Fox, 2000), infants of mothers with high levels of depressive symptoms did not show the normative increase in RSA between 3 and 6 months (Field, Pickens, Fox, & Nawrocki, 1995).

Similarly, recent research has found that maternal sensitivity moderated the relation between purported genetic risk for behavior problems and change in infants' RSA regulation across the first year of life (Propper et al., 2008). At 3 and 6 months of age, infants carrying the A_1^+ allele of the dopamine receptor D2 gene, which has been associated with novelty seeking, aggression, and impulse control disorders in adolescents and adults, failed to show expected and normative RSA withdrawal in response to mothers' disengagement during the SFP. By 12 months of age, infants with the A_1^+ allele whose mothers were rated as being higher in sensitivity at earlier ages now displayed RSA withdrawal when separated from mothers, reaching levels that were comparable to infants who were not at genetic risk. Infants with the A_1^+ allele whose mothers were rated lower in sensitivity failed to withdraw RSA in the expected situations at each age.

Together findings from these two studies suggest that the emotional quality of the early caregiving environment may influence trajectories of RSA development during periods of rapid physiological change.

Parenting may affect developing RSA regulation through support for infants' abilities to activate and to inhibit RSA appropriately during social interactions. Porges' (2007) polyvagal theory proposes that vagal regulation is associated with social affiliation behaviors that assist the individual in maintaining optimal arousal via parasympathetic influence on cardiac functioning. When there is little need to regulate arousal, RSA is activated or augmented, contributing to a slowing down of heart rate, whereas RSA is inhibited or withdrawn when the environment requires active (rather than passive) participation and/or requires active coping and self-regulatory behaviors.

Research has examined infants' RSA regulation during social interactions using the SFP, an experimental procedure that affords the opportunity to observe infants' RSA during periods of social engagement and during social disengagement when infants are presented with the challenge of self-regulation. According to polyvagal theory (Porges, 2007), the expected pattern would be RSA augmentation (higher RSA) during interactive episodes when infants can rely on parents to structure and support social interaction, and RSA withdrawal (lower RSA) during the still-face episode when parents' support is withdrawn. Studies have confirmed this normative pattern of RSA regulation across episodes of the SFP (Bazhenova, Plonskaia, & Porges, 2001; Moore & Calkins, 2004; Moore, Hill-Soderlund, Propper, Calkins, Mills-Koonce, & Cox, 2009; Weinberg & Tronick, 1996). This pattern of RSA augmentation and withdrawal may reflect infants' increased need for regulation in response to social disengagement and supports theories that infants rely on parents to structure social interaction and support regulation.

Consistent with this, infants who failed to show the expected response of RSA withdrawal when mothers disengaged from social interaction during the SFP, also showed an atypical pattern of responses when interacting with their mothers during normal play and reunion episodes (Moore & Calkins, 2004). Although their baseline levels of RSA did not differ from those of other infants, during the normal play episode of the SFP, they showed greater RSA withdrawal, less positive affect, and the dyad showed lower behavioral synchrony, suggesting that infants may have been less likely or able to rely on mothers' support for regulation of emotion and social interaction and may have relied on self-regulation to a greater degree than other infants (Moore & Calkins, 2004). The reliance on self-regulation may also explain why they did not show RSA withdrawal during the still-face episode. If the infants were accustomed to relying on self-regulation whether or not their mothers were interacting with them, they may not have shown the expected RSA withdrawal in response to mothers' inattention and unavailability.

As a whole, research examining infants' baseline RSA and RSA reactivity during parent-infant interaction supports the theory that a sensitive care-giving environment may promote the development of effective RSA regulation. Higher conflict between parents in a family may diminish parents' abilities to respond sensitively to their infants' needs for support in regulating behavior and emotions, obliging the child to become more dependent on self-regulation (e.g., Davies & Cummings, 2006; Donovan, Leavitt, & Walsh, 1998).

Parent Conflict and Infants' RSA Regulation

Only one published study of which we are aware has examined infants' RSA in relation to parent conflict (Porter, Wouden-Miller, Silva, & Porter, 2003). Results of that study showed that 6-month-old infants in families reporting higher marital conflict showed lower baseline RSA, suggesting a decreased capacity for effective regulation. In addition, infants with lower baseline RSA also showed less behavioral regulation as assessed by the Behavior Rating Scales (Bayley, 1993). In contrast, Gottman and Katz (1989) found that preschool-aged children in families reporting greater parent conflict showed higher RSA than other children. Gottman and

Katz (1989) proposed that this unexpected finding represented a temporary adaptation that, over the course of development, could overburden children's capacities for self-regulation.

The Current Study

The current study examined 6-month-old infants' mean levels of RSA and change in RSA, i.e., RSA reactivity, across episodes of the SFP in relation to parent conflict. The overall prediction was that higher levels of parent conflict would be related to atypical patterns of RSA regulation. The first hypothesis was that greater parent conflict would be related to lower mean levels of baseline RSA and to lower RSA across episodes of the SFP. This prediction was consistent with empirical work with infants (Porter et al., 2003) and with older children (El-Sheikh, et al., 2001).

The second hypothesis addressed RSA reactivity (i.e., change in RSA) in response to mothers' disengagement during the still-face episode of the SFP. In prior research, infants typically showed significant RSA withdrawal when mothers disengaged from social interaction (Moore & Calkins, 2004; Weinberg & Tronick, 1996) and those who did not showed less optimal interaction with their mothers in the normal play and reunion episodes (Moore & Calkins, 2004). We predicted that higher parent conflict would be related to a lesser degree of RSA withdrawal than typical in response to mothers' still-face, suggesting either ineffective RSA regulation or lack of reactivity to mothers' disengagement.

The third hypothesis was that infants in families reporting higher levels of parent conflict would show a lesser degree of expected RSA activation during contexts in which infants were interacting with mothers (normal play and reunion episodes). This hypothesis was based on previous research finding that infants who failed to show expected RSA withdrawal in response to mothers' disengagement (see second hypothesis) failed to show expected RSA activation when they were interacting with their mothers (Moore & Calkins, 2004).

Methods

Participants

Approximately half of participants were recruited from birth records and the other half from a separate, longitudinal study of child health and development (Durham Child Health and Development Study, DCHD), which also recruited participants from birth records and postings at birth and parenting classes. All participants were recruited from the same, urban area, thus the two recruiting methods were comparable. Seventy-five mothers who delivered full-term, healthy infants and who were married and/or cohabiting or had at least weekly contact with infants' fathers in the presence of the infant agreed to participate. All mothers signed informed consent for their and their infants' participation and study procedures were approved by the Institutional Review Board.

For analyses, forty-eight infants had complete physiological data across baseline and all three episodes of the SFP. Additional information regarding missing RSA data is provided below. Of the 48 infants, 61% were male, 51% were Euro American, and infants ranged in age from 6 to 8.5 months ($M = 6.81$, $SD = .68$). Thirty-eight of the 45 mothers who reported their marital status were cohabiting. Thirty-one percent of mothers reported family incomes below the poverty level. Of those families reporting incomes below poverty level, 36% were Euro American and 50% were non-cohabiting. There were no significant differences in demographic variables between the full sample and the subsample with complete RSA data.

Procedures

Conflict between parents—Each mother completed a 23 item self-report instrument describing the quality of partner relationships (Braiker & Kelly, 1979). Four subscales can be derived from the measure; for the current study only the Conflict subscale, composed of five items, was used. Sample items include: “How often do you and your partner argue with one another?” “When you and your partner argue, how serious are the problems or arguments?” “To what extent do you communicate negative feelings (anger, dissatisfaction, frustration) toward your partner?” Responses to items are scaled from 1 (Not at all) to 9 (Very much). Possible total scores range from 5 to 45. Validity of the instrument is supported by evidence of the instrument’s sensitivity to marital change across the transition to parenthood and by correlations between husbands’ and wives’ reports ranging from .24 to .62 across subscales (Belsky, Lang, & Rovine, 1985). Internal consistency in previous research on marital conflict and parenting ranged across subscales from .61 to .70 with a mean of .76 and test-retest reliability over a 12 month period ranged from .51 to .81 (Belsky & Hsieh, 1998; Belsky et al., 1985).

In this sample, internal consistency for the Conflict subscale was good ($\alpha = .759$) and total scores ranged from 5 to 34 ($M = 15.75$, $SD = 6.89$), suggesting mild to moderate levels of conflict. The mean and range were comparable to those obtained using this same instrument in an independent study on marital conflict and infants’ RSA functioning (Porter et al., 2003).

Still-Face Paradigm—To examine infants’ responses to social engagement and disengagement in relation to level of parent conflict in the home, infants were observed in the SFP. Infants’ behaviors were videotaped and their physiological responses were measured. Prior to the SFP, the experimenter attached heart rate monitoring equipment as described below. Mothers placed infants in an infant seat and sat in a chair directly in front of the infants. Mothers were given verbal instructions for the SFP, and then were asked to sit quietly and to review the same instructions in written form for 3 minutes. During this period, baseline cardiac data was collected for infants. The baseline episode was designed to minimize stimulation for infants.

For the normal play episode mothers were instructed to play with their babies as they normally would. Immediately after the 2-minute normal play episode, following standard SFP procedures, mothers were told to turn away from their infants for 15 seconds, then to turn back toward their infants for the still-face episode. Mothers were instructed to look at their infants for 2 minutes without responding in any way with facial or vocal expressions and were assured that the examiner would stop the procedure if the infant became too distressed. A 2-minute reunion episode followed the still-face in which mothers were instructed to respond to their babies in any way they felt was appropriate. The episodes of the SFP were video recorded and a time code was added to the videotape.

Coding infants’ and mothers’ affective behaviors—Infants’ and mothers’ behaviors were coded by trained coders naïve to hypotheses of the current study. Infants’ and mothers’ behaviors were coded by independent coders. Facial affect was coded at 1-s intervals as positive, neutral, or negative. If coders were unable to see infants’ or mothers’ faces, affect was coded as missing. Coders were initially trained to reliability using a large pool of videorecorded SFP interactions. To ensure that coders maintained inter-observer reliability in the current study, fifteen percent of the interactions were selected randomly and coded by a second coder. Agreement was calculated as coders observing the same behavior within one second of each other and quantified using kappa to correct for chance agreement ($\kappa = .89$ for infants, .83 for mothers).

Following previous research (Campbell et al., 1995; Moore & Calkins, 2004), to determine the amount of time during the SFP that infants and mothers spent in positive and negative affective states, the time series of the affect codes were aggregated individually for each infant and each parent. The total number of seconds in which each affective state code occurred was calculated for infants for each episode of the SFP and expressed as percentages of valid (not missing) time. Because mothers rarely displayed negative affect, percentages of mothers' negative affect were not computed and because mothers' affect was constrained to be neutral during the still-face episode, aggregate scores for mothers were computed only for the normal play and reunion episodes. Following previous research (e.g., Moore & Calkins, 2004), synchrony between mothers' and infants' behaviors was computed as the Pearson correlation between the second-by-second time series of mothers' and infants' affect.

Cardiac vagal tone—The researcher placed two disposable pediatric electrodes on the child's chest while the infant was seated in an infant seat on a table next to the mother. The electrodes were connected to a preamplifier, the output of which was transmitted to a monitor configured to collect heart interbeat intervals (IBIs; Mini Logger 2000; Mini-Mitter Corp., Bend, OR). A data file containing the IBIs for the entire period of collection was transferred to a laptop computer for later artifact editing and analysis using MXEdit software (Delta Biometrics, Bethesda, MD). Editing the files consisted of scanning the data for outlier points relative to adjacent data and replacing those points by dividing them or summing them so that they would be consistent with the surrounding data. Data files that required editing of more than 2% of the data (e.g., 12 data points in a 5-minute period) were not included in the analyses. Porges's (1985) method of calculating RSA was used. RSA was calculated every 30-s for the 3-min baseline period and every 15-sec during each of the 2-min episodes of the SFP (normal play, still-face, reunion).

Porges' method applies an algorithm to the sequential IBI data. The algorithm uses a moving 21-point polynomial to detrend periodicities in heart period (HP) slower than RSA. Then, a band-pass filter extracts the variance of HP within the frequency band of spontaneous respiration in young children (.24 – 1.04). Although lower frequency bands may be studied, research with young children has consistently examined this band and identified associations to child functioning (Huffman et al., 1998; Porges, 1996; Stifter & Fox, 1991). This estimate of RSA is derived by calculating the natural log of this variance and is reported in units of $\ln(\text{msec})^2$. RSA was calculated every 30-s for the 3-min baseline period and every 15-sec during each of the 2-min episodes of the SFP (normal play, still-face, reunion). These epoch durations are typical and valid for studies of short duration tasks with infants (Bar-Haim et al., 2000; Huffman et al., 1998). The mean RSA of the 15-s or 30-s epochs within each episode was used in subsequent analyses. Larger values of RSA indicated greater vagal tone, which putatively describes a state that does not require active regulation of emotion.

Change in RSA—To measure the degree of RSA reactivity, following prior research (e.g., Donzella, Gunnar, Krueger, & Alwin, 2000; Moore & Calkins, 2004; Quigley & Stifter, 2006), a set of change scores was computed (Δ RSA). Although change from baseline to a challenge task has traditionally been used as a measure of RSA reactivity, because the SFP presents a series of distinct but contiguous social contexts, the most relevant measure of RSA reactivity was change from the preceding episode. Thus, change scores were computed to measure Δ RSA from baseline to the normal play episode, from normal play to the still-face episode, and from the still-face to the reunion episode, by subtracting episode RSA from RSA in the previous episode or from baseline RSA in the case of change from baseline to normal play. Positive values of Δ RSA represented an increase in RSA (RSA activation) and negative values of Δ RSA represented a decrease in RSA (RSA withdrawal).

Missing data—RSA data were available for 52 infants in the baseline condition, 50 in the normal play episode, 51 in the still-face episode, and 48 in the reunion episode, with complete data across baseline and all episodes of the SFP available for 47 infants. One infant was missing only the baseline data point so this was replaced with the mean value, resulting in a sample size of $N = 48$ with complete data to test study hypotheses regarding RSA. RSA data were missing due to infants becoming too distressed to complete the SFP ($N = 6$), technical problems, such as electrodes falling off or other equipment failure ($N = 4$), editing of artifacts required for more than 2% of the episode, and invalid data indicated by a standard deviation across epochs greater than 1.00 for RSA (high degree of variability over the course of the episode) most likely due to movement artifact or infants' pulling or chewing on wires or ($N = 17$). This amount of missing physiological data is typical of studies collecting IBI from infants as they are often non-compliant with procedures, are physically active, and complete data were required across a number of assessment points (e.g., Calkins, 1997; Stifter & Jain, 1996). There were no differences in infant age, infant sex, behavioral measures, or parent conflict between infants with complete RSA data and those with missing data. Of the 48 infants who had complete RSA data across the SFP, four had missing behavioral data because either the infants became distressed and mothers stood to soothe the baby, blocking the camera view ($N = 2$) or technical problems with the video equipment ($N = 2$).

Results

Preliminary Analyses

Demographic variables—All preliminary and main analyses were conducted on the subsample of participants with complete RSA across baseline and the SFP ($N = 48$). One-way analyses of variance were conducted to examine differences in behavioral and physiological responses in relation to infant sex and ethnicity. Female infants were more positive than male infants during the still-face episode ($F(1, 43) = 7.45, p < .05$), with females showing positive expressions 8% of the time and males showing positive expressions 4% of the time. Male infants showed higher baseline RSA than female infants ($F(1, 47) = 4.99, p < .05$; $M_s = 3.73$ and 3.14 for males and females respectively). Therefore infant sex was included in main analyses. There was no relation between infant sex and parent conflict.

Euro American infants showed more negative affect in the still-face episode than African American infants ($F(1, 43) = 3.52, p < .05$), with Euro American infants showing negative expressions 26% of the time and African American infants showing negative expressions 15% of the time. There were no differences between African American and Euro American infants for RSA. However, because of the difference in negative affect in the still-face episode, suggesting different levels of distress, and because two independent studies with larger samples than this pilot study have reported that African American infants had higher baseline RSA and higher RSA during testing with the Bayley scales (C. Stifter, personal communication, July 22, 2007) and during the SFP (Moore et al., 2009), ethnicity was included in main analyses. There was no relation between infant ethnicity and parent conflict.

Level of parent conflict was unrelated to family income (below or above poverty) and did not differ between parents who lived together and those who did not. One-way analyses of variance revealed that infants' RSA and behavioral responses did not differ between cohabiting and non-cohabitating families. Because it is possible that infants from cohabitating and non-cohabitating families may be exposed to different rates and types of parent conflict, we conducted main analyses two ways: for all infants with complete RSA data across the SFP ($N = 48$) and excluding infants of parents who were not cohabitating ($N = 7$) or did not respond to the question about marital status ($N = 3$). Similar results were obtained for both sets of analyses; therefore results are reported for the full subsample with complete RSA data.

There was no relation between family income and infants' behavioral or RSA measures. There were no significant correlations between infant age and behavioral and RSA measures.

Correlations among behavioral and physiological responses—Higher baseline RSA was correlated with greater negative affect in the still-face episode ($r(44) = .36, p < .05$) but was unrelated to Δ RSA (RSA reactivity change scores). Δ RSA was correlated with negative affect in the normal play episode ($r(44) = -.33, p < .05$), such that higher levels of negative affect were associated with greater RSA withdrawal from baseline. Higher RSA in the still-face episode was correlated with greater negative affect in the preceding normal play episode ($r(44) = .29, p < .05$). Consistent with this, Δ RSA between normal play and still-face was correlated with negative affect ($r(44) = .34, p < .05$) and positive affect ($r(44) = -.32, p < .05$) in the normal play episode, indicating that infants who were more negative and less positive when interacting with their mothers showed less RSA withdrawal when their mothers became still-faced.

Correlations among mothers' behavior and infants' behavioral and physiological responses—Mothers' and infants' positive affect was correlated in the reunion episode only ($r(44) = .41, p < .01$). Mothers' positive affect was uncorrelated with infants' RSA or Δ RSA. Dyadic synchrony was uncorrelated with infants' RSA or Δ RSA.

Infants' Responses across the SFP

On average, infants showed normative patterns of behavioral and RSA responses across the SFP that have been found in prior research (Table 1). They were less positive during the still-face episode relative to the normal play and reunion episodes and more negative in the still-face relative to the normal play episode. Consistent with other research (e.g., Weinberg & Tronick, 1996), they showed more negative affect during the reunion episode than during normal play and a carryover of negative affect from the still-face episode. Consistent with previous research (e.g., Moore & Calkins, 2004; Weinberg & Tronick, 1996), RSA decreased significantly between the normal play and the still-face episodes. RSA did not increase significantly between the still-face and reunion episodes but RSA in the reunion episode was not significantly different from RSA in normal play, suggesting a carryover of physiological responses similar to that observed with behavioral responses.

Parent Conflict and Infants' Responses across the SFP

To examine whether parent conflict was related to mean levels of infants' positive and negative affect during episodes of the SFP, separate repeated measures general linear models were tested with parent conflict, infant sex, and infant ethnicity as independent variables and episode (normal play, still-face, reunion) as the within-subjects factor. There were no effects of parent conflict on infants' positive or negative affect during episodes of the SFP. There were also no effects of parent conflict on mothers' positive affect or on dyadic synchrony in either the normal play or reunion episodes.

To test the first study hypothesis regarding the relation between parent conflict and mean levels of infants' RSA at baseline and during each episode of the SFP, a repeated measures general linear model was tested with parent conflict, infant sex, and infant ethnicity as independent variables, and episode (baseline and each episode of the SFP) as the within-subjects factor. There was a main effect for parent conflict ($F(1, 43) = 5.22, p < .05, \eta^2 = .11$, see Figure 1). The interaction between episode and parent conflict was not significant ($F(3, 41) = 2.32, p < .09, \eta^2 = .15$). To probe and to illustrate the main effect, infants were categorized into three groups based on the interval conflict score (low conflict = -1 SD below the mean; average conflict = mean levels; high conflict = $+1$ SD above the mean) and the values of RSA were plotted by level of conflict (low, average, high; see Figure 1). Pairwise comparisons identified

significant differences in RSA for infants in families highest in parent conflict compared to all other infants (all $p < .05$), indicating that mean levels of RSA were lower at the highest levels of parent conflict during baseline as well as during each episode of the SFP.

RSA reactivity—To address hypotheses regarding the degree of change in RSA in the SFP in relation to parent conflict, i.e., RSA reactivity, we tested repeated measures general linear models with Δ RSA as the dependent repeated measure, parent conflict, infant sex, and infant ethnicity as independent variables, and episode (Δ RSA from baseline to normal play, Δ RSA from normal play to still-face, Δ RSA from still-face to reunion) as the within-subjects factor. There was a significant main effect for parent conflict ($F(1, 43) = 5.46, p < .05, \eta^2 = .14$), with higher parent conflict related to a lesser degree of Δ RSA across episodes. Pairwise comparisons identified a significant difference between the lowest ($-1 SD$) and highest levels ($+1 SD$) of parent conflict ($p < .05$) in degree of Δ RSA. The interaction between episode and parent conflict was not significant, indicating that the degree of RSA reactivity was lower for infants in the highest conflict families compared to the lowest conflict families across episodes of the SFP.

Correlations between parent conflict and infants' behavioral and RSA

responses—To further describe relations between parent conflict and infants' responses, we computed correlation coefficients among parent conflict scores, positive and negative affect, RSA at baseline and during each episode of the SFP, and the Δ RSA change scores. Results were consistent with analyses of mean levels of RSA and Δ RSA. Parent conflict was uncorrelated with infants' behavioral responses. Parent conflict was negatively correlated with RSA in each episode of the SFP (r 's = $-.30, -.30, -.35$, all $p < .05$ for normal play, still-face, and reunion episodes respectively). Of the Δ RSA reactivity scores, the correlation between parent conflict and change from baseline to the normal play episode approached significance ($r(48) = -.28, p = .058$), suggesting that higher conflict was associated with greater RSA withdrawal from baseline to the first interactive episode of the SFP.

As post hoc analyses, the Sobel test and bootstrapping methods (Preacher & Hayes, 2004) were used to test possible indirect effects of parent conflict on infant behavior via RSA, given that conflict was related to RSA and RSA was related to infants' behavior (in a few cases) even though conflict was unrelated to behavior. None of the tests were significant. In addition, relations were examined between parent conflict and mothers' positive affect and dyadic synchrony in the normal play and reunion episodes. No significant relations were found.

Discussion

To understand better the impact that parent conflict may have on infants' developing physiological regulation, this study examined the relation of parent conflict to infants' RSA regulation while interacting with their mothers. Based on prior research, we hypothesized that infants in families with higher levels of conflict would show lower levels of RSA at baseline and during episodes of the SFP, that they would show a lesser degree of RSA withdrawal in response to mothers' disengagement in the still-face episode, and that they would show an atypical response of RSA withdrawal when interacting with their mothers during the SFP.

The first hypothesis, that greater parent conflict would be related to lower levels of RSA, was confirmed, consistent with one previous study with infants (Porter et al., 2003) and with a larger body of research with older children (e.g., El-Sheikh et al., 2001; Katz & Gottman, 1995, 1997). Given that low RSA is generally found to be linked to disorders that involve problems with regulating behavior and emotion (Beauchaine et al., 2007), these findings suggested that the robust association between parent conflict and behavior problems later in development may be mediated by ineffective RSA regulation that could be evident as early as 6 months of age.

The second question addressed whether parent conflict was related to the degree of RSA reactivity (i.e., change in RSA) in response to mothers' disengagement in the still-face episode. Mothers' disengagement is a challenge task that has been found typically to elicit RSA withdrawal in infants (Moore & Calkins, 2004; Weinberg & Tronick, 1996), a purported indicator that infants are engaging in effective self-regulation in response to their mothers' unavailability. We found that infants from the highest conflict families did show RSA withdrawal in response to the still-face but to a significantly lesser degree than infants in the lowest conflict families. Note that in this study, RSA reactivity to mothers' still-face was measured as change in RSA from the previous normal play episode rather than change from baseline, although change from baseline to a challenge task has traditionally been used as a measure of RSA reactivity. Because the SFP presents a series of distinct but contiguous social contexts, we believed that the most relevant measure of RSA reactivity was change from the preceding episode.

Because we measured RSA reactivity to mothers' still-face as a function of change from RSA during normal play, results regarding RSA reactivity to mothers' disengagement should be interpreted in the context of RSA during the preceding normal play episode. The third study hypothesis that infants in families reporting higher parent conflict would show a lesser degree of expected RSA activation during episodes in which infants were interacting with mothers (the normal play and reunion episodes) was supported. The normative response of infants during interactive episodes of the SFP is RSA activation (Moore & Calkins, 2004; Moore et al., 2009; Weinberg & Tronick, 1996). However, in the current study, infants in the highest conflict families showed RSA withdrawal rather than RSA activation when interacting with their mothers during the normal play episode and significantly lower levels of activation during the reunion episode than infants in families with the lowest levels of parent conflict. This is consistent with previous research finding that 3-month-old infants who failed to show expected RSA withdrawal in response to mothers' disengagement also failed to show RSA activation when interacting with their mothers (Moore & Calkins, 2004).

Together with the finding that high parent conflict was associated with a lesser degree of RSA withdrawal when mothers disengaged from interacting with their infants between the normal play and still-face episodes, the finding that high parent conflict was associated with either RSA withdrawal or low levels of RSA activation during the interactive episodes suggest that infants in the highest conflict families may have been less physiologically reactive to mothers' disengagement because they were already showing a reliance on self-regulatory mechanisms. RSA withdrawal or low levels of RSA activation while interacting with parents may suggest greater self-regulation and we theorized that this may be due to the impact that parent conflict may have on diminishing parents' abilities to respond sensitively to their infants' needs for support in regulating arousal.

Taken together, our findings suggested that infants in families with relatively higher parent conflict may have a diminished capacity for RSA regulation in response to challenge, based on their lower overall RSA and attenuated degree of RSA reactivity in all episodes of the SFP, and may have adapted to rely on self-regulatory mechanisms even when interacting with their mothers, based on their failure to show significant RSA activation during interactive episodes. Alternative explanations, such as infants in highest conflict families were less distressed by the still-face or more distressed during normal play or reunion episodes were not supported by analyses of the behavioral data.

Whether infants in higher conflict families are likely to have negative or positive outcomes or to show ineffective RSA regulation later in development is unclear. At 6 months of age, they appeared to have adapted to rely on self-regulation even when interacting with their mothers and this could suggest a precocious ability to self-regulate that may serve them well in

environments where parents cannot be depended on to reliably provide support for regulating arousal, perhaps due to being absorbed or distracted by coping with conflict with the other parent. This would be consistent with the biological sensitivity to context theory (Boyce & Ellis, 2005), which proposes a developmental plasticity of stress-reactivity systems that evolved to facilitate successful adaptation to the specific environments in which children are raised, and with the physiological toughness theory (Dienstbier, 1989; Dienstbier & Zillig, 2002), which proposes that experiencing intermittent and mild to moderate stress has a beneficial effect on some biological systems.

On the other hand, an over-reliance on self-regulation that is less effective (indicated by overall low RSA) may have cumulative negative effects over time. Regulatory fatigue has been demonstrated in adults over short periods of time in conditions that require self-control (Muraven & Baumeister, 2000; Muraven & Slessareva, 2003), suggesting that chronic activation of self-regulatory mechanisms could detract from resources that could be utilized to promote optimal social and cognitive development. Thus, as Gottman and Katz (1989) suggested, a pattern of RSA regulation that may be an effective adaptation in early childhood, with time could become a maladaptive regulatory response.

It is important to note that the highest levels of conflict reported in this sample, which was not selected for marital distress or conflict, were moderate levels, compared to very little or almost no conflict. Therefore, findings apply to relatively normative levels of parent conflict, given that rates of parent conflict are highest during infancy and preschool and typically subside thereafter (Belsky & Rovine, 1990). In families with more intense or frequent conflict or conflict that reaches levels of physical violence, we might expect there to be more severe and lasting negative effects on children's functioning.

In addition, research findings linking RSA reactivity and behavioral problems have been mixed. Although lack of expected RSA withdrawal in response to challenge situations during infancy generally has been thought to indicate ineffective regulation (Calkins, 1997; Huffman, et al., 1998; Porges et al., 1994; Stifter & Corey, 2001; Suess et al., 1994), *greater* than normal RSA withdrawal was found to be associated with more anger in three- to five-year-olds (Donzella et al., 2000), and studies of older children with disruptive behavior disorders have found that low levels of RSA overall, rather than atypical RSA reactivity, are functionally related to problem behaviors (Beauchaine et al., 2007). Therefore, a lesser degree of expected RSA withdrawal, relative to other children, in response to a challenge situation may not necessarily mean ineffective regulation, and normative levels of change in RSA may not necessarily indicate effective regulation. However, because baseline RSA reflects the system's capacity for effective reactivity, low baseline RSA is typically associated with a lesser degree of RSA reactivity (Salomon, Matthews, & Allen, 2000). Therefore the findings of the current study that infants in families with the highest levels of conflict showed lower baseline RSA and lower levels of RSA across episodes of the SFP than other infants suggest that these infants may be at risk for later problems in behavioral and emotional regulation.

Limitations

There are several limitations of the study to be considered when interpreting results. First, the sample size in this study precluded the power to identify significant interactions between context (episode) and level of conflict. Some findings approached significance in theoretically consistent patterns and effect sizes of the findings were moderate. Although significant findings for RSA reactivity were those comparing the highest and the lowest levels of parent conflict, visual inspection of the profiles of change in RSA across the SFP suggested that the highest level of parent conflict may have differed from average levels of conflict, in the same way that mean levels of RSA did.

Another limitation of the study was that there was no independent measure of parenting. Thus we do not know whether infants in higher conflict families showed RSA withdrawal or diminished RSA activation when interacting with their mothers because of less sensitive and responsive parenting. The theoretical basis of the hypotheses was that parent conflict could affect RSA during infancy because caregivers' abilities to provide sensitive support for infants' developing physiological regulation may be undermined by parent conflict. Although we were unable to measure quality of parenting in this study, the SFP afforded the opportunity to observe infants' RSA when mothers' support for regulation was available (normal play and reunion episodes) and when mothers' support was unavailable (still-face episode). In addition, mothers' affective behaviors were observed during the interactive episodes as was dyadic behavioral synchrony, but both were unrelated to either parent conflict or to infants' RSA regulation.

Irrespective of parent conflict, in the current study, we found that infants who were more negative and less positive when interacting with their mothers in the normal play episode showed diminished RSA reactivity to their mothers' disengagement, consistent with prior research finding that infants who showed RSA withdrawal when interacting with mothers were less positive and showed less behavioral dyadic synchrony with their mothers, and also failed to show RSA withdrawal during the still-face episode (Moore & Calkins, 2004). This pattern of findings suggests that when interactions between mothers and their infants are less positive and less mutually responsive, infants show greater self-regulation of RSA while interacting with their mothers and are less physiologically reactive to mothers' disengaging from interaction. An example of this pattern of interaction that we have observed is where a mother engaged her baby insensitively by poking roughly at his cheeks, to which the baby responded by turning his face away. The mother then used her hands to turn the baby's face back towards her and the baby became negative and fussy and tried to turn away again. However, as soon as this mother adopted the still-face, the infant relaxed and began to look quietly at his mother.

The developmental theory guiding the current research, that physiological regulation may mediate the relation between parent conflict and later behavior disorders, could not be assessed in the current study. Parent conflict was not directly related to infant behavior during the SFP, but parent conflict was related to RSA, and RSA was related, in a few episodes of the SFP, to infant behavior. Various tests of indirect effects of parent conflict on infant behaviors via RSA did not result in significant findings. This could be a function of sample size, although the bootstrapping method provides reliable results with small samples (Preacher & Hayes, 2004).

One possible explanation for not finding associations between parent conflict and infant behavior and for finding relatively few significant associations between infants' RSA and behavior is that at 6 months of age, infants' behaviors are not consistently linked to their physiological responses (e.g., Gunnar, Mangelsdorf, Larson, & Hertzgaard, 1989; Weinberg and Tronick, 1996; Zelenko, Kraemer, Huffman, Gschwendt, Pageler, & Steiner, 2005). Therefore, associations between parent conflict and children's behavior may emerge later in development as children's physiological and behavioral responses consolidate into predictable patterns. Longitudinal research is needed to examine these developmental trajectories.

Another important limitation of the study is that there were not measures of the extent to which infants were directly exposed to parent conflict, although this is also a problem in the larger literature on marital conflict. A few of the families included in analyses were not cohabiting. It could be the case that even if children of parents who are not cohabiting are exposed to conflict less often that the conflict they do see is more intense; perhaps the parents are not cohabiting because of conflict, and, given that the noncustodial parent may be more likely to be in contact with the custodial parent only to visit with the child, it may be that conflict between the two occurs more often in the child's presence, whereas in cohabiting couples conflict may occur at other times. Questions about frequency of and exposure to conflict are very important

ones that should be addressed in future research. Given the small sample size, we could not analyze differences in relation to cohabiting, so these families were included in reported results after analyses excluding non-cohabiting families resulted in similar findings.

The duration of parent conflict and whether it is chronic may be another important variable in understanding developmental consequences of parent conflict. The prevailing theory is that children are sensitized to parent conflict through repeated exposure (Davies & Cummings, 2006). However, El Sheikh and Whitson (2006) note that there is “scant” evidence regarding the effects of exposure over time, mainly because the work has not yet been done. We propose that timing in terms of developmental stage of exposure to parent conflict may be a particularly important variable for understanding the effects of parent conflict on developing biological systems. Exposure during sensitive periods may be sufficient to set a child on a particular trajectory of risk, regardless of whether or not conflict persists over time. Developmental research is needed to identify whether such sensitive periods exist and, if so, whether they coincide with critical shifts in other domains. For example, given that RSA is thought to mediate social affiliation (e.g., Porges, 2007) and to be a function of positive engagement with the environment (e.g., Bazhenova et al., 2001), does the onset of social smiling between 3–4 months of age coincide with important developmental transitions in RSA functioning? In future research, moderators such as timing, chronicity, frequency, and degree of exposure to conflict will be critical factors to measure. Of note, these parameters have also been relatively neglected in the larger body of research on marital conflict and older children.

There are several caveats when interpreting research findings on infants’ vagal tone. It is important to take into account individual differences in reactivity; some individuals respond with parasympathetic (e.g., vagal tone) activation, some with sympathetic activation, and others with activation of both systems (Berntson & Cacioppo, 2007; Quigley & Stifter, 2006) and infants may also show patterns of individual differences that are not apparent when studying only RSA. Therefore, conclusions regarding whether the infants were regulating and, if so, how effectively they regulated, are tentative. Because there are methodological limitations on the ability to measure both parasympathetic and sympathetic responses in early infancy (Fox, Schmidt, Henderson, & Marshall, 2007), researchers may need to rely on behavioral indicators of reactivity. Two important directions for future research would be to identify groups of infants with distinct patterns of behavioral and physiological responses across the SFP or other challenge tasks and to follow the developmental trajectories of these groups of children with an emphasis on outcomes in early and middle-childhood, when most of the research on the consequences of parent conflict has been conducted.

In summary, the current study suggested that moderate levels of parent conflict are associated with low levels of RSA, a physiological risk factor for later problems with behavioral and emotion regulation, as early as 6 months of age. Parent conflict may also result in an adaptation of greater reliance on self-regulation rather than on parents to provide structure and support for infants’ regulation. Whether this adaptation is protective or is temporary requires further study.

Acknowledgments

This research was supported by a grant from the National Institute of Child Health and Development to the author (RO3HD043181). Portions of this research were presented at the May, 2004 XIVth Biennial International Conference for Infant Studies, Chicago, IL.

References

- Bar-Haim Y, Marshall PJ, Fox NA. Developmental changes in heart period and high-frequency heart period variability from 4 months to 4 years of age. *Developmental Psychobiology* 2000;37:44–56. [PubMed: 10937660]
- Bayley, N. Bayley Scales of Infant Development. 2. San Antonio, TX: The Psychological Corporation; 1993.
- Bazhenova OV, Plonskaia O, Porges SW. Vagal reactivity and affective adjustment in infants during interaction challenges. *Child Development* 2001;72:1314–1326. [PubMed: 11699673]
- Beauchaine TP, Gatzke-Kopp L, Mead HK. Polyvagal theory and developmental psychopathology: Emotion dysregulation and conduct problems from preschool to adolescence. *Biological Psychology* 2007;74:174–184. [PubMed: 17045726]
- Belsky J, Hsieh K. Patterns of marital change during the early childhood years: Parent personality, coparenting, and division-of-labor correlates. *Journal of Family Psychology* 1998;12:511–528.
- Belsky J, Lang M, Rovine M. Stability and change in marriage across the transition to parenthood: A second study. *Journal of Marriage & the Family* 1985;47:855–865.
- Belsky J, Rovine M. Patterns of marital change across the transition to parenthood: Pregnancy to three years postpartum. *Journal of Marriage & the Family* 1990;52:5–19.
- Berntson, GG.; Cacioppo, JT. Integrative physiology: Homeostasis, allostasis, and the orchestration of systemic physiology. In: Cacioppo, J.; Tassinary, L.; Berntson, G., editors. *Handbook of Psychophysiology*. Cambridge, UK: Cambridge University Press; 2007. p. 433–452.
- Boyce WT, Ellis BJ. Biological sensitivity to context: I. An evolutionary-developmental theory of the origins and functions of stress reactivity. *Development and Psychopathology* 2005;17:271–301. [PubMed: 16761546]
- Braiker, H.; Kelly, H. Conflict in the development of close relationship. In: Burgess, R.; Huston, T., editors. *Social exchange and developing relationships*. New York: Academic Press; 1979.
- Calkins SD. Cardiac vagal tone indices of temperamental reactivity and behavioral regulation in young children. *Developmental Psychobiology* 1997;31:125–135. [PubMed: 9298638]
- Campbell SB, Cohn JF, Meyers T. Depression in first-time mothers: Mother-infant interaction and depression chronicity. *Developmental Psychology* 1995;31:349–357.
- Cox, MJ.; Paley, B.; Harter, K. Interparental conflict and parent-child relationships. In: Grych, JH.; Fincham, FD., editors. *Interparental conflict and child development: Theory, research, and applications*. New York, NY: Cambridge University Press; 2001. p. 249–272.
- Cummings, EM.; Davies, P. Children and marital conflict: The impact of family dispute and resolution. New York: Guilford Press; 1994.
- Davies, PT.; Cummings, EM. Interpersonal discord, family process, and developmental psychopathology. In: Cicchetti, D.; Cohen, DJ., editors. *Developmental Psychopathology: Vol. 3: Risk, Disorder, and Adaptation*. 2. New York: Wiley & Sons; 2006. p. 86–128.
- Dienstbier RA. Arousal and physiological toughness: Implications for mental and physical health. 1989;96:84–100.
- Dienstbier, RA.; Zillig, LM. Toughness. In: Snyder, CR.; Lopez, SJ., editors. *Handbook of positive psychology*. New York, NY: Oxford University Press; 2002. p. 515–527.
- Donovan WL, Leavitt LA, Walsh RO. Conflict and depression predict maternal sensitivity to infant cries. *Infant Behavior and Development* 1998;21:505–517.
- Donzella B, Gunnar MR, Krueger WK, Alwin J. Cortisol and vagal tone responses to competitive challenge in preschoolers: Associations with temperament. *Developmental Psychobiology* 2000;37:209–220. [PubMed: 11084602]
- El-Sheikh M, Harger J, Whitson SM. Exposure to interparental conflict and children's adjustment and physical health: The moderating role of vagal tone. *Child Development* 2001;72:1617–1636. [PubMed: 11768136]
- El-Sheikh M, Whitson SA. Longitudinal relations between marital conflict and child adjustment: Vagal regulation as a protective factor. *Journal of Family Psychology* 2006;20:30–39. [PubMed: 16569087]

- Fantuzzo J, Boruch R, Beriama A, Atkins M. Domestic violence and children: Prevalence and risk in five major U.S. cities. *Journal of the American Academy of Child & Adolescent Psychiatry* 1997;36:116–122. [PubMed: 9000789]
- Feldman R, Greenbaum CW, Yirmiya N. Mother-infant affect synchrony as an antecedent of the emergence of self-control. *Developmental Psychology* 1999;35:223–231. [PubMed: 9923477]
- Field T, Pickens J, Fox NA, Nawrocki T. Vagal tone in infants of depressed mothers. *Development & Psychopathology* 1995;7:227–231.
- Fox, NA.; Schmidt, LA.; Henderson, HA.; Marshall, PJ. Developmental psychophysiology: Conceptual and methodological issues. In: Cacioppo, J.; Tassinary, L.; Berntson, G., editors. *Handbook of Psychophysiology*. Cambridge, UK: Cambridge University Press; 2007. p. 453-481.
- Gottman JM, Katz LF. The effects of marital discord on young children's peer interaction and health. *Developmental Psychology* 1989;25:273–381.
- Gottman JM, Katz LF. Patterns of marital conflict predict children's internalizing and externalizing behaviors. *Developmental Psychology* 1993;29:940–950.
- Gunnar MR, Mangelsdorf S, Larson M, Hertsgaard L. Attachment, temperament, and adrenocortical activity in infancy: A study of psychoendocrine regulation. *Developmental Psychology* 1989;25:355–363.
- Huffman LC, Bryan YE, Del Carmen R, Pedersen FA, Doussard-Roosevelt JA, Porges SW. Infant temperament and cardiac vagal tone: Assessments at twelve weeks of age. *Child Development* 1998;69:624–635. [PubMed: 9680676]
- Katz, LF. Physiological processes and mediators of the impact of marital conflict on children. In: Grych, JH.; Fincham, FD., editors. *Interparental conflict and child development: Theory, research, and applications*. New York, NY: Cambridge University Press; 2001. p. 188-212.
- Katz LF, Gottman JM. Patterns of marital conflict predict children's internalizing and externalizing behavior. *Developmental Psychology* 1993;29:940–950.
- Katz LF, Gottman JM. Vagal tone protects children from marital conflict. *Development and Psychopathology* 1995;7:83–92.
- Katz LF, Gottman JM. Buffering children from marital conflict and dissolution. *Journal of Clinical Child Psychology* 1997;26:157–171. [PubMed: 9169376]
- Moore GA, Calkins SD. Infants' vagal regulation in the still-face paradigm is related to dyadic coordination of mother-infant interaction. *Developmental Psychology* 2004;40:1068–1080. [PubMed: 15535757]
- Moore GA, Hill-Soderlund AL, Propper CB, Calkins SD, Mills-Koonce WR, Cox MJ. Mother-infant vagal regulation in the face-to-face still-face paradigm is moderated by maternal sensitivity. *Child Development* 2009;80:209–223. [PubMed: 19236402]
- Muraven M, Baumeister RF. Self-regulation and depletion of limited resources: Does self-control resemble a muscle? *Psychological Bulletin* 2000;126:247–259. [PubMed: 10748642]
- Muraven M, Slessareva E. Mechanisms of self-control failure: Motivation and limited resources. *Personality and Social Psychology Bulletin* 2003;29:894–906. [PubMed: 15018677]
- Panksepp J. The long-term psychobiological consequence of infant emotions: Prescriptions for the twenty-first century. *Infant Mental Health Journal* 2001;22:132–173.
- Pollack SD. Early adversity and mechanisms of plasticity: Integrating affective neuroscience with developmental approaches to psychopathology. *Development and Psychopathology* 2005;17:735–752. [PubMed: 16262990]
- Porges, SW. Method and apparatus for evaluating rhythmic oscillations in a periodic physiological response system. 4,510,944. 1985.
- Porges SW. The polyvagal perspective. *Biological Psychology* 2007;74:116–143. [PubMed: 17049418]
- Porges SW, Doussard-Roosevelt JA, Portales AL, Suess PE. Cardiac vagal tone: Stability and relation to difficulties in infants and 3-year-olds. *Developmental Psychobiology* 1994;27:289–300. [PubMed: 7926281]
- Porter CL, Wouden-Miller M, Silva S, Porter AE. Marital harmony and conflict: Links to infants' emotional regulation and cardiac vagal tone. *Infancy* 2003;4:297–307.

- Preacher KJ, Hayes AF. SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behavior Research Methods, Instruments, and Computers* 2004;36:717–731.
- Propper C, Moore GA. The influence of parenting on infant emotionality: A multi-level psychobiological perspective. *Developmental Review* 2006;26:427–460.
- Propper CB, Moore GA, Mills-Koonce WR, Halpern C, Hill A, Calkins SD, et al. Gene-environment contributions to the development of vagal functioning: An examination of DRD2 and maternal sensitivity. *Child Development* 2008;78:1378–1395.
- Quigley KS, Stifter CA. A comparative validation of sympathetic reactivity in children and adults. *Psychophysiology* 2006;43:357–365. [PubMed: 16916431]
- Salomon K, Matthews KA, Allen MT. Patterns of sympathetic and parasympathetic reactivity in a sample of children and adolescents. *Psychophysiology* 2000;37:842–849. [PubMed: 11117464]
- Schore, AN. The self-organization of the right brain and the neurobiology of emotional development. In: Lewis, MD.; Granic, I., editors. *Emotion, development, and self-organization*. Cambridge: Cambridge University Press; 2000. p. 155-185.
- Siegel DJ. Toward an interpersonal neurobiology of the developing mind: Attachment relationships, “mindsight,” and neural integration. *Infant Mental Health Journal* 2001;22:67–94.
- Stifter CA, Corey JM. Vagal regulation and observed social behavior in infancy. *Social Development* 2001;10:189–201.
- Stifter CA, Jain A. Psycho physiological correlates of infant temperament: Stability of behavior and autonomic patterning from 5 to 18 months. *Developmental Psychobiology* 1996;29:379–391. [PubMed: 8732809]
- Suess PE, Porges SW, Plude DJ. Cardiac vagal tone and sustained attention in school-age children. *Psychophysiology* 1994;31:17–22. [PubMed: 8146250]
- Tronick EZ, Als H, Adamson L, Wise S, Brazelton B. The infants’ response to entrapment between contradictory messages in face-to-face interaction. *American Academy of Child Psychiatry* 1978;1:1–13.
- Weinberg KM, Tronick EZ. Infants’ affective reactions to the resumption of maternal interaction after the still-face. *Child Development* 1996;67:905–914. [PubMed: 8706534]
- Zelenko M, Kraemer H, Huffman L, Gschwendt M, Pageler N, Steiner H. Heart rate correlates of attachment status in young mothers and their infants. *Journal of the American Academy of Child and Adolescent Psychiatry* 2005;44:470–476. [PubMed: 15843769]

Figure 1.
Levels of RSA across baseline and episodes of the SFP by parent conflict.

Figure 2.

Infants' RSA Reactivity (Δ RSA) in episodes of the SFP by parent conflict.

Note. Positive values indicate RSA activation, negative values indicate RSA withdrawal.

Table 1
Mean Values of Infants' Behavioral and Physiological Responses across the SFP

Variable	Normal Play		Still-Face		Reunion		F	df	η^2
	M (SD)		M (SD)		M (SD)				
Positive Affect	.24 ^a (0.20)		.05 ^b (0.06)		.23 ^a (0.17)		17.91 ^{***}	2, 40	.31
Negative Affect	.07 ^a (0.10)		.20 ^b (0.20)		.16 ^b (0.22)		9.75 ^{**}	2, 40	.20
RSA	3.68 ^a (1.08)		3.41 ^b (0.96)		3.61 ^{ab} (1.09)		3.22 [†]	3, 44	.13

Note. RSA = Respiratory Sinus Arrhythmia. Values with different superscripts within rows differ significantly.

[†] $p < .10$.
** $p < .01$.
*** $p < .001$.