Published in final edited form as:

J Org Chem. 2009 October 2; 74(19): 7595–7597. doi:10.1021/jo9016502.

Efficient Methodology for the Synthesis of 3-Amino-1,2,4-triazoles

Romain Noël, Xinyi Song, Rong Jiang, Michael J. Chalmers, Patrick R. Griffin, and Theodore M. Kamenecka*

Department of Molecular Therapeutics and Translational Research Institute, The Scripps Research Institute, Scripps Florida, 130 Scripps Way No. A2A, Jupiter, Florida 33458

Abstract

A general and efficient method for the preparation of 3-amino-1,2,4-triazoles has been developed. The desired 3-amino-1,2,4-triazoles (**1**) were prepared in good overall yield via two convergent routes. The key intermediate within both routes is substituted hydrazinecarboximide derivative **2**.

Triazole heterocycles occupy a central position in modern heterocyclic chemistry, principally because this heterocyclic ring is an important recognition element in biologically active molecules. Consequently new and efficient methods for the preparation of this important heterocyclic ring system are of contemporary interest. 3-Amino-1,2,4-triazole and its derivatives have also been the subject of numerous studies because of the many reported applications in the fields of medicinal and agrochemistry. For example, 3-aminotriazole derivatives are inhibitors of catalase¹ and histidine² bio-synthesis. Sufotidine bismuth citrate, a complex with an aminotriazole motif that is a histamine H₂-receptor antagonist, is used in the treatment of duodenal and gastric ulceration and other conditions where histamine is a known mediator.³ 3-Aminotriazoles have also been found effective for the treatment of chronic bronchial asthma,⁴ used as herbicides,⁵ and patented as neuropeptide Y receptor ligands.⁶ Aminotriazoles have also been described as potent CRF1 receptor antagonists,⁷ as well as inhibitors of methionine aminopeptidase-2.⁸

Numerous methods have been developed for the synthesis of 1,5-disubstituted 3-amino-1,2,4-triazoles.⁹ However, few of these describe direct synthesis of the triazole lacking substitution at the 5-position. Methods do exist to synthesize 5-amino-substituted analogues which are then diazotized to remove the amino group, but this is laborious as it requires an additional step for every analogue synthesized.¹⁰ Additionally, other published routes to 5-unsubstituted triazoles suffer some disadvantages including the following: (1) not atom economical,¹¹ (2) low yielding^{11a,12} and (3) limited scope with regards to substitution of the 3-amino group¹³ or in the N-1 position¹⁴ (the substituent in this position was added in an additional step). In some cases, chemistry is limited to symmetric bisaryl substitution patterns, which may not find general use.¹⁵ In other approaches, there are only a few examples given. Hence, the breadth and applicability of these methods to synthesize a variety of substituted (alkyl, aryl, primary or secondary alkyl or aryl amine) 3-amino-1,2,4-triazoles is lacking.¹¹

As part of our medicinal chemistry research program, we required a robust facile synthesis of 3-aminotriazole derivatives (devoid of C-5 substitution) **1** wherein we could vary the R¹, R², and R³ groups. Herein we report a convergent and convenient method for the preparation of these derivatives.

We envisioned that 3-aminotriazole **1** could be obtained by cyclization of hydrazinecarboximidamide derivative **2** with a formic acid equivalent (Scheme 1). This precursor could be prepared by two different methods. One route begins with thiourea **3** and the other with hydrazinecarbothioamide **4** wherein the R¹/R²/R³ groups are introduced just prior to cyclization. This convergent synthesis facilitates the study of structure–activity relationships for **1** depending on the starting material **3** or **4**. To illustrate the potential of this approach, we decided to optimize the chemistry using commercially available derivatives: 1-phenylthiourea **3a** (R² = Ph and R³ = H) and 1-phenyl-3-thiosemicarbazide **4a** (R¹ = Ph).

Thiourea **3a** was converted to the known sulfonic acid **5** following the reported literature procedure (Scheme 2).¹⁶ Oxidation of thiourea **3a** with hydrogen peroxide in the presence of sodium molybdate dehydrate gave **5** in excellent yield.¹⁷ The reaction of **5** with a variety of commercially available aryl- or alkyl-hydrazines afforded intermediates **2a** which when treated with trimethyl orthoformate provided the desired 3-aminotriazoles **1**.

Our initial reaction of **5** with phenylhydrazine at room temperature (without base) showed good conversion to the expected compound **2a** (R¹ = Ph). Because of the aqueous solubility of this intermediate the crude reaction mixture was simply concentrated in vacuo, and taken to the next step. Heating of this intermediate in trimethyl orthoformate at 140 °C for 14 h produced the 3-aminotriazole **1a** in 66% isolated yield (Table 1, entry 1).

The same procedure was then applied to the other hydrazines in Table 1. In the cases where R¹ is a substituted phenyl ring (Table 1, entries 2 and 3), the 3-aminotriazoles were obtained with moderate yields. When the reaction was performed with 2-hydrazinopyridine (Table 1, entry 4), the conversion of the first step was very slow at room temperature, but proceeded more efficiently at 80 °C. With *tert*-butylhydrazine hydrochloride (Table 1, entry 5), the first step did not occur even at 80 °C and starting material was recovered. However, the reaction did progress smoothly in the presence of an external base. Of the different bases tried (pyridine, 1,4,6-collidine, diisopropylethylamine, cesium carbonate, sodium hydroxide, triethylamine, and DBU) triethylamine proved to give the highest overall yields for the two steps. Base was required in all reactions starting with hydrazine hydrochloride salts (Table 1, entries 3, 5, and 6). Other (free base) aliphatic hydrazines (Table 1, entries 7 and 8) reacted smoothly at room temperature without additive to give the desired product in good yield. Although the yields are modest in some cases, the two-step one-pot method provided rapid access to the desired compounds not readily available via other synthetic methods.

To investigate substitution of the 3-position of triazole **1**, we envisioned an alternative route featuring 1-phenyl-3-semithiocarbazide **4a** as the starting material in place of 1-phenylthiourea **3a** (Scheme 3). The synthesis of sulfonic acid derivative **6** had not been reported in the literature, but was readily prepared in 93% yield from **4a** following the same procedure described to synthesize **5**.¹⁷ The reaction of **6** with a variety of commercially available aryl- or alkylamines led to intermediates **2b** which when treated with trimethyl orthoformate afforded the desired 3-aminotriazoles **1**. A study was carried out to optimize the reaction conditions.

Treatment of **6** with aniline at 120 °C for 30 min (without base) gave a very low conversion to intermediate **2b**. The reaction was then tried in the presence of different bases. When the reaction was performed with DBU, triethylamine, sodium hydroxide, or cesium carbonate, we noted that the formation of byproducts was considerably increased in relation to the conversion to **2b**. However, we found that conversion to **2b** was dramatically improved in the presence of pyridine. We also tried the reactions in different solvents (DMF, EtOH, CH₃CN) with the cleanest conversions occurring in acetonitrile. The cyclization of intermediates **2b** was performed as before at 140 °C for 14 h to afford the desired products **1** in good yields for the two steps.

As reported in Table 2, the 3-aminotriazoles **1** were obtained in all cases in good yields. In examples where the reaction was performed with aniline derivatives (Table 2, entries 1 and 2), the yields are slightly lower perhaps reflecting the reduced reactivity of these nucleophiles toward intermediate **6**. It is noteworthy that the reactions are compatible with primary and secondary amines as well as anilines. Moreover, the reactions are compatible with several functional groups such as olefins, acetals, carbamates, and esters.

In conclusion, we have developed an efficient and convergent method to synthesize differently substituted 3-amino-1,2,4-triazoles **1** from readily available starting materials. To vary the substitution at the N-1 position, the synthesis of aniline derived sulfonic acid intermediate **5** is followed by addition of hydrazine and then cyclization. To vary the 3-amino substituents, the synthesis of sulfonic acid **6** is followed by amine addition and then cyclization. The yields are good for the two-step, one-pot protocol and mild enough to accommodate a range of functional groups. Our ongoing efforts toward the synthesis of biologically active compounds containing the 3-aminotriazole motif using this methodology will be reported elsewhere.

Experimental Section

Representative Experimental Procedure for the Preparation of Compound 1: General Procedure A

A mixture of **5** (240 mg, 1.2 mmol) and phenylhydrazine (108 mg, 1.0 mmol) in anhydrous acetonitrile (1 mL) was stirred at room temperature for 1.5 h. The reaction mixture was then concentrated to a solid, which was added with trimethyl orthoformate (1 mL) and heated overnight at 140 °C in a sealed tube. The resulting mixture was cooled to room temperature, filtered through a short pad of silica gel, and flushed with 20% MeOH in CH₂Cl₂. The filtrate was concentrated and purified by preparative HPLC to provide the desired product **1a** (156 mg, 66%) as a solid.

General Procedure B

A mixture of **6** (100 mg, 0.465 mmol), aniline (37 μ L, 0.406 mmol), and pyridine (72 μ L, 0.890 mmol) in anhydrous acetonitrile (470 μ L) was stirred at 120 °C for 30 min in a sealed tube. The reaction mixture was then concentrated. The residue was diluted with methyl orthoformate (1.2 mL) and heated overnight at 140 °C in a sealed tube. The resulting mixture was partitioned

between a saturated solution of NaH-CO₃ (5 mL) and EtOAc (3 × 20 mL). The combined organic layers were washed with brine (10 mL), dried over anhydrous MgSO₄, filtered, and concentrated in vacuo. The residue was purified by flash chromatography to provide the desired product **1a** (41 mg, 43%) as a solid. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 6.83–6.87 (m, 1H), 7.26–7.30 (m, 2H), 7.33–7.37 (m, 1H), 7.52–7.56 (m, 2H), 7.64–7.66 (m, 2H), 7.84–7.87 (m, 2H), 9.07 (s, 1H), 9.44 (s, 1H); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ 116.0, 118.1, 119.5, 126.6, 128.7, 129.7, 136.9, 140.8, 141.5, 160.8; MS (ESI) [M + H]⁺ 237.19

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Professor William R. Roush (The Scripps Research Institute, Scripps Florida) is gratefully acknowledged for proofreading the manuscript. The efforts of P.R.G. and M.J.C. were supported by the National Institutes of Health (NIH) Molecular Library Screening Center Network (MLSCN) grant U54MH074404 (Hugh Rosen, Principal Investigator).

References

- (1). Khadir A, Verreault J, Averill DA. Arch. Biochem. Biophys 1999;370:163–175. [PubMed: 10510274]
- (2). Ventura L, PerezGonzales JA, Ramon D. FEMS Microbiol. Lett 1997;149:207–212. [PubMed: 9141661]
- (3). Clitherow, JW. EP 367484. 1989.
- (4). Naito Y, Akahoshi F, Takeda S, Okada T, Kajii M, Nishimura H, Sigiura M, Fukaya C, Kagitani Y. J. Med. Chem 1996;39:3019–3029. [PubMed: 8709136]
- (5). Hartmann M, Bauer H-J, Wermann K. Biocide Polym 1985:195.
- (6). Fauchere, J-L.; Ortuno, J-C.; Duhault, J.; Boutin, JA.; Levens, N. EP 1044970. 2000.
- (7). Lowe RF, Nelson J, Dang TN, Crowe PD, Pahuja A, McCarthy JR, Grigoriadis DE, Conlon P, Saunders J, Chen C, Szabo T, Chen TK, Bozigian H. J. Med. Chem 2005;48:1540–1549. [PubMed: 15743196]
- (8). Marino JP, Fisher PW, Hofmann GA, Kirkpatrick RB, Janson CA, Johnson RK, Ma C, Mattern M, Meek TD, Ryan MD, Schulz C, Smith WW, Tew DG, Tomazek TA, Veber DF, Xiong WC, Yamamoto Y, Yamashita K, Yang G, Thompson SK. J. Med. Chem 2007;50:3777–3785. [PubMed: 17636946]
- (9). (a) Temple, C, Jr.. Triazoles 1,2,4. Montgomery, JA., editor. John Wiley and Sons; New York: 1981. p. 130(b) Polya, JB. Comprehensive Heterocyclic Chemistry. Potts, KT., editor. Vol. 5. Pergamon Press; Oxford, UK: 1984. p. 733 (c) Katritzky AR, Rogovoy BV, Vvedensky VY, Kovalenko K, Steel PJ, Markov VI, Forood B. Synthesis 2001;6:897–893. and references cited therein. (d) Makara GM, Ma Y, Margarida L. Org. Lett 2002;4:1751–1754. and references cited therein. [PubMed: 12000290]
- (10). Davies, RJ.; Arnost, MJ.; Bemis, GW.; Forster, CJ.; Grey, R., Jr.; Ledford, B.; Marhefka, C.; Messersmith, D.; Pierce, AC.; Salituro, F.; Wang, J. WO 2005013982. 2005.
- (11). (a) Miyamoto Y, Yamazaki C. J. Heterocycl. Chem 1989;26:763–767. (b) Miyamoto Y, Yamazaki C. J. Heterocycl. Chem 1989;26:327–332.
- (12). Frenna V, Vivona N, Consiglio G, Spinelli D, Mezzina E. J. Chem. Soc., Perkin Trans. 2 1993;7:1339–1343.
- (13). (a) El-Gendy Z, Abdel-Rahman RM, Abdel-Malik MS. Indian J. Chem., Sect. B 1989;28B(6):479–485. (b) Abdel-Rahman RM, Ghareib M. Indian J. Chem., Sect. B 1987;26B(5):496–500.
- (14). Brenchley, G.; Charrier, J-D.; Durrant, S.; Knegt, R.; Ramaya, S.; Sadiq, S. WO 2007139795. 2007.
- (15). Baker W, Ollis WD, Poole VD. J. Chem. Soc 1950:3389–3394.

- (16). Maryanoff CA, Stanzione RC, Plampin JN, Mills JE. *J. Org. Chem* 1986;51:1882–1884.
- (17). While crude **5** or **6** could be used directly, better yields were obtained if **5** or **6** were purified on silica gel.

Scheme 1. Retrosynthesis

Scheme 2. Synthesis of 3-Amino-1,2,4-triazole from 3a

Reagents and conditions: (a) $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ (0.05 equiv), 30% H_2O_2 (5 equiv), NaCl (0.4 equiv), H_2O , 0 °C to rt; (b) R^1NHNH_2 (1 equiv), without or with triethylamine (2.2 equiv), CH_3CN , rt or 80 °C, 1.5 h; (c) $\text{HC}(\text{OMe})_3$, 140 °C, overnight.

Scheme 3. Synthesis of 3-Amino-1,2,4-triazole from 4a

Reagents and conditions: (a) Na₂MoO₄·2H₂O (0.05 equiv), 30% H₂O₂ (5 equiv), NaCl (0.4 equiv), H₂O, 0 °C to rt; (b) R²R³NH (1 equiv), pyridine (2.2 equiv), CH₃CN, 120 °C, 30 min; (c) HC(OMe)₃, 140 °C, overnight.

11

11

Entry	R ¹
2
3
4
5
6

Entry	R ¹
7
8

^a Reaction condition of the first step.

^b Isolated yield over two steps.

^c Without additive.

^d With 2.2 equiv of triethylamine.

TABLE 2
Preparation of 1 from 6

Entry
1

Entry

2

Entry
3
4
5

Entry

6

Entry
7
8
9

Entry

10

^aIsolated yield over two steps.