Published in final edited form as:

Aust N Z J Psychiatry. 2010 January ; 44(1): 4–19. doi:10.3109/00048670903393597.

Prevalence Estimates of Combat-Related PTSD: A Critical Review

Lisa K. Richardson, M.App.Psy.¹, B. Christopher Frueh, Ph.D.², and Ronald Acierno, Ph.D.³

¹ School of Psychology, Murdoch University, Perth, WA, Australia

² The Menninger Clinic and Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas, USA

³ Veterans Affairs Medical Center and Department of Psychiatry and Behavioral Sciences, Medical University of South Carolina, Charleston, South Carolina, USA

Abstract

Objective—To provide a critical review of prevalence estimates of combat-related PTSD among military personnel and veterans, and of the relevant factors that may account for the variability of estimates within and across cohorts, including methodological and conceptual factors accounting for differences in prevalence rates across nations, conflicts/wars, and studies.

Method—We examined MEDLINE and PsycINFO databases for literature on combat-related PTSD. The following terms were used independently and in combinations in this search: PTSD, combat, veterans, military, epidemiology, prevalence.

Results—The point prevalence of combat-related PTSD in US military veterans since the Vietnam War ranges from about 2 – 17%. Studies of recent conflicts suggest that combat-related PTSD afflicts between 4 – 17% of US Iraq War veterans, but only 3 – 6% of returning UK Iraq War veterans. Thus, the prevalence range is narrower and tends to have a lower ceiling among combat veterans of non-US Western nations. Variability in prevalence is likely due to differences in sampling strategies; measurement strategies; inclusion and measurement of the *DSM-IV* clinically significant impairment criterion; timing and latency of assessment and potential for recall bias; and combat experiences. Prevalence rates are also likely affected by issues related to PTSD course, chronicity, and comorbidity; symptom overlap with other psychiatric disorders; and sociopolitical and cultural factors that may vary over time and by nation.

Conclusions—The disorder represents a significant and costly illness to veterans, their families, and society as a whole. However, further carefully conceptualized research is needed to advance our understanding of disorder prevalence, as well as associated information on course, phenomenology, protective factors, treatment, and economic costs.

Keywords

posttraumatic stress disorder (PTSD); prevalence; combat; veterans; military

Despite a large body of literature on combat-related posttraumatic stress disorder (PTSD) there remains a lack of clarity regarding the prevalence of the disorder among military personnel and veterans. Even when there is consistency in issue or phenomenon, sufficient heterogeneity exists in the methodology of the studies that the conclusions drawn are often different, and data are available to support almost any position [1]. The 2007 debate in *Science*, and then later in

the *Journal of Traumatic Stress*, following Dohrenwend et al.'s [2] re-evaluation of the prevalence of PTSD among Vietnam veterans documented by Kulka et al., [3] in the National Vietnam Veterans Readjustment Survey (NVVRS), demonstrates an interesting range of perspectives that experts in this field hold about the same study [4–15]. Data from this one study have been interpreted to indicate a 15.2% [3], 9.1% [2], and 5.4% [11,13] point prevalence of PTSD among Vietnam veterans when measured in the late 1980s – a potential overestimation of nearly 300%. A recent article by Dobbs (2009) [16] in *Scientific American* and numerous media articles, radio and television stories, and blogs further reinforce that the issue of PTSD prevalence remains heatedly debated and continues to polarise the traumatology field and even politics in the U.S. This is partly due to the fact that the stakes are huge: major decisions about VA funding, research allocations, disability payments, and even society's perspective on war itself hinge on these data.

Recent studies of returning Operation Iraqi Freedom (OIF) and Operation Enduring Freedom (OEF) veterans [17–23] have generated numerous concerns regarding the long term mental health consequences of combat exposure. Historically, military conflicts have differed in duration, intensity, and public credibility of purpose [24] and the question regarding the similarity or difference of the psychological effects of combat across different combatant cohorts remains unanswered. Our goal is to provide a critical review of prevalence estimates of combat-related PTSD among military personnel and veterans, and of the relevant factors that may account for the variability of estimates within and across cohorts, including methodological and conceptual factors accounting for differences in prevalence rates across nations, conflicts/wars, and studies. It is not the aim of this paper to identify a single point prevalence estimate. Indeed, it is difficult to narrow the range of point prevalence for combat-related PTSD due to the significant heterogeneity of methods and samples used across studies, even within the same conflict/war cohort. The direct comparison between US and other NATO military forces across the same conflicts offers a unique approach to demonstrating how these factors interact with sociopolitical and cultural factors and may manifest differently across different samples and studies. Because of factors unique to the peacekeeping experience compared to active combat, the issue of PTSD prevalence in peacekeepers will not be included in this review.

For this review we examined MEDLINE and PsycINFO databases for literature on combat-related PTSD. The following terms were used independently and in combinations in this search: PTSD, combat, veterans, military, epidemiology, prevalence. This review is not intended to be exhaustive of all information pertaining to combat-related PTSD, a large and diverse body of literature.

Prevalence of Combat-Related PTSD in US Samples

The point prevalence of combat-related PTSD reported across studies of US combat veterans ranges from about 2–17%; and lifetime prevalence about 6–31% [2,3,17–19,25–37]. Point prevalence rates from veterans of the Vietnam War ranges from 2.2% – 15.2% [2,3,25–28]. Among Persian Gulf War veterans, PTSD rates reported are between 1.9% – 13.2% [30–39] and from veterans of the current conflict in the Middle East, PTSD point prevalence is reported from 4%–17.1% [17–19]. See Table 1 for a summary of studies.

US Combat veterans demonstrate a two- to four-fold increase in prevalence of PTSD compared to US civilians, depending on the subject cohort and research methodology used. Estimates of the point prevalence of PTSD among community samples of adults in the US is 5 – 6% [40, 41], with overall lifetime prevalence of the disorder estimated at 7.8%, [42]. Overall PTSD prevalence in Veterans Affairs (VA) primary care clinics is 11.5% [43]. The numbers of veterans seeking VA mental healthcare for PTSD has grown dramatically over the past ten years [44]; a large proportion of which is due to newly diagnosed Vietnam veterans [16], with

lowest rates among veterans of WWII [45]. Given that the VA serves a relatively small percentage of veterans nationally, and their patient population has a lower income and higher illness burden than the general population of veterans, one might expect that PTSD prevalence would be higher than among the entire population of veterans.

The Vietnam War—The debate regarding the prevalence of combat related PTSD was fuelled to a large extent by the publication of contrasting findings of two major US studies of Vietnam Veterans conducted in the late 1980's. The NVVRS [3] used multiple self-report instruments in addition to the Structured Clinical Interview for *DSM-III-R* PTSD (SCID-PTSD) to estimate PTSD prevalence in a community sample of 1,200 Vietnam veterans and Vietnam era non-serving veterans and civilians. The NVVRS estimated prevalence of lifetime PTSD among male Vietnam theatre veterans was 30.9%, and point rates at 15.2%. Against the expectations of post-combat psychological burden at the time, the high rates of PTSD obtained in the NVVRS represented a significant and unanticipated mental health problem [2,12,46, 47]. These elevated rates remain among the most frequently cited in the combat-related PTSD literature, although they have been criticised in relation to a number of theoretical and methodological factors [4,7,8,10–13,48]. Multiple revisions of the original NVVRS data on the basis of definitions of clinical severity have obtained lower prevalence rates [2,10].

Just prior to the release of the NVVRS findings, results from the Vietnam Experience Study (VES) were published [25]. Using a modified version of the DIS-PTSD (Diagnostic Interview Schedule for PTSD), this study compared rates of PTSD in a random sample of serving and non-serving Vietnam veterans ($n = 4,462$) matched in terms of level of education, employment, income, marital status, and satisfaction with personal relationships. The VES reported 14.7% of Vietnam veterans experienced combat-related PTSD at some time during or after military service, and 2.2% had the disorder during the month before the examination. This was dramatically lower than prevalence estimates of the NVVRS. According to the NVVRS researchers, the validity of the PTSD diagnoses made in the VES using the DIS-PTSD is unsupported in the literature [14]. Further discussion of the impact on prevalence estimates of different PTSD assessment instruments is included below.

Also conducted in the late 1980's, the findings of the methodologically rigorous Vietnam Era Twin Registry studies [49], received relatively little attention. In this study ($n = 6,744$ MZ & DZ pairs), monozygotic twins who were discordant for service in Vietnam were examined. Of the twins who had served in Vietnam, 16.8% had PTSD in contrast to 5% of twins who had not served in Vietnam. The prevalence of PTSD in the twins exposed to high rates of combat exposure was 9 times higher than in their non-combat siblings. Subsequent analyses from this study have shown a significant influence of genetic heritability on symptom expression of PTSD [50,51]. This Twin study is now in the early stages of being replicated and extended almost twenty years later [52].

Operation Desert Storm (The 1991 Gulf War)—In later war cohorts the estimates of PTSD prevalence are also mixed. The 8-month, 1990 conflict in the Persian Gulf was unique in US Armed Forces history because 17% of the 694,000 personnel were from Reserve and National Guard Units. For the 40 days of air warfare and 5 days of ground combat, there were fewer than 200 battle deaths and relatively few casualties inflicted upon US military personnel. However, unlike previous veterans, Gulf War veterans were subject to a variety of potentially harmful environmental exposures, both natural and man-made [33]. Researchers generally report lower rates of PTSD among Gulf War veterans than among military personnel from other wars [32,33,38,39,53].

To investigate mental health outcomes from the Gulf war, an epidemiological study (The National Health Survey of Gulf War Era Veterans and Their Families) was conducted on a

randomly chosen sample of 15,000 Gulf War and 15,000 non-war veterans selected from the total number of US troops who were deployed in the Persian Gulf area during the Gulf War [26,33,34,36]. In 1995, A total of 11,441 Gulf veterans (75%) and 9,476 non-Gulf veterans (64%) selected for the first wave of evaluation, participated. Five to seven years post-deployment, 10.1% of deployed Gulf War troops (compared to 4.2% of non-deployed era veterans) were estimated to have PTSD during the month prior to completing the self-administered PCL (PTSD checklist), and general health symptom questionnaire [34]. Toomey et al. [36] evaluated a subset ($n = 2,189$) of the same Gulf veterans using retrospective self-report and face-to-face psychological examinations conducted about 10 years after deployment (i.e., 1998–2001). The war-onset prevalence of PTSD was estimated at 6.3% for deployed veterans and 1.1% in the non-Gulf deployed era veterans. Ten years after the end of the Gulf War, 1.8% of deployed veterans who had war-onset PTSD remained symptomatic. Within this cohort, the retrospective estimates of the prevalence of PTSD decreased over time. In contrast, Wolfe, Proctor, et al., [39] reported a prospective PTSD prevalence of 3–7% among New England army veterans of the Persian Gulf War within 5 days of return from deployment, increasing at follow-up 18–24 months later to 8% – 16%; a phenomenon also noted by Gray, Bolton & Litz [54].

A curious outlier, the Iowa Persian Gulf Study Group [30] reported PTSD prevalence estimates of 1.9% in a ($n=3,969$) sub-sample of GW veterans. Such an anomalous result has been suggested to be the consequence of a study participation bias and response bias which may have occurred given the contemporary media emphasis on reporting the less specific, somatically focused symptoms of “Gulf War Syndrome” at the time [53–55]. The variable course of the disorder, coupled with methodological, exposure and cohort differences have influenced the heterogeneity of PTSD prevalence rates among Gulf war veterans, as they have done in Vietnam veterans.

OIF & OEF—Using the self-report PCL, Hoge, Castro, Messer et al. [18] evaluated the psychological sequelae of current US Army & Marine commitments to Iraq (OIF) and Afghanistan (OEF). In this study, PTSD was present in 18.1% – 19.9% of returning OIF soldiers, 11.5% of returning OEF soldiers, and 9.4% of soldiers prior to deployment, but when stricter criterion measuring symptom severity were included these rates dropped to 12.2% – 12.9%, 6.2% and 5% respectively. Unlike earlier studies, Hoge et al.’s [18] study estimated combat-related PTSD while fighting continued, and incorporated cross-sectional pre-deployment data. Such contemporaneous data collection may reduce the potential impact of a wide range of influencing factors such as media contagion effects, social influence or peer contamination, recall bias, participation bias, and exaggeration or falsification due to compensation or litigation claims. In a later study of deployed US personnel ($n = 303,950$), the presence of PTSD was measured by scoring 2 or more on the four item PTSD scale to obtain estimates of PTSD at 9.8% for OIF, 4.7% for OEF, and 2.1% for other locations [17]. In this study, prior trauma exposure was not controlled for and false-positives likely since such measures are designed to err in that direction.

In a population survey of 103,788 veterans active from 2001–2005 and newly registered with VA, 13% of the study population were diagnosed with PTSD [19]. Rates were highest in the veterans aged 18–24 years and lowest in veterans aged 40+ years. Another large scale study of 50,184 active military duty and Reserve/National Guard personnel reported an overall prevalence of new-onset PTSD based on the self-report PCL of 7.6 – 8.7% among those who were both deployed and reporting combat exposures, compared to 1.4 – 2.1% of deployers not reporting combat exposures and 2.3 – 3.0% of non-deployers [56]. Findings revealed higher rates of PTSD among personnel who were younger, less educated, enlisted, current smokers, and problem drinkers; with no effect found for race or service component (i.e., active duty vs.

Reserve/National Guard). Findings highlight the importance of specific combat exposures over mere war-zone deployment as contributing to new onset PTSD.

Detracting from the value of these recent studies are concerns that results 1) are based entirely on self-report checklists, which have differential validity from structured psychiatric interviews [57]; 2) may not incorporate information regarding functional impairment (required for *DSM-IV* diagnoses); and 3) do not inquire about exposure to prior trauma. Failing to take such factors into account may result in overestimated rates of combat-related PTSD [2,10,58,59].

Prevalence of Combat-Related PTSD among Samples from Other Western Nations

The prevalence estimates in studies of combat veterans of Western nations other than the US have smaller ranges and lower ceilings (see Table 1). The 2002 Canadian Forces Mental Health Survey of Regular Forces reported lifetime prevalence estimates of 7.2% and current estimates (i.e., within the last year) at 2.7%. For those who have completed three or more operational tours, the figures rise to 10.3% in their lifetime, and 4.7% over the previous year [60].

In Australian samples, PTSD prevalence in Australian Defense Force (ADF) Vietnam veterans has been reported at around 21% lifetime and 12% current [61]. Ten to 15 years post Gulf-War deployment, 5.4% of 1,871 ADF Gulf-War Veterans have current PTSD [62]. Both estimates were obtained through structured clinical interviews. Estimates of PTSD in ground forces of the ADF serving in Iraq & Afghanistan are as yet unpublished however, Royal Australian Navy (RAN) estimates of PTSD in the 1,739 sailors deployed to the Middle Eastern Area of Operations (MEAO) between 2001–2005 was 1.6% in total (corrected) as measured by scores >50 on the PCL [63]. From UK based research, Lee et al. [64] reported 12% prevalence of PTSD, based on a psychiatrists' assessment among a consecutive sample of UK Gulf war veterans ($n = 3,000$) attending Veterans Medical Service between 1993 and 2001. Jones et al. [65] mailed the PCL and GHQ-12 to a random sample of UK veterans ($n = 2,873$; including all branches) deployed after 1999. They reported that 53% of their sample had deployment experience and 2.5% of their total sample had PTSD in 2002. In the largest population based study of UK combat veterans thus far, Hotopf et al., [20] reported health outcomes in a random sample of UK armed forces personnel deployed to the 2003 War in Iraq ($n = 4,722$) and personnel not deployed ($n = 5,550$), who were demographically similar in all other ways. Participants completed a 28-page questionnaire on various health concerns, general well-being and alcohol consumption as well as PTSD symptoms. Findings for current rates of PTSD were 4% of the sample. In a later study of the same sample, differences in prevalence rates between OIF deployed regular UK army personnel and OIF deployed UK army reservists was 4% to 6% respectively [21]. Iversen et al. [22] also report 3.7% prevalence PTSD among UK veterans ($n = 4,662$) of Iraq.

The findings of PTSD prevalence in the 2.5 – 6% range are in contrast to Hoge et al.'s [18] finding of 17.1% among US veterans, when using an identical measure and case definition. In all UK-based studies, personnel from regular forces who had a combat role and/or contact with the enemy were more likely to have had PTSD symptoms than those deployed in other roles [20,22,23,65]. One possible reason for the disparity between US and UK samples lies with differences in the frequency, duration and intensity of combat exposure during deployment. However, other UK based research argues that it is premature to conclude that significant deleterious effects are an inevitable outcome of going to war [20,66]. Differences between UK and US estimates may also be related to pre- and post-exposure factors including childhood vulnerability, sociopolitical variables and military training [21,67–69].

Methodological and Conceptual Factors

The 2–17% point prevalence for PTSD represents significant heterogeneity in numbers of affected veterans, even within the same war cohort. It has been suggested that the sources of variance are largely the results of methodological and conceptual factors that differ across studies, as is demonstrated in the progressive scaling down of estimates based on the NVVRS [70]. In the next section we address more generally a number of methodological and conceptual factors that differ across studies and may help to explain the wide range of combat-related PTSD prevalence across these studies.

Sampling Strategies

Sample Size—Over time the sample sizes on which prevalence has been determined have increased dramatically. Perhaps the most frequently cited point-prevalence of PTSD is from the NVVRS, which is based on only 1,200 subjects compared with Seal et al.'s [19] sample 103,788 OIF/OEF veterans. To what extent were the NVVRS' 1,200 (and then the smaller sub-sample over-represented by high combat exposure and probable PTSD [26]) genuinely representative of the 2.7 million Vietnam veterans? Despite the complex statistical randomisation procedures used, such figures may not logically appear representative, or even at face-value, seem adequate to extrapolate to such a large population. On the other hand, studies with very large samples rely on self-report measures over clinical interviews and are thus, methodologically less rigorous.

Selection/Participation Bias—Maximising response rates is important for survey research to be considered valid and reliable. Given that avoidance of trauma cues is a symptom of the disorder, those with PTSD may have avoided participation in epidemiological studies. The mass pre- and post-deployment psychiatric screening programs seen in the recent US studies are likely to eliminate such issues from US studies, however other western nations do not necessarily conduct mass-screening of their armed forces and this issue may remain of interpretive concern. Participants who are voluntarily assessed and remain anonymous may have fewer disincentives for candid self appraisal; whereas those who are compulsorily screened may have more incentives to minimize distress including stigma from disclosure, avoidance of delays to post-deployment reunion, and the possibility of long-term consequences to future employment opportunities in and outside the military [17]. Alternatively, combat veterans who are ill may assign more salience to memories of combat, whereas those who are well may minimize combat exposure [53,55]. It is possible that veterans who most strongly associate their symptoms with their war experience may be prone to inflate the relationship between these variables [71]. Consequently, due to the demonstrated relationship between affective state and recall, perception and cognitive processing [71–73], the potential for sampling bias among those who do participate in research studies may increase. Thus far, research suggests that the influence of low participation rates [70] or non-responders [34,66, 73,74] is unlikely to affect such studies' larger outcomes.

Comparison Groups—Appropriate comparison groups are difficult to identify in military research, as comparisons between military and civilian groups are often unsuitable. The military may attract high risk-takers, exhibit a “healthy warrior bias,” and/or lack the representative racial and socioeconomic diversity of the general population [53,75]; yet often age-matched civilian peers serve as comparison groups in military mental health research [65].

As a comparison group, reservists can also be distinct from regular military personnel. They may not maintain the same levels of fitness, have minimal deployment experience, be deployed to roles for which they are untrained, be assigned dangerous combat duties which they did not

expect, and may face unique social and personal stressors when deploying or returning home [20,21,76]. In the past, soldiers who were physically or mentally ill were unlikely to be deployed [65,73,77]. Yet, armed forces are relying on larger numbers of reserve personnel whose pre-deployment health status may be less assured, and there may be a possible association between combat-related psychiatric disorders and reservist status [20,21,32,78, 79]. Research has demonstrated that elite forces and officers, who are typically better trained and more cohesive, have better fitness and higher motivation, and frequently have lower levels of PTSD than lower ranks, general personnel and reservists [22,56,66]. Findings from research on Gulf War veterans demonstrate differences in levels of post-deployment mental health and psychosocial problems among National Guard and Reserve personnel involved in the 1990 Persian Gulf War compared to active-duty troops [32,34,35]. On the other hand, one large recent study found no PTSD prevalence differences between active duty personnel and reservists [56].

Exposure to combat risk has been one explanation for the differences in PTSD estimates observed between the US and other nations' combat forces. Personnel from different nations are deployed to different areas, which may differ in risk levels and combat intensity. However, it is unlikely that non-US forces are not exposed to significant risk. In fact, several studies report that high risk and significant exposure occurs for personnel from all nations deployed to war zones [22,73], and as such, the differential exposure argument alone is insufficient to explain PTSD prevalence differences. Contrasting differences in prevalence results from US and UK force samples, Hotopf & Wessely [53] report that unlike US combat forces deployed to the Persian Gulf war, in most cases even non-Gulf war deployed veterans in UK studies had been deployed to other combat zones (e.g., Falkland Islands, Northern Ireland, Somalia, Kosovo), and therefore exposed to risk previously. They and others have argued that because of differences in pre-, and post-military experiences, not combat exposure alone, it may be inappropriate to make too much of such comparisons [21,65,76,80].

Measurement Strategies

Estimates of combat-related PTSD prevalence depend heavily on how symptoms and impairment are measured [81–83], the subsequent reliability and validity of the instruments themselves, and their lay versus expert administrators [2]. The VES study was criticised for using lay interviewers to administer a single abbreviated measure of PTSD (the abbreviated Diagnostic Interview Schedule; DIS-PTSD) to estimate prevalence, compared to the multicomponent assessment process administered by experienced clinicians, used in the NVVRS [2,84]. The DIS-PTSD measure has been argued to underestimate the rate of PTSD in civilian and military personnel [6,42]. However, the authors of the NVVRS acknowledged that “when instrumentation and scoring procedures matched to those used by CDC were used with NVVRS data, the resulting estimates of current prevalence were statistically indistinguishable from those published from the CDC VES (p. 556)” [84]. Clearly the type of measures used, their purpose for use as either screening or diagnostic tools, the anonymity or voluntary nature of their administration, in addition to the timing of administration, are all important mediating factors which contribute to the variability observed and the subsequent concern regarding the overarching definition of the PTSD construct [16,85].

Structured Psychiatric Interviews vs. Self-Report Instruments—Differences in PTSD prevalence have also been attributed to whether data were obtained via “self-report” measures or “structured psychiatric interviews,” as there is good data that the former overestimate PTSD. This is, at least in part, because few self-report measures attempt to assess the extent of functional impairment. In a sample of war-zone deployed Dutch soldiers, PTSD prevalence from structured psychiatric interviews was 41% lower than estimates from self-report questionnaires [58]. Similar findings were noted in a post-9/11 sample of NYC adults,

where it was found that 4.1% obtained a score indicative of “probable PTSD” on a self-report instrument (the PCL) whereas less than half that (1.7%) met full symptom criteria on a structured psychiatric interview for PTSD [57]. The PCL is a commonly used self-report measurement tool to screen for PTSD and a clinical cut-off of ≥ 50 is typical (e.g. [18,32]. However some studies have demonstrated the overestimation of PTSD by the PCL with cut-off ≥ 50 [62,65]. Other studies have reported lower estimates of PTSD when assessed using structured clinical interviews (i.e. Clinician-Administered PTSD Scale (CAPS)) compared to rates obtained via self-report measures (i.e. PCL). They consider the former the “gold standard” for diagnosing mental disorders, and that the estimates obtained through structured interview are therefore more accurate [34]. Thus, studies based entirely on self-report measures that do not evaluate clinical impairment [17,18] may represent significant overestimates of PTSD prevalence.

The *DSM-IV* Clinically Significant Impairment Criterion—The *DSM-IV*’s introduction of the requirement that a trauma exposed individual must appraise their exposure with horror, fear or helplessness, and that they must be significantly impaired as a consequence of their exposure for a diagnosis of PTSD to be made, has ultimately decreased the prevalence rate of PTSD and is a major explanatory factor for discordant prevalence estimates from studies using *DSM-III-R* criteria which did not require the “clinically significant impairment” criterion for a diagnosis of PTSD. Studies have demonstrated that PTSD rates are reduced substantially when adjusted for PTSD symptoms that are unrelated to functional impairment [2,18,58]. Some assert that Dohrenwend et al.’s [2] GAF score of 7 as a cut-off for positive PTSD is sufficiently conservative to capture genuine cases of impaired veterans [8,9], while others disagree [4,7, 13]. McNally [11,13] argues that the NVVRS PTSD prevalence drops from 9.1% to 5.4% (a 65% reduction of the original estimate) when a “clinically significant functional impairment” criterion (e.g., score of 1–6 on a 10-point global assessment of functioning) is applied to Dohrenwend et al.’s [2] reanalyses of the NVVRS data. Data from a large community-sample of civilians show that inclusion of the clinical significance criterion in *DSM-IV* reduces the conditional probability of PTSD, given exposure to trauma, by about 30% [83]. This particular issue however is much disputed due to its centrality to the debate surrounding the construct of PTSD as a whole [16,46,86] and further discussion is beyond the scope of this paper.

Timing and Latency of Assessment and Potential for Recall Bias—The delay between combat exposure and the assessment of symptoms is a significant issue [20,87]. The validity and reliability of PTSD prevalence estimates may be compromised by: a) selection bias associated with normal military personnel attrition; b) poor recall; c) intervening influences of the media and popular opinion; d) systemic influences, such as VA disability incentives; e) the influence of emotional and psychological state on memory and cognition; and f) other sociopolitical and cultural factors that may influence reporting.

In the epidemiological studies conducted with Vietnam veterans, data was not collected until 10–20 years after the war-zone deployment and relied on retrospective self-reports of combat experiences and symptoms [47]. Concern regarding the reporting of combat exposure has been minimized in the NVVRS and largely mitigated for other community samples via the recent reanalyses by Dohrenwend et al. [2], which included a careful examination of archival records and supported the general reliability of self-reported combat exposures in a community sample of Vietnam veterans. Similarly, although Gulf War veterans’ health problems began surfacing in the early 1990s, the first large scale research into the health consequence of participation in this war did not occur until five years after the war ended in both the US [88] and the UK [53].

Recall biases are a direct threat to validity of war-related research. Typically, studies linking combat exposure to PTSD have relied upon retrospective self reports of veterans [74,89].

Prevalence studies treat such accounts as reliable and accurate despite the presupposition that respondents' clinical state was altered by the trauma [90]. In fact, research shows current clinical state, self-perceived health status, and salience of the memory all affect how traumatic experiences are recalled [39,55,71,72,91,92]. Because perception of illness can affect recall or interpretation of events leading to illness, veterans who feel unwell may be more likely to recall experiencing combat exposures during deployment [39,55,93]. Koenen et al. [94] examined consistency in combat exposure reports and their relation to PTSD symptoms in Vietnam Veteran American Legionnaires ($n = 1,462$) who responded to two mailed surveys, 14 years apart (1984, 1998). Combat exposure reports were highly reliable (test–retest correlation = 0.87). However, changes in exposure reporting were related to changes in PTSD symptoms; specifically re-experiencing symptoms. Other research demonstrates 88% of Gulf War veterans remembered traumatic events differently than how they had originally reported them one-month after their war service and that increased reporting was positively associated with reports of PTSD symptoms [71].

Over time, the delay between exposure to combat and time of assessment has been reduced. Compared to the data collected on Vietnam veterans 10–20 years after the end of the war, Hoge et al. [17,18] and Seal et al. [19] report the findings of combat exposure and symptom presentation immediately post-deployment, while the fighting continues and increasingly, also have access to pre-deployment data [18] Rona et al. [23] and Iversen et al. [22] also collected post-deployment data longitudinally and within one-month post-return. Such approaches should result in stronger validity of conclusions. However, timing of assessment is a complex issue in that data indicate prevalence of mental health problems may increase significantly at 120-days post-deployment relative to immediate post-deployment assessments [16,84]. While a long delay between exposure and assessment is clearly undesirable, immediate assessment post-deployment may also under-report the prevalence of PTSD due to the observation that soldiers may feel relief and optimism about their imminent return home, thereby masking immediate distress, or may purposefully minimise distress to avoid delaying their return and subsequent leave entitlements [87].

Finally, research demonstrates that attrition rates from the military after active duty for health and socioeconomic reasons are approximately 5% per year from UK armed forces which translates to large numbers of veterans being lost from research sampling when post-deployment studies are delayed [53]. Hoge et al. [95] report that within 6-months after hospitalization, 47% of soldiers who were hospitalized for the first time in 1996 with a mental disorder as a primary diagnosis, left military service. In comparison, only 12% of personnel hospitalized for non-psychiatric medical illnesses left the service in the same period. Therefore in cross-sectional studies such as those cited above, it is possible the samples may reflect the self-selecting out of the unwell, the homeless, and the substance using—all factors which may be associated with psychiatric distress and disorder, thereby obscuring genuine prevalence rates.

Pre-, Peri-, and Post-Deployment Factors

Factors associated with combat are assumed to be critical mediators in the expression of PTSD, including deployment-specific factors such as intensity and duration of combat. Risk factors for PTSD also include features of personal vulnerability, other war zone experiences, and a wide range of socio-cultural and situational factors. Pre-, peri-, and post-deployment factors interact to influence symptom development and adaptation, and significantly affect PTSD prevalence and course [22,23,76,96,97].

Pre-Deployment Factors—The expression of PTSD is partially mediated by genetic factors [50,94,98–101]. Other relevant pre-deployment factors include age and education at time of

deployment, gender, race, early conduct problems, intelligence, childhood adversity, family history of psychiatric disorder, pre-deployment psychological and physical health, poor social support after a trauma, and personality pathology [23,102]. Our understanding of the relative contribution of each of these factors is still undeveloped.

Peri-Deployment Factors—Relevant peri-deployment factors include characteristics of the combat arena, intensity and duration of fighting, logistical/military support, and unit characteristics, and each combat deployment in the same cohort appears to have unique types of hazards [20,53] suggesting that deployment-specific stressors may contribute to the differences between groups.

A dose-response relationship between trauma intensity and PTSD severity has been hypothesized and has some empirical support [2,17,18,20,22,23,73]. For example, one study found that combat infantry units deployed to Iraq ($n = 2,530$) reported significantly greater combat exposure than those deployed to Afghanistan ($n = 3,671$), and accordingly the percentage who met screening criteria for major depression, generalized anxiety, or PTSD was significantly higher after duty in Iraq (15.6% – 17.1%) than after duty in Afghanistan (11.2%), or before deployment to Iraq (9.3%) [18]. Prevalence of PTSD was higher after deployment to Iraq (12.2 – 19.9%) than before (5.0 – 9.4%), and it increased in a linear manner as exposure to fire-fights increased (e.g., no fire-fights = 4.5% PTSD prevalence; to > 5 fire-fights = 19.3%). Similar findings have been replicated in a later US ($n = 303,905$) population-based study [17]. Iversen et al. [22] have also reported a strong association between time spent in a forward area and PTSD in UK soldiers deployed to Iraq. Smith et al. [56] also found that specific combat exposures, rather than mere war-zone deployment, were associated with PTSD in a population-based cohort of US military personnel ($n = 50, 184$).

The fact that most personnel exposed to combat do not develop PTSD, and that the dose-response relationship between combat and pathology is not always linear, obviously suggests the development and course of PTSD is not solely determined by the features of the combat itself [20,22,59,66]. In fact, PTSD is highly dependent on cognitive appraisals of threat [103], and the risk of PTSD appears to be substantially reduced by high unit morale and good leadership in the war-zone [22,23] and the perception of support and group cohesion among military personnel [23,66].

Post-Deployment Factors—Post-military factors also appear to contribute to course, presentation, and chronicity of PTSD [76]. These post-military factors have become central to the prevalence question, particularly in light of the latency between exposure and assessment seen in earlier research. Relevant post-military factors include individual symptoms, circumstances, stressors, and social support, as well as the post-deployment social and political environment.

Sociopolitical and Cultural Factors

Historical, Contextual, and Socioeconomic Factors—The diagnosis of PTSD, first added to the *DSM* in 1980, was largely the result of attempts to account for the challenging impairment presented by Vietnam veterans at the time of homecoming [46,59]. In the immediate post-Vietnam era, compensation for significant functional impairment was difficult to obtain other than for observable physical injuries and entrée to VA medical services were possible only via a “war-related” disorder [47]. Thus, the development of the clinical conceptualization of PTSD has been heavily influenced by socio-economic and political factors [46,104–106]. More recently there has been concern that healthcare systems and disability policies encourage psychiatric illness. In fact, some argue that the US VA system has failed to benefit from the lessons of 20th century military psychiatry regarding social expectations and

incentives [46,59,90,107–112]. Certainly recent administrative trends regarding PTSD disability claims are troubling. Among the relatively small subset of veterans seeking mental health treatment in VA clinics, most (up to 94%) also concurrently apply for PTSD disability benefits [113]. Further, the number of veterans receiving VA disability payments for PTSD increased 79.5% from 1999 to 2004, while all other disabilities increased only 12.2% during that same period [114]. Different compensation and support structures for veterans of non-US armed forces may partly explain different estimates of PTSD prevalence [16].

A particular concern is that veterans' disability-seeking status may influence their clinical presentation in healthcare settings. Some veterans seeking evaluation in US VA medical centers may exaggerate, misrepresent or falsify their combat experiences [48,109,115], and some have created fictitious military service altogether [107,116–118]. Studies also show that veterans evaluated for PTSD in VA clinical settings may malingering or exaggerate psychiatric symptoms, especially when disability compensation is at stake [59,110,113,119–121]. Freeman et al. [120] found that 53% of veterans in a PTSD treatment-seeking clinical sample appeared to be malingering psychopathology on a structured forensic psychiatric interview. For a trenchant review of the effects of government disability pensions and benefits on veterans' psychiatric functioning see Jones and Wessely [110].

The effect of disability-seeking for subjective ailments like PTSD on the behavior of claimants remains a significant issue in research and practice, and a matter of discussion among a number of US policy agencies [114,122]. Researchers conducting clinical trials have been encouraged to control for or exclude compensation-seeking veterans from clinical research [117,120,123]. However, the extent to which secondary gain may influence prevalence rates is unclear and outcomes between clinical and community samples are not necessarily generalizable [1,13]. While certain VA system policies or incentives may have an influence on treatment-seeking veterans (a small subset of all veterans), they may have relatively little impact on the larger veteran population represented in community samples. Thus, it is difficult to estimate how healthcare or other government policies might shape larger attitudes and reporting styles related to PTSD in community samples by influencing social attitudes and expectations.

Media and Popular Culture Influences—Media stories and societal expectations influence symptom reporting and effect the validity of any retrospectively determined relationship between the trauma and psychiatric symptoms [55,62,89,92]. Research into significant historical events and memory recall has demonstrated that frequently repeated media-based narratives may reinforce inaccurate memories and beliefs about the event [53,92,124] and the impact on help-seeking following a Hollywood glamorization of a psychologically tortured veteran can only be speculated [125]. Wessely and Jones [47] cite the Vietnam War as an example where the public's changing opinion surrounding the war was likely to have influenced “a gradual ‘inflation’ of traumatic memories to fit with the changing views (p. 12).” See also Burkett & Whitley [107], Shephard [46], Jones & Wessely [110], McHugh & Triesman [126], and Satel & Frueh [105] for further discussion.

PTSD Onset and Course

The vast majority of combat veterans do not ever develop PTSD. Further, studies that estimate both lifetime and current prevalence of PTSD typically find that roughly half of the veterans who had PTSD at some point in the past do not meet diagnostic criteria for current PTSD [2,3,25,34,61,85]. On the other hand, retrospective studies in samples of Holocaust survivors [127], Korean war veterans [128], Lebanon War veterans [129] and World War II veterans [130–132] indicate that the duration of “chronic PTSD” can span an entire adult lifetime; up to 50 years after combat exposure [133]. The phenomenon of “delayed-onset” PTSD remains somewhat controversial. While there is some empirical evidence to support the phenomenon

[54], a recent review of delayed-onset PTSD found the phenomena in the absence of prior symptoms was rare [134]. In fact, one recent study found that delayed-onset PTSD was extremely rare among veterans more than one year after combat exposure, and there was no evidence of PTSD symptom onset six or more years after combat [135].

Symptom Overlap with Other Psychiatric Disorders

While Criterion A “the traumatic event” [136], is a unique requirement for a *DSM-IV* PTSD diagnosis compared to other psychological disorders, PTSD is classified as an anxiety disorder and conceptually shares a significant overlap between symptoms of other mood and anxiety disorders. This overlap may partially account for the high co-morbidity evident in PTSD [121]. High comorbidity of PTSD with psychiatric and medical diagnoses have been well documented, including high rates of co-morbid substance abuse, anxiety, mood, and personality disorders [3,30,43,95,132,137–139]. The high degree of symptom overlap between PTSD and other Axis I psychiatric disorders has been suggested to indicate a problem with the conceptual foundation of the disorder, one of a number of challenges raised recently regarding the question of whether PTSD is actually a distinct and unique form of psychopathology [48, 86,126,140,141].

Conclusions

The point prevalence of combat-related PTSD in studies of US military veterans ranges from about 2 – 17%, and the range tends to have a lower ceiling and is more narrow among veterans of other Western nations. There is considerable variability in rates of prevalence due to differences in sampling strategies; measurement strategies (e.g., use of structured psychiatric interviews versus self-report measures); inclusion and measurement of the *DSM-IV* clinically significant impairment criterion; latency of assessment and potential for recall bias; and the combat experience (including deployment factors, and pre-, peri-, and post-deployment factors). Prevalence rates are also likely affected by issues related to PTSD course, chronicity, and comorbidity; symptom overlap with other psychiatric disorders; and sociopolitical and cultural factors. Regardless of the “true” prevalence of combat-related PTSD, the disorder is associated with severe functional impairment (both occupational and social), high co-morbidity with other psychiatric disorders, high medical comorbidity, and/or reduced quality of life for the veterans who suffer from it. Thus, it represents a significant and costly illness to veterans, their families, and society as a whole. Further methodologically rigorous research is needed to better elucidate our understanding of the prevalence, course, phenomenology, protective factors, and costs associated with combat-related PTSD.

Acknowledgments

This work was partially supported by Australian-American Fulbright Commission Postgraduate Award to Ms. Richardson, Grant MH074468 from the National Institute of Mental Health to Dr. Frueh, and awards from the McNair and Menninger Foundations to Dr. Frueh. We thank Steve Herrera for his technical assistance. All views and opinions expressed herein are those of the authors and do not necessarily reflect those of our respective institutions or the U.S. Department of Veterans Affairs.

References

1. Kaylor JA, King DW, King LA. Psychological effects of military service in Vietnam: a meta-analysis. *Psychol Bull* 1987;102(2):257–271. [PubMed: 3659223]
2. Dohrenwend BP, Turner JB, Turse NA, Adams BG, Koenen KC, Marshall R. The psychological risks of Vietnam for U.S. veterans: A revisit with new data and methods. *Science* 2006;313:979–982. [PubMed: 16917066]

3. Kulka, RA.; Schlenger, WE.; Fairbank, JA.; Hough, RL.; Jordan, BK.; Marmar, CR., et al. Trauma and the Vietnam War generation: Report of findings from the National Vietnam Veterans Readjustment Study. New York: Brunner/Mazel; 1990.
4. Buckley TC. PTSD and Vietnam veterans. *Science* 2007;315:185.
5. Dohrenwend BP, Turner JB, Turs N, Adams BG, Koenan KC, Marshall R. PTSD and Vietnam Veterans. *Science* 2007;315:186–187.
6. Dohrenwend BP, Turner JB, Turs N, Marshall R. Continuing controversy over the psychological risks of Vietnam for U.S. veterans. *J Trauma Stress* 2007;20:449–465. [PubMed: 17721952]
7. Frueh BC. PTSD and Vietnam Veterans. *Science* 2007;315(5809):184–187. author reply 184–187. [PubMed: 17225296]
8. Kilpatrick DG. PTSD and Vietnam Veterans. *Science* 2007;315(5809):184–187. author reply 184–187. [PubMed: 17225294]
9. Kilpatrick DG. Confounding the critics: the Dohrenwend and colleagues reexamination of the National Vietnam Veteran Readjustment Study. *J Trauma Stress* 2007;20(4):487–493. [PubMed: 17721965]
10. McNally RJ. Psychology. Psychiatric casualties of war. *Science* 2006;313(5789):923–924. [PubMed: 16917047]
11. McNally RJ. PTSD and Vietnam veterans. *Science* 2007;315:185–186.
12. McNally RJ. Revisiting Dohrenwend et al. 's revisit of the National Vietnam Veterans Readjustment Study. *J Trauma Stress* 2007;20(4):481–486. [PubMed: 17721958]
13. McNally RJ. Can we solve the mysteries of the National Vietnam Veterans Readjustment Study? *J Anxiety Disord* 2007;21:192–200. [PubMed: 17045777]
14. Schlenger WE, Kulka RA, Fairbank JA, Hough RL, Jordan BK, Marmar CR, et al. The psychological risks of Vietnam: the NVVRS perspective. *J Trauma Stress* 2007;20(4):467–479. [PubMed: 17721970]
15. Vermetten E, Bremner JD, Skelton L, Spiegel D. PTSD and Vietnam Veterans. *Science* 2007;315(5809):184. [PubMed: 17218506]
16. Dobbs D. The post-traumatic stress trap. *Sci Am* 2009;300(4):64–69. [PubMed: 19363922]
17. Hoge CW, Auchterlonie JL, Milliken CS. Mental health problems, use of mental health services, and attrition from military service after returning from deployment to Iraq or Afghanistan. *JAMA* 2006;295(9):1023–1032. [PubMed: 16507803]
18. Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL. Combat duty in Iraq and Afghanistan: Mental health problems and barriers to care. *NEJM* 2004;351:13–22. [PubMed: 15229303]
19. Seal KH, Bertenthal D, Miner CR, Sen S, Marmar C. Bringing the war back home: mental health disorders among 103,788 US veterans returning from Iraq and Afghanistan seen at Department of Veterans Affairs facilities. *Arch Intern Med* 2007;167(5):476–482. [PubMed: 17353495]
20. Hotopf M, Hull L, Fear NT, Browne T, Horn O, Iversen A, et al. The health of UK military personnel who deployed to the 2003 Iraq war: a cohort study. *Lancet* 2006;367(9524):1–12. [PubMed: 16399129]
21. Browne T, Hull L, Horn O, Jones M, Murphy D, Fear NT, et al. Explanations for the increase in mental health problems in UK reserve forces who have served in Iraq. *Br J Psychiatry* 2007;190:484–489. [PubMed: 17541107]
22. Iversen AC, Fear NT, Ehlers A, Hacker Hughes J, Hull L, Earnshaw M, et al. Risk factors for post-traumatic stress disorder among UK Armed Forces personnel. *Psychol Med* 2008;38(4):511–522. [PubMed: 18226287]
23. Rona RJ, Hooper R, Jones M, Iversen AC, Hull L, Greenberg N, et al. Vulnerability, combat exposure and psychological symptoms in the Iraq war: A longitudinal study of UK Armed Forces. 2008 Manuscript submitted for publication.
24. Larson GE, Highfill-McRoy RM, Booth-Kewley S. Psychiatric diagnoses in historic and contemporary military cohorts: combat deployment and the healthy warrior effect. *Am J Epidemiol* 2008;167(11):1269–1276. [PubMed: 18436536]
25. Centers for Disease Control. Vietnam Experience Study: Health status of Vietnam veterans: psychosocial characteristics. *JAMA* 1988;259:2701–2707. [PubMed: 2833630]

26. Eisen SA, Griffith KH, Xian H, Scherrer JF, Fischer ID, Chantarujikapong S, et al. Lifetime and 12-month prevalence of psychiatric disorders in 8,169 male Vietnam War era veterans. *Mil Med* 2004;169(11):896–902. [PubMed: 15605939]
27. Schlenger WE, Kulka RA, Fairbank JA, Hough RL, Jordan BK, Marmar CR, et al. The prevalence of post-traumatic stress disorder in the Vietnam generation: a multimethod, multisource assessment of psychiatric disorder. *J Trauma Stress* 1992;5:333–363.
28. Thompson WW, Gottesman I, Zalewski C. Reconciling disparate prevalence rates of PTSD in large samples of US male Vietnam veterans and their controls. *BMC Psychiatry* 2006;6:19–29. [PubMed: 16670009]
29. Barrett DH, Doebbeling CC, Schwartz DA, Voelker MD, Falter KH, Woolson RF, et al. Posttraumatic stress disorder and self-reported physical health status among U.S. Military personnel serving during the Gulf War period: a population-based study. *Psychosomatics* 2002;43(3):195–205. [PubMed: 12075034]
30. Engel CC Jr, Ursano R, Magruder C, Tartaglione R, Jing Z, Labbate LA, et al. Psychological conditions diagnosed among veterans seeking Department of Defense Care for Gulf War-related health concerns. *J Occup Environ Med* 1999;41(5):384–392. [PubMed: 10337608]
31. Gray GC, Reed RJ, Kaiser KS, Smith TC, Gastanaga VM. Self-reported symptoms and medical conditions among 11,868 Gulf War-era veterans: the Seabee Health Study. *Am J Epidemiol* 2002;155(11):1033–1044. [PubMed: 12034582]
32. Iowa Persian Gulf Study Group. Self-reported illness and health status among Gulf War veterans. A population-based study. *JAMA* 1997;277:238–245. [PubMed: 9005274]
33. Kang HK, Mahan CM, Lee KY, Magee CA, Murphy FM. Illnesses among United States veterans of the Gulf War: a population-based survey of 30,000 veterans. *J Occup Environ Med* 2000;42(5):491–501. [PubMed: 10824302]
34. Kang HK, Natelson BH, Mahan CM, Lee KY, Murphy FM. Post-traumatic stress disorder and chronic fatigue syndrome-like illness among Gulf War veterans: a population-based survey of 30,000 veterans. *Am J Epidemiol* 2003;157(2):141–148. [PubMed: 12522021]
35. Stretch RH, Marlowe DH, Wright KM, Bliese PD, Knudson KH, Hoover CH. Post-traumatic stress disorder symptoms among Gulf War veterans. *Mil Med* 1996;161(7):407–410. [PubMed: 8754714]
36. Toomey R, Kang HK, Karlinsky J, Baker DG, Vasterling JJ, Alpern R, et al. Mental health of US Gulf War veterans 10 years after the war. *Br J Psychiatry* 2007;190:385–393. [PubMed: 17470952]
37. Wolfe J, Erickson DJ, Sharkansky EJ, King DW, King LA. Course and predictors of posttraumatic stress disorder among Gulf War veterans: a prospective analysis. *J Consult Clin Psychol* 1999;67(4):520–528. [PubMed: 10450622]
38. Barrett DH, Gray GC, Doebbeling BN, Clauw DJ, Reeves WC. Prevalence of symptoms and symptom-based conditions among Gulf War veterans: current status of research findings. *Epidemiol Rev* 2002;24(2):218–227. [PubMed: 12762094]
39. Wolfe J, Proctor SP, Erickson DJ, Heeren T, Friedman MJ, Huang MT, et al. Relationship of psychiatric status to Gulf War veterans' health problems. *Psychosom Med* 1999;61(4):532–540. [PubMed: 10443762]
40. Norris FH. Epidemiology of trauma: Frequency and impact of different potentially traumatic events on different demographic groups. *J Consult Clin Psychol* 1992;60:409–418. [PubMed: 1619095]
41. Resnick HS, Kilpatrick DG, Dansky BS, Saunders BE, Best CL. Prevalence of civilian trauma and posttraumatic stress disorder in a representative national sample of women. *J Consult Clin Psychol* 1993;61:984–991. [PubMed: 8113499]
42. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB. Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 1995;52:1048–1060. [PubMed: 7492257]
43. Magruder KM, Frueh BC, Knapp RG, Davis L, Hamner MB, Martin RH, et al. Prevalence of posttraumatic stress disorder in Veterans Affairs primary care clinics. *Gen Hosp Psychiatry* 2005;27:169–179. [PubMed: 15882763]
44. Rosenheck RA, Fontana AF. Recent trends in VA treatment of post-traumatic stress disorder and other mental disorders. *Health Aff (Millwood)* 2007;26(6):1720–1727. [PubMed: 17978391]

45. Frueh BC, Grubaugh AL, Acierno A. Age differences in posttraumatic stress disorder, psychiatric disorders, and healthcare service use among veterans in Veterans Affairs primary care clinics. *Am J Geriatr Psychiatry* 2007;15:660–672. [PubMed: 17670996]
46. Shephard, B. *A war of nerves: Soldiers and psychiatrists in the twentieth century*. Cambridge, Massachusetts: Harvard University Press; 2001.
47. Wessely S, Jones E. Psychiatry and the 'lessons of Vietnam': What were they, and are they still relevant? *War & Society* 2004;22:89–103.
48. McNally RJ. Progress and controversy in the study of posttraumatic stress disorder. *Annu Rev Psychol* 2003;54:229–252. [PubMed: 12172002]
49. Goldberg J, Eisen SA, True WR, Henderson WG. A twin study of the effects of the Vietnam conflict on alcohol drinking patterns. *Am J Public Health* 1990;80(5):570–574. [PubMed: 2327534]
50. True WR, Rice J, Eisen SA, Heath AC, Goldberg J, Lyons MJ, et al. A twin study of genetic and environmental contributions to liability for posttraumatic stress symptoms. *Arch Gen Psychiatry* 1993;50(4):257–264. [PubMed: 8466386]
51. Chantarujikapong SI, Scherrer JF, Xian H, Eisen SA, Lyons MJ, Goldberg J, et al. A twin study of generalized anxiety disorder symptoms, panic disorder symptoms and post-traumatic stress disorder in men. *Psychiatry Res* 2001;103(2–3):133–145. [PubMed: 11549402]
52. Magruder, KM. Personal Communication. 2007.
53. Hotopf M, Wessely S. Can epidemiology clear the fog of war? Lessons from the 1990–91 Gulf War. *Int J Epidemiol* 2005;34(4):791–800. [PubMed: 15911546]
54. Gray MJ, Bolton EE, Litz BT. A longitudinal analysis of PTSD symptom course: delayed-onset PTSD in Somalia peacekeepers. *J Consult Clin Psychol* 2004;72(5):909–913. [PubMed: 15482050]
55. Wessely S, Unwin C, Hotopf M, Hull L, Ismail K, Nicolaou V, et al. Stability of recall of military hazards over time: Evidence from the Persian Gulf War of 1991. *Br J Psychiatry* 2003;183:314–322. [PubMed: 14519609]
56. Smith TC, Ryan MA, Wingard DL, Slymen DJ, Sallis JF, Kritz-Silverstein D. New onset and persistent symptoms of post-traumatic stress disorder self reported after deployment and combat exposures: prospective population based US military cohort study. *BMJ* 2008;336(7640):366–371. [PubMed: 18198395]
57. Ruggiero KJ, Rheingold AA, Resnick HS, Kilpatrick DG, Galea S. Comparison of two widely used PTSD-screening instruments: Implications for public mental health planning. *J Trauma Stress* 2006;19:699–707. [PubMed: 17075907]
58. Engelhard IM, van den Hout MA, Weerts J, Arntz A, Hox JJ, McNally RJ. Deployment-related stress and trauma in Dutch soldiers returning from Iraq. Prospective study. *Br J Psychiatry* 2007;191:140–145. [PubMed: 17666498]
59. Wessely S. War stories: Invited commentary on.. Documented combat exposure of US veterans seeking treatment for combat-related post-traumatic stress disorder. *Br J Psychiatry* 2005;186(6):473–475.
60. Survey, CCH., editor. Statistics Canada. The Canadian Forces 2002 Supplement of the Statistics Canada. 2003.
61. O'Toole BI, Marshall RP, Grayson DA, Schureck RJ, Dobson M, Ffrench M, et al. The Australian Vietnam veterans health study: III. Psychological health of Australian Vietnam veterans and its relationship to combat. *Int J Epidemiol* 1996;25:331–340. [PubMed: 9119558]
62. Ikin JF, Sim MR, Creamer MC, Forbes AB, McKenzie DP, Kelsall HL, et al. War-related psychological stressors and risk of psychological disorders in Australian veterans of the 1991 Gulf War. *Br J Psychiatry* 2004;185:116–126. [PubMed: 15286062]
63. Rayner SPS. Prevalence of psychological trauma in operationally deployed Navy personnel: a baseline surveillance report. *ADF Health* 2005;6:81–84.
64. Lee HA, Gabriel R, Bolton JP, Bale AJ, Jackson M. Health status and clinical diagnoses of 3000 UK Gulf War veterans. *J R Soc Med* 2002;95(10):491–497. [PubMed: 12356969]
65. Jones M, Rona RJ, Hooper R, Wessely S. The burden of psychological symptoms in UK Armed Forces. *Occup Med (Lond)* 2006;56(5):322–328. [PubMed: 16720583]

66. Hughes JH, Cameron F, Eldridge R, Devon M, Wessely S, Greenberg N. Going to war does not have to hurt: preliminary findings from the British deployment to Iraq. *Br J Psychiatry* 2005;186:536–537. [PubMed: 15928366]
67. Iversen AC, Fear NT, Simonoff E, Hull L, Horn O, Greenberg N, et al. Influence of childhood adversity on health among male UK military personnel. *Br J Psychiatry* 2007;191:506–511. [PubMed: 18055954]
68. Schnurr PP, Lunney CA, Sengupta A. Risk factors for the development versus maintenance of posttraumatic stress disorder. *J Trauma Stress* 2004;17(2):85–95. [PubMed: 15141781]
69. McFarlane AC, Creamer M. Current knowledge about psychological trauma: A response to Milton. *ADF Health* 2006;7:78–82.
70. Stimpson NJ, Thomas HV, Weightman AL, Dunstan F, Lewis G. Psychiatric disorder in veterans of the Persian Gulf War of 1991. Systematic review. *Br J Psychiatry* 2003;182:391–403. [PubMed: 12724242]
71. Southwick SM, Morgan CA, Nicolaou AL, Charney DS. Consistency of memory for combat-related traumatic events in veterans of Operation Desert Storm. *Am J Psychiatry* 1997;154:173–177. [PubMed: 9016264]
72. Roemer L, Litz BT, Orsillo SM, Ehlich PJ, Friedman MJ. Increases in retrospective accounts of war-zone exposure over time: the role of PTSD symptom severity. *J Trauma Stress* 1998;11(3):597–605. [PubMed: 9690197]
73. Unwin C, Blatchley N, Coker W, Ferry S, Hotopf M, Hull L, et al. Health of UK servicemen who served in Persian Gulf War. *Lancet* 1999;353(9148):169–178. [PubMed: 9923871]
74. Iverson A, Dyson C, Smith N, Greenberg N, Walwyn R, Unwin C, et al. 'Goodbye and goodluck': The mental health needs and treatment experiences of British ex-service personnel. *Br J Psychiatry* 2005;186:480–486. [PubMed: 15928358]
75. Warner JT, Asch BJ. The record and prospects of the all volunteer military in the United States. *J Econ Perspect* 2001;15:169–192.
76. Friedman MJ, Schnurr P, McDonagh-Coyle A. Posttraumatic stress disorder in the military psychiatr. *Clin North Am* 1994;17:265–277.
77. Rona RJ, Fear NT, Hull L, Greenberg N, Earnshaw M, Hotopf M, et al. Mental health consequences of overstretch in the UK armed forces: first phase of a cohort study. *BMJ* 2007;335(7620):603. [PubMed: 17664192]
78. Smith TC, Corbeil TE, Ryan MA, Heller JM, Gray GC. In-theatre hospitalizations of US and allied personnel during the 1991 Gulf War. *Am J Epidemiol* 2004;159:1064–1076. [PubMed: 15155291]
79. Turner MA, Kiernan MD, McKechnie AG, Finch PJ, McManus FB, Neal LA. Acute military psychiatric casualties from the war in Iraq. *Br J Psychiatry* 2005;186:476–479. [PubMed: 15928357]
80. Killgore WD, Stetz MC, Castro CA, Hoge CW. The effects of prior combat experience on the expression of somatic and affective symptoms in deploying soldiers. *J Psychosom Res* 2006;60(4):379–385. [PubMed: 16581362]
81. Breslau N. Epidemiologic studies of trauma, posttraumatic stress disorder, and other psychiatric disorders. *Can J Psychiatry* 2002;47(10):923–929. [PubMed: 12553127]
82. Breslau N, Kessler RC. The stressor criterion in DSM-IV posttraumatic stress disorder: An empirical investigation. *Biol Psychiatry* 2001;50:699–704. [PubMed: 11704077]
83. Breslau N, Alvarado GF. The clinical significance criterion in DSM-IV post-traumatic stress disorder. *Psychol Med* 2007;37(10):1437–1444. [PubMed: 17445282]
84. Kulka RA, Schlenger WE, Fairbank JA, Jordan BK, Hough RL, Mamar CR, et al. Assessment of posttraumatic stress disorder in the community: Prospects and pitfalls from recent studies of Vietnam veterans. *Psychol Assess* 1991;3:547–560.
85. Ehlers A, Mayou RA, Bryant B. Psychological predictors of chronic posttraumatic stress disorder after motor vehicle accidents. *J Abnorm Psychol* 1998;107(3):508–519. [PubMed: 9715585]
86. Spitzer RL, First MB, Wakefield JC. Saving PTSD from itself in DSM-V. *J Anxiety Disord* 2007;21:233–241. [PubMed: 17141468]
87. Bliese PDW, Kathleen M, Adler AB, Thomas JL, Hoge CW. Timing of postcombat mental health assessments. *Psychological Services* 2007;4 (3):141–148.

88. Statement of Donna Heivilin Director of Planning and Reporting National Security and International Affairs Division. Gulf War Illness Federal Research Strategy Needs Reexamination 1998.
89. McCauley LA, Joos SK, Spencer PS, Lasarev M, Shuell T. Strategies to assess validity of self-reported exposures during the Persian Gulf War. Portland Environmental Hazards Research Center. Environ Res 1999;81(3):195–205. [PubMed: 10585015]
90. Mossman D. Veterans affairs disability compensation: a case study in counter therapeutic jurisprudence. Bull Am Acad Psychiatry Law 1996;24(1):27–44. [PubMed: 8891320]
91. Koenen KC. Genetics of posttraumatic stress disorder: Review and recommendations for future studies. J Trauma Stress 2007;20(5):737–750. [PubMed: 17955543]
92. McCauley LA, Joos SK, Lasarev MR. Gulf War unexplained illnesses: persistence and unexplained nature of self-reported symptoms. Environ Res 1999;81(3):215–223. [PubMed: 10585017]
93. Barsky A. Amplification, somatization, and the somatoform disorders. Psychosomatics 1992;33(1):28–34. [PubMed: 1539100]
94. Koenen K, Stellman SD, Dohrenwend BP, Sommer JF Jr, Stellman J. The consistency of combat exposure reporting and course of PTSD in Vietnam War veterans. J Trauma Stress 2007;20(1):3–13. [PubMed: 17345644]
95. Hoge CW, Lesikar SE, Guevara R, Lange J, Brundage JF, Engel CC, et al. Mental disorders among U.S. Military personnel in the 1990s: Association with high levels of health care utilization and early military attrition. Am J Psychiatry 2002;159:1576–1583. [PubMed: 12202280]
96. Bowman ML. Individual differences in posttraumatic distress: problems with the DSM-IV model. Can J Psychiatry 1999;44(1):21–33. [PubMed: 10076738]
97. Friedman MJ. Acknowledging the psychiatric cost of war. N Engl J Med 2004;351(1):75–77. [PubMed: 15229311]
98. Jang KL, Taylor S, Stein MB, Yamagata S. Trauma exposure and stress response: exploration of mechanisms of cause and effect. Twin Res Hum Genet 2007;10(4):564–572. [PubMed: 17708697]
99. Segman RH, Cooper-Kazaz R, Macciardi F, Goltser T, Halfon Y, Dobrobrski T, et al. Association between the dopamine transporter gene and posttraumatic stress disorder. Mol Psychiatry 2002;7(8):903–907. [PubMed: 12232785]
100. Stein MB, Jang KL, Taylor S, Vernon PA, Livesley WJ. Genetic and environmental influences on trauma exposure and posttraumatic stress disorder symptoms: a twin study. Am J Psychiatry 2002;159(10):1675–1681. [PubMed: 12359672]
101. Xian H, Chantarujikapong SI, Scherrer JF, Eisen SA, Lyons MJ, Goldberg J, et al. Genetic and environmental influences on posttraumatic stress disorder, alcohol and drug dependence in twin pairs. Drug Alcohol Depend 2000;61(1):95–102. [PubMed: 11064187]
102. Brewin CR, Andrews B, Valentine JD. Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. J Consult Clin Psychol 2000;68:748–766. [PubMed: 11068961]
103. Ehlers A, Clark DM. A cognitive model of posttraumatic stress disorder. Behav Res Ther 2000;38(4):319–345. [PubMed: 10761279]
104. Mezey G, Robbins I. Usefulness and validity of post-traumatic stress disorder as a psychiatric category. BMJ 2001;323(7312):561–563. [PubMed: 11546705]
105. Satel, SL.; Frueh, BC. Sociopolitical aspects of psychiatry. Psychiatry, Cto, editor. Baltimore, MD: Lippincott, Williams, & Wilkins; In Press
106. Yehuda R, McFarlane AC. Conflict between current knowledge about posttraumatic stress disorder and its original conceptual basis. Am J Psychiatry 1995;152(12):1705–1713. [PubMed: 8526234]
107. Burkett, BG.; Whitley, G. Stolen valor: How the Vietnam generation was robbed of its heroes and history. Dallas, Texas: Verity Press; 1998.
108. Durodie B. Risk and the social construction of ‘Gulf War Syndrome’. Philos Trans R Soc Lond B Biol Sci 2006;361(1468):689–695. [PubMed: 16687271]
109. Frueh BC, Grubaugh AL, Elhai JD, Buckley TC. US Department of Veterans Affairs disability policies for posttraumatic stress disorder: Administrative trends and implications for treatment, rehabilitation, and research. American Journal of Public Health 2007;97:2143–2145. [PubMed: 17971542]

110. Jones E, Wessely S. A paradigm shift in the conceptualization of psychological trauma in the 20th century. *J Anxiety Disord* 2007;21(2):164–175. [PubMed: 17064877]
111. Shephard B. “Pitiless psychology”: The role of prevention in British military psychiatry in the Second World War. *Hist Psychiatry* 1999;10:491–510. [PubMed: 11624331]
112. Summerfield D. The invention of post-traumatic stress disorder and the social usefulness of a psychiatric category. *Br Med J* 2001;322:95–98. [PubMed: 11154627]
113. Frueh BC, Elhai JD, Gold PB, Monnier J, Magruder KM, Keane TM, et al. Disability compensation seeking among veterans evaluated for posttraumatic stress disorder. *Psychiatr Serv* 2003;54:84–91. [PubMed: 12509672]
114. Department of Veterans Affairs Office of Inspector General. Review of State Variances in VA disability Compensation Payments (#05-00765-137). Washington, D.C.: 2005.
115. Rosen GM, Taylor S. Pseudo-PTSD. *J Anxiety Disord* 2007;21(2):201–210. [PubMed: 17084063]
116. Baggeley M. ‘Military Munchausen’s’: assessment of factitious claims of military service in psychiatric patients. *Psychiatr Bull* 1998;22(3):153–154.
117. Frueh BC, Elhai JD, Grubaugh AL, Monnier J, Kashdan TB, Sauvageot JA, et al. Documented combat exposure of US veterans seeking treatment for combat-related post-traumatic stress disorder. *Br J Psychiatry* 2005;186:467–472. discussion 473–465. [PubMed: 15928355]
118. Sparr LF, Pankratz LD. Factitious posttraumatic stress disorder. *Am J Psychiatry* 1983;140:1016–1019. [PubMed: 6869583]
119. Dean, ET. Shook over hell: Post-traumatic stress, Vietnam, and the Civil War. Cambridge, Massachusetts: Harvard University Press; 1997.
120. Freeman T, Powell M, Kimbrell TA. Measuring symptom exaggeration in veterans with chronic posttraumatic stress disorder. *Psychiatry Res* 2008;158:374–380. [PubMed: 18294699]
121. Frueh BC, Hamner MB, Cahill SP, Gold PB, Hamlin K. Apparent symptom overreporting among combat veterans evaluated for PTSD. *Clin Psychol Rev* 2000;20:853–885. [PubMed: 11057375]
122. Institute of Medicine of the National Academies. Treatment of posttraumatic stress disorder: An assessment of the evidence. Washington, DC: The National Academies Press; 2007.
123. Charney DS, Davidson JRT, Friedman M, Judge R, Keane T, McFarlane S, et al. A consensus meeting on effective research practice in PTSD. *CNS Spectrums* 1998;3:7–10.
124. Sturken, M. Tangled Memories: The Vietnam War, the AIDS Epidemic, and the Politics of Remembering. Berkeley: University of California Press; 1997.
125. Bowman J. Spielberg Stress Disorder. *Am Spect* 2006;39(1):64–65.
126. McHugh PR, Treisman G. PTSD: A problematic diagnostic category. *J Anxiety Disord* 2007;21:211–222. [PubMed: 17085011]
127. Yehuda R, Tischler L, Golier JA, Grossman R, Brand SR, Kaufman S, et al. Longitudinal assessment of cognitive performance in Holocaust survivors with and without PTSD. *Biol Psychiatry* 2006;60(7):714–721. [PubMed: 17008144]
128. Ikin JF, Sim MR, McKenzie DP, Horsley KW, Wilson EJ, Moore MR, et al. Anxiety, post-traumatic stress disorder and depression in Korean War veterans 50 years after the war. *Br J Psychiatry* 2007;190:475–483. [PubMed: 17541106]
129. Solomon Z, Mikulincer M. Trajectories of PTSD: A 20-Year Longitudinal Study. *Am J Psychiatry* 2006;163(4):659–666. [PubMed: 16585441]
130. Bramsen I, van der Ploeg HM. Use of medical and mental health care by World War II survivors in the Netherlands. *J Trauma Stress* 1999;12:243–261. [PubMed: 10378164]
131. Port CL, Engdahl B, Frazier P. A longitudinal and retrospective study of PTSD among older prisoners of war. *Am J Psychiatry* 2001;158(9):1474–1479. [PubMed: 11532734]
132. Schnurr PP, Spiro AI, Vielhauer MJ, Findler MN, Hamblen JL. Trauma in the lives of older men: Findings from the Normative Aging Study. *Journal of Clinical Geropsychology* 2002;8:175–187.
133. Schnurr PP, Friedman MJ, Foy DW, Shea MT, Hsieh FY, Lavori PW, et al. Randomized trial of trauma-focused group therapy for posttraumatic stress disorder. *Arch Gen Psychiatry* 2003;60:481–489. [PubMed: 12742869]
134. Andrews B, Brewin CR, Philpott R, Stewart L. Delayed-onset posttraumatic stress disorder: a systematic review of the evidence. *Am J Psychiatry* 2007;164(9):1319–1326. [PubMed: 17728415]

135. Frueh BC, Grubaugh AL, Yeager DE, Magruder KM. Delayed-onset posttraumatic stress disorder among patients in Veterans Affairs primary care clinics. *Br J Psychiatry* 2009;194:515–520. [PubMed: 19478290]
136. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4. Washington, DC: American Psychiatric Association; 1994.
137. Browne MW. An overview of analytic rotation in exploratory factor analysis. *Multivariate Behavioral Research* 2001;36:111–150.
138. Kashdan TB, Collins RL, Elhai JD. Social anxiety and positive outcome expectancies on risk-taking behaviors. *Cognitive Therapy and Research* 2006;30:749–761.
139. Keane TM, Wolfe J. Comorbidity in post-traumatic stress disorder: An analysis of community and clinical studies. *J Appl Soc Psychol* 1990;20:1776–1788.
140. Elhai JD, Grubaugh AL, Kashdan TB, Frueh BC. Empirical examination of a proposed refinement to DSM-IV posttraumatic stress disorder symptom criteria using the National Comorbidity Survey Replication data. *J Clin Psychiatry* 2008;69:597–602. [PubMed: 18294026]
141. Rosen GM, Lilienfeld SO. Posttraumatic stress disorder: an empirical evaluation of core assumptions. *Clin Psychol Rev* 2008;28(5):837–868. [PubMed: 18329146]
142. Goss-Gilroy Inc. Health Study of Canadian Forces Personnel Involved in the 1991 Conflict in the Persian Gulf. Ottawa: 1998.

Table 1

Prevalence Estimates of Combat-Related PTSD

Country	Study Year	Study Design	Participants (Sample Size)	Assessment Instruments	(Lifetime) Current Prevalence
US	NVRS, 1990	retrospective, assessed 11–12 yrs post-return	Vietnam veterans, males (1,200)	MISS-PTSD (SR), MMPI-PTSD (SR), SCID-III-R (CI); PTSD from DSM-III-R	(30.9%) 15.2%
US	VES, 1988	retrospective, assessed 11–12 yrs post-return, telephone	Vietnam era, males (2,490 VV; 1,972 NVV)	MMPI-PTSD (SR), DIS-3A (CI)	(15.2%) 2.2%
US	Eisen et al., 2004 [26]	retrospective, assessed 17yrs post-return	Vietnam veterans, males, twin registry (8,169)	DIS-III-R (based on DSM-III-R criteria)	(10%) 4.5%
US	Dohrenwend et al., 2006 [2]	retrospective review	Original NVRS sample data (1,200)	as per NVRS	(18.7%) 9.1%
US	Thompson et al., 2006 [28]	retrospective review	Original NVRS and VES sample data (1,200; 2,490)	as per NVRS & VES	NVRS=2.9% VES=2.6%
US	Goldberg et al., 1990 [49]	retrospective, 15yrs post deployment, MZ twin pairs, males postal/telephone	Vietnam veterans, males, twin registry (2,092)	PCL (SR)	16.8% VV, 5.0% NVV
Australia	O'Toole et al., 1996 [61]	retrospective assessed 20–25 yrs post-deployment	Vietnam veterans, males (641)	MISS-PTSD (SR) AUSCID (SR)	(20.9%) 11.6%
US	Iowa Persian Gulf Study Group, 1997 [32]	retrospective, assessed 4–5 yrs post deployment, cross sectional, telephone	1990 PGW veterans (1,896); era veterans not deployed to PGW (1,799)	PCL-M (SR)	PGWV=1.9% Era Non-GWV=0.8%
US	Barrett et al., 2002 [29]	as per Iowa Persian Study group sample, Review of data	1990 PGW veterans (1,889); era veterans not deployed to PGW (1,793)	PCL-M (SR)	1.09%
US	Kang et al., 2003 [34]	retrospective, cross sectional, postal, assessed 5–6 yrs post-deployment	1990 PGW veterans (1,441); era veterans not deployed to PGW (9,476)	PCL- (SR)	PGW =10.1%; Era Non-GW =4.2%
US	Wolfe, Proctor et al., 1999 [39]	retrospective, cross sectional, assessed 3–6 yrs post	1990 PGW veterans (204), sampled from Louisiana & Massachusetts bases; era veterans not deployed to PGW (48)	CAPS (CI) SCID (CI) MISS-PTSD (SR)	PGWV= Approx 6.3% (approx 7.35%); Era Non-GWV = not reported

Country	Study Year	Study Design	Participants (Sample Size)	Assessment Instruments	(Lifetime) Current Prevalence
US	Wolfe, Erikson et al., 1999 [37]	prospective, assessed 5 days post deployment (T1) and recontacted 2 yrs later (T2)	1990 PGW veterans, males & females, from Massachusetts base (2,942)	MISS-PTSD (SR)	T1 GWV= 3%; T2 GWV = 8%
US	Gray et al., 2002 [31]	based on Seabee Health Study 1997–1999 Retrospective, cross sectional, postal	1990 PGW veterans, Seabees compared to US era veterans not deployed to Gulf (3,831 GWV; 4,933 deployed elsewhere Era Seabee; 3,104 non-deployed Seabees)	PTSD screen from DSM-IV (SR)	GWV= 3.04%, Era= 0.61; Non- deployed = 0.71
US	Stretch et al., 1993 [35]	retrospective, cross sectional assessed 2 yrs post-deployments, postal	1990 PGW veterans compared to non- deployed era veterans residing in Hawaii/Pennsylvania (1,524 GWV; 2,512 NGV)	WRAIR algorithm	8.6% GWV, 1.6% NGV
Canada	Goss-Gilroy et al., 1998 [142]	retrospective, assessed 6yrs post- deployment, postal	1990 PGW, males & females, compared to deployed elsewhere (3,113 GWV; 3,439 NGV)	PCL-R (SR)	OR=2.69 (95% CI 1.59–4.26)
UK	Unwin et al., 1999 [73]	retrospective, cross sectional postal	1990 PGW veterans, males compared to non- deployed era veterans, compared to era veterans deployed to Bosnia peacekeeping (3,284 GWV; 1,815 Bosnian vets; 2,408 era vets)	MISS-PTSD	GWV= 13.2%, Bosnia deployed = 4.7%, Era=4.1%
Canada	Statistics Canada, 2003 [60]	retrospective, interview, random selection from total deployed forces	Canadian Regular Forces, males & females (5,000 regulars, 3,000 reservists)	CIDI (CI)	(7.2%) 2.8%; reserves (4.7) 1.2%; >3 tours (10.3%); > 3 tours 4.7%
UK	Toomey et al., 2007 [36]	regular UK personnel, postal, assessed T1 = 3–4 yrs post- deployment and T2 = 10 yrs later	1990 PGW veterans (subset of Kang et al 2000) (4,762)	CAPS (CI) PCL-C (SR)	3.7%
UK	Lee et al., 2002 [64]	consecutive serving + former serving, cross sectional	Post-1993, all branch UK regular forces veterans (3,000)	Clinician Assessment	12%
Australia	Ikin et al., 2004 [62]	retrospective, mailed plus interview, all deployed ADF personnel to GW	1990 PGW veterans (1,381)	CIDI (CI)	5.4%
UK	Jones et al., 2006 [65]	retrospective, postal, random sample	active, past-deployed, all units (2,873)	PCL (SR), GHQ-12 (SR)	2.5%

Country	Study Year	Study Design	Participants (Sample Size)	Assessment Instruments	(Lifetime) Current Prevalence
UK	Hotopf et al., 2006 [20]	retrospective, random, postal and sub-sample in-person	2003 Iraq War (4,722 OIF; 5,500 OIF era)	PCL-C (SR)	4%
UK	Brown et al., 2007 [21]	postal, random	as per Hotopf et al., 2006 (3,936 regulars, 796 reservists)	PCL-C (SR), GHQ-12 (SR)	4% = regulars, 6% = reservists
UK	Iversen et al., 2008 [22]	retrospective regular personnel, postal	OIF (4,762)	PCL-C (SR)	3.7%
US	Hoge et al., 2004 [18]	cross-sectional, longitudinal, pre- post	OIF/OEF active deployed (2,530 pre deployment; 3,671 post-deployment)	routine post- deployment (SR)	post-OIF = 15.6%(army), 17.1%(marine), post-OEF=11.2%
US	Hoge et al., 2006 [17]	cross sectional, pre-post, Army & Marines	OIF/OEF/other active deployed (303,905)	routine post- deployment (SR)	OIF=9.8%; OEF = 4.7%; Other = 2.1%
US	Seal et al., 2007 [19]	prospective, 2001–2005, all deployed personnel	OEF/OIF veterans enrolled at VA (103,788)	VA diagnosis on files (CI)	13%
US	Smith et al., 2008 [56]	prospective cohort analysis, deployed personnel	OIF/OEF/other active duty and Reserve/National Guard deployed (50,184)	PCL-C (SR)	7.6–8.7%

Note: US = United States; UK = United Kingdom; NVVRS = National Vietnam Veterans Readjustment Survey; MISS-PTSD = Mississippi Scale for posttraumatic stress disorder; SR = self report; MMPI-PTSD = Minnesota Multiphasic Personality Inventory, PTSD subscale; SCID = Structured Clinical Interview for DSM; CI = clinical interview; DSM = Diagnostic and Statistical Manual; VES = Vietnam Experience Study; VV = Vietnam veteran; NVV = Vietnam era veteran, did not serve in Vietnam; DIS = Diagnostic Interview Schedule; MZ = monozygotic twins; PCL = PTSD Checklist; AUSCID = Australian Version Structured Clinical Interview for DSM-III-R; PGW = Persian Gulf War; GW = Gulf War era veteran, did not serve in Gulf War; ADF = Australian Defence Force; CAPS = Clinician Administered PTSD Scale; PCL-C = PTSD Checklist-Civilian; CIDJ = Composite International Diagnostic Interview; GHQ-12 = General Health Questionnaire – 12; OIF = Operation Iraqi Freedom; OEF = Operation Enduring Freedom (Afghanistan); VA = Department of Veterans Affairs; WRAIR = Walter Reed Army Institute of Research.