Continuing the search for the engram: examining the mechanism of fear memories

Sheena A. Josselyn, PhD

Program in Neurosciences and Mental Health, The Hospital for Sick Children, and the Department of Physiology and Institute of Medical Sciences, University of Toronto, Toronto, Ont.

The goal of my research is to gain insight using rodent models into the fundamental molecular, cellular and systems that make up the base of memory formation. My work focuses on fear memories. Aberrant fear and/or anxiety may be at the heart of many psychiatric disorders. In this article, I review the results of my research group; these results show that particular neurons in the lateral amygdala, a brain region important for fear, are specifically involved in particular fear memories. We started by showing that the transcription factor CREB (cAMP/Ca²⁺ response element binding protein) plays a key role in the formation of fear memories. Next, we used viral vectors to overexpress CREB in a subset of lateral amygdala neurons. This not only facilitated fear memory formation but also “drove” the memory into the neurons with relatively increased CREB function. Finally, we showed that selective ablation of the neurons overexpressing CREB in the lateral amygdala selectively erased the fear memory. These findings are the first to show disruption of a specific memory by disrupting select neurons within a distributed network.

A fundamental goal of neuroscience is to understand how memories are encoded and stored in the brain. Ensembles (or groups) of neurons are thought to serve as the physical representation of memory (the memory trace or “engram,” a term first coined by Richard Semon in 1921¹). However, identifying the precise neurons that constitute a memory trace has been challenging neuroscientists since Karl Lashley began his famous search for the elusive engram in the 1920s. In his studies, Lashley trained rats in various tasks (such as traversing a maze to find a food reward) and lesioned various portions of different cortical regions either before or after training. After 30 years of work, he summarized his findings in a seminal paper entitled “In search of the engram.”²

Lashley observed that cortical lesions disrupted performance in the maze and that the degree of disruption was roughly proportional to the amount of cortex removed but not to the location of the lesion. From this, he concluded that all cortical areas can substitute for each other as far as learning is concerned (the principle of equipotentiality) and that the cortex tends to act as a whole in that the amount, rather than the location, of cortical tissue removed correlated with performance in the maze (the principle of mass action). These

findings led Lashley to conclude that memories are not localized, but rather distributed, in the cortex. In his words,

“...this series of experiments has yielded a good bit of information about what and where the memory trace is not. It has discovered nothing directly of the real nature of the memory trace. I sometimes feel, in reviewing the evidence of the localization of the memory trace, that the necessary conclusion is that learning is just not possible. It is difficult to conceive of a mechanism that can satisfy the conditions set for it. Nevertheless, in spite of such evidence against it, learning sometimes does occur.”²

Lashley’s failure to find the engram may have been because of his choice of memory task (learning a maze is a complex task, and this sort of memory likely relies on many brain regions) and brain region examined (regions other than the cortex may be involved in memory).

Since Lashley’s studies, though, there has been progress in “finding the engram.” Both cellular imaging and electrophysiological studies³⁻⁷ have detected neurons whose activity is correlated with memory encoding or expression or both, suggesting that these active neurons make up the memory trace. For example, John Guzowski’s laboratory used a technique

Correspondence to: Dr. S.A. Josselyn, The Hospital for Sick Children, 555 University Ave., Toronto ON M5G 1X8; fax 406 813-7717; sheena.josselyn@sickkids.ca

J Psychiatry Neurosci 2010;35(4):221-8.

Submitted Jan. 22, 2010; Revised Apr. 7, 2010; Accepted Apr. 12, 2010.

DOI: 10.1503/jpn.100015

they developed, referred to as catFISH (compartmental analysis of temporal activity by fluorescence in situ hybridization),⁸ and found that some of the same neurons in the dorsal hippocampus of rats were active when the rats were re-exposed to a unique environment. More recently, Mark Mayford's laboratory used a sophisticated transgenic mouse approach that allowed them to tag active neurons.⁵ They showed that some of the same amygdala neurons that were active during fear learning were also active during the retrieval of that fear memory. Furthermore, the number of reactivated neurons in the amygdala was correlated with the strength of the retrieved fear memory.

These findings suggest that specific neurons are involved in a memory. However, these correlative studies do not address whether these active neurons are essential components of the memory trace. A direct test of this hypothesis would require specifically disrupting only these activated neurons while leaving their neighbours intact and then determining whether subsequent memory expression is blocked. Establishing such a causal role for particular neurons in a memory has been difficult because the neuronal ensembles that make up this memory trace are thought to be sparsely distributed,⁹ and the ability of current pharmacological, genetic and lesioning techniques to target specific subsets of neurons within a brain region is limited.

In my laboratory's "search for the engram," we used auditory fear conditioning as our memory task. In this Pavlovian paradigm, a neutral tone is paired with a mildly aversive footshock. One tone-shock pairing is sufficient to produce a robust and long-lasting fear memory that can be quantified by measuring the percentage of time mice spend freezing (an adaptive defensive response) when the tone is subsequently replayed. Automated measures have been established to easily and reliably quantify freezing.¹⁰ We chose this task because genetic, biochemical, electrophysiological and behavioural studies from many laboratories, including those of Mike Davis, Michael Fanselow, Joe LeDoux and Steven Maren, established that lateral amygdala is required for auditory-conditioned fear memories^{11–15} (but see¹⁶). Now we knew where to look in the brain for a fear memory trace. The next step was to target lateral amygdala neurons that are active following fear training or testing.

To target neurons whose activity is correlated with memory, Jin-Hee Han, Adelaide Yiu and Hwai-Lin (Liz) Hsiang in my laboratory, together with our collaborators Steven Kushner, Bruno Bontempi and Paul Frankland, took advantage of our recent findings that increasing the level of the transcription factor CREB (cAMP/Ca²⁺ responsive element binding protein) in a small portion of lateral amygdala neurons enhances auditory fear memory under certain training conditions.¹⁷ Furthermore, we found that neurons with relatively increased CREB levels are more likely than their neighbours to be active following fear memory training or testing. This suggested to us that these neurons with high CREB levels outcompete their neighbours for inclusion in the fear memory trace.

The idea that CREB is involved in memory is not new. Indeed, studies by Tim Tully's and Eric Kandel's laboratories showed that CREB is necessary for memory formation in

invertebrates.^{18–22} Parallel studies in rodents have also showed that CREB is important for memory formation^{23–34} (but see³⁵).

I first began examining the effects of increasing CREB function on memory when I was a postdoctoral fellow at Yale in Mike Davis's laboratory. Eric Nestler's laboratory was located across the hall at the Connecticut Center for Mental Health, and Bill Carlezon, a postdoctoral fellow in his laboratory at the time, was manipulating CREB function using replication-defective herpes simplex viral (HSV) vectors to determine the effects of CREB on cocaine-induced drug seeking.³⁶ Although there were many ways to decrease CREB function, there were relatively few ways to transiently enhance its function in rodents. The HSV system was an ideal tool to study the effects of enhancing CREB on memory. This system, which allows expression of a transgene in a particular brain region at a particular time, was largely pioneered by my colleague, Rachael Neve, now at the Massachusetts Institute of Technology.

Herpes simplex viral vectors offer many advantages over other viral vector systems.³⁷ First, HSV infects adult (nondividing) neurons (rather than glia) with relatively high efficiency.^{38,39} Second, the DNA from HSV remains episomal, thus avoiding potentially confounding effects of integration into the host DNA. Third, these vectors offer a large capacity for the insertion of foreign DNA. In our studies, we drove transgene expression with the HSV immediate-early gene *IE4/5* such that transgene expression peaked 2–5 days following infusion and declined within 7 days.^{40,41} In this way, we could examine the effects of acutely increasing CREB function on fear memory in the lateral amygdala. In collaboration with Bill Carlezon and Eric Nestler, Mike Davis and I showed that increasing CREB levels in the lateral amygdala enhanced memory for fear-potentiated startle, another fear memory task,¹² in rats that had been given weak training (consisting of several training trials massed together without intervening rest periods, which has been shown to produce weak memory).⁴² What intrigued us about our finding was that we observed robust memory enhancement despite the fact that we increased CREB levels in a relatively small percentage of lateral amygdala neurons (our HSV vectors infected roughly 15% of lateral amygdala neurons). This disconnect between the small number of neurons infected but large behavioural results continued to motivate my experiments.

Later, when I was a postdoctoral fellow in Alcino Silva's laboratory, Steven Kushner, Alcino and I had long conversations about how we might make sense of these findings. We definitely believed the data. Mike Davis taught me to always follow the data (to borrow from Shakira, data, like hips, don't lie). Plus, our basic effect was replicated by 2 additional groups using the same HSV vectors.^{43,44} We reasoned that one way to explain the data was that the low number of neurons with increased levels of CREB were somehow outcompeting their neighbours for inclusion in the memory trace. When I began my laboratory in Toronto, Jin-Hee Han, a talented postdoctoral fellow in my laboratory, continued these studies.

First, we revamped our viral vectors to allow infected neurons to be easily visualized by fusing the transgenes with green fluorescent protein (GFP). Tagging the N-terminus of

CREB with GFP does not interfere with the functional activity of CREB.^{37,40,45,46} In our first studies, we used 3 main vectors: HSV-CREB expressing wild-type CREB (CREB^{WT}); HSV-mCREB (the mutant CREB vector) expressing a mutated version of CREB that cannot be phosphorylated at the key Ser133 residue (CREB^{S133A}); and HSV-GFP (control, Cntrl vector) expressing GFP or LacZ.

Our overall aim was to determine whether neurons infected with the CREB^{WT} vector were preferentially recruited to the lateral amygdala fear memory trace. We reasoned that if competition between neurons for inclusion in the memory trace occurred during learning, the easiest way to see this would be to “stack” the experimental deck by maximizing the difference between the neurons infected with the CREB^{WT} vector and their noninfected neighbours. Thus, we first infused our CREB^{WT} vector into the lateral amygdala of mice with a targeted disruption of the 2 main isoforms of CREB, CREB^{αδ-/-} mice.⁴⁷ The CRE-DNA binding is virtually abolished (by > 90%),^{48,49} and the levels of CREB protein are dramatically reduced (roughly 85%–90% reduction compared with controls) in the brains of CREB^{αδ-/-} mice.^{48,50} These CREB-deficient mice have previously been shown to be impaired in auditory fear conditioning.^{23,28} Importantly, we replicated this memory deficit and showed that infusing our Cntrl vector into the lateral amygdala did not change this. However, infusing the CREB^{WT} vector into the lateral amygdala of these mice completely rescued the auditory fear deficit; the CREB-deficient mice now froze at the same levels as their wild-type littermate control mice. Furthermore, under weak training conditions, we found that increasing CREB function in a similar small fraction of lateral amygdala neurons enhanced memory in wild-type mice (similar to my initial findings using fear-potentiated startle in rats⁴²). Together these findings suggested that these neurons with increased CREB function were preferentially recruited to the trace supporting the fear memory.

We were inspired to explore this possibility further by John Guzowski's elegant studies using the catFISH technique. Specifically, Jin-Hee Han, Adelaide Yiu and Christina (Christy) Cole in my laboratory, together with our collaborators, Steven Kushner, John Guzowski and Alcino Silva, attempted to visualize the fear memory trace by taking advantage of the unique transcriptional time course of the activity-dependent gene *Arc* (activity-regulated cytoskeleton-associated protein; Arg3.1).^{6,8} Under basal conditions, neurons contain very low levels of *Arc* RNA. However, neuronal activation (the type of activation that is associated with learning or long-term potentiation) produces a rapid but transient burst in *Arc* RNA synthesis (within about 3–5 min of activation).⁸ This RNA is then delivered to dendrites within roughly 20 minutes of activation. Therefore, the localization of *Arc* RNA (either in the nucleus, cytoplasm or both) can serve as an activity marker for that particular neuron, with *Arc* RNA localized to the nucleus being a molecular marker of a neuron that was active 5 minutes ago. We used this technique to identify neurons that were activated by fear memory training or testing and asked whether these neurons were also the ones we infected with the CREB^{WT} vector.

We observed that neurons infected with the CREB^{WT} vector were 3 times more likely than their noninfected neighbours to be *Arc* positive in wild-type mice and 10 times more likely in CREB-deficient mice. In contrast, neurons infected with the dominant-negative CREB vector were 12 times less likely than their neighbours to be *Arc* positive. This was consistent with our behavioural data showing that disrupting CREB function (using the CREB^{S133A} vector) in roughly 20% of lateral amygdala neurons had no effect on memory in wild-type mice, perhaps because the remaining neurons (over 80%) with normal levels of CREB were sufficient for normal memory.¹⁷

Together with many control studies, these findings suggest that neurons with relatively higher CREB function are preferentially recruited to the memory trace. Would this mean that increasing CREB levels in all lateral amygdala neurons would enhance fear memory? We hypothesized that it would not, because we think that competition between eligible neurons is critical in memory enhancement. We reasoned that raising CREB levels in all neurons would not produce a stronger memory because all of the neurons would again be equal (and the signal-to-noise ratio would not be enhanced). As Gore Vidal put it, it is not enough to succeed; others must fail. The same may apply to neuronal competition underlying memory formation.

Our findings suggested that neurons with higher levels of CREB “win” the competition between neurons for inclusion in a memory trace. My laboratory then attempted to use this property to determine if selectively ablating just these neurons after training disrupts the expression of an established fear memory. If so, this would indicate that these neurons are critical components of the elusive memory trace. We used several techniques to do this but had no luck. It turns out that these neurons, like Steven Segal, are hard to kill, perhaps because CREB is a survival factor.⁵¹ Then, Steven Kushner, an avid reader of scientific journals, came across a paper in *Nature Methods* by Ari Waisman and colleagues⁵² describing a novel transgenic mouse used for cell lineage ablation studies based on diphtheria toxin (DT). We decided to try to kill neurons overexpressing CREB using this approach.

First, a bit of background on DT. *Corynebacterium diphtheriae* produces DT, a potent toxin that, once it binds to its receptor (diphtheria toxin receptor; DTR) and is internalized into a mammalian cell, efficiently blocks protein synthesis to cause rapid apoptotic cell death.^{53–55} For cell death to occur using the DT-based system, both key components (DT and DTR) are required. Interestingly, mice do not normally express a functional DTR.^{56–58} So, Waisman and colleagues⁵² engineered a mouse that expresses a simian DTR transgene (driven by the ubiquitous Rosa promoter; Fig. 1). However, expression of DTR is dependent on the cre recombinase-mediated removal of a transcriptional STOP cassette (upstream of the DTR transgene they placed a floxed STOP cassette that silences DTR expression until the STOP cassette is removed by cre recombinase-mediated recombination). Systemic injection of DT anytime thereafter induced apoptosis, but only in the cells that have undergone cre-mediated recombination and express DTR. A single internalized catalytically active fragment of DT (DT-A)

is sufficient to kill a cell, indicating the sensitivity of DT-induced cell ablation.⁵⁹ Importantly, neither high doses of DT in wild-type mice nor the expression of DTR alone (without DT) induces apoptosis, indicating the specificity of the system.^{60,61} Because DT readily crosses the blood-brain barrier,⁶² these inducible DTR (iDTR) mice have been used to induce cell death in the brain.^{52,60}

Many strategies can be used to delete defined cells, including the expression of cytotoxic proteins,⁶³ antibodies⁶⁴ or chemicals.⁶⁵ However, these systems, and indeed earlier versions of the DT system, were not suited for our studies because they lacked temporal control over cell death and/or are prone to “leakiness” (in that cell death is not temporally or spatially limited to the targeted cell population).⁶³ The iDTR mice were designed to address these drawbacks. Because DT induces cell death by apoptosis, it is thought to produce less inflammation than necrosis-induced cell death.⁶⁶ In this way, the iDTR mice were designed to minimize the “bystander effect,” in which the death of a targeted cell directly or indirectly affects neighbouring neurons.

Jin-Hee Han, together with other members of my laboratory and our collaborators Steven Kushner and Paul Frankland, took advantage of the iDTR transgenic mice to selectively induce cell death in the neurons that we hypothesized were preferentially involved in the memory trace (neurons infected with the CREB^{WT} vector). To this end, we re-engineered our viral vectors to include cDNA for cre recombinase. Our first vector expressed both CREB and cre recombinase (CREB-cre vector). When injected into the lateral amygdala of the iDTR mouse, we expected that a small portion of neurons would have increased levels of CREB and that a subsequent injection of DT would ablate just these neurons. Recombination

and DTR expression can only occur in neurons expressing cre recombinase, allowing us to persistently tag infected neurons for subsequent ablation. This tagging is critical because of the relatively brief time course of transgene expression using HSV.⁴⁰ Because this technique produces a small lesion of the lateral amygdala that may affect memory on its own, we needed a control vector. So we added cre recombinase to our Cntrl vector. Our thinking was that this Cntrl-cre vector would ablate a similar number of lateral amygdala neurons but that these neurons would be randomly located in the lateral amygdala and not involved in the memory trace (because they did not overexpress CREB). As a first step in this study, Jin-Hee confirmed that this iDTR/vector system selectively induced cell death by examining 2 markers of apoptosis (activated caspase 3 and terminal deoxynucleotidyl transferase dUTP nick end labeling [TUNEL]). We quantified the number of cells with activated caspase 3 and those that were positive for TUNEL and found that significant apoptotic cell death was only observed in the experimental groups (iDTR mice infused with either the CREB-cre or Cntrl-cre vectors). Importantly, we found that the CREB-cre and Cntrl-cre vectors produced a similar level of cell death. We were excited because we thought that this system, which allowed us to temporally ablate tagged neurons, would bring us one step closer to finding the engram.

To examine the effects of post-training ablation of neurons overexpressing CREB, we performed a series of behavioural tests. First, we found that increasing CREB levels in a population of lateral amygdala neurons (by microinjecting the CREB-cre vector) enhanced memory following weak training in iDTR mice. This finding replicated our earlier finding using the CREB^{WT} vector (without cre recombinase) in wild-type mice.¹⁷ This memory enhancement, however, was completely reversed after systemic administration of DT (to delete just those neurons infected with the CREB-cre vector; Fig. 2). Importantly, the reversal of this memory enhancement was not observed in the control groups that lacked a key component of the DT-killing system (either cre recombinase or DT), consistent with the lack of cell death in these control groups. This was important because it showed that the memory reversal was not due to fear memory extinction caused by the second memory test.

Next, we showed that deleting neurons infected with the CREB-cre vector (but not a similar portion of random neurons infected with the Cntrl-cre vector) also impaired the expression of a strong memory. Therefore, even though increasing CREB function in a portion of lateral amygdala neurons did not further enhance memory in mice that were trained using a strong protocol, these neurons are nevertheless important in the fear memory. In both of these experiments, however, we administered DT after a fear memory test. This memory test, however, may reactivate the fear memory and trigger a second wave of consolidation (referred to as reconsolidation).⁶⁷ Karim Nader, Joe LeDoux and several other groups have shown that, similar to initial consolidation, reconsolidation requires protein synthesis.⁶⁷ Because DT disrupts protein synthesis, we wanted to examine whether the memory disruption that we observed in the previous experiments was due to a

Fig. 1: Schematic diagram depicting the strategy for selectively ablating neurons using inducible diphtheria toxin receptor (iDTR) transgenic mice. The mice express a simian DTR transgene. Expression of the DTR transgene is suppressed by a STOP cassette, which is floxed and removed when a neuron is infected with herpes simplex virus expressing cre recombinase (cAMP/Ca²⁺ response element binding protein [CREB]-cre vector or control-cre vector). Only neurons that have undergone cre-induced recombination will constitutively express DTR on the cell surface. At any point thereafter, systemic injection of diphtheria toxin (DT) induces apoptosis only in cells expressing DTR. In this way, only neurons infected with CREB-cre vector or Cntrl-cre vector will be ablated.

disruption in reconsolidation. Therefore, we examined additional groups of mice but did not reactivate the memory (we did not give the mice a memory test) before DT administration. Instead, we administered DT at the same time but in the homecage. Again, we observed that mice microinjected with the CREB-cre vector and administered DT showed impaired memory whereas similarly treated Cntrl-cre mice did not.

If neurons overexpressing CREB during training (neurons with the CREB-cre vector) are critically involved in the memory trace, deleting this subpopulation should permanently block memory expression. To examine the persistence of memory loss, we trained mice using a strong protocol, administered DT in the homecage and assessed memory 2, 5 and 12 days later. The memory loss observed in the CREB-

Fig. 2: Neurons with relatively increased CREB (cAMP/Ca²⁺ response element binding protein) function are essential for memory recall. (A) Schematic diagram of an experiment designed to test whether neurons with increased CREB function at the time of training are required for subsequent memory recall. (i) A subset of lateral amygdale (LA) neurons are infected with a CREB-cre vector (labelled green via green fluorescent protein [GFP]), which increases CREB in these neurons and induces them to undergo cre-induced recombination to express diphtheria toxin receptor (DTR). (ii) Mice are trained. A subset of LA neurons have high CREB levels, but because diphtheria toxin (DT) has not been administered, there is no cell death. (iii) Mice are tested. Neurons involved in the memory trace are depicted in red. Neurons with increased CREB out-compete their neighbours for inclusion in the fear memory trace (labelled yellow [green+red]). (iv) Mice receive systemic injections of DT to induce cell death only in the cells that express DTR (have undergone cre-induced recombination and, therefore, must have been infected with the CREB-cre vector). Mice are tested again. Neurons that previously had high CREB (infected) are now ablated. (B) Results from this experiment indicate that selectively ablating neurons with increased CREB at the time of training impairs subsequent memory recall. The CREB-cre group shows that increasing CREB function in a portion of neurons at the time of training enhances memory (Test 1) and subsequent ablation of these particular neurons completely reverses this enhancement (Test 2). The Cntrl-cre group shows that GFP does not enhance memory and that selectively ablating these cells does not change this. The off-target group shows that increasing CREB function in neurons outside the LA does not enhance memory. The CREB alone (no cre) group shows that increasing CREB in the LA enhances memory (Test 1) and that DT injection (with no recombination) does not reverse this enhancement Test 2). The mice that received phosphate-buffered saline but no DT (PBS, no DT) show that CREB enhances memory in inducible DTR mice, and a systemic injection of PBS (rather than DT) does not change this on the second test day.

cre vector mice was long lasting; mice showed low freezing during the tone over these repeated tests. In contrast, memory remained robust over repeated tests in similarly treated Cntrl-cre mice. Therefore, we found no evidence of memory recovery in mice in which neurons overexpressing CREB were deleted, indicating that the memory was not just transiently suppressed.

To determine whether the memory loss could be due to a nonspecific impairment of lateral amygdala function (the mice did have a small lesion of the lateral amygdala), we retrained these mice. Following retraining, both groups of mice (CREB-cre and Cntrl-cre mice) showed equally high levels of freezing during the tone. That is, the post-training ablation of neurons that were overexpressing CREB (and cre recombinase) at the time of training produced a memory deficit, but these mice were capable of learning a tone-shock association (and freezing). Similarly, we also found that ablating neurons with the CREB-cre vector before fear conditioning did not impair subsequent memory formation. In this experiment, we infused the CREB-cre vector and systemically administered DT (to kill the neurons overexpressing CREB) before strong training and tested fear memory daily for 3 days. We found that even though a subpopulation of neurons was ablated before training, the mice acquired memory normally. Furthermore, this memory was neither more prone to extinction nor more fragile than a memory acquired without this killing (iDTR mice microinjected with the CREB-cre vector but systemically administered phosphate-buffered saline instead of DT). Together, these findings indicate that ablation of neurons that were overexpressing CREB at the time of memory-encoding blocks the memory of that particular learning event while leaving subsequent learning intact. Much like the data from wild-type mice infused with the mCREB vector (which showed normal memory), the high portion of remaining (noninfected) neurons seemed sufficient to encode a new memory. Finally, we examined the effects of ablating neurons that were infected with the CREB-cre vector after training. In this experiment, we reasoned that there would be no memory disruption because the CREB-infected neurons would not be part of the memory trace (because training had already taken place). This is exactly what we observed. Therefore, the memory loss induced by ablating neurons overexpressing CREB is robust, persistent and specific.

Our results show that the neurons with increased CREB levels at the time of fear learning are critical to the stability of that memory because selectively ablating just these neurons after training blocks this fear memory. This indicates that these neurons themselves are essential for later memory expression; they are not simply creating a local environment that promotes memory formation (such as releasing trophic factors). Fear learning may generate a broad memory trace that encompasses more lateral amygdala neurons than affected by our treatment or multiple memory traces throughout the brain. However, deleting just the neurons overexpressing CREB at the time of training produces amnesia, which suggests that these neurons play an essential role in what is likely a broader fear neuronal network.

Our results established a causal link between the activity of

a defined subpopulation of neurons and expression of a fear memory, thereby identifying a key component of the memory trace. They also indicate that neurons with relatively high CREB levels are selectively recruited to a fear memory trace. But what is the mechanism underlying this preferential selection? Why are the neurons that overexpress CREB so special? One possibility is that increasing CREB alters the intrinsic excitability of a neuron. Intrinsic excitability (the propensity of a neuron to fire action potentials in response to an input) is determined by the distribution and properties of ion channels (e.g., Na^+ , K^+ , Ca^{2+}) in the plasma membrane. Indeed, increasing CREB function stimulates transcription of a voltage-dependent Na^+ subunit (1 β subunit) and inhibits transcription of a voltage-dependent K^+ channel subunit (Kv1.4).⁶⁸ Hebb proposed that links between 2 cells are strengthened if both cells are active simultaneously.⁶⁹ That is, coincident firing of the presynaptic neuron and depolarization of the postsynaptic neuron is necessary for Hebbian plasticity. A postsynaptic neuron that is more excitable than its neighbour might therefore be more likely to be depolarized and subsequently “fire together” and “wire together” with the presynaptic neuron. In this way, neurons with increased intrinsic excitability may be “primed” for learning and more likely to outcompete their neighbours for inclusion in a fear memory trace. Unlike synaptic plasticity, which involves changes at the level of the individual synapse, intrinsic excitability or plasticity involves changes at the level of the entire neuron,⁷⁰ thus making it an attractive mechanism for the effects of CREB on neuronal competition during memory formation.

To investigate this, my laboratory collaborated with Mike Salter's laboratory at The Hospital for Sick Children. We examined the electrophysiological characteristics of principal neurons in the lateral amygdala infected with CREB^{WT} and Cntrl vectors in acute slices from wild-type mice. We were inspired by work from Eric Nestler's and Robert Malenka's laboratories⁷¹ showing that medium aspiny neurons in the nucleus accumbens infected with a similar CREB vector showed increased excitability (increased evoked action potential firing, decreased rheobase), whereas disrupting CREB (using the dominant-negative CREB^{S133A} vector) decreased intrinsic excitability without affecting passive membrane properties.

We found a similar effect in lateral amygdala neurons. Our initial results showed that neurons with the CREB^{WT} vector were more excitable than both their noninfected neighbours and those infected with the Cntrl vector.⁷² Specifically, neurons with the CREB^{WT} vector fired more action potentials in response to the same input stimulus than neurons with the Cntrl vector or the noninfected neighbouring neurons. Importantly, we observed no change in passive membrane properties (average resting membrane potential, input conductance) or action potential waveforms between the groups. This increase in neuronal excitability is in agreement with findings using viral vectors^{73,74} and transgenic mice^{75–78} to increase CREB levels. Recently, the effects of increasing CREB function on the electrophysiological properties of lateral amygdala neurons has been replicated and extended by Alcino Silva's group.⁷⁹

Francis Crick wrote that a crucial step in understanding the

mechanisms underlying memory is to interfere with defined neuronal populations in intact neural circuits.⁸⁰ He emphasized that it was necessary that such manipulations target specific neurons in time and space. With the advent of modern genetic, molecular and imaging tools, this is becoming possible.⁸¹ Our studies have shown that interfering (retrograde deletion) with a defined neuronal population (lateral amygdala neurons overexpressing CREB) in intact circuits disrupts memory expression. This may be the first example of the disruption of a specific memory within a distributed network. We believe that this takes us one step closer to finding and even manipulating the elusive engram.

Acknowledgments: In this review, I outlined my scientific journey. Of course, I did not take this journey alone, as all the work described here has been a team effort. I mentioned many of the important people along the way. I would also like to specifically acknowledge certain individuals whose contributions were vital. First, my postdoctoral mentors, Mike Davis and Alcino Silva. Next, the members of my laboratories (both past and present), especially Jin-Hee Han, Adelaide Yiu, Christy Cole and Hwai-Lin (Liz) Hsiang. Finally, I thank my collaborators, Eric Nestler, Bill Carlezon, Rachael Neve and John Guzowski, and especially Steven Kushner and Paul Frankland. I would like to thank the people in my lab and my many collaborators for making this work possible. I would also like to thank Paul W. Frankland for suggestions on this manuscript. This work was supported by Canadian Institutes of Health Research (MOP74650), EJLB Foundation and grants from the National Sciences and Engineering Research Council of Canada.

Competing interests: None declared.

References

- Schacter DL. *Forgotten, neglected pioneers: Richard Semon and the story of memory*. Philadelphia (PA): Psychology Press; 2001.
- Lashley K. In search of the engram. *Symp Soc Exp Biol* 1950;4:454-82.
- Quirk GJ, Repa C, LeDoux JE. Fear conditioning enhances short-latency auditory responses of lateral amygdala neurons: parallel recordings in the freely behaving rat. *Neuron* 1995;15:1029-39.
- Berger TW, Rinaldi PC, Weisz DJ, et al. Single-unit analysis of different hippocampal cell types during classical conditioning of rabbit nictitating membrane response. *J Neurophysiol* 1983;50:1197-219.
- Reijmers LG, Perkins BL, Matsuo N, et al. Localization of a stable neural correlate of associative memory. *Science* 2007;317:1230-3.
- Guzowski JF, Timlin JA, Roysam B, et al. Mapping behaviorally relevant neural circuits with immediate-early gene expression. *Curr Opin Neurobiol* 2005;15:599-606.
- Hall J, Thomas KL, Everitt BJ. Cellular imaging of zif268 expression in the hippocampus and amygdala during contextual and cued fear memory retrieval: selective activation of hippocampal CA1 neurons during the recall of contextual memories. *J Neurosci* 2001;21:2186-93.
- Guzowski JF, McNaughton BL, Barnes CA, et al. Environment-specific expression of the immediate-early gene *Arc* in hippocampal neuronal ensembles. *Nat Neurosci* 1999;2:1120-4.
- Perez-Orive J, Mazor O, Turner GC, et al. Oscillations and sparsening of odor representations in the mushroom body. *Science* 2002;297:359-65.
- Anagnostaras SG, Josselyn SA, Frankland PW, et al. Computer-assisted behavioral assessment of Pavlovian fear conditioning in mice. *Learn Mem* 2000;7:58-72.
- LeDoux JE. Emotion circuits in the brain. *Annu Rev Neurosci* 2000;23:155-84.
- Davis M. The role of the amygdala in fear and anxiety. *Annu Rev Neurosci* 1992;15:353-75.
- Fanselow MS, Gale GD. The amygdala, fear, and memory. *Ann N Y Acad Sci* 2003;985:125-34.
- Maren S, Quirk GJ. Neuronal signalling of fear memory. *Nat Rev Neurosci* 2004;5:844-52.
- Rumpel S, LeDoux J, Zador A, et al. Postsynaptic receptor trafficking underlying a form of associative learning. *Science* 2005;308:83-8.
- McGaugh JL. Memory consolidation and the amygdala: a systems perspective. *Trends Neurosci* 2002;25:456.
- Han JH, Kushner SA, Yiu AP, et al. Neuronal competition and selection during memory formation. *Science* 2007;316:457-60.
- Yin JC, Del Vecchio M, Zhou H, et al. CREB as a memory modulator: induced expression of a dCREB2 activator isoform enhances long-term memory in *Drosophila*. *Cell* 1995;81:107-15.
- Yin JC, Wallach JS, Del Vecchio M, et al. Induction of a dominant negative CREB transgene specifically blocks long-term memory in *Drosophila*. *Cell* 1994;79:49-58.
- Kaang BK, Kandel ER, Grant SG. Activation of cAMP-responsive genes by stimuli that produce long-term facilitation in Aplysia sensory neurons. *Neuron* 1993;10:427-35.
- Dash PK, Hochner B, Kandel ER. Injection of the cAMP-responsive element into the nucleus of Aplysia sensory neurons blocks long-term facilitation. *Nature* 1990;345:718-21.
- Bartsch D, Casadio A, Karl KA, et al. CREB1 encodes a nuclear activator, a repressor, and a cytoplasmic modulator that form a regulatory unit critical for long-term facilitation. *Cell* 1998;95:211-23.
- Bourtchuladze R, Frenguelli B, Blendy J, et al. Deficient long-term memory in mice with a targeted mutation of the cAMP-responsive element-binding protein. *Cell* 1994;79:59-68.
- Kogan JH, Frankland PW, Blendy JA, et al. Spaced training induces normal long-term memory in CREB mutant mice. *Curr Biol* 1997;7:1-11.
- Kida S, Josselyn SA, Peña de Ortiz S, et al. CREB required for the stability of new and reactivated fear memories. *Nat Neurosci* 2002;5:348-55.
- Pittenger C, Huang YY, Paletzki RF, et al. Reversible inhibition of CREB/ATF transcription factors in region CA1 of the dorsal hippocampus disrupts hippocampus-dependent spatial memory. *Neuron* 2002;34:447-62.
- Guzowski JF, McGaugh JL. Antisense oligodeoxynucleotide-mediated disruption of hippocampal cAMP response element binding protein levels impairs consolidation of memory for water maze training. *Proc Natl Acad Sci U S A* 1997;94:2693-8.
- Frankland PW, Josselyn SA, Anagnostaras SG, et al. Consolidation of CS and US representations in associative fear conditioning. *Hippocampus*. 2004;14:557-69.
- Graves L, Dalvi A, Lucki J, et al. Behavioral analysis of CREB alphas mutation on a B6/129 F1 hybrid background. *Hippocampus* 2002;12:18-26.
- Lamprecht R, Hazvi S, Dudai Y. cAMP response element-binding protein in the amygdala is required for long- but not short-term conditioned taste aversion memory. *J Neurosci* 1997;17:8443-50.
- Impey S, Smith DM, Obrietan K, et al. Stimulation of cAMP response element (CRE)-mediated transcription during contextual learning. *Nat Neurosci* 1998;1:595-601.
- Josselyn SA, Kida S, Silva AJ. Inducible repression of CREB function disrupts amygdala-dependent memory. *Neurobiol Learn Mem* 2004;82:159-63.
- Josselyn SA, Nguyen PV. CREB, synapses and memory disorders: past progress and future challenges. *Curr Drug Targets CNS Neurol Disord* 2005;4:481-97.
- Restivo L, Tafi E, Ammassari-Teule M, et al. Viral-mediated expression of a constitutively active form of CREB in hippocampal neurons increases memory. *Hippocampus* 2009;19:228-34.
- Balschun D, Wolfer DP, Gass P, et al. Does cAMP response element-binding protein have a pivotal role in hippocampal synaptic plasticity and hippocampus-dependent memory? *J Neurosci* 2003;23:6304-14.
- Carlezon WA Jr, Thome J, Olson VG, et al. Regulation of cocaine reward by CREB. *Science* 1998;282:2272-5.
- Neve RL, Neve KA, Nestler EJ, et al. Use of herpes virus amplicon vectors to study brain disorders. *Biotechniques* 2005;39:381-91.
- Fink DJ, DeLuca NA, Goins WF, et al. Gene transfer to neurons using herpes simplex virus-based vectors. *Annu Rev Neurosci* 1996;19:265-87.
- Carlezon WA Jr, Nestler EJ, Neve RL. Herpes simplex virus-mediated gene transfer as a tool for neuropsychiatric research. *Crit Rev Neurobiol* 2000;14:47-67.
- Barrot M, Olivier JD, Perrotti LI, et al. CREB activity in the nucleus

- accumbens shell controls gating of behavioral responses to emotional stimuli. *Proc Natl Acad Sci U S A* 2002;99:11435-40.
41. Carlezon WA Jr, Neve RL. Viral-mediated gene transfer to study the behavioral correlates of CREB function in the nucleus accumbens of rats. *Methods Mol Med* 2003;79:331-50.
 42. Josselyn SA, Shi C, Carlezon WA Jr, et al. Long-term memory is facilitated by cAMP response element-binding protein overexpression in the amygdala. *J Neurosci* 2001;21:2404-12.
 43. Jasnow AM, Shi C, Israel JE, et al. Memory of social defeat is facilitated by cAMP response element-binding protein overexpression in the amygdala. *Behav Neurosci* 2005;119:1125-30.
 44. Wallace TL, Stellitano KE, Neve RL, et al. Effects of cyclic adenosine monophosphate response element binding protein overexpression in the basolateral amygdala on behavioral models of depression and anxiety. *Biol Psychiatry* 2004;56:151-60.
 45. Chao JR, Ni YG, Bolaños CA, et al. Characterization of the mouse adenylyl cyclase type VIII gene promoter: regulation by cAMP and CREB. *Eur J Neurosci* 2002;16:1284-94.
 46. Olson VG, Zabetian CP, Bolaños CA, et al. Regulation of drug reward by cAMP response element-binding protein: evidence for two functionally distinct subregions of the ventral tegmental area. *J Neurosci* 2005;25:5553-62.
 47. Hummler E, Cole TJ, Blendy JA, et al. Targeted mutation of the CREB gene: compensation within the CREB/ATF family of transcription factors. *Proc Natl Acad Sci U S A* 1994;91:5647-51.
 48. Walters CL, Blendy JA. Different requirements for cAMP response element binding protein in positive and negative reinforcing properties of drugs of abuse. *J Neurosci* 2001;21:9438-44.
 49. Pandey SC, Mittal N, Silva AJ. Blockade of cyclic AMP-responsive element DNA binding in the brain of CREB delta/alpha mutant mice. *Neuroreport* 2000;11:2577-80.
 50. Walters CL, Kuo YC, Blendy JA. Differential distribution of CREB in the mesolimbic dopamine reward pathway. *J Neurochem* 2003;87:1237-44.
 51. Finkbeiner S. CREB couples neurotrophin signals to survival messages. *Neuron* 2000;25:11-4.
 52. Buch T, Heppner FL, Tertilt C, et al. A Cre-inducible diphtheria toxin receptor mediates cell lineage ablation after toxin administration. *Nat Methods* 2005;2:419-26.
 53. Dorland RB, Middlebrook JL, Leppla SH. Receptor-mediated internalization and degradation of diphtheria toxin by monkey kidney cells. *J Biol Chem* 1979;254:11337-42.
 54. Honjo T, Nishizuka Y, Hayaishi O. Diphtheria toxin-dependent adenosine diphosphate ribosylation of aminoacyl transferase II and inhibition of protein synthesis. *J Biol Chem* 1968;243:3553-5.
 55. Breitman ML, Clapoff S, Rossant J, et al. Genetic ablation: targeted expression of a toxin gene causes microphthalmia in transgenic mice. *Science* 1987;238:1563-5.
 56. Eidels L, Proia RL, Hart DA. Membrane receptors for bacterial toxins. *Microbiol Rev* 1983;47:596-620.
 57. Stenmark H, Olsnes S, Sandvig K. Requirement of specific receptors for efficient translocation of diphtheria toxin A fragment across the plasma membrane. *J Biol Chem* 1988;263:13449-55.
 58. Middlebrook JL, Dorland RB. Response of cultured mammalian cells to the exotoxins of *Pseudomonas aeruginosa* and *Corynebacterium diphtheriae*: differential cytotoxicity. *Can J Microbiol* 1977;23:183-9.
 59. Yamaizumi M, Mekada E, Uchida T, et al. One molecule of diphtheria toxin fragment A introduced into a cell can kill the cell. *Cell* 1978;15:245-50.
 60. Gropp E, Shanabrough M, Borok E, et al. Agouti-related peptide-expressing neurons are mandatory for feeding. *Nat Neurosci* 2005;8:1289-91.
 61. Saito M, Iwawaki T, Taya C, et al. Diphtheria toxin receptor-mediated conditional and targeted cell ablation in transgenic mice. *Nat Biotechnol* 2001;19:746-50.
 62. Wrobel CJ, Wright DC, Dedrick RL, et al. Diphtheria toxin effects on brain-tumor xenografts. Implications for protein-based brain-tumor chemotherapy. *J Neurosurg* 1990;72:946-50.
 63. Brockschneider D, Pechmann Y, Sonnenberg-Riethmacher E, et al. An improved mouse line for Cre-induced cell ablation due to diphtheria toxin A, expressed from the Rosa26 locus. *Genesis* 2006;44:322-7.
 64. Woodruff MF, Anderson NA. Effect of lymphocyte depletion by thoracic duct fistula and administration of antilymphocytic serum on the survival of skin homografts in rats. *Nature* 1963;200:702.
 65. Franklin RJ, Crang AJ, Blakemore WF. The role of astrocytes in the remyelination of glia-free areas of demyelination. *Adv Neurol* 1993;59:125-33.
 66. Kanduc D, Mittelman A, Serpico R, et al. Cell death: apoptosis versus necrosis [review]. *Int J Oncol* 2002;21:165-70.
 67. Nader K, Schafe GE, Le Doux JE. Fear memories require protein synthesis in the amygdala for reconsolidation after retrieval. *Nature* 2000;406:722-6.
 68. McClung CA, Nestler EJ. Regulation of gene expression and cocaine reward by CREB and DeltaFosB. *Nat Neurosci* 2003;6:1208-15.
 69. Hebb DO. *The organization of behavior, a neuropsychological theory*. New York (NY): Wiley; 1949.
 70. Schrader LA, Anderson AE, Varga AW, et al. The other half of Hebb: K⁺ channels and the regulation of neuronal excitability in the hippocampus. *Mol Neurobiol* 2002;25:51-66.
 71. Dong Y, Green T, Saal D, et al. CREB modulates excitability of nucleus accumbens neurons. *Nat Neurosci* 2006;9:475-7.
 72. Moran TD, Han JH, Josselyn SA, et al. Lasting synaptic potentiation in the lateral amygdala of CREB^{ad+/+} mice and its inhibition by anisomycin. *Society for Neuroscience Abstracts* 2006;536.7.
 73. Han MH, Bolaños CA, Green TA, et al. Role of cAMP response element-binding protein in the rat locus ceruleus: regulation of neuronal activity and opiate withdrawal behaviors. *J Neurosci* 2006;26:4624-9.
 74. Huang YH, Lin Y, Brown TE, et al. CREB modulates the functional output of nucleus accumbens neurons: a critical role of N-methyl-D-aspartate glutamate receptor (NMDAR) receptors. *J Biol Chem* 2008;283:2751-60.
 75. Lopez de Armentia M, Jancic D, Olivares R, et al. cAMP response element-binding protein-mediated gene expression increases the intrinsic excitability of CA1 pyramidal neurons. *J Neurosci* 2007;27:13909-18.
 76. Barco A, Alarcon JM, Kandel ER. Expression of constitutively active CREB protein facilitates the late phase of long-term potentiation by enhancing synaptic capture. *Cell* 2002;108:689-703.
 77. Viosca J, Lopez de Armentia M, Jancic D, et al. Enhanced CREB-dependent gene expression increases the excitability of neurons in the basal amygdala and primes the consolidation of contextual and cued fear memory. *Learn Mem* 2009;16:193-7.
 78. Jancic D, Lopez de Armentia M, Valor LM, et al. Inhibition of cAMP response element-binding protein reduces neuronal excitability and plasticity, and triggers neurodegeneration. *Cereb Cortex* 2009;19:2535-47.
 79. Zhou Y, Won J, Karlsson MG, et al. CREB regulates excitability and the allocation of memory to subsets of neurons in the amygdala. *Nat Neurosci* 2009;12:1438-43.
 80. Crick F. Neural networks and REM sleep. *Biosci Rep* 1988;8:531-5.
 81. Reijmers L, Mayford M. Genetic control of active neural circuits. *Front Mol Neurosci* 2009;2:27.

Correction

In the May issue of the *Journal of Psychiatry and Neuroscience*, the degrees of authors Martin Roy and Philippe-Olivier Harvey should both have been listed as PhD.

Reference

Roy M, Harvey PO, Berlim MT, et al. Medial prefrontal cortex activity during memory encoding of pictures and its relation to symptomatic improvement after citalopram treatment in patients with major depression. *J Psychiatry Neurosci* 2010;35(3):152-62.