Published in final edited form as:

Nat Neurosci. 2008 April ; 11(4): 417–419. doi:10.1038/nn2050.

Redefining the central serotonergic system based on genetic lineage

Patricia Jensen^{1,†}, Anna F. Farago^{1,†}, Rajeshwar B. Awatramani^{2,†}, Michael M. Scott³, Evan S. Deneris³, and Susan M. Dymecki^{1,*}

¹ Department of Genetics, Harvard Medical School, 77 Avenue Louis Pasteur, Boston, MA 02115

² Department of Neurology and Center for Genetic Medicine, Northwestern University Feinberg Medical School, 7-113 Lurie, 303 E. Superior St., Chicago, IL 60611

³ Department of Neurosciences, Case Western Reserve University, School of Medicine, Room E653, 10900 Euclid Avenue, Cleveland, OH 44106-4975

Abstract

Central serotonin (5-HT, 5-hydroxytryptamine)-producing neurons are heterogeneous, differing in location, morphology, neurotoxin sensitivity and associated developmental disorders such as sudden infant death syndrome, fetal alcohol syndrome, and autism. Yet the molecular underpinnings associated with such heterogeneity and differential disease vulnerability are largely unknown, as are molecular markers capable of identifying physiological subtypes of 5-HT neurons. Here we redefine subtypes within the mature 5-HT system based on genetic programs differentially enacted and differentially required in progenitor cells. We present a molecular framework for the 5-HT neural system that, having genetic lineages as its basis, is likely to have physiological relevance and will provide a means for selectively accessing 5-HT neurons for in vivo manipulations.

Abnormalities in 5-HT-producing neurons are increasingly implicated in a broad spectrum of developmental disorders, including sudden infant death syndrome¹, fetal alcohol syndrome², and autism (reviewed in ³). Each disorder differs in clinical feature, and mounting evidence suggests that different 5-HT neuron subtypes are selectively affected. Heterogeneity within the 5-HT neuron population is further demonstrated by differences in anatomical distribution, cell morphology and axonal trajectory, neurotoxin sensitivity and physiological properties (reviewed in ⁴). Mechanisms that determine these differences are largely unknown and presently few molecular markers have been identified which are capable of distinguishing individual 5-HT neuron subtypes. Such knowledge is central to understanding etiological differences among 5-HT neuron disorders and for gaining genetic access to select 5-HT neuron subgroups for experimental study.

While markers capable of distinguishing mature 5-HT neuron subtypes are wanting, at hand are markers that, when viewed in combinations, can resolve 5-HT progenitor cells into discrete subsets. From these subsets may arise physiologically relevant groupings of mature 5-HT neurons; this is because developmental programs that define the fate and function of neurons are often set in motion by the action of factors differentially expressed among their antecedent

*to whom correspondence should be addressed: Telephone: 617-432-4812, Facsimile: 617-432-7595, dymecki@genetics.med.harvard.edu.

[†]these authors contributed equally

Competing Interests Statement

The authors declare that they have no competing financial interests.

progenitor cells. 5-HT progenitor cells reside in the embryonic hindbrain in bilateral territories flanking the floor plate and spanning much of the anteroposterior (AP) extent of the hindbrain. This progenitor territory can be subdivided along the AP axis into molecularly distinct subsets based on the broader partitioning of the hindbrain into segments (rhombomeres) with distinguishing gene expression profiles (reviewed in ⁵). Thus, aspects of 5-HT neuron subtype identity may be determined through the action of rhombomere(r)-specific genetic programs on resident 5-HT progenitor and precursor cell subsets. We have set out to deconstruct the 5-HT neural system based on rhombomere-defined 5-HT sublineages. Our investigations have begun with studies of 5-HT progenitor cells situating in r1, r2 or r3.

Our approach extends the recently developed paradigm of intersectional and subtractive genetic fate mapping^{6,7} (and reviewed in ⁸) through the generation of (1) a novel, broadly applicable dual recombinase-responsive indicator allele, *RC::Fela* (Fig. 1a,b) that provides enhanced single-cell resolution by comparison with our previously generated intersectional alleles^{6,7}; and (2) a highly efficient Flpe recombinase driver line, *ePet::Flpe* (Fig. 1d,e; Supplementary Fig. 1a–q), capable of mediating recombination in 5-HT precursors defined by expression of the ETS-domain transcription factor Pet-1 – thus, in most if not all 5-HT neurons (Supplementary Fig. 1a–q). Placing *RC::Fela* and *ePet::Flpe* in combination with a cre driver line active in a specific rhombomere allows for determining which mature 5-HT neurons arise from that specific rhombomere (with its associated unique molecular code); moreover, it presents a means to gain genetic access for further manipulation of just that 5-HT neuron subset in isolation – such cannot be achieved using single recombinase-based strategies.

To study the subset of 5-HT neurons arising from r1, we partnered *RC::Fela* and *ePet::Flpe* with an *En1-cre* knock-in allele (*En^{Cki}*)⁹ – while the latter allele is active in midbrain progenitor cells in addition to the entirety of r1 (Fig. 1f, data not shown, and ⁹) this is not confounding given that 5-HT neurons do not arise from midbrain^{4,10}. In triple transgenic animals, we show that the intersectional marker, nuclear-β-galactosidase (nβgal), is efficiently activated and can be used to trace r1-derived 5-HT precursor cells; in other words, those 5-HT cells having a history of both *En1* and *Pet-1* expression (Fig. 1g, blue cells; Fig. 1h–l, red cells). In these same triply transgenic embryos, 5-HT precursors derived from progeny situated caudal to r1 (caudal to *En1*) expressed enhanced green fluorescent protein (eGFP) as a lineage tracer instead of nβgal, as these cells underwent only Flpe-mediated recombination of *RC::Fela* (Fig. 1h,i; the “subtractive” green population). No cells were co-labeled with both nβgal and eGFP (Fig. 1h,i), demonstrating that nβgal was not produced at detectable levels from the Flpe/non-Cre-recombined (eGFP⁺) *RC::Fela* allele; thus, intersectional and subtractive populations can be distinguished unequivocally (Fig. 1h,i).

Upon tracking r1 (*En1-cre*)-derived 5-HT precursors into mature postnatal stages, we found that they comprise entirely 5-HT nuclei B7, B6, and even B4, despite its relatively caudal position (Fig. 2h and data not shown – red cells). r1 (*En1-cre*)-derived 5-HT neurons were also found populating ventral pontine 5-HT nuclei (B9, B8 and B5)(Fig. 2g,i,j – red cells), along with contributions from progenitor cells situated caudal to r1(*En1* territory). These findings extend the work of Zervas et al¹⁰ and Simon et al¹¹: (1) they reveal an unpredicted developmental and genetic heterogeneity among B9, B8, and B5; (2) they substantiate that the rostral 5-HT system should not only include B9-B5 but also B4 (a point variably represented in the literature); and (3) they support the notion that the requirement for *En1* to generate B7 and B6 neurons (and likely all r1 5-HT-descendants) is autonomous to the parental 5-HT progenitor cells and not secondary to alterations in the 5-HT progenitor cell environment within r1.

Upon tracking r2-derived 5-HT precursor cells into postnatal brain, by partnering *Rse2::cre*⁷ with *RC::Fela* and *ePet::Flpe*, we found that they contribute to nuclei B9, B8 and B5 (Fig.

2,1,n,o – red cells), intermingling with non-r2-derived 5-HT neurons (Fig. 2,1,n,o – green cells). Consistent with the r1-derived fate maps, r2 5-HT progenitor cells do not contribute to nuclei B7, B6, and B4 (Fig. 2m and data not shown) – instead these nuclei derive entirely from r1 (Fig. 2h and data not shown). Similar results were found upon incorporation of *Egr2::cre* (r3/r5)¹² as the cre driver: contribution to nuclei B9, B8 and B5 (Fig. 2q,s,t – red cells), intermingling with r1- and r2-derived 5-HT neurons (Fig. 2q,s,t – green cells), and no contribution to B7, B6, and B4 (Fig. 2r and data not shown). Because the *Egr2::cre* driver marks as the intersectional (nβgal+) population those 5-HT neurons arising from either r3 or r5, we cannot rule out a contribution from r5 to the B9, B8, and B5 nuclei. Arguing against such an r5 contribution, however, is that the intervening rhombomere, r4, is thought not to produce 5-HT neurons (reviewed in ⁴), creating a gap separating development of the B9-B4 nuclei (the “rostral” 5-HT neural system) from the B3-B1 nuclei (the “caudal” 5-HT neural system). Thus, the simplest interpretation of our r3/r5 5-HT fate map is that the nβgal+ cells (red cells) seen in B9, B8 and B5 (Fig. 2q,s,t) arise from r3 while the nβgal+ cells seen in the nucleus B3 (Fig. 2u) come from r5; the latter cells are dispersed among eGFP+ cells likely deriving from r6-r7. Collectively, the above findings (summarized in Fig. 2a-f) reveal that the distribution of 5-HT neurons derived from distinct rhombomere-defined gene expression domains differs from the anatomically defined groupings of mature 5-HT neurons¹³.

The transcription factor *Nkx2.2*, expressed throughout the 5-HT primordium, has been reported necessary for the generation of most 5-HT neuronal precursors except a rostral cohort (reviewed in ⁴), suggesting that some rostral 5-HT progenitors, despite expressing *Nkx2.2*, do not require it for 5-HT neuron production. Towards localizing this *Nkx2.2*-independent subset of 5-HT progenitor cells, we examined the distribution of 5-HT neurons in the postnatal *Nkx2.2*^{-/-} brain stem using our 5-HT fate map as a basis for interpretation. We found that in *Nkx2.2*^{-/-} mutants, 5-HT neurons comprising B6 and B7 are present in qualitatively appropriate number and density (Fig. 3d, compared to 3a). Additionally, we found that neurons comprising the B4 nucleus as well as a subset of neurons within B8, B5, and B9 are also present (Fig. 3e, compared to 3b, and data not shown). This distribution is highly similar to that of the r1-derived 5-HT fate map, suggesting that it is largely 5-HT progenitors situated caudal to r1 that require *Nkx2.2* activity. Unexpectedly, we also found a cohort of 5-HT neurons in the B1-B3 areas in mutant tissue (Fig. 3f, compared to 3c, and data not shown). Given our fate map showing that these most caudal nuclei do not receive contributions from r1 progenitor cells, it is possible that there exists a subset of caudal progenitors, in addition to those within r1, that are capable of producing 5-HT progeny in the absence of *Nkx2.2* activity. Alternatively, in the absence of *Nkx2.2*, r1-derived 5-HT progenitors may expand their distribution to include aspects of the B1-B3 complex, permitting a skeletal distribution of 5-HT neurons in all nuclei. To distinguish between these possibilities, we mapped in *Nkx2.2*^{-/-} mutants the fate of r1 (*En1-cre*)-derived (nβgal+) and non-r1-derived (eGFP+) 5-HT neurons (Fig. 3j-l). In the vicinity of B1-B3 in mutant tissue, we detected eGFP+ but not nβgal+ cells (Fig. 3l), suggesting that these 5-HT neurons do not arise from r1 but rather there exists a subset of caudal cells that, like r1 5-HT progenitor cells, can produce 5-HT neurons without *Nkx2.2* activity. The precise location and molecular signature of these newly identified progenitor cells remains unknown and of great interest.

By developing and applying intersectional and subtractive gene activation technologies, we have shown that 5-HT neuron subtypes with distinct molecular histories and requirements can be isolated in situ. From our findings, we define a novel molecular framework for the 5-HT system. Because this model is based on genetic lineage, we predict it to have functional and pathophysiological relevance. Further, the tools presented here provide a powerful template from which select 5-HT lineages can be manipulated in vivo in virtual isolation – such capabilities open new avenues for considering 5-HT neuron functions and the etiological differences among 5-HT neuron disorders.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Dr. A. Joyner for the *En1-cre* mice, Dr. Sylvie Schneider-Maunoury for the *Egr2-cre*, Dr. L. Sussel for the *Nkx2.2* mutant mice, and Drs. C. Cepko and L. Murtaugh for plasmids. We also thank Dr. F. Farley and J. J. Mai for their technical assistance, as well as Drs. H. Kinney, E. Nattie, G. Richerson, and K. Commons and members of the Dymecki and Kinney laboratories for their input and critical reading. A.F.F. and S.M.D. were supported by grants from the NIH (P01 HD036379, R01 DK 067826, and R21 DA023643), P.J. was supported by a grant from the Charles H. Hood Foundation (Boston, MA), R.B.A. was supported by a grant from the Charles King Trust Foundation, and E.S.D. and M.M.S. were supported by R01 MH62723.

References

1. Paterson DS, et al. *Jama* 2006;296:2124–32. [PubMed: 17077377]
2. Riikonen RS, et al. *Biol Psychiatry* 2005;57:1565–72. [PubMed: 15953494]
3. Pardo CA, Eberhart CG. *Brain Pathol* 2007;17:434–47. [PubMed: 17919129]
4. Cordes SP. *Clin Genet* 2005;68:487–94. [PubMed: 16283875]
5. Lumsden A, Krumlauf R. *Science* 1996;274:1109–15. [PubMed: 8895453]
6. Farago AF, et al. *Neuron* 2006;50:205–18. [PubMed: 16630833]
7. Awatramani R, et al. *Nat Genet* 2003;35:70–5. [PubMed: 12923530]
8. Dymecki SM, Kim JC. *Neuron* 2007;54:17–34. [PubMed: 17408575]
9. Kimmel RA, et al. *Genes Dev* 2000;14:1377–89. [PubMed: 10837030]
10. Zervas M, et al. *Neuron* 2004;43:345–57. [PubMed: 15294143]
11. Simon HH, et al. *Mol Cell Neurosci* 2005;28:96–105. [PubMed: 15607945]
12. Voiculescu O, et al. *Genesis* 2000;26:123–6. [PubMed: 10686605]
13. Wallace JA, et al. *Brain Res Bull* 1982;9:117–29. [PubMed: 6756548]
14. Scott MM, et al. *J Neurosci* 2005;25:2628–36. [PubMed: 15758173]
15. Lidov HG, Molliver ME. *Brain Res Bull* 1982;8:389–430. [PubMed: 6178481]

a Dual recombinase-responsive indicator allele

b Intersectional fate mapping strategy

Figure 1. Intersectional and subtractive cell marking distinguishes r1 (*En1-cre*)- from non-r1-derived 5-HT precursors

(a) *RC::Fela* dual recombinase responsive indicator allele. *CAG* promoter/enhancer elements followed by, 5' to 3', an *FRT*-flanked *PGK-neo^R* cassette with two consecutive SV40 polyadenylation (pA) sequences; a *loxP*-flanked *eGFP* cassette with five consecutive bovine growth hormone pA sequences; and a *nlacZ* sequence followed by a single pA sequence was targeted to the *Rosa26* locus. (b) Strategy for single (*Flp-FRT*) and dual (*Flp-FRT* and *cre-loxP*) recombinase-based fate mapping. Tubes represent the neural tube at early (left) to late (right) developmental stages. Top row illustrates early transient expression of *GeneA*. Middle row illustrates expression of *GeneA::Flpe* and *GeneB::cre* transgenes. Bottom row illustrates

coupling of *GeneA::Flpe* and *GeneB::cre* transgenes with dual recombinase-responsive indicator allele. Cells expressing Flpe recombinase activate expression of eGFP (the subtractive population, green); cells expressing Flpe and Cre recombinase activate expression of nβgal (the intersectional population, red). Reporter molecule activation is permanent, and expression is retained in all descendent cells (bottom right). (c) Cartoon schematic illustrates sagittal section of embryonic day (E) 12.5 brain. (d, e) E12.5 *ePet::Flpe* transgenic embryo. A 40kb *ePet1* enhancer region¹⁴ coupled with β-globin minimal promoter followed by an *intron-Flpe* cassette⁷ was used to generate the *ePet::Flpe* transgene. Adjacent sagittal sections, and horizontal sections inset, show expression of *Pet1* (d) and *Flpe* (e) mRNA by in situ hybridization detectable in the mantle zone (mz) immediately ventral to the ventricular zone (vz). (f) E11.5 *En1-cre; indicator* double transgenic embryo, and horizontal section inset, showing *En1* descendants as βgal⁺ by X-gal detection. (g-l) *En1-cre; ePet::Flpe; RC::Fela* triply transgenic animals. E12.5 sagittal sections (g-i) and P21 coronal sections (j-l) stained to detect nβgal (the r1 (*En1-cre*)-derived intersectional 5-HT population) by X-gal detection (g) and immunofluorescence (h, i, k, l), and eGFP (the non-r1-derived 5-HT subtractive population) by immunofluorescence (h, i, k, l).

Figure 2. The organization of 5-HT neurons as defined by embryonic origin and developmental gene expression profile is different from that based on anatomical architecture

(a) Shown are cartoon schematics of the embryonic hindbrain (left) and the mature brainstem (right): dark blue represents the 5-HT progenitor cells in r1 (left) or r1-derived mature 5-HT neurons (right); yellow, r2 5-HT progenitors (left) or descendants (right); light blue, r3; pink, presumed r5; gray, r6-r7. The mature sagittal diagram is compressed along the mediolateral axis. The B1-B9 abbreviations refer to the names given to nuclei defined anatomically, some of which are also referred to as the dorsal raphe nucleus (DR) or median raphe nucleus (MR). Dashed lines represent coronal sections (b-f) presented rostral to caudal, top to bottom. r1-derived 5-HT neurons (dark blue) populate the B7, B6 and B4 groups in their entirety, as well

as aspects of the B9, B8 and B5 nuclei. r2-derived 5-HT neurons (yellow) populate the B9, B8 and B5 nuclei intermingled with both the r1-derived (dark blue) and r3-derived (light blue) 5-HT neurons. Presumed r5-derived 5-HT neurons (pink) populate the B3 nucleus intermingled with more caudally (r6-r7-) derived 5-HT neurons (gray). Boxes highlight images of coronal sections from various triple transgenic P21 brainstem stained to detect nβgal (red) and eGFP (green) by immunofluorescence: **(g-k)** En1::cre (r1); ePet::Flpe; RC::Fela; **(l-p)** Rse2::cre (r2); ePet::Flpe; RC::Fela; **(q-u)** Egr2::cre (r3/r5); ePet::Flpe; RC::Fela,. 5-HT neurons with a combined history of rhombomere specific::cre and ePet::Flpe activity are identified as nβgal⁺ (the intersectional population, in other words that population derived from a particular rhombomere), while cells with a history of only ePet::Flpe activity are eGFP⁺ only (the subtractive population, in other words those neurons derived from the remaining pool of 5-HT progenitor cells).

Figure 3. r1 (*En1-cre*)- and non-r1-derived 5-HT neurons populate the postnatal *Nkx2.2*^{-/-} mutant brainstem

(**a-f**) Coronal brainstem sections from P0.5 *Nkx2.2*^{-/-} animals (**d-f**) and wild-type littermates (*Nkx2.2*^{+/+}, **a-c**) stained to detect 5-HT. Adjacent images are taken from approximately the same axial levels. At the level of the B4, B6 and B7 nuclei (shown is B7), there is a comparable distribution of 5-HT positive neurons in *Nkx2.2*^{+/+} and *Nkx2.2*^{-/-} sections. In more ventral and caudal territories there is a clear diminution of the B9, B8, B5, B3, B2 and B2 nuclei (shown is B8 and B3). (**g-l**) Coronal sections from P0.5 *En1-cre; ePet::Flpe; RC::Fela; Nkx2.2*^{-/-} animals (**g-i**) and wild-type littermates (*En1-cre; ePet::Flpe; RC::Fela; Nkx2.2*^{+/+}, **j-l**) stained to detect nβgal (the r1 (*En1-cre*)-derived intersectional 5-HT population) and eGFP (the non-r1-derived 5-HT subtractive population) by immunofluorescence. (**j-l**) In *Nkx2.2*^{-/-} brains the rostral 5-HT nuclei (B4-B9) receives contributions exclusively from r1-derived 5-HT progenitors (**j,k**) and the caudal 5-HT nuclei (B1-3) is populated by non-r1-derived 5-HT progenitors (**l**).