Published in final edited form as:

Neurobiol Learn Mem. 2009 February ; 91(2): 155–165. doi:10.1016/j.nlm.2008.09.006.

Some Surprising Findings on the Involvement of the Parietal Lobe in Human Memory

Ingrid R. Olson^{1,2} and Marian Berryhill^{1,2}

¹ Department of Psychology, Temple University

² Center for Cognitive Neuroscience, University of Pennsylvania

Abstract

The posterior parietal lobe is known to play some role in a far-flung list of mental processes: linking vision to action (saccadic eye movements, reaching, grasping), attending to visual space, numerical calculation, and mental rotation. Here we review findings from humans and monkeys that illuminate an untraditional function of this region: memory. Our review draws on neuroimaging findings that have repeatedly identified parietal lobe activations associated with short-term or working memory and episodic memory. We also discuss recent neuropsychological findings showing that individuals with parietal lobe damage exhibit both working memory and long-term memory deficits. These deficits are not ubiquitous; they are only evident under certain retrieval demands. Our review elaborates on these findings and evaluates various theories about the mechanistic role of the posterior parietal lobe in memory. The available data point towards the conclusion that the posterior parietal lobe plays an important role in memory retrieval irrespective of elapsed time. The two models that are best supported by existing data are the Attention to Memory Model and the Subjective Memory Model. We conclude by formalizing several open questions that are intended to encourage future research.

Keywords

parietal lobe; Balint's syndrome; short-term memory; working memory; episodic memory; spatial processing; retrieval; episodic buffer; meta-memory

If you were to peruse any textbook on memory or neuroscience, you would be hard-pressed to find the terms “memory” and “parietal lobe” together. How then, do we explain the large number of neuroimaging findings reporting parietal lobe activations to various mnemonic demands? The sheer volume of these findings raises the question of whether the parietal lobe plays a functional role in mnemonic processing that has been overlooked.

To address this question, we review evidence linking the parietal lobe to memory. We focus on visual short-term or working memory (WM) and episodic memory for the simple reason that there is now sufficient material in these literatures to provide some nascent consensus. We note that links between verbal WM and parietal lobe function has recently been reviewed elsewhere (Buchsbaum and D'Esposito, 2008).

Corresponding author: Ingrid R. Olson, Department of Psychology, Temple University, 1701 N. 13th Street, Philadelphia, PA 19122, Telephone: 215-204-7318, Email: iolson@temple.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Our first piece of evidence that the parietal lobe may have some role in memory comes from white matter tractography revealing close anatomical linkages between the parietal lobe and frontal and medial temporal lobe areas. Subsequent sections describe experimental evidence from spatial WM, object WM and episodic memory studies. We explore various hypotheses describing the mechanism of parietal involvement in memory and conclude with a series of open questions meriting further research.

Anatomy and Connectivity of the Posterior Parietal Cortex

Because in vivo axon tracing techniques cannot be applied to humans, much of our knowledge of the connectivity of the parietal lobe is drawn from work in non-human primates. This presents us with a significant intellectual hurdle, since the extent of inter-species homology is unclear (Culham and Kanwisher, 2001; Glover, 2004), but see (Rushworth, Behrens, and Johansen-Berg, 2006). The human parietal lobe is nearly 20 times larger than that of the macaque. This ratio is markedly higher than the same comparison made between human and macaque temporal (9 times larger), or occipital (2 times larger) cortices (Van Essen, Lewis, Drury, Hadjikhani, Tootell, Bakircioglu, and Miller, 2001). The size difference is partially explained by the expansion of the human inferior parietal lobe. Indeed, there appears to be no equivalent of the human supramarginal gyrus (BA 39) in monkeys (Karnath, Ferber, and Himmelbach, 2001). Despite this problem, there are some parietal regions that appear to share anatomical and functional similarities between macaques and humans (Culham and Valyear, 2006) and more recent non-invasive functional connectivity studies are beginning to provide information about human parietal connectivity (Rushworth et al., 2006).

Beginning anteriorly, the human parietal lobe is located immediately posterior to the central sulcus, where the somatosensory cortices (Brodmann areas (BA) 1, 2, and 3) occupy the postcentral sulcus. More posterior regions, referred to herein as the posterior parietal cortex (PPC) are traditionally divided into the superior parietal lobe (SPL; BA 5 and 7), the medial parietal lobe (mesial extent of BA 7), and the inferior parietal lobe (IPL; BA 39 and 40)¹. The SPL is posterior to somatosensory cortex and above the intraparietal sulcus (IPS), excluding medial parietal cortex. The medial parietal lobe consists primarily of the precuneus, which is buried in the interhemispheric fissure. Recently some authors have defined the medial parietal cortex very inclusively, including the limbic posterior cingulate and retrosplenial cortices in their definitions of this region (Vincent, Snyder, Fox, Shannon, Andrews, Raichle, and Buckner, 2006). Here, we hold to the more traditional definition of medial parietal cortex. The IPL is located below the horizontal portion of the IPS, anterior to the superior occipital lobe, and consists of the angular (BA 39) and the supramarginal gyri (BA 40).

Inferior Parietal Lobe and Medial Temporal Lobe Connectivity

In 1957, Scoville and Milner reported that bilateral damage to the human medial temporal lobe causes chronic anterograde amnesia (Scoville and Milner, 1957). We now know that within the medial temporal lobe (MTL), the hippocampus plays a central role in spatial and episodic memory formation. Monkey tractography studies have identified two large axons tracts connecting the IPL to the medial temporal lobe. First, portions of the inferior longitudinal fasciculus connect the angular gyrus with the parahippocampal gyrus (Seltzer and Pandya, 1984). Second, the cingulum bundle, which lies within the cingulate cortex, connects lateral and medial regions of the posterior IPL (PG and Opt²) to the parahippocampal gyrus (Seltzer

¹An alternative anatomical nomenclature divides the posterior parietal lobe into dorsal and ventral regions. The dorsal parietal lobe corresponds to the lateral and medial parts of BA7, the ventral parietal lobe corresponds to the inferior parietal lobe.

²The anatomical terms PG, PF, Opt, 7a, and 7b are used primarily by monkey researchers following von Economo's terminology (von Economo, C., & Koskinas, G. N. (1925). *Die Cytoarchitektonik der Hirnrinde des erwachsenen Menschen*. Berlin: Verlag von Julius Springer.

and Pandya, 1984). Connections also exist between entorhinal cortex and IPL area 7 in the monkey (Wellman and Rockland, 1997; Insausti and Amaral, 2008). Most interestingly, there are connections between area CA1 in the anterior portion of the hippocampus and area 7a, in the posterior IPL of the monkey (also known as PG or Opt) (Rockland and Van Hoesen, 1999; Clower, West, Lynch, and Strick, 2001) and 7b, found in the anterior IPL (also known as area PF) of the monkey (Rockland and Van Hoesen, 1999). Information flows predominantly from CA1 to 7a, suggesting that memory functions of the hippocampus modulate processing in area 7a (Clower et al., 2001) (see Figure 1A). Connections have also been identified between the presubiculum and area 7a (Cavada and Goldman-Rakic, 1989; Ding, Van Hoesen, and Rockland, 2000).

Although these findings are largely based on investigations of monkey anatomy, there is some evidence that the human inferior parietal lobe has similar patterns of MTL connectivity. Rushworth and colleagues, using a diffusion weighted tractography method, reported a pattern of connectivity between a region of the parahippocampal gyrus just lateral to the hippocampus and the angular gyrus that resembled the inferior longitudinal fasciculus (see Figure 1C). The angular gyrus in the IPL was the only lateral parietal region with a high probability of connection to the parahippocampus (Rushworth et al., 2006). In addition, resting state fluctuations in the BOLD signal show correlations between the hippocampus and medial-lateral PPC (Vincent et al., 2006); see also (Greicius and Menon, 2004; Takahashi, Ohki, and Kim, 2008).

Superior and Inferior Parietal-Frontal Connectivity

Portions of the frontal lobe are known to play a critical supporting role in mnemonic functions. In the monkey, several large fiber tracts connect both inferior and superior parietal regions to the prefrontal cortex. The dominant white matter tract connecting these regions is the superior longitudinal fasciculus (SLF). The SLF can be divided into three subcomponents from the most superior to the most inferior: SLF I, SLF II, SLF III (Makris, Kennedy, McInerney, Sorensen, Wang, Caviness, and Pandya, 2005; Schmahmann, Pandya, Wang, Dai, D'Arceuil, de Crespigny, and Wedeen, 2007) (see Figure 1B). The SLF I connects medial and dorsal portions of the SPL (part of areas PE, PG, 31) to dorsal premotor and prefrontal regions (BA 6 and 9, supplementary motor area) (Schmahmann and Pandya, 2006; Schmahmann et al., 2007) and is most likely important for higher order motor behavior (Petrides and Pandya, 2006). The SLF II unites the posterior IPL (area PO in the IPS, PG, Opt) with the mid- and dorso-lateral prefrontal cortices (BA 6, 8, 9, 46) (Schmahmann and Pandya, 2006; Schmahmann et al., 2007). The fronto-occipital fasciculus links a broad swath of medial (areas PO, PG medial) and posterior IPL regions (areas PG lateral, Opt, DP) with a band of regions in dorsolateral prefrontal cortex (areas 6D, 8Ad, 8B, 9, 46d) (Schmahmann and Pandya, 2006). SLF III connects the anterior inferior parietal lobe (region PF/PO of the IPS) and the anterior intraparietal area within the IPS, with ventral premotor, ventral prefrontal, and dorsolateral prefrontal cortex (areas 6, 44, 9/46) (Schmahmann and Pandya, 2006; Schmahmann et al., 2007). The function of these tracts is unknown, although it has been suggested that they are important for visual attention and working memory (Cavada and Goldman-Rakic, 1989; Schmahmann et al., 2007). Last, a separate axon tract, the cingulum bundle (temporal aspects only) links posterior IPL (area PG, Opt) to the supplementary motor area and the dorsolateral prefrontal cortex (Mori, 2005). The cingulum bundle is considered the "dorsal limbic pathway" as it has branches that extend not only throughout the length of the cingulate, but also to the hippocampus and parahippocampus (Seltzer and Pandya, 1984). Thus, via the cingulum bundle, the posterior IPL is connected both to the dorsolateral prefrontal cortex and to the hippocampus.

Working Memory for Spatial Attributes

Over fifty years ago MacDonald Critchley remarked “*Bilateral disease of the brain, and especially of the parieto-occipital regions, may be followed by the most conspicuous spatial disorders, especially entailing visual disorientation.*” (Critchley, 1953, p. 354). Of the many spatial processes linked to parietal lobe function, spatial WM is a more recent addition. This type of memory allows one to remember spatial information over short delay periods— where one looked a few moments ago, for instance. There are several sources of evidence linking the parietal lobe to spatial working memory: neurophysiological, neuropsychological, brain imaging, and transcranial magnetic stimulation (TMS).

“Memory cells” are cells that display persistent activity during the delay period of a WM task. The discharge of memory cells is higher during the delay period than during baseline periods between trials. Such cells were first found in the dorsolateral prefrontal cortex (PFC) of monkeys performing spatial WM tasks (Fuster and Alexander, 1970; Niki and Watanabe, 1976). Since their discovery, memory cells have also been found in other brain regions including the lateral intraparietal cortex in the PPC (Gnadt and Andersen, 1988; Mazzoni, Bracewell, Barash, and Andersen, 1996). PPC memory cells also exhibit delay activity during spatial WM tasks (Gnadt and Andersen, 1988; Fuster, 1990; Constantinidis and Steinmetz, 1996; reviewed in Constantinidis, 2006; Constantinidis and Procyk, 2004; Constantinidis and Wang, 2004) that is comparable to delay activity observed in PFC neurons (Fuster, 1984). Selective cooling of areas 5 and 7 disrupts spatial memory (Quintana and Fuster, 1993).

In accord with these findings, it has been known for several decades that damage to the right PPC in humans leads to impaired spatial WM. Spatial WM deficits subsequent to PPC damage are observed across a range of tasks and stimuli (see Table 1) such as old/new recognition of simultaneously or sequentially presented locations (Husain, Mannan, Hodgson, Wojciulik, Driver, and Kennard, 2001; Malhotra, Jager, Parton, Greenwood, Playford, Brown, Driver, and Husain, 2005; Pisella, Berberovic, and Mattingley, 2004; Berryhill and Olson, 2008b), or recall, by pointing, to a series of locations (Baldo and Dronkers, 2006; De Renzi, Faglioni, and Previdi, 1977; De Renzi and Nichelli, 1975; Hanley, Young, and Pearson, 1991; Malhotra et al., 2005).

It is important to bear in mind that IPL lesions, near the temporoparietal junction frequently result in hemispatial neglect (Vallar and Perani, 1986). Hemispatial neglect is a heterogeneous disorder (Shimozaki, Kingstone, Olk, Stowe, and Eckstein, 2006), reviewed in (Ellis, Della Sala, and Logie, 1996; Heilman, Valenstein, and Watson, 2000; Driver and Vuilleumier, 2001; Halligan, Fink, Marshall, and Vallar, 2003; Heilman, Jeong, and Finney, 2004; Pisella and Mattingley, 2004; Buxbaum, 2006; Hillis, 2006) characterized by a failure to attend, react to, or to perceive stimuli presented in the contralesional visual field. Spatial WM deficits are known to occur in patients with hemispatial neglect. For instance, several recent studies of neglect patients required subjects to remember sequentially presented locations arrayed in a vertical line. The results showed that neglect patients had impaired spatial WM performance (Malhotra, Mannan, Driver, and Husain, 2004; Malhotra et al., 2005; Parton, Malhotra, Nachev, Ames, Ball, Chataway, and Husain, 2006). When the eye-movements of neglect patients were assessed in visual search tasks, it was shown that they tend to revisit previously foveated items as if they were new items, indicative of a spatial WM impairment (Husain et al., 2001; Wojciulik, Husain, Clarke, and Driver, 2001). After neglect resolves (or even if it never existed), spatial WM deficits remain present (see Table 1), indicating that the spatial WM deficits are not attributable to inattentiveness to portions of space. Converging evidence for the importance of the right PPC in spatial WM is offered by several TMS studies showing that transient inactivation of these regions impairs spatial WM (Kessels, van Zandvoort, Postma, Kappelle, and de Haan, 2000; Oliveri, Turriziani, Carlesimo, Koch, Tomaiuolo,

Panella, and Caltagirone, 2001; Koch, Oliveri, Torriero, Carlesimo, Turriziani, and Caltagirone, 2005).

Neuroimaging studies have overwhelmingly corroborated these findings (Smith and Jonides, 1997; 1998; D'Esposito, Aguirre, Zarahn, Ballard, Shin, and Lease, 1998; Ungerleider, Courtney, and Haxby, 1998; Hartley and Speer, 2000;). More recent event-related fMRI findings suggest that the PPC is involved specifically in maintaining spatial information, as shown by sustained delay-period activity (reviewed in Curtis, 2006; Linden, 2007) and possibly in WM manipulation (Postle, Ferrarelli, Hamidi, Feredoes, Massimini, Peterson, Alexander, and Tononi, 2006).

The PPC appears to work in concert with the PFC to solve the computations necessary for accurate spatial WM. The firing patterns of these regions is intrinsically linked: when cell populations in the dorsolateral PFC and lateral PPC cells were simultaneously recorded during a delayed saccade task, the neural activity patterns were shown to match (Chafee and Goldman-Rakic, 1998; Quintana and Fuster, 1999). Selective lesions to these regions in the macaque leads to a similar disruption in saccadic memory – saccades become less accurate as the delay period lengthens (reviewed in Curtis, 2006). Similarly, cortical cooling of either region while recording single-units in the other region showed that these regions have a strong interdependency during spatial WM tasks (Chafee and Goldman-Rakic, 2000). In addition, TMS to either the PPC or dorsolateral PFC during spatial WM maintenance slows reaction times (Oliveri et al., 2001; Koch et al., 2005). Both regions are robustly activated during spatial WM tasks (Jonides, Smith, Koeppe, Awh, Minoshima, and Mintun, 1993; Smith and Jonides, 1998); reviewed in (Ungerleider et al., 1998; Wager and Smith, 2003; Passingham and Sakai, 2004; Naghavi and Nyberg, 2005; D'Esposito, 2007).

In spite of these similarities, there are differences between the neural profiles associated with the PPC and the PFC. The mnemonic activity of PPC neurons, specifically neurons in area 7a, appears to be closely aligned with bottom-up attentional activity as PPC neurons fire to any salient stimuli, even when it is behaviorally irrelevant and PPC neurons begin firing before PFC neurons under bottom-up attentional conditions, suggesting that this region detects and registers the spatial location of visually presented stimuli (Constantinidis and Steinmetz, 2005; reviewed in Constantinidis, 2006). In contrast, the mnemonic activity of lateral PFC and frontal eye field neurons may be more closely aligned with top-down attention as frontal neurons begin to fire before lateral intraparietal neurons under top-down attentional conditions (Buschman and Miller, 2007).

Working Memory for Objects or Object Features

In contrast to the strong converging evidence linking the PPC to spatial WM functions, the evidence for functional links between the PPC and object WM is weaker. Neurons in macaque lateral intraparietal sulcus are responsive to object shape during the remember periods of delayed-match-to sample tasks. Neuronal selectivity can be independent of eye movements, reaching, or object manipulation – or in other words, independent from spatial processing (Serenio and Maunsell, 1998; Sereno and Amador, 2006). Also, lateral intraparietal neurons exhibit delay activity to target color when it is behaviorally relevant (Toth and Assad, 2002).

In humans, BOLD activity in the IPS has been shown to titrate with the number of shapes and colors that can be held in memory over brief amount of time, suggesting that this region has some role in governing WM capacity (Todd and Marois, 2004; 2005; Macoveanu, Klingberg, and Tegner, 2006; Xu and Chun, 2006). This activity is not due to the number of items viewed at encoding, but rather to the number of items accurately maintained in WM, lending support to the idea that the function of the IPS is mnemonic, not perceptual (Todd and Marois, 2004). Converging evidence for these findings can be found in a small number of neuropsychological

studies. We required subjects with or without right PPC damage to remember four sequentially presented colors, abstract shapes, or common objects over a short delay period and then make an old/new recognition decision. The stimuli were presented sequentially and at central fixation in order to minimize spatial demands. The results showed that right PPC damage was associated with diminished WM performance across stimulus categories (Berryhill and Olson, 2008b; for deficits of face WM see Warrington and James, 1967). A subsequent study tested patients with bilateral parietal lobe damage on exactly the same task. A similar pattern of deficits was observed (Berryhill and Olson, 2008a).

Unfortunately, the results of other studies cast some doubt on the robustness of these findings. Several patients with right PPC damage and neglect were tested on an object WM task and no deficits were observed (Pisella et al., 2004). Another study found that right PPC patients were impaired on object delay-match-to-sample tasks only when the probed dimension (object or spatial) was unpredictable (Finke, Bublak, and Zihl, 2006), hinting that the right PPC's role in WM may be attentional rather than mnemonic. One of the only TMS studies relevant to this question administered single TMS pulses during the maintenance period of an object WM task, but found no effect of unilateral or bilateral PPC stimulation on performance (Oliveri et al., 2001).

As a whole, the evidence suggests that object WM relies to some degree on an intact parietal lobe although several unanswered questions remain (see Box 1, question 1). The fact that there are discrepant findings is troubling, and the reason for this remains elusive, although we have observed that the manner in which memory is probed leads to different patterns of performance after parietal lobe damage. This finding hints that retrieval processes may be key to understanding the functionality of the PPC (Berryhill and Olson, 2008a; Berryhill, Phuong, Picasso, Cabeza, and Olson, 2007; Drowos, Berryhill, and Olson, 2008, accepted pending revisions). We expand on this topic in the next section.

Mechanistic Role of the Parietal Lobe in Working Memory

The mechanistic role of the parietal lobe in WM is currently unknown. The neuroimaging literature has focused on WM maintenance and two hypothetical roles of the PPC in WM that have emerged are in the manipulation of information during WM maintenance (Chamod and Petrides, 2007; Wendelken, Bunge, and Carter, 2008), and in total information accumulation during WM maintenance (Xu, 2007). In contrast, the neuropsychological literature has offered the hypothesis that the PPC's role in WM is in instantiating particular retrieval processes (Berryhill and Olson, 2008a).

Information Manipulation Hypothesis

Several investigators have proposed that the mnemonic function of the PPC is specifically associated with the manipulation of information held in WM (Chamod and Petrides, 2007; Wendelken et al., 2008). For instance, in one recent study subjects were required to remember letters and colors under several different conditions. In the *low* and *high* memory load conditions, the task was to remember the alphanumeric items in the memory set and respond whether two probe items had been present in the preceding memory set. In the *organization* condition, the task was to remember whether the probe items had been previously shown in the same color. In their second experiment, the organization task was to spatially reorganize the items according to arrows. In both studies, the authors observed frontal activity (DLPFC, VLPFC) during the delay period associated with WM load. However, superior parietal activity (BA7) was only observed when WM maintenance required manipulation (Wendelken et al., 2008). Similar SPL regions are activated during other tasks demanding item reorganization in WM such as order sequencing tasks (Marshuetz and Smith, 2006; Marshuetz, Smith, Jonides, DeGutis, and Chenevert, 2000; Wager and Smith, 2003), and mathematical calculation tasks

(Dehaene, Molko, Cohen, and Wilson, 2004; Dehaene, Spelke, Pinel, Stanescu, and Tsivkin, 1999). One problem with this interpretation is it does not account for the large number of WM studies in which the task does not include a manipulation requirement, yet SPL activations are observed. Also, the manipulation hypothesis falls prey to the “difficulty explanation”, that greater PPC activations in the manipulation condition are simply due to the greater difficulty of that task compared to the baseline condition.

Information Load Hypothesis

Behavioral studies have shown that on average, most people can maintain approximately four objects in visual WM (Luck and Vogel, 1997) and that memory capacity varies with the complexity (Alvarez and Cavanagh, 2004), or with the similarity of these items (Olson and Jiang, 2002; Awh, Barton, and Vogel, 2007). Interestingly, portions of the parietal lobe appear to be sensitive to memory capacity limitations. In one study, subjects observed an array of colored squares and after a brief delay, were shown a probe image and made a same/different judgment. In a whole brain analysis, only one region titrated with memory capacity: the superior IPS. Note that superior IPS activity was not modulated by set size but rather, the number of objects that subjects could accurately maintain in memory such that neural activity leveled off as memory capacity reached asymptote. Additional analyses revealed that activity correlated with individual performance on the object WM task (Todd and Marois, 2004; 2005). This finding and those of others (Xu and Chun, 2006) suggest that one role of the superior IPS in WM is to maintain a finite number of objects over short delay periods.

An alternative explanation is that the superior IPS maintains visual features, rather than discrete objects, over a short period of time. Note that the number of objects and visual features are highly correlated in most studies using visual stimuli. To understand their individual contributions, Xu (2007) dissociated feature and object information. In her study, subjects were required to remember stimulus arrays with the same number of objects but different numbers of features by varying shape and/or color. In a separate task, the number of objects was varied, but the number of features remained constant. The results showed that the superior IPS exhibited greater changes in signal strength when there were more visual features present. However there was no difference in the superior IPS when the number of objects was varied, holding the number of features constant (Xu, 2007; see also Xu and Chun, 2006). This finding provides evidence for the view that the superior IPS represents the sum total of perceptual information stored in WM.

These findings predict that superior IPS damage will diminish visual WM capacity, as a function of the number of visual features of the items. We tested this prediction in two studies by requiring patients with unilateral or bilateral parietal lobe damage to remember visual stimuli that varied in the number of features: color patches, novel shapes, and common objects. The first part of the prediction was supported: visual WM capacity was diminished after PPC damage. However, the second part of the prediction was *not* borne out in these data: there was no effect of stimulus complexity or number of features on patient performance (Berryhill and Olson, 2008a; b), suggesting that visual feature load does not fully explain the mechanistic function of the PPC in WM.

Retrieval Process Hypothesis

In a series of WM studies, we examined the role of the PPC in WM while manipulating the retrieval process. The performance of two patients with bilateral parietal damage was compared to that of matched controls (Berryhill and Olson, 2008a). In the first pair of experiments, order WM was tested with either recall or recognition retrieval probes. Subjects observed four sequentially presented items (colors, shapes, or objects) and after a brief delay, the same items were repeated. The recognition task was to make a same/different judgment regarding the order

of the four items. In the recall version, a single probe item appeared after the delay and the task was to provide the ordinal position (1–4) of that image. The patients exhibited impaired WM performance when probed by old/new recognition, but intact WM performance when probed by recall. In a second pair of studies, the task was to remember the identity of four common objects over a brief delay, and then to make either an old/new judgment or to verbally name the objects that had been shown. The same pattern of preserved recall accompanied by impaired recognition was observed (see Figure 2). These findings cannot be attributed to deficient encoding or maintenance processes because these demands were held constant; only the retrieval process varied. This finding leads us to suggest that the PPC's role in WM is specifically linked to retrieval processes. If true, this function coincides with a wealth of findings on retrieval in the episodic memory literature, discussed next.

Episodic Memory

Findings linking episodic memory to the parietal lobe recently have attracted considerable attention from the cognitive neuroscience community. Several excellent reviews of this topic are available and we direct interested readers to these sources (Wagner, Shannon, Kahn, and Buckner, 2005; Cabeza, 2008; Cabeza, Ciaramelli, Olson, and Moscovitch, 2008; Simons and Mayes, 2008; Vilberg and Rugg, 2008, see also Uncapher & Wagner, this volume). To summarize: parietal activity is greater when an old stimulus is correctly identified as old, compared to when a new item is correctly identified as new. The benefit afforded stimuli retrieved as old even extends to items that were *incorrectly* perceived as old (Wheeler and Buckner, 2003; Kahn, Davachi, and Wagner, 2004). Separable regions in the lateral parietal lobe, a superior and an inferior region, show complementary patterns of activity; the superior area appears to reflect top-down influence in mediating low-confidence responses and the inferior region reflects bottom-up influence mediating high-confidence responses (Kim and Cabeza, 2007).

Until recently, these findings were viewed with skepticism because they suffered from an absence of converging evidence from other techniques. A key piece of converging evidence was recently provided by a study of patients with bilateral parietal lobe lesions (lesions are shown in Figure 2a). In this study, patients and matched controls were asked to recollect various autobiographical memories. In condition 1, they freely recalled events from their lifetime in as much detail as possible. In condition 2, they answered specific questions about these memories. The results showed that parietal lobe damage decreased the vividness and amount of detail freely recalled. However, when patients were probed for specific details pertaining to their memories, they performed normally (Berryhill et al., 2007; see also Hunkin, Parkin, Bradley, Burrows, Aldrich, Jansari, and Burdon-Cooper, 1995). We recently tested the same patients on the Deese-McDermott Roediger (DRM) false memory paradigm and found that they exhibited normal false memory when tested by recall and impaired levels when tested by old/new recognition (see Figure 2). Interestingly, they also exhibited an abnormally low level of “remember” responses compared to “know” response (Drowos et al., 2008, accepted pending revisions) (see Figure 2).

Mechanistic Role of the Parietal Lobe in Episodic Memory

A large number of fMRI studies have reported that portions of the PPC are modulated as a function of episodic memory retrieval demands. We do not discuss these findings in detail as they have been thoroughly reviewed elsewhere (Naghavi and Nyberg, 2005; Wagner et al., 2005; Cabeza, 2008). Suffice it to say that these findings come from a number of different laboratories using different tasks and stimuli, providing a compelling case for PPC involvement in episodic memory retrieval. Discerning what exactly that role might be is the subject of much current investigation. Suggested roles include serving as an episodic buffer, a type of WM

(Baddeley, 2000; Vilberg, Moosavi, and Rugg, 2006; Vilberg and Rugg, 2008), providing an assessment of memory signal strength, as a mnemonic accumulator (Wagner et al., 2005), in directing internal attention (Cabeza, 2008; Cabeza et al., 2008; Ciaramelli, Grady, and Moscovitch, 2008), or as a gauge of memory subjectivity or confidence (Ally, Simons, McKeever, Peers, and Budson, 2008; Simons, Peers, Mazuz, Berryhill, and Olson, 2008-submitted).

Our episodic memory findings, described earlier, also point towards disrupted memory retrieval processes following PPC damage because episodic memory is impaired under some, but not all, retrieval conditions (Berryhill et al., 2007; Drowos et al., 2008, accepted pending revisions). Preserved memory on several episodic memory tasks, in studies conducted by ourselves and others (Davidson, Anaki, Ciaramelli, Cohn, Kim, Murphy, Troyer, Moscovitch, and Levine, 2008; Simons, Peers, Hwang, Ally, Fletcher, and Budson, 2008; Simons et al., 2008-submitted) indicate that memory retrieval per se is not diminished by parietal lobe damage, but what is damaged is a particular subprocess that tapped in certain retrieval tasks.

One proposed hypothesis that appears to offer a happy marriage between the WM findings reviewed earlier, and episodic memory findings is that portions of the PPC serve as the interface between WM and episodic memory retrieval – as the episodic buffer (Baddeley, 2000). According to this hypothesis, the retrieval of greater amounts of integrated multi-modal information causes subjects to have confidence in their retrieval judgments and increases the maintenance demands on the episodic buffer. This hypothesis can explain many of the findings in the visual WM literature, such as increased BOLD activations with recollected rather than familiar memories (Wheeler and Buckner, 2004; Yonelinas, Otten, Shaw, and Rugg, 2005; Daselaar, Fleck, and Cabeza, 2006; reviewed in Skinner and Fernandes, 2007), and decreased WM capacity after PPC damage (Berryhill and Olson, 2008a; b). However, it fails to explain why PPC damage leads to different levels of memory performance depending on the probe task (Cabeza et al., 2008).

A second proposal suggests that portions of the PPC assess the strength of a memory signal in order to determine whether a stimulus was previously viewed. This mnemonic accumulator hypothesis (Wagner et al., 2005) can be easily understood as an analogy to signal detection theory in which evidence is accrued and measured against some criterion level. This view is unable to explain why different memory tasks produce different response patterns in the parietal cortex, as some sort of signal estimate should be resistant to the type of information retrieved (reviewed in Cabeza, 2008).

A third possibility is that the PPC's role in memory retrieval parallels its role in attention (Wagner et al., 2005; Cabeza, 2008; Cabeza et al., 2008). To understand how this may work it is important to understand that attention and memory are tightly intertwined at nearly every stage of mnemonic processing (reviewed in Naghavi and Nyberg, 2005; Awh, Vogel, and Oh, 2006; Chun and Turk-Browne, 2007). They are linked at encoding because top-down attention to any given item improves the perception of features, such as contrast (Carrasco, Ling, and Read, 2004). Selective attention acts as a gatekeeper, filtering which aspects of the environment are encoded into memory, and which items are not. Attention and memory are also intertwined during memory maintenance as demonstrated by the fact that distraction during delay periods disrupts memory performance. Last, attention may be needed for accurate retrieval because divided attention tasks can cause interference and failures of retrieval even for items that were encoded with full attention (Fernandes and Moscovitch, 2000). As originally conceived, the attention to internal representation hypothesis suggested that the parietal lobe focused attention on items held in memory (Wagner et al., 2005). This hypothesis has been expanded from a purely goal driven, or top-down, attentional modulation to include spontaneous, or bottom-up, attentional modulation, which maps onto dorsal and ventral parietal regions (Cabeza, 2008).

The most recent version of this expanded view, the *attention to memory model* (AtoM) suggests that memory goals can be allocated deliberately (top-down) or can be captured by a memory cue (bottom-up) (see Figure 3). Explicit directions provide a goal to guide internal memory retrieval towards a particular memory. This process is sustained by a region in the BA 7 of the superior parietal lobe. In contrast, a memory may be spontaneously evoked by an environmental stimulus in a bottom-up fashion. When this happens, the MTL connections with inferior parietal lobe regions cause the superior parietal region to redirect attention to the bottom-up driven recollection. This hypothesis can account for much of the reviewed episodic memory literature, including our findings in which bottom-up, spontaneous retrieval of autobiographical memories was impaired in patients with bilateral ventral parietal damage, but top-down, cued retrieval was normal (Berryhill et al., 2007).

The last hypothesis we discuss, the subjective memory hypothesis, proposes that the parietal lobe is responsible for the subjective experience of confidence and vividness in one's retrieved memories (Ally et al., 2008). This hypothesis finds support in a small number of neuroimaging findings (Chua, Schacter, Rand-Giovannetti, and Sperling, 2006; Durate, Henson, and Graham, 2008), in statements made by parietal patients that they had diminished vividness and confidence in their memory abilities (Ally et al., 2008; Davidson et al., 2008), and in neuropsychological findings showing decreased levels of memory confidence (Davidson et al., 2008; Drowos et al., 2008, accepted pending revisions; Simons et al., 2008-submitted), and memory vividness (Berryhill et al., 2007). Notably, several years ago Hunkin and colleagues described patient, DH, who had parieto-occipital damage due to a closed head injury. This patient had semantic memories for his lifetime but he felt that he lacked "genuine" memories of this life. He claimed that he did not feel that he had truly experienced these memories: "*He said it was as if he had read a book about his life but he did not confuse these facts with real memories any more than one would confuse a story in a book with real life*" (Hunkin et al., 1995).

Such reports are reminiscent of findings in the hemispatial neglect literature suggesting that portions of the parietal lobe (right inferior) play a role in the subjective states of awareness. Perhaps it is the case that the inferior parietal lobe also plays an analogous role in the subjective awareness, and hence confidence, of memories.

Conclusions

The functionality of the PPC, like that of the frontal lobe, cannot be characterized as simply sensory, motor, or cognitive. Parietal functions that were identified generations ago in lesion studies - spatial attention, visual guidance of action (eye movements, grasping), calculation, reading - do not appear to have an underlying categorical structure that provides for easy analysis. Complicating the situation, in this paper we suggest that an additional function should be added to this list: memory retrieval.

Our review indicates that (1) The PPC plays a critical role in spatial working memory; (2) The PPC may also have a role in object WM, although this role remains less substantiated; (3) The PPC's role in WM may be most strongly linked to memory retrieval (Berryhill and Olson, 2008a); (4) The PPC plays an important role in episodic memory retrieval, although encoding functions may also exist (see Uncapher & Wagner, this volume). Links between the posterior parietal lobe and memory retrieval were first noted in fMRI studies of episodic memory. More recently, lesion studies indicate that PPC damage can cause selective retrieval difficulties in the context of episodic retrieval tasks. (5) Many mechanistic accounts of the PPC's role in episodic memory have been proposed. Two accounts have emerged that find agreement with both neuroimaging and lesion data are the attention to memory (AtoM) hypothesis (Cabeza et al., 2008) and the subjective memory hypothesis (Ally et al., 2008).

The goal of this review is in part to raise interest in these remaining questions and to provoke researchers to develop clearly testable hypotheses (see Box 1). The prospect of answering these questions is certain to drive the field of memory research in the years to come.

Box 1. Questions for Future Research

1. Given that a hallmark of parietal functioning is spatial selectivity, is object WM truly independent from spatial processing functions of the PPC? According to one view, whenever we encode objects, we automatically encode and maintain stimulus location, even when location is task irrelevant (Jiang, Olson, and Chun, 2000). In support of this finding, macaque research indicates that PPC coding is largely retinotopic even in object WM tasks and that IPS neurons are more sensitive to remembered locations than remembered objects (Sabes, Breznen, and Andersen, 2002). The object attributes to which IPS neurons are most sensitive – shape and size – are inherently spatial in nature. However, the inter-species generalizability of these findings is poorly understood.
2. Are the parietal mechanisms for attention and memory encoding one and the same? According to one view, PPC activations during WM tasks simply represent attention to objects (Cowan, 2001; Majerus, Bastin, Poncelet, Van der Linden, Salmon, Collette, and Maquet, 2007). In support of this view, a recent fMRI study probed set-size effects during an object WM task and two control tasks of perceptual attention, finding similar IPS activity and limitations during all tasks (Mitchell and Cusack, in press). Because the attention task had no WM requirements, the authors suggest that the activity reflects the processing of attended items.
3. Can different mnemonic processes be linked to different parietal lobe regions? Episodic memory *retrieval* has been associated with IPL activations, while visual working memory *maintenance* has been associated with IPS activations. Expected laterality differences – left PPC for verbal material, right PPC for visual material – have seldom been reported in fMRI studies, although they are commonly reported in neuropsychological studies.
4. Our neuropsychological findings showed a dissociation between WM recall and recognition such that parietal lobe damage only adversely affected recognition. A similar dissociation has not been reported in long-term memory tasks and indeed, several studies have reported normal levels of recognition memory after parietal lobe damage (Davidson et al., 2008; Simons et al., 2008; Simons et al., 2008-submitted). Are recall and recognition differentially reliant on the parietal lobe at different time scales?

Acknowledgments

We would like to thank Roberto Cabeza for helpful discussions. We also thank Alexandra Giardino. This research was supported by RO1 MH071615 to I. Olson and Ruth Kirschstein NRSA F32 NS059093 to M. Berryhill.

References

- Ally BA, Simons JS, McKeever JD, Peers PV, Budson AE. Parietal contributions to recollection: Electrophysiological evidence from aging and patients with parietal lesions. *Neuropsychologia* 2008;46:1800–1812. [PubMed: 18402990]
- Alvarez GA, Cavanagh P. The capacity of visual short-term memory is set by visual information load and by number of objects. *Psychological Science* 2004;15:106–111. [PubMed: 14738517]

- Awh E, Barton B, Vogel EK. Visual working memory represents a fixed number of items regardless of complexity. *Psychological Science* 2007;18:622–628. [PubMed: 17614871]
- Awh E, Vogel EK, Oh SH. Interactions between attention and working memory. *Neuroscience* 2006;139:201–208. [PubMed: 16324792]
- Baddeley A. The episodic buffer: a new component of working memory? *Trends in Cognitive Science* 2000;4:417–423.
- Baldo JV, Dronkers NF. The role of inferior parietal and inferior frontal cortex in working memory. *Neuropsychology* 2006;20:529–538. [PubMed: 16938015]
- Berryhill ME, Olson IR. Is the posterior parietal lobe involved in working memory retrieval? Evidence from patients with bilateral parietal lobe damage. *Neuropsychologia* 2008a;46:1775–1786. [PubMed: 18439630]
- Berryhill ME, Olson IR. The right parietal lobe is critical for visual working memory. *Neuropsychologia* 2008b;46:1767–1774. [PubMed: 18308348]
- Berryhill ME, Phuong L, Picasso L, Cabeza R, Olson IR. Parietal lobe and Episodic Memory: Bilateral Damage Causes Impaired Free Recall of Autobiographical Memory. *Journal of Neuroscience* 2007;14415–14423. [PubMed: 18160649]
- Buchsbaum BR, D'Esposito M. The search for the phonological store: from loop to convolution. *Journal of Cognitive Neuroscience* 2008;20:762–778. [PubMed: 18201133]
- Buschman TJ, Miller EK. Top-down versus bottom-up control of attention in the prefrontal and posterior parietal cortices. *Science* 2007;315:1860–1862. [PubMed: 17395832]
- Buxbaum LJ. On the right (and left) track: Twenty years of progress in studying hemispatial neglect. *Cognitive Neuropsychology* 2006;23:184–201.
- Cabeza R. Role of lateral posterior parietal regions in episodic memory retrieval: The dual attention hypothesis. *Neuropsychologia* 2008;46:1813–1827. [PubMed: 18439631]
- Cabeza R, Ciaramelli E, Olson IR, Moscovitch M. Parietal cortex and memory: an attentional account. *Nature Reviews Neuroscience* 2008;9:613–625.
- Carrasco M, Ling S, Read S. Attention alters appearance. *Nature Neuroscience* 2004;7:308–313.
- Cavada C, Goldman-Rakic PS. Posterior parietal cortex in rhesus monkey: I. Parcellation of areas based on distinctive limbic and sensory corticocortical connections. *Journal of Comparative Neurology* 1989;287:393–421. [PubMed: 2477405]
- Chafee MV, Goldman-Rakic P. Inactivation of parietal and prefrontal cortex reveals interdependence of neural activity during memory-guided saccades. *J Neurophysiol* 2000;83:1550–1566. [PubMed: 10712479]
- Chafee MV, Goldman-Rakic PS. Matching patterns of activity in primate prefrontal area 8a and parietal area 7ip neurons during a spatial working memory task. *Journal of Neurophysiology* 1998;79:2919–2940. [PubMed: 9636098]
- Chamod AS, Petrides M. Dissociable roles of the posterior parietal and the prefrontal cortex in manipulation and monitoring processes. *Proceeding of the National Academy of Sciences U S A* 2007;104:14837–14842.
- Chua EF, Schacter DL, Rand-Giovannetti E, Sperling RA. Understanding metamemory: Neural correlates of the cognitive process and subjective level of confidence in recognition memory. *Neuroimage* 2006;29:1150–1160. [PubMed: 16303318]
- Chun MM, Turk-Browne NB. Interactions between attention and memory. *Current Opinion in Neurobiology* 2007;17:177–184. [PubMed: 17379501]
- Ciaramelli E, Grady CL, Moscovitch M. Top-down and bottom-up attention to memory: A hypothesis (AtoM) on the role of the posterior parietal cortex in memory retrieval. *Neuropsychologia* 2008;46:1828–1851. [PubMed: 18471837]
- Clower DM, West RA, Lynch JC, Strick PL. The inferior parietal lobule is the target of output from the superior colliculus, hippocampus, and cerebellum. *Journal of Neuroscience* 2001;21:6283–6291. [PubMed: 11487651]
- Constantinidis C. Posterior parietal mechanisms of visual attention. *Reviews in Neuroscience* 2006;17:415–427.

- Constantinidis C, Procyk E. The primate working memory networks. *Cognitive, Affective, and Behavioral Neuroscience* 2004;4:444–465.
- Constantinidis C, Steinmetz MA. Neuronal activity in posterior parietal area 7a during the delay periods of a spatial memory task. *Journal of Neurophysiology* 1996;76:1352–1355. [PubMed: 8871242]
- Constantinidis C, Steinmetz MA. Posterior parietal cortex automatically encodes the location of salient stimuli. *Journal of Neuroscience* 2005;25:233–238. [PubMed: 15634786]
- Constantinidis C, Wang XJ. A neural circuit basis for spatial working memory. *Neuroscientist* 2004;10:553–565. [PubMed: 15534040]
- Cowan N. The magical number 4 in short-term memory: A reconsideration of mental storage capacity. *Behavioral & Brain Sciences* 2001;24:87–114. [PubMed: 11515286]
- Critchley, M. *The Parietal Lobes*. London: Edward Arnold; 1953.
- Culham JC, Kanwisher NG. Neuroimaging of cognitive functions in human parietal cortex. *Current Opinion in Neurobiology* 2001;11:157–163. [PubMed: 11301234]
- Culham JC, Valyear KF. Human parietal cortex in action. *Current Opinion in Neurobiology* 2006;16:205–212. [PubMed: 16563735]
- Curtis CE. Prefrontal and parietal contributions to spatial working memory. *Neuroscience* 2006;139:173–180. [PubMed: 16326021]
- D'Esposito M. From cognitive to neural models of working memory. *Philosophical Transactions of the Royal Society of London B* 2007;362:761–772.
- D'Esposito M, Aguirre GK, Zarahn E, Ballard D, Shin RK, Lease J. Functional MRI studies of spatial and nonspatial working memory. *Brain Research, Cognitive Brain Research* 1998;7:1–13.
- Daselaar SM, Fleck MS, Cabeza R. Triple dissociation in the medial temporal lobes: recollection, familiarity, and novelty. *Journal of Neurophysiology* 2006;96:1902–1911. [PubMed: 16738210]
- Davidson PS, Anaki D, Ciaramelli E, Cohn M, Kim AS, Murphy KJ, Troyer AK, Moscovitch M, Levine B. Does lateral parietal cortex support episodic memory? Evidence from focal lesion patients. *Neuropsychologia* 2008;46:1743–1755. [PubMed: 18313699]
- De Renzi E, Faglioni P, Previdi P. Spatial memory and hemispheric locus of lesion. *Cortex* 1977;13:424–433. [PubMed: 604009]
- De Renzi E, Nichelli P. Verbal and non-verbal short-term memory impairment following hemispheric damage. *Cortex* 1975;11:341–354. [PubMed: 1222578]
- Dehaene S, Molko N, Cohen L, Wilson AJ. Arithmetic and the brain. *Current Opinions in Neurobiology* 2004;14:218–224.
- Dehaene S, Spelke E, Pinel P, Stanescu R, Tsivkin S. Sources of mathematical thinking: behavioral and brain-imaging evidence. *Science* 1999;284:970–974. [PubMed: 10320379]
- Della Sala S, Gray C, Baddeley A, Allamano N, Wilson L. Pattern span: a tool for unwinding visuo-spatial memory. *Neuropsychologia* 1999;37:1189–1199. [PubMed: 10509840]
- Ding SL, Van Hoesen G, Rockland KS. Inferior parietal lobule projections to the presubiculum and neighboring ventromedial temporal cortical areas. *Journal of Comparative Neurology* 2000;425:510–530. [PubMed: 10975877]
- Driver J, Vuilleumier P. Perceptual awareness and its loss in unilateral neglect and extinction. *Cognition* 2001;79:39–88. [PubMed: 11164023]
- Drowos D, Berryhill ME, Olson IR. Memory retrieval deficits and the parietal lobe. *Neuropsychology*. 2008 accepted pending revisions.
- Durate A, Henson RN, Graham KS. The effects of aging on the neural correlates of subjective and objective recollection. *Cerebral Cortex* 2008;18:2169–2180. [PubMed: 18165281]
- Ellis AX, Della Sala S, Logie RH. The bailiwick of visuo-spatial working memory: evidence from unilateral spatial neglect. *Brain Research Cognitive Brain Research* 1996;3:71–78. [PubMed: 8713547]
- Ferber S, Danckert J. Lost in space--the fate of memory representations for non-neglected stimuli. *Neuropsychologia* 2006;44:320–325. [PubMed: 15946711]
- Fernandes MA, Moscovitch M. Divided attention and memory: evidence of substantial interference effects at retrieval and encoding. *Journal of Experimental Psychology, General* 2000;129:155–176. [PubMed: 10868332]

- Finke K, Bublak P, Zihl J. Visual spatial and visual pattern working memory: neuropsychological evidence for a differential role of left and right dorsal visual brain. *Neuropsychologia* 2006;44:649–661. [PubMed: 16111725]
- Fuster JM. Behavioral electrophysiology of the prefrontal cortex. *Trends Neurosci* 1984;7:408–414.
- Fuster JM. Inferotemporal units in selective visual attention and short-term memory. *Journal of Neurophysiology* 1990;64:681–697. [PubMed: 2230917]
- Fuster JM, Alexander GE. Delayed response deficit by cryogenic depression of frontal cortex. *Brain Research, Cognitive Brain Research* 1970;20:85–90.
- Glover S. Separate visual representations in the planning and control of action. *Behavioral Brain Sciences* 2004;27:3–24. discussion 24–78.
- Gnadt JW, Andersen RA. Memory related motor planning activity in posterior parietal cortex of macaque. *Experimental Brain Research* 1988;70:216–220.
- Greicius MD, Menon V. Default-mode activity during a passive sensory task: uncoupled from deactivation but impacting activation. *Journal of Cognitive Neuroscience* 2004;16:1484–1492. [PubMed: 15601513]
- Halligan PW, Fink GR, Marshall JC, Vallar G. Spatial cognition: evidence from visual neglect. *Trends in Cognitive Science* 2003;7:125–133.
- Hanley JR, Young AW, Pearson NA. Impairment of the visuo-spatial sketch pad. *Quarterly Journal of Experimental Psychology A* 1991;43:101–125.
- Hartley AA, Speer NK. Locating and fractionating working memory using functional neuroimaging: storage, maintenance, and executive functions. *Microsc Res Tech* 2000;51:45–53. [PubMed: 11002352]
- Heilman KM, Jeong Y, Finney GR. Spatial cognition. *Neurology* 2004;63:1994–1996. [PubMed: 15596737]
- Heilman KM, Valenstein E, Watson RT. Neglect and related disorders. *Seminars in Neurology* 2000;20:463–470. [PubMed: 11149702]
- Hillis AE. Neurobiology of unilateral spatial neglect. *Neuroscientist* 2006;12:153–163. [PubMed: 16514012]
- Hunkin NM, Parkin AJ, Bradley VA, Burrows EH, Aldrich FK, Jansari A, Burdon-Cooper C. Focal retrograde amnesia following closed head injury: a case study and theoretical account. *Neuropsychologia* 1995;33:509–523. [PubMed: 7617158]
- Husain M, Mannan S, Hodgson T, Wojciulik E, Driver J, Kennard C. Impaired spatial working memory across saccades contributes to abnormal search in parietal neglect. *Brain* 2001;124:941–952. [PubMed: 11335696]
- Insausti R, Amaral DG. Entorhinal cortex of the monkey: IV. Topographical and laminar organization of cortical afferents. *Journal of Comparative Neurology* 2008;509:608–641. [PubMed: 18551518]
- Jiang Y, Olson IR, Chun MM. Organization of visual short-term memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition* 2000;26:683–702.
- Jonides J, Smith EE, Koeppe RA, Awh E, Minoshima S, Mintun MA. Spatial working memory in humans as revealed by PET. *Nature* 1993;363:623–625. [PubMed: 8510752]
- Kahn I, Davachi L, Wagner AD. Functional-neuroanatomic correlates of recollection: implications for models of recognition memory. *Journal of Neuroscience* 2004;24:4172–4180. [PubMed: 15115812]
- Karnath HO, Ferber S, Himmelbach M. Spatial awareness is a function of the temporal not the posterior parietal lobe. *Nature* 2001;411:950–953. [PubMed: 11418859]
- Kessels RP, Jaap Kappelle L, de Haan EH, Postma A. Lateralization of spatial-memory processes: evidence on spatial span, maze learning, and memory for object locations. *Neuropsychologia* 2002;40:1465–1473. [PubMed: 11931950]
- Kessels RP, van Zandvoort MJ, Postma A, Kappelle LJ, de Haan EH. The Corsi Block-Tapping Task: standardization and normative data. *Appl Neuropsychol* 2000;7:252–258. [PubMed: 11296689]
- Kim H, Cabeza R. Differential contributions of prefrontal, medial temporal, and sensory-perceptual regions to true and false memory formation. *Cerebral Cortex* 2007;17:2143–2150. [PubMed: 17110592]

- Koch G, Oliveri M, Torriero S, Carlesimo GA, Turriziani P, Caltagirone C. rTMS evidence of different delay and decision processes in a fronto-parietal neuronal network activated during spatial working memory. *Neuroimage* 2005;24:34–39. [PubMed: 15588594]
- Linden DE. The working memory networks of the human brain. *Neuroscientist* 2007;13:257–267. [PubMed: 17519368]
- Luck SJ, Vogel EK. The capacity of visual working memory for features and conjunctions. *Nature* 1997;390:279–281. [PubMed: 9384378]
- Macoveanu J, Klingberg T, Tegner J. A biophysical model of multiple-item working memory: a computational and neuroimaging study. *Neuroscience* 2006;141:1611–1618. [PubMed: 16777342]
- Majerus S, Bastin C, Poncelet M, Van der Linden M, Salmon E, Collette F, Maquet P. Short-term memory and the left intraparietal sulcus: focus of attention? Further evidence from a face short-term memory paradigm. *Neuroimage* 2007;35:353–367. [PubMed: 17240164]
- Makris N, Kennedy DN, McInerney S, Sorensen AG, Wang R, Caviness VS Jr, Pandya DN. Segmentation of subcomponents within the superior longitudinal fascicle in humans: a quantitative, in vivo, DT-MRI study. *Cerebral Cortex* 2005;15:854–869. [PubMed: 15590909]
- Malhotra P, Jager HR, Parton a, Greenwood R, Playford ED, Brown MM, Driver J, Husain M. Spatial working memory capacity in unilateral neglect. *Brain* 2005;128:424–435. [PubMed: 15644422]
- Malhotra P, Mannan S, Driver J, Husain M. Impaired spatial working memory: one component of the visual neglect syndrome? *Cortex* 2004;40:667–676. [PubMed: 15505977]
- Mannan SK, Mort DJ, Hodgson TL, Driver J, Kennard C, Husain M. Revisiting previously searched locations in visual neglect: role of right parietal and frontal lesions in misjudging old locations as new. *Journal of Cognitive Neuroscience* 2005;17:340–354. [PubMed: 15811244]
- Markowitsch HJ, Kalbe E, Kessler J, von Stockhausen HM, Ghaemi M, Heiss WD. Short-term memory deficit after focal parietal damage. *Journal of Clinical and Experimental Neuropsychology* 1999;21:784–797. [PubMed: 10649534]
- Marshuetz C, Smith EE. Working memory for order information: multiple cognitive and neural mechanisms. *Neuroscience* 2006;139:195–200. [PubMed: 16359810]
- Marshuetz C, Smith EE, Jonides J, DeGutis J, Chenevert TL. Order information in working memory: fMRI evidence for parietal and prefrontal mechanisms. *Journal of Cognitive Neuroscience* 2000;12 (Suppl 2):130–144. [PubMed: 11506653]
- Mazzoni P, Bracewell RM, Barash S, Andersen RA. Motor intention activity in the macaque's lateral intraparietal area. I. Dissociation of motor plan from sensory memory. *Journal of Neurophysiology* 1996;76:1439–1456. [PubMed: 8890265]
- Mitchell DJ, Cusack R. Flexible, capacity-limited activity of posterior parietal cortex in perceptual as well as visual short-term memory tasks. *Cerebral Cortex* 18:1788–1798. in press.
- Mori, S.; Wakana, S.; Nague-Poetscher, LM.; van Zijl, PCM. Atlas of human white matter. Amsterdam: Elsevier; 2005.
- Naghavi HR, Nyberg L. Common fronto-parietal activity in attention, memory, and consciousness: shared demands on integration? *Consciousness and Cognition* 2005;14:390–425. [PubMed: 15950889]
- Niki H, Watanabe M. Prefrontal unit activity and delayed response: relation to cue location versus direction of response. *Brain Research, Cognitive Brain Research* 1976;105:79–88.
- Nys GM, van Zandvoort MJ, van der Worp HB, Kappelle LJ, de Haan EH. Neuropsychological and neuroanatomical correlates of perseverative responses in subacute stroke. *Brain* 2006;129:2148–2157. [PubMed: 16870885]
- Oliveri M, Turriziani P, Carlesimo GA, Koch G, Tomaiuolo F, Panella M, Caltagirone C. Parieto-frontal interactions in visual-object and visual-spatial working memory: evidence from transcranial magnetic stimulation. *Cerebral Cortex* 2001;11:606–618. [PubMed: 11415963]
- Olson IR, Jiang Y. Is visual short-term memory object based? *Perception & Psychophysics* 2002;64:1055–1067. [PubMed: 12489661]
- Parton A, Malhotra P, Nachev P, Ames D, Ball J, Chataway J, Husain M. Space re-exploration in hemispatial neglect. *Neuroreport* 2006;17:833–836. [PubMed: 16708024]
- Passingham D, Sakai K. The prefrontal cortex and working memory: physiology and brain imaging. *Current Opinion in Neurobiology* 2004;14:163–168. [PubMed: 15082320]

- Petrides M, Pandya DN. Efferent association pathways originating in the caudal prefrontal cortex in the macaque monkey. *Journal of Comparative Neurology* 2006;498:227–251. [PubMed: 16856142]
- Pisella L, Berberovic N, Mattingley JB. Impaired working memory for location but not for colour or shape in visual neglect: a comparison of parietal and non-parietal lesions. *Cortex* 2004;40:379–390. [PubMed: 15156795]
- Pisella L, Mattingley JB. The contribution of spatial remapping impairments to unilateral visual neglect. *Neuroscience and Biobehavioral Reviews* 2004;28:181–200. [PubMed: 15172763]
- Postle BR, Ferrarelli F, Hamidi M, Ferdoes E, Massimini M, Peterson M, Alexander A, Tononi G. Repetitive transcranial magnetic stimulation dissociates working memory manipulation from retention functions in the prefrontal, but not posterior parietal, cortex. *Journal of Cognitive Neuroscience* 2006;18:1712–1722. [PubMed: 17014375]
- Postma A, Sterken Y, de Vries L, de Haan EH. Spatial localization in patients with unilateral posterior left or right hemisphere lesions. *Experimental Brain Research* 2000;134:220–227.
- Quintana J, Fuster JM. Spatial and temporal factors in the role of prefrontal and parietal cortex in visuomotor integration. *Cerebral Cortex* 1993;3:122–132. [PubMed: 8490318]
- Quintana J, Fuster JM. From perception to action: temporal integrative functions of prefrontal and parietal neurons. *Cerebral Cortex* 1999;9:213–221. [PubMed: 10355901]
- Rockland KS, Van Hoesen GW. Some temporal and parietal cortical connections converge in CA1 of the primate hippocampus. *Cerebral Cortex* 1999;9:232–237. [PubMed: 10355903]
- Rushworth MF, Behrens TE, Johansen-Berg H. Connection patterns distinguish 3 regions of human parietal cortex. *Cerebral Cortex* 2006;16:1418–1430. [PubMed: 16306320]
- Sabes PN, Breznen B, Andersen RA. Parietal representation of object-based saccades. *Journal of Neurophysiology* 2002;88:1815–1829. [PubMed: 12364508]
- Schmahmann, JD.; Pandya, DN. *Fiber pathways of the brain*. Oxford: Oxford University Press; 2006.
- Schmahmann JD, Pandya DN, Wang R, Dai G, D'Arceuil HE, de Crespigny AJ, Wedeen VJ. Association fibre pathways of the brain: parallel observations from diffusion spectrum imaging and autoradiography. *Brain* 2007;130:630–653. [PubMed: 17293361]
- Scoville WB, Milner B. Loss of recent memory after bilateral hippocampal lesions. *Journal of Neurosurgery and Psychiatry* 1957;20:11–21.
- Seltzer B, Pandya DN. Further observations on parieto-temporal connections in the rhesus monkey. *Experimental Brain Research* 1984;55:301–312.
- Sereno AB, Amador SC. Attention and memory-related responses of neurons in the lateral intraparietal area during spatial and shape-delayed match-to-sample tasks. *Journal of Neurophysiology* 2006;95:1078–1098. [PubMed: 16221750]
- Sereno AB, Maunsell JH. Shape selectivity in primate lateral intraparietal cortex. *Nature* 1998;395:500–503. [PubMed: 9774105]
- Shimozaki S, Kingstone A, Olk B, Stowe R, Eckstein M. Classification images of two right hemisphere patients: a window into the attentional mechanisms of spatial neglect. *Brain Research, Cognitive Brain Research* 2006;1080:26–52.
- Simons JS, Mayes AR. What is the parietal lobe contribution to human memory? *Neuropsychologia* 2008;46:1739–1742. [PubMed: 18501933]
- Simons JS, Peers PV, Hwang DY, Ally BA, Fletcher PC, Budson AE. Is the parietal lobe necessary for recollection in humans? *Neuropsychologia* 2008;46:1185–1191. [PubMed: 17850832]
- Simons JS, Peers PV, Mazuz Y, Berryhill ME, Olson IR. Dissociation between memory accuracy and memory confidence following bilateral parietal lesions. 2008 submitted.
- Skinner EI, Fernandes MA. Neural correlates of recollection and familiarity: A review of neuroimaging and patient data. *Neuropsychologia* 2007;45:2163–2179. [PubMed: 17445844]
- Smith EE, Jonides J. Working memory: a view from neuroimaging. *Cognitive Psychology* 1997;33:5–42. [PubMed: 9212720]
- Smith EE, Jonides J. Neuroimaging analyses of human working memory. *Proceedings of the National Academy of Sciences U S A* 1998;95:12061–12068.
- Takahashi E, Ohki K, Kim DS. Dissociated Pathways for Successful Memory Retrieval from the Human Parietal Cortex: Anatomical and Functional Connectivity Analyses. *Cerebral Cortex*. 2008

- Todd JJ, Marois R. Capacity limit of visual short-term memory in human posterior parietal cortex. *Nature* 2004;428:751–754. [PubMed: 15085133]
- Todd JJ, Marois R. Posterior parietal cortex activity predicts individual differences in visual short-term memory capacity. *Cognitive, Affective, and Behavioral Neuroscience* 2005;5:144–155.
- Toth LJ, Assad JA. Dynamic coding of behaviourally relevant stimuli in parietal cortex. *Nature* 2002;415:165–168. [PubMed: 11805833]
- Ungerleider LG, Courtney SM, Haxby JV. A neural system for human visual working memory. *Proceeding of the National Academy of Sciences U S A* 1998;95:883–890.
- Vallar G, Perani D. The anatomy of unilateral neglect after right-hemisphere stroke lesions. A clinical/CT-scan correlation study in man. *Neuropsychologia* 1986;24:609–622. [PubMed: 3785649]
- van Asselen M, Kessels RP, Neggers SF, Kappelle LJ, Frijns CJ, Postma A. Brain areas involved in spatial working memory. *Neuropsychologia* 2006;44:1185–1194. [PubMed: 16300806]
- Van Essen DC, Lewis JW, Drury HA, Hadjikhani N, Tootell RB, Bakircioglu M, Miller MI. Mapping visual cortex in monkeys and humans using surface-based atlases. *Vision Res* 2001;41:1359–1378. [PubMed: 11322980]
- Vilberg KL, Moosavi RF, Rugg MD. The relationship between electrophysiological correlates of recollection and amount of information retrieved. *Brain Research, Cognitive Brain Research* 2006;1122:161–170.
- Vilberg KL, Rugg MD. Memory retrieval and the parietal cortex: a review of evidence from event-related fMRI. *Neuropsychologia* 2008;46:1787–1799. [PubMed: 18343462]
- Vincent JL, Snyder AZ, Fox MD, Shannon BJ, Andrews JR, Raichle ME, Buckner RL. Coherent spontaneous activity identifies a hippocampal-parietal memory network. *Journal of Neurophysiology* 2006;96:3517–3531. [PubMed: 16899645]
- von Economo, C.; Koskinas, GN. *Die Cytoarchitektonik der Hirnrinde des erwachsenen Menschen*. Berlin: Verlag von Julius Springer; 1925.
- Vuilleumier P, Sergent C, Schwartz S, Valenza N, Girardi M, Husain M, Driver J. Impaired perceptual memory of locations across gaze-shifts in patients with unilateral spatial neglect. *Journal of Cognitive Neuroscience* 2007;19:1388–1406. [PubMed: 17651010]
- Wager TD, Smith EE. Neuroimaging studies of working memory: a meta-analysis. *Cognitive, Affective, and Behavioral Neuroscience* 2003;3:255–274.
- Wagner AD, Shannon BJ, Kahn I, Buckner RL. Parietal lobe contributions to episodic memory retrieval. *Trends in Cognitive Science* 2005;9:445–453.
- Warrington EK, James M. An experimental investigation of facial recognition in patients with unilateral cerebral lesions. *Cortex* 1967;3:317–326.
- Wellman BJ, Rockland KS. Divergent cortical connections to entorhinal cortex from area TF in the macaque. *Journal of Comparative Neurology* 1997;389:361–376. [PubMed: 9414000]
- Wendelken C, Bunge SA, Carter CS. Maintaining structured information: an investigation into functions of parietal and lateral prefrontal cortices. *Neuropsychologia* 2008;46:665–678. [PubMed: 18022652]
- Wheeler ME, Buckner RL. Functional dissociation among components of remembering: control, perceived oldness, and content. *Journal of Neuroscience* 2003;23:3869–3880. [PubMed: 12736357]
- Wheeler ME, Buckner RL. Functional-anatomic correlates of remembering and knowing. *Neuroimage* 2004;21:1337–1349. [PubMed: 15050559]
- Wojciulik E, Husain M, Clarke K, Driver J. Spatial working memory deficit in unilateral neglect. *Neuropsychologia* 2001;39:390–396. [PubMed: 11164877]
- Xu Y. The role of the superior intraparietal sulcus in supporting visual short-term memory for multifeature objects. *Journal of Neuroscience* 2007;27:11676–11686. [PubMed: 17959810]
- Xu Y, Chun MM. Dissociable neural mechanisms supporting visual short-term memory for objects. *Nature* 2006;440:91–95. [PubMed: 16382240]
- Yonelinas AP, Otten LJ, Shaw KN, Rugg MD. Separating the brain regions involved in recollection and familiarity in recognition memory. *Journal of Neuroscience* 2005;25:3002–3008. [PubMed: 15772360]

Parietal-Medial Temporal Lobe Connections

Parietal-Frontal Lobe Connections

Parietal-Medial Temporal, Frontal Lobe Connections

Figure 1. Parietal-Medial Temporal Lobe Connections. **A.** Hippocampal area CA1 (black dots in left-most panel) projects to area 7a (demarked with an oval in the right-most panel) in the lateral intraparietal area (LIP), as shown by retrograde viral tracing in the cebus monkey (Clower et al., 2001). The middle figures provide anatomical orientation for monkey area 7a. **B.** Parietal-Frontal Lobe Connections. The three branches of the macaque superior longitudinal fasciculus (SLF I–III) are shown from two perspectives: on the left, the lateral view and on the right the coronal view demonstrating the medial connections and the lateral connections (Schmahmann et al., 2007). **C.** Human Tractography. The left panel shows the results of a human diffusion weighted tractography study in which broad connections between inferior parietal regions and the whole of the parahippocampal gyrus (left panel), and connections between superior parietal

and frontal areas (right panel) were found (Rushworth et al., 2006). Figures taken with kind permission from the publishers.

Figure 2.

The effects of parietal lobe lesions on working memory and episodic memory. **A.** Working memory and episodic memory was investigated in two patients with bilateral parietal lobe lesions. MRI images show the lesions outlined in red. **B.** Three experiments spanning short and long-term forms of memory probed memory with two different response tasks. In order and object WM (left), recall (Rec) performance was preserved whereas recognition (Recog) performance was impaired (Berryhill and Olson, 2008a). In a long-term memory (LTM) task, performance was assessed with the DRM paradigm, and the same pattern of preserved recall and impaired recognition performance was observed (Drowos et al., 2008, accepted pending revisions). Data are shown as standardized scores in which performance greater than 1.96 standard deviations is considered impaired.

Attention to Memory (AtoM) Model

Figure 3.

The Attention to Memory (AtoM) Model proposes two loci of parietal involvement in memory retrieval that map onto two forms of attentional demands (Cabeza et al., 2008). Top-down attention is controlled and sustained by neurons in the superior, or dorsal parietal cortex (DPC). The DPC directs attention towards a retrieval goal, in this case, the memory of a 1st birthday party. In contrast, bottom-up attention can be captured by environmental stimuli or by sensory cues, such as the image of the birthday cake. In this case, the medial temporal lobes (MTL) cause the inferior, or ventral parietal cortex (VPC) to signal the DPC to attend to the spontaneously retrieved memory. The waveforms reflect the variability in retrieval demands.

Table 1

Neuropsychological studies of spatial working memory as assessed by recall or recognition in patients with putative parietal lobe damage. Note that in many studies precise cerebral localization was not provided. Highlighted text in the Lesion Site column indicates which patient population, if any, was impaired. Abbreviations: Bi = bilateral, Corsi = Corsi block-tapping task, hemi = hemisphere, inf = inferior, L = left, R = Right, PPC = posterior parietal cortex, PFC = prefrontal cortex, 2AFC = two-alternative forced-choice.

Authors	Task	Lesion Sites	Spared	Impaired
(De Renzi and Nichelli, 1975)	Recall: Corsi	L hemi, R hemi		√
(De Renzi et al., 1977)	Recall: Corsi	R hemi, R hemi with visual field defect		√
(Hanley et al., 1991)	Recall: Corsi	R hemi , (case study)		√
(Markowitsch, Kalbe, Kessler, von Stockhausen, Ghaemi, and Heiss, 1999)	Recall: Corsi	L angular gyrus, Case	√	
(Della Sala, Gray, Baddeley, Allamano, and Wilson, 1999)	Recall: Pattern Span, Corsi	R hemi, L hemi, Bi hemi	√	
(Kessels et al., 2000)	Recall: Corsi	R hemi , L hemi, Bi hemi, subcortical		√
(Postma, Sterken, de Vries, and de Haan, 2000)	Recall: Perceptual Localization (2AFC)	R hemi, L hemi		√
(Kessels, Jaap Kappelle, de Haan, and Postma, 2002)	Recall: Object- Location Conjunction, Locations, Corsi, Maze learning	L hemi, R hemi , Bi hemi, Anterior (frontal, temporal), Posterior (occipital, parietal)	√* Corsi block	√*conjunc tion task
(Pisella et al., 2004)	Old/new recognition	R PPC + neglect , non- PPC + neglect		√
(Malhotra et al., 2005)	Recall Old/new Recognition: Vertical Corsi	R PPC + neglect ; R PPC non-neglect		√
(van Asselen, Kessels, Neggers, Kappelle, Frijns, and Postma, 2006)	Recall: Corsi	R PFC, L PFC, R PPC, L PPC	√	
(Nys, van Zandvoort, van der Worp, Kappelle, and de Haan, 2006)	Recall: Corsi	R hemi + neglect, L hemi + neglect, Bi hemi + neglect ,		√
(Baldo and Dronkers, 2006)	Recall: Corsi	L inf PPC , L inf frontal		√
(Berryhill and Olson, 2008b)	Old/new recognition: Locations	R PPC		√
(Ferber and Danckert, 2006)	Old/new recognition: Vertical Locations	R PPC + neglect , R PPC		√
(Vuilleumier, Sergent, Schwartz, Valenza, Girardi, Husain, and Driver, 2007)	Old/new recognition	R PPC + neglect		√
(Husain et al., 2001)	Visual search revisiting	R PPC + neglect , R PPC		√
(Parton et al., 2006)	Visual search revisiting	R PPC + neglect , R PPC		√
(Mannan, Mort, Hodgson, Driver, Kennard, and Husain, 2005)	Visual search revisiting	R PPC (by lesion location) + neglect		√