Published in final edited form as:

Inorganica Chim Acta. 2000 August ; 306(1): 30–37. doi:10.1016/S0020-1693(00)00144-4.

Synthesis and characterization of complexes of the {ReO}³⁺ core with SNS and S donor ligands

Frank J. Femia^a, Xiaoyuan Chen^a, Kevin P. Maresca^b, Timothy M. Shoup^b, John W. Babich^b, and Jon Zubieta^{a,*}

^a Department of Chemistry, Syracuse University, Syracuse, NY 13244, USA

^b Biostream Inc., 160 Second Street, Cambridge, MA 02142, USA

Abstract

The reaction of [ReOCl₃(PPh₃)₂] with *N,N*-bis(2-mercaptoethyl)benzylamine and 4-bromobenzenethiol allowed for the isolation of [ReO{η³-(SCH₂CH₂)₂N(CH₂C₆H₅)}-(η¹-C₆H₄Br-4-S)] (1). The reaction of [ReOCl₃(PPh₃)₂] with [(HSCH₂CH₂)₂N(CH₂C₅H₄N)] and the appropriate thiol in chloroform treated with triethylamine has led to the isolation of a series of neutral rhenium complexes of the type [ReO{η³-(SCH₂CH₂)₂N(CH₂C₅H₄N)}(η¹-C₆H₄X-4-S)] (X = Br (2), Cl (3), F (4), and OCH₃ (5)) and [ReO{η³-(SCH₂CH₂)₂N(CH₂C₅H₄N)}(η¹-C₆H₄OCH₃-4-CH₂S)] (6). Likewise, under similar reaction conditions, the use of the related tridentate ligand, [(HSCH₂CH₂)₂N(CH₂CH₂C₅H₄N)], has led to the isolation of a series of rhenium complexes of the type [ReO{η³-(SCH₂CH₂)₂N(CH₂CH₂C₅H₄N)}(η¹-C₆H₄X-4-S)] (X=Br (7), Cl (8), OCH₃ (9)), as well as [ReO{η³-(SCH₂CH₂)₂N(CH₂CH₂C₅H₄N)}(η¹-C₆H₄Cl-4-CH₂S)]·0.5CH₃(CH₂)₄CH₃ (10). These compounds are extensions of the ‘3+1’ approach to the synthesis of materials with the {MO}³⁺ core (M=Tc and Re), which have applications in nuclear medicine. The ligands chosen allow systematic exploration of the consequences of *para*-substitution on the monodentate thiolate ligand [S] and of derivatization of the substituent R on the tridentate aminodithiol ligand [SNS] of the type (HSCH₂CH₂)₂NR. Such modifications can influence lipophilicity, charge, size and molecular weight of the complex and consequently the biodistribution.

Keywords

Crystal structure; Rhenium complexes; Oxo complexes; SNS-donor ligand complexes; S-donor ligand complexes

1. Introduction

The significant contemporary interest in the coordination chemistry of rhenium derives from its analogies to technetium chemistry, as well as its expanding role in therapeutic nuclear chemistry [1]. The radionuclides of Re, the Group 7 congener of technetium, are β-emitters with properties which make them suitable candidates for therapeutic applications.

*Corresponding author. Tel.: 31-315-443 2547; fax: 31-315-443 4070. jazubiet@syr.edu (J. Zubieta).

5. Supplementary material

All atomic and thermal parameters and all interatomic angles are available from the authors upon request. Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as publication Nos. CCDC 139416–139424. Copies of the data can be obtained free of charge on application to The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: +44-1223-336033; deposit@ccdc.cam.ac.uk or www: http://www.ccdc.cam.ac.uk).

Rhenium-186 is particularly attractive [2–4] because of its half-life (90 h) and strong β -emission ($\beta_{\max}$, 1070 keV), capable of delivering high radiation doses to tissues. Rhenium-188, also a β -emitting radionuclide ($\beta_{\max}$, 2120 keV), with a half-life of 17 h and available from a $^{188}\text{W}/^{188}\text{Re}$ radionuclide generator at no carrier added levels [5], has become a logical choice for the development of radio-pharmaceuticals for use in the treatment of cancer [1].

The synthesis of novel compounds, which allow the systematic exploration of size, shape, molecular weight, and charge of the metal–thiolate complexes with respect to their biodistribution patterns, provides a driving force for advances in the radiopharmaceutical chemistry of rhenium. One approach to the systematic design of Re(V) complexes with the $\{\text{ReO}\}^{3+}$ core exploits the ‘3+1’ concept of ligand addition to a Group 7 metal-oxo species [6–16]. The methodology is based on forming a tridentate ‘3’ and a monodentate ‘1’ mixed-thiolate core. The technique uses tridentate dithiolates with the $(\text{S-X-S})^{2-}$ ($\text{X}=\text{NR}$, O , S) donor set and a -2 charge in combination with a monothiol ligand of -1 charge disposed about a Group 7 metal-oxo core, so as to occupy four coordination sites on the metal with a total formal charge of -3 . The ‘3+1’ ligand combination preserves the metal-oxo core and the formal $+5$ oxidation state of the metal, thus conforming to the metal coordination and oxidation preferences.

Over the past decade, thiol based tridentate ligands have received significant attention in the design of new Group 7 metal-oxo complexes [8–11,17–31]. One such series of compounds belongs to the class of $[\text{SSS}]/[\text{S}]$ ‘3+1’ mixed ligand complexes [10,11,17–21]. Numerous groups have synthesized neutral rhenium and technetium compounds containing 2-mercaptoethyl sulfide as the tridentate in combination with various monothiols, to produce complexes such as $[\text{ReO}\{(\text{SCH}_2\text{CH}_2)_2\text{S}\}\{\text{SCH}_2\text{C}_6\text{H}_5\}]$ [11], $[\text{ReO}\{(\text{SCH}_2\text{CH}_2)_2\text{S}\}\{\text{SCH}_2\text{CH}(\text{OH})\text{CH}(\text{OH})\text{CH}_2\text{SH}\}]$ [17], $[\text{ReO}\{(\text{SCH}_2\text{CH}_2)_2\text{S}\}\{\text{S}(\text{CH}_2)_3\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{C}_6\text{H}_5\}]$ [18] and $[\text{ReO}\{(\text{SCH}_2\text{CH}_2)_2\text{S}\}\{\text{SCH}_2\text{CH}(\text{CH}_2\text{CH}_3)\text{-4-HOC}_6\text{H}_5\}]$ [19]. Not only have such neutral compounds reported but also several cationic complexes such as $[\text{ReO}\{(\text{SCH}_2\text{CH}_2)_2\text{S}\}\{\text{C}_5\text{H}_4\text{NH-2-S}\}][\text{Br}]$ [20].

Another subclass of compounds includes the class of $[\text{SOS}]/[\text{S}]$ ‘3+1’ mixed ligand complexes [11,21,22]. In most instances of this family of compounds, the tridentate 2-mercaptoethylether seems to be the tridentate of choice, and has been used in the design of such neutral compounds as $[\text{ReO}\{(\text{SCH}_2\text{CH}_2)_2\text{O}\}\{\text{SC}_6\text{H}_5\}]$ [11] and $[\text{ReO}\{(\text{SCH}_2\text{CH}_2)_2\text{O}\}\{\text{SCH}_2\text{-4-CH}_3\text{OC}_6\text{H}_5\}]$ [22], a cationic species $[\text{ReO}\{(\text{SCH}_2\text{CH}_2)_2\text{O}\}\{\text{C}_5\text{H}_4\text{NH-2-S}\}]$ [22], and several binuclear complexes, of which $[\text{Tc}_2\text{O}_2\{(\text{SCH}_2\text{CH}_2)_2\text{O}\}_2\{\text{SCH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{S}\}]$ [21] and $[\text{Re}_2\text{O}_2\{(\text{SCH}_2\text{CH}_2)_2\text{O}\}_2\{\text{SCH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{S}\}]$ [22] are representatives.

A third series of ‘3+1’ complexes includes the $[\text{ONS}]/[\text{S}]$ mixed ligand complexes [8,9,20,23–27]. One subclass of this family incorporates the tridentate Schiff base $[\text{HOC}_6\text{H}_4\text{C}(\text{H})\text{NC}_6\text{H}_4\text{SH}]$ and includes complexes such as $[\text{TcO}\{\text{OC}_6\text{H}_4\text{C}(\text{H})\text{NC}_6\text{H}_4\text{S}\}\{\text{SC}_6\text{H}_5\}]$ [9], $[\text{TcO}\{\text{OC}_6\text{H}_4\text{C}(\text{H})\text{NC}_6\text{H}_4\text{S}\}\{n\text{-SC}_8\text{H}_{17}\}]$ [8], and a cationic species $[\text{ReO}\{\text{OC}_6\text{H}_4\text{C}(\text{H})\text{NC}_6\text{H}_4\text{S}\}\{\text{C}_5\text{H}_4\text{NH-2-S}\}][\text{Br}]$ [23]. Another tridentate ligand type found in the $[\text{ONS}]/[\text{S}]$ series is represented by $[\text{C}_5\text{H}_3\text{N-2-CH}_2\text{OH-6-CH}_2\text{SH}]$, which allows isolation of both neutral compounds, such as $[\text{ReO}\{\text{C}_5\text{H}_3\text{N-2-CH}_2\text{O-6-CH}_2\text{S}\}\{\text{S-4-ClC}_6\text{H}_4\}]$ [27], and cationic species, $[\text{ReO}\{\text{C}_5\text{H}_3\text{N-2-CH}_2\text{O-6-CH}_2\text{S}\}\{\text{C}_5\text{H}_4\text{NH-2-S}\}][\text{Br}]$ [20].

The vast majority of the ‘3+1’ Group 7 metal-oxo complexes belong to the class $[\text{SN(R)S}]/[\text{S}]$, where the $-\text{R}$ group of the tridentate may be exploited to provide a variety of derivatives and secondary functionality [19,28–31]. Examples include

[ReO{C₅H₃N(CH₂S)₂}{S-4-C₆H₄OH}] [19], [ReO{C₂H₅N(CH₂CH₂S)₂}{S-4-Cl-C₆H₄}] [28], [ReO{C₂H₅SCH₂CH₂N(CH₂CH₂S)₂}{S-4-CH₃OC₆H₄}] [29], a binuclear complex [Re₂-O₂{C₂H₅N(CH₂CH₂S)₂}₃] [28] and a cationic species, [ReO{C₅H₃N(CH₂S)₂}{C₅H₄NH-2-S}][Cl] [27].

In an effort to expand the '3+1' chemistry to the oxo-rhenium core, our work has concentrated on the reactions of the tridentate thiol, [(HSCH₂CH₂)₂N-(CH₂C₆H₅)], and the potentially tetradentate ligands-[(HSCH₂CH₂)₂N(CH₂C₅H₄N)] and [(HSCH₂CH₂)₂-N(CH₂CH₂C₅H₄N)] with [ReOCl₃(PPh₃)₂], so as to prepare families of compounds with mixed thiol ligands, and allowing the systematic exploration of the *para*-substituted benzenethiol and benzylmercaptan series. This study continues our development of the coordination chemistry of the Group 7 metal rhenium [17,20,22,23,27], in order to provide new materials for radiolabeling of chemotactic peptides [32]. We report the synthesis and structural characterization of a series of oxo-rhenium '3+1' complexes, namely [ReO{η³-(SC-H₂CH₂)₂N(CH₂C₆H₅)}(η¹-C₆H₄Br-4-S)] (1), [ReO{η³-(SCH₂CH₂)₂N(CH₂C₅H₄N)}(η¹-C₆H₄X-4-S)] (X=Br (2), Cl (3), F (4), and OCH₃ (5)), [ReO{η³-(SC-H₂CH₂)₂N(CH₂C₅H₄N)}(η¹-C₆H₄OCH₃-4-CH₂S)] (6), [ReO{η³-(SCH₂CH₂)₂N(CH₂CH₂C₅H₄N)}(η¹-C₆H₄-X-4-S)] (X=Br (7), Cl (8), OCH₃ (9)), and [ReO{η³-(SCH₂CH₂)₂N(CH₂CH₂C₅H₄N)}(η¹-C₆H₄Cl-4-CH₂S)]·0.5CH₃(CH₂)₄CH₃ (10).

2. Experimental

2.1. General considerations

NMR spectra were recorded on a Bruker DPX 300 (¹H 300.10 MHz) spectrometer in CDCl₃ (δ 7.27). IR spectra were recorded as KBr discs with a Perkin-Elmer series 1600 FT IR. Elemental analysis for carbon, hydrogen, and nitrogen were carried out by Oneida Research Services, Whitesboro, NY. Ammonium perhenate (Aldrich), triethylamine (Aldrich), triphenylphosphine (Aldrich), benzylamine (Aldrich), 2-(aminomethyl)pyridine (Aldrich), ethylene sulfide (Aldrich), 2-(2-aminoethyl)pyridine (Aldrich), 4-bromobenzenethiol (Aldrich), 4-chlorobenzenethiol (Aldrich), 4-fluorobenzenethiol (Aldrich), 4-methoxybenzenethiol (Aldrich), 4-chlorobenzyl mercaptan (Aldrich), 4-methoxybenzyl mercaptan (Lancaster) were used as received without further purification. All other solvents and reagents were purchased from Aldrich and used as received, unless otherwise stated. The [ReOCl₃(PPh₃)₂] [33] and *N,N*-bis(2-mercapto-ethyl)benzylamine [34] were synthesized according to published procedures.

2.2. Syntheses

2.2.1. [(HSCH₂CH₂)₂N(CH₂C₅H₄N)]—A similar procedure was employed as for the preparation of *N,N*-bis(2-mercaptoethyl)benzylamine. ¹H NMR (CDCl₃, ppm): 1.72 (s, 2H), 2.61 (m, 4H), 2.78 (m, 4H), 3.81 (s, 2H), 7.18 (m, 1H), 7.36, (d, 1H), 7.67 (m, 1H), 8.52 (m, 1H).

2.2.2. [(HSCH₂CH₂)₂N(CH₂CH₂C₅H₄N)]—A similar procedure was employed as for the preparation of *N,N*-bis(2-mercaptoethyl)benzylamine. ¹H NMR (CDCl₃, ppm): 1.75 (s, 2H), 2.10–2.90 (m, 12H), 7.18 (m, 1H), 7.20, (d, 1H), 7.60 (t, 1H), 8.50 (d, 1H).

2.2.3. General synthetic procedure—To a solution of [ReOCl₃(PPh₃)₂] (0.0822 g, 0.0986 mmol) in 1 ml of chloroform was added dropwise a solution of 1 equiv. (0.0986 mmol) of monothiol (the HSR donor ligand) and 1 equiv. of the dithiol [(HSCH₂CH₂)₂NR, R=CH₂C₆H₅, CH₂C₅H₄N, (CH₂)₂C₅H₄N] (0.0986 mmol) in 1 ml of chloroform. The solution remained olive green until the addition of triethylamine (0.08 ml, 0.574 mmol) whereupon it immediately changed from olive to forest green. The solution was stirred and

refluxed for an additional 30 min and then evaporated to dryness. Crystals for each of the compounds were grown as indicated.

2.2.3.1. Preparation of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-C}_6\text{H}_5)\}(\eta^1\text{-C}_6\text{H}_4\text{Br-4-S})]$ (1): X-ray quality crystals were grown by slow diffusion of pentane in a solution of the compound in methylene chloride (yield: 0.021 g, 34.6%). *Anal.* Calc. for $\text{C}_{17}\text{H}_{19}\text{NOS}_3\text{BrRe}$ (mol. wt. 615.62): C, 33.2; H, 3.11; N, 2.28. Found: C, 32.5; H, 3.49; N, 2.16%.

2.2.3.2. Preparation of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-C}_5\text{H}_4\text{N})\}(\eta^1\text{-C}_6\text{H}_4\text{Br-4-S})]$ (2): X-ray quality crystals were grown by slow diffusion of pentane in a solution of the compound in methylene chloride (yield: 0.027 g, 44.4%). IR (KBr, cm^{-1}): 1654 (m), 1560 (m), 1508 (w), 1267 (m), 1080 (m), 1008 (m), 948 (s), 813 (w). ^1H NMR (CDCl_3 , ppm): 2.73 (m, 2H), 2.85 (m, 2H), 3.52 (m, 2H), 3.85 (m, 2H), 5.03 (s, 2H), 7.26 (m, 2H), 7.38 (m, 2H), 7.53 (m, 2H), 7.80 (m, 1H), 8.71 (m, 1H).

2.2.3.3. Preparation of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-C}_5\text{H}_4\text{N})\}(\eta^1\text{-C}_6\text{H}_4\text{Cl-4-S})]$ (3): X-ray quality crystals were grown by slow diffusion of diethylether into a methylene chloride solution of **3** (yield: 0.019 g, 33.9%). IR (KBr, cm^{-1}): 1654 (m), 1560 (m), 1508 (w), 1266 (m), 1085 (m), 1012 (m), 949 (s), 818 (w). ^1H NMR (CDCl_3 , ppm): 2.72 (m, 2H), 2.83 (m, 2H), 3.52 (m, 2H), 3.86 (m, 2H), 5.03 (s, 2H), 7.35 (m, 2H), 7.41 (m, 2H), 7.73 (m, 2H), 7.82 (m, 1H), 8.71 (m, 1H). *Anal.* Calc. for $\text{C}_{16}\text{H}_{18}\text{N}_2\text{OS}_3\text{ClRe}$ (mol. wt. 572.15): C, 33.6; H, 3.17; N, 4.90. Found: C, 34.1; H, 3.55; N, 4.07%.

2.2.3.4. Preparation of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-C}_5\text{H}_4\text{N})\}(\eta^1\text{-C}_6\text{H}_4\text{F-4-S})]$ (4): X-ray quality crystals were grown by slow diffusion of pentane into a methylene chloride solution of **4** (yield: 0.025 g, 45.6%). IR (KBr, cm^{-1}): 1654 (m), 1560 (w), 1508 (w), 1263 (s), 1154 (m), 1014 (m), 942 (s), 833 (w). ^1H NMR (CDCl_3 , ppm): 2.69 (m, 2H), 2.83 (m, 2H), 3.53 (m, 2H), 3.85 (m, 2H), 5.03 (s, 2H), 7.09 (m, 2H), 7.39 (m, 2H), 7.63 (m, 2H), 7.80 (m, 1H), 8.71 (m, 1H).

2.2.3.5. Preparation of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-C}_5\text{H}_4\text{N})\}(\eta^1\text{-C}_6\text{H}_4\text{OCH}_3\text{-4-S})]$ (5): X-ray quality crystals were grown by slow diffusion of pentane into a solution of the compound in methylene chloride (yield: 0.023 g, 41.1%). IR (KBr, cm^{-1}): 1654 (m), 1560 (w), 1508 (w), 1283 (s), 1172 (m), 1028 (m), 941 (s), 828 (w). ^1H NMR (CDCl_3 , ppm): 2.71 (m, 2H), 2.84 (m, 2H), 3.55 (m, 2H), 3.82 (m, 2H), 3.88 (s, 3H), 5.07 (s, 2H), 6.95 (m, 2H), 7.37 (m, 2H), 7.61 (m, 2H), 7.79 (m, 1H), 8.72 (m, 1H).

2.2.3.6. Preparation of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-C}_5\text{H}_4\text{N})\}(\eta^1\text{-C}_6\text{H}_4\text{OCH}_3\text{-4-CH}_2\text{S})]$ (6): X-ray quality crystals were grown by slow diffusion of diethylether into a methylene chloride solution of **6** (yield: 0.027 g, 47.1%). IR (KBr, cm^{-1}): 1654 (m), 1560 (w), 1509 (w), 1248 (s), 1104 (m), 1032 (m), 948 (s), 819 (w). ^1H NMR (CDCl_3 , ppm): 2.61 (m, 2H), 3.01 (m, 2H), 3.53 (m, 2H), 3.81 (s, 3H), 3.87 (m, 2H), 4.89 (s, 2H), 5.00 (s, 2H), 6.84 (m, 2H), 7.34 (m, 2H), 7.41 (m, 2H), 7.76 (m, 1H), 8.71 (m, 1H). *Anal.* Calc. for $\text{C}_{18}\text{H}_{23}\text{N}_2\text{O}_2\text{S}_3\text{Re}$ (mol. wt. 581.76): C, 37.2; H, 3.98; N, 4.82. Found: C, 36.8; H, 3.73; N, 4.55%.

2.2.3.7. Preparation of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-CH}_2\text{C}_5\text{H}_4\text{N})\}(\eta^1\text{-C}_6\text{H}_4\text{Br-4-S})]$ (7): X-ray quality crystals were grown by slow diffusion of pentane into a methylene chloride solution of **7** (yield: 0.031 g, 49.9%). IR (KBr, cm^{-1}): 1654 (m), 1578 (w), 1438 (w), 1273 (s), 1095 (m), 1029 (m), 943 (s), 846 (w). ^1H NMR (CDCl_3 , ppm): 2.76 (m, 2H), 2.91 (m, 2H), 3.31 (t, 2H), 3.42 (m, 2H), 3.68 (m, 2H), 4.19 (m, 2H), 7.24 (m, 2H), 7.32 (m, 2H), 7.51 (m, 2H), 7.67 (m, 1H), 8.56 (m, 1H). *Anal.* Calc. for $\text{C}_{17}\text{H}_{20}\text{N}_2\text{OS}_3\text{BrRe}$ (mol. wt. 630.64): C, 32.4; H, 3.20; N, 4.44. Found: C, 32.9; H, 3.36; N, 4.38%.

2.2.3.8. Preparation of $[\text{ReO}(\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-CH}_2\text{C}_5\text{H}_4\text{N))}(\eta^1\text{-C}_6\text{H}_4\text{Cl-4-S)] (8)}$: Crystals were grown by slow diffusion of pentane into a solution of the compound in methylene chloride (yield: 0.024 g, 41.5%). IR (KBr, cm^{-1}): 1654 (m), 1574 (w), 1472 (w), 1268 (s), 1090 (m), 1009 (m), 947 (s), 816 (w). ^1H NMR (CDCl_3 , ppm): 2.79 (m, 2H), 2.89 (m, 2H), 3.31 (t, 2H), 3.42 (m, 2H), 3.68 (m, 2H), 4.20 (m, 2H), 7.22 (m, 2H), 7.33 (m, 2H), 7.55 (m, 2H), 7.68 (m, 1H), 8.57 (m, 1H). *Anal.* Calc. for $\text{C}_{17}\text{H}_{20}\text{N}_2\text{OS}_3\text{ClRe}$ (mol. wt. 586.19): C, 34.8; H, 3.44; N, 4.78. Found: C, 33.6; H, 3.21; N, 2.84%.

2.2.3.9. Preparation of $[\text{ReO}(\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-CH}_2\text{C}_5\text{H}_4\text{N))}(\eta^1\text{-C}_6\text{H}_4\text{OCH}_3\text{-4-S)] (9)}$: X-ray quality crystals were grown by slow diffusion of diethylether into a solution of the compound in methylene chloride (yield: 0.023 g, 40.1%). IR (KBr, cm^{-1}): 1654 (m), 1577 (m), 1492 (w), 1283 (s), 1171 (m), 1024 (m), 938 (s), 826 (w). ^1H NMR (CDCl_3 , ppm): 2.76 (m, 2H), 2.90 (m, 2H), 3.30 (t, 2H), 3.40 (m, 2H), 3.65 (m, 2H), 3.83 (s, 3H), 4.18 (m, 2H), 6.93 (m, 2H), 7.22 (m, 2H), 7.54 (m, 2H), 7.67 (m, 1H), 8.56 (m, 1H).

2.2.3.10. Preparation of $[\text{ReO}(\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{-CH}_2\text{C}_5\text{H}_4\text{N))}(\eta^1\text{-C}_6\text{H}_4\text{Cl-4-CH}_2\text{S)]} \cdot 0.5\text{CH}_3(\text{CH}_2)_4\text{CH}_3$ (10): X-ray quality crystals were grown by slow diffusion of hexane into a methylene chloride solution of **10** (yield: 0.028 g, 44.2%). IR (KBr, cm^{-1}): 1650 (m), 1589 (w), 1487 (w), 1263 (s), 1090 (m), 1009 (m), 948 (s), 831 (w). ^1H NMR (CDCl_3 , ppm): 2.71 (m, 2H), 3.01 (m, 2H), 3.30 (t, 2H), 3.43 (m, 2H), 3.69 (m, 2H), 4.15 (m, 2H), 4.83 (s, 2H), 7.19–7.27 (m, 4H), 7.33 (m, 2H), 7.65 (m, 2H), 8.56 (m, 1H).

2.3. X-ray crystallography

All data were collected on a Bruker SMART diffractometer system using graphite monochromated Mo K α radiation ($\lambda(\text{Mo K}\alpha)=0.71073 \text{ \AA}$). All the data collections were carried out at low temperature (87–93 K). The crystal parameters and other experimental details of the data collections are summarized in Table 1. A complete description of the details of the crystallographic methods is given in Supporting Information. Data was corrected for Lorentz and polarization effects, and absorption corrections were made using SAD-ABS [35]. The structures were solved by direct methods [36]. Neutral atom scattering factors were taken from Cromer and Waber [37] and anomalous dispersion corrections were taken from those of Creagh and McAuley [38]. All calculations were performed using SHELXTL-96 [36]. Non-hydrogen atoms were refined anisotropically. No anomalies were encountered in the refinements of any of the structures. Atomic positional parameters for the structures have been deposited with the Cambridge Structural Database (Section 5).

3. Results and discussion

The synthesis of *N,N*-bis(2-mercaptoethyl)benzylamine employs the route of Schröder and coworkers [34], in which 4 equiv. of ethylene sulfide are reacted with 1 equiv. of benzylamine at 65°C for 48 h. A similar procedure was employed in the synthesis of $[(\text{HSCH}_2\text{-CH}_2)_2\text{N(CH}_2\text{C}_5\text{H}_4\text{N})]$ and $[(\text{HSCH}_2\text{CH}_2)_2\text{N(CH}_2\text{CH}_2\text{-C}_5\text{H}_4\text{N})]$ starting from the appropriate amines, 2-(aminomethyl)pyridine and 2-(2-aminoethyl)pyridine, respectively.

The compounds of this study were all prepared readily from the reactions of the appropriate ligands with $[\text{ReOCl}_3(\text{PPh}_3)_2]$. While $[\text{Bu}_4\text{N}][\text{ReOBr}_4(\text{OPPh}_3)]$ also proved to be an effective starting material, there was no advantage to its use in terms of yield or product purity. Since the synthesis of $[\text{Bu}_4\text{N}][\text{ReOBr}_4(\text{OPPh}_3)]$ is relatively involved and of low yield, the more easily prepared and purified starting material $[\text{ReOCl}_3(\text{PPh}_3)_2]$ was used exclusively for the synthesis of **1–10**. The green solid is indefinitely air and moisture stable making it suitable for probing the reactivities of a variety of ligand types. Consequently, the compounds of this study were synthesized in a straightforward fashion from the reaction of $[\text{ReOCl}_3(\text{PPh}_3)_2]$ with one equivalent of the tridentate dithiol ligand (or '3' ligand) and 1

equiv. of the monodentate thiol ligand (or '1' ligand) in chloroform. An addition of excess triethylamine produces a forest green solution, which upon workup yields crystalline products **1–10**.

The infrared spectrum of **2–10** are characterized by a series of ligand bands in the 1008–1654 cm^{-1} range and a strong band in the 938–949 cm^{-1} range attributed to $\nu(\text{Re}=\text{O})$. The ^1H NMR spectra of compounds **2–6** are unexceptional. All display a series of multiplets ranging from 6.84 to 8.72 ppm assignable to the eight aromatic protons, as well as four multiplets at 2.69, 2.87, 3.53, and 3.85 ppm assignable to each CH_2 moiety comprising the backbone of the tridentate ligand ($^-\text{SCH}_2\text{CH}_2\text{N}(\text{R})\text{CH}_2\text{CH}_2\text{S}^-$). All four of the multiplets integrate to two protons; however, individual assignments of the resonances are tenuous at best, due to the multiplicity of the peaks and the flexibility of the ligand in solution, as noted by others [6]. In addition, compounds **2–6** also displayed a single proton resonance integrating to two protons at 5.03 ppm corresponding to the methylene spacer of the tridentate backbone. Furthermore, $[\text{ReO}\{\eta^3-(\text{SCH}_2\text{CH}_2)_2\text{N}(\text{CH}_2-\text{C}_5\text{H}_4\text{N})\}(\eta^1-\text{C}_6\text{H}_4\text{OCH}_3-4-\text{S})]$ (**5**) and $[\text{ReO}\{\eta^3-(\text{SCH}_2-\text{CH}_2)_2\text{N}(\text{CH}_2\text{C}_5\text{H}_4\text{N})\}(\eta^1-\text{C}_6\text{H}_4\text{OCH}_3-4-\text{CH}_2\text{S})]$ (**6**) exhibited an additional singlet at approximately 3.88 and 3.81 ppm, respectively, which is attributed to the three protons of the methoxy group. The monodentate benzylmercaptan of compound **6** displayed methylene protons at approximately 4.89 ppm as a single peak, as anticipated from Sadtler spectra [39].

Similarly, the ^1H NMR spectra for compounds **7–10** contain multiplets at 2.76, 2.93, 3.42, and 3.68 ppm assignable to each CH_2 moiety comprising the backbone of the tridentate ligand, as well as series of multiplets ranging from 6.93 to 8.57 ppm assignable to the eight aromatic protons. Compounds **7–10** also displayed a triplet at 3.31 ppm and a multiplet at 4.18 ppm each integrating to two protons and corresponding to the ethylene spacer of the tridentate ligand. In addition, the monodentate benzylmercaptan of compound **10** displayed methylene protons at approximately 4.83 ppm as a single peak, as anticipated from Sadtler spectra [38]. $[\text{ReO}\{\eta^3-(\text{SCH}_2\text{CH}_2)_2\text{N}(\text{CH}_2\text{CH}_2\text{C}_5\text{H}_4\text{N})\}(\eta^1-\text{C}_6\text{H}_4\text{OCH}_3-4-\text{S})]$ (**9**) exhibited an additional singlet at approximately 3.83 ppm, which is attributed to the three protons of the methoxy group. In no case does the ^1H NMR data suggest the presence of *syn/anti* isomers with respect to the amine substituent R and the oxo-group orientations. The crystal structure data, likewise, confirm that all compounds adopt the *syn* geometry exclusively.

As shown in Fig. 1, the coordination geometry around the metal for $[\text{ReO}\{\eta^3-(\text{SCH}_2\text{CH}_2)_2\text{N}(\text{CH}_2\text{C}_6\text{H}_5)\}(\eta^1-\text{C}_6\text{H}_4\text{Br}-4-\text{S})]$ (**1**) is distorted trigonal bipyramidal, with the oxo group and the sulfur atoms of the [SNS] ligand defining the equatorial plane and the axial positions occupied by the sulfur atom of the monothiol and the nitrogen atom of the [SNS] ligand. The angles between the atoms in the equatorial plane are close to the ideal 120° (Table 2), but a value of $157.91(15)^\circ$ observed for the $\text{N}(1)-\text{Re}(1)-\text{S}(3)$ angle indicates a distortion of the trigonal bipyramid, which can also be seen in the value of the trigonality index [40], $\tau=0.61$. The bond distances in the coordination environment of the metal are characteristic for this type of complexation [31]. The $\text{Re}=\text{O}$ bond length is found to be $1.699(5) \text{ \AA}$ well within the range observed for other $\{\text{ReO}\}^{3+}$ complexes [24,31,41–52]. Likewise, the $\text{Re}-\text{N}$ bond length [$\text{Re}-\text{N}(1)=2.208(6) \text{ \AA}$] is typical for the $\text{Re}-\text{N}_{\text{amine}}$ single bond in which the nitrogen atom exhibits sp^3 hybridization [24,31,41–52]. However, in the case of the metal–sulfur bond lengths, those in the equatorial plane are similar [$\text{Re}-\text{S}(1)=2.2794(18)$, $\text{Re}-\text{S}(2)=2.2824(18) \text{ \AA}$], while for the $\text{Re}-\text{S}$ axial bond there is an expected lengthening [$\text{Re}-\text{S}(3)=2.3097(19) \text{ \AA}$] [24,31,41–52]. The rhenium atom is situated $0.0817 \text{ \AA}$, above the equatorial plane of the trigonal bipyramid in the direction of the monothiol. This distance is within the range reported for five-coordinate $\text{Re}/\text{Tc}(\text{V})$ complexes with similar distorted trigonal bipyramidal geometry [31].

The compounds $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{C}_5\text{H}_4\text{N)}\}(\eta^1\text{-C}_6\text{H}_4\text{X-4-S})]$ (X = Br (**2**), Cl (**3**), F (**4**), OCH₃ (**5**)), $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{C}_5\text{H}_4\text{N)}\}(\eta^1\text{-C}_6\text{H}_4\text{O-CH}_3\text{-4-CH}_2\text{S})]$ (**6**), $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{CH}_2\text{-C}_5\text{H}_4\text{N)}\}(\eta^1\text{-C}_6\text{H}_4\text{X-4-S})]$ (X = Br (**7**), Cl (**8**), OCH₃ (**9**)), and $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{CH}_2\text{C}_5\text{H}_4\text{N)}\}(\eta^1\text{-C}_6\text{H}_4\text{Cl-4-CH}_2\text{S})]$ · 0.5CH₃(CH₂)₄CH₃ (**10**) are structurally similar to **1** with the exception of the *para*-substituent of the benzenethiol or benzylmercaptan, as well as the substitution on the nitrogen of the tridentate. The core geometry of the five-coordinate oxo-rhenium complexes containing the tridentate ligand, $[(\text{HSCH}_2\text{CH}_2)_2\text{N(CH}_2\text{C}_5\text{H}_4\text{N})]$, is shown in Fig. 2 for $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{C}_5\text{H}_4\text{N)}\}(\eta^1\text{-C}_6\text{H}_4\text{Br-4-S})]$ (**2**) and is seen to consist of distorted trigonal bipyramidal geometry with the oxo group and the sulfur atoms of the [SNS] ligand defining the equatorial plane and the axial positions occupied by the sulfur atom of the monothiol and the nitrogen atom of the [SNS] ligand. The angles between the atoms in the equatorial plane are close to the ideal 120° (Table 2), but a value of 163.35(17)° observed for the N(1)–Re(1)–S(3) angle indicates a distortion of the trigonal bipyramid, which can also be seen in the value of the trigonality index [51], $\tau=0.74$. The Re=O bond length is found to be 1.695(6) Å and the Re–N bond length [Re–N(1)=2.221(7) Å] is typical for the Re–N_{amine} single bond in which the nitrogen atom exhibits sp³ hybridization. The metal–sulfur bond lengths in the equatorial plane are unexceptional [Re–S(1)=2.270(2), Re–S(2)=2.281(2) Å], while the Re–S(3) axial bond length is 2.313(2) Å, a lengthening phenomena noted by others [24,31,41–52]. The rhenium atom is situated 0.0977 Å, above the equatorial plane of the trigonal bipyramid in the direction of the monothiol.

Similarly, the core geometry of the five-coordinate oxo-rhenium complexes containing the $[(\text{HSCH}_2\text{-CH}_2)_2\text{N(CH}_2\text{CH}_2\text{C}_5\text{H}_4\text{N})]$ tridentate is shown in Fig. 3 for $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{CH}_2\text{C}_5\text{H}_4\text{N)}\}(\eta^1\text{-C}_6\text{H}_4\text{Br-4-S})]$ (**7**) and is also seen to consist of distorted trigonal bipyramidal geometry. The oxo group and the sulfur atoms of the [SNS] ligand define the equatorial plane, while the axial positions are occupied by the sulfur atom of the monothiol and the nitrogen atom of the [SNS] ligand. The angles between the atoms in the equatorial plane are close to the ideal 120° (Table 2), likewise, a value of 159.42(10)° observed for the N(1)–Re(1)–S(3) angle indicates a distortion of the trigonal bipyramid, which can also be seen in the value of the trigonality index [40], $\tau=0.60$. The Re=O and Re–N_{amine} bond lengths are typical for this particular class of compounds and are found to be 1.694(3) and 2.217(4) Å respectively [24,31,39–49,53]. As in the previous cases, the metal–sulfur bond lengths in the equatorial plane are unexceptional [Re–S(1)=2.2758(12), Re–S(2)=2.2918(11) Å], while the Re–S(3) axial bond length is slightly lengthened to 2.3144(11) Å. The rhenium atom is situated 0.0811 Å above the equatorial plane of the trigonal bipyramid in the direction of the monothiol.

In all the oxo-rhenium benzenethiolates and benzylmercaptans (compounds **2–7**, **9**, **10**) the equatorial plane is defined by the two sulfur donors of the tridentate [SNS] ligand and the oxo group, while the sulfur atom of the monothiol and the nitrogen atom of the [SNS] ligand occupy the axial positions. The Addison τ indices for the rhenium mixed thiolate compounds of this study are listed in Table 3, revealing that compounds **1–7**, **9** and **10** actually tend to lie closer to the distorted trigonal bipyramidal structure, a result that is similar to other compounds that are members of the oxo-rhenium ‘3+1’ [SNS]/[S] mixed-thiolate family [28–31].

In all cases, one equatorial position is occupied by a terminal oxo-group with a typical Re=O distance ranging from approximately 1.69–1.71 Å. The two other equatorial positions are occupied by the two sulfur donors of the [SNS] tridentate. In the complexes of this study, the displacement of Re site from the equatorial plane in the direction of the monothiol group is approximately 0.0753–0.1087 Å as shown in Table 4. The Re–N_{amine} distances range from 2.21 to 2.23 Å while the sulfur distances are all fairly unexceptional, ranging from 2.27

to 2.32 Å well within the range of those reported previously [52–54]. Tables 2 and 4 compare the relevant bond angles and lengths for compounds **1–7**, **9** and **10**, respectively.

4. Conclusions

Exploitation of the thiolate chemistry of the oxo-rhenium(V) core has allowed the preparation of a series of mononuclear oxo-rhenium-mixed thiolate complexes. We have demonstrated that the action of the tridentate [SN(R)S] aminodithiol ligand on a suitable {ReO}³⁺ precursor in the presence of a monodentate ligand gives rise to stable, neutral oxo-rhenium complexes of the ‘3+1’ type. This route appears to be a quite general, facile method for moderate yield synthesis of materials with the [ReO{SN(R)S}(S)] core. The geometry and charge of the monodentate thiol ligand may be manipulated easily to produce complexes of various size, shape and charges for clinical development.

Acknowledgments

This work was supported by a grant from the Department of Energy, Office of Health and Environmental Research (D2-FG02-99ER62791).

References

1. Schubiger PA, Alberto R, Smith A. *Bioconjug Chem* 1996;7:165. [PubMed: 8983338]
2. Weininger J, Ketring AR, Deutsch E, Maxon HR, Goeckler WR. *J Nucl Med* 1983;24:125.
3. Wessels BW, Rogus RD. *Med Phys* 1984;11:638. [PubMed: 6503879]
4. Yorke ED, Beaumier P, Wessels B, Fritzberg A, Morgan A. *Nucl Med Biol* 1991;18:827.
5. Vanderheyden JL, Fritzberg AR, Rao TN, Kasina S, Srinivasan A, Reno JM, Morgan AC. *J Nucl Med* 1987;28:656.
6. Pietzsch HJ, Spies H, Hoffman S, Stach J. *Inorg Chim Acta* 1989;161:15.
7. Spies H, Pietzsch HJ, Sayhe R, Hoffman S. *Isotopenpraxis* 1990;26:159.
8. Pietzsch HJ, Spies H, Hoffman S, Scheller D. *Appl Radiat Isot* 1990;41:185.
9. Bandoli G, Mazzi U, Pietzsch HJ, Spies H. *Acta Crystallogr, Sect C* 1992;48:1422.
10. Fietz T, Spies H, Pietzsch HJ, Leibnitz P. *Inorg Chim Acta* 1995;231:233.
11. Spies H, Fietz T, Pietzsch HJ, Johannsen B, Leibnitz P, Reck G, Scheller D, Klostermann K. *J Chem Soc, Dalton Trans* 1995:2277.
12. Pelecanou M, Pirmettis IC, Nock BA, Papadopoulos MS, Chiotellis E, Stassinopoulou CI. *Inorg Chim Acta* 1998;281:148.
13. Megalla S, Plossl K, Kung MP, Chumpradit S, Stevenson DA, Frederick D. *Bioconjug Chem* 1996;7:421. [PubMed: 8853455]
14. Megalla S, Plossl K, Kung MP, Liable-Sands LM, Rheingold AL. *J Am Chem Soc* 1995;117:11037.
15. Mastrostamatis SG, Papadopoulos MS, Pirmettis IC, Paschali E, Varvarigou AD, Stassinopoulou CI, Raptopoulou CP, Terzis A, Chiotellis E. *J Med Chem* 1994;37:3212. [PubMed: 7932548]
16. Pirmettis IC, Papadopoulos MS, Chiotellis E. *J Med Chem* 1997;40:2539. [PubMed: 9258360]
17. Maresca KP, Bonavia GH, Babich JW, Zubieta J. *Inorg Chim Acta* 1999;284:252.
18. Johannsen B, Scheunemann M, Spies H, Brust P, Wober J, Syhre R, Pietzsch HJ. *Nucl Med Biol* 1996;23:429. [PubMed: 8832697]
19. Skaddan MB, Katzenellenbogen JA. *Bioconjug Chem* 1999;10:119. [PubMed: 9893973]
20. Maresca KP, Femia FJ, Bonavia GH, Babich JW, Zubieta J. *Inorg Chim Acta* 2000;297:98.
21. Pietzsch HJ, Spies H, Hoffmann S. *Inorg Chim Acta* 1989;165:163.
22. Maresca KP, Femia FJ, Babich JW, Zubieta J. *Inorg Chem Commun* 1998;1:209.
23. Femia FJ, Chen X, Babich JW, Zubieta J. *Inorg Chim Acta* 2000;300–302:517.
24. Tisato F, Refosco F, Mazzi U, Bandoli G, Nicolini M. *J Chem Soc, Dalton Trans* 1987:1693.

25. Bandoli G, Mazzi U, Wilcox B, Jurisson S, Deutsch E. *Inorg Chim Acta* 1984;95:217.
26. Duatti A, Marchi A, Rossi R, Magon L, Deutsch E, Bertoalsi V, Bellucci F. *Inorg Chem* 1988;27:4208.
27. Chen X, Femia FJ, Babich JW, Zubieta J. *Inorg Chim Acta* 2000;306:113–116.
28. Papadopoulos M, Pirmettis I, Tsoukalas C, Nock B, Maina T, Raptopoulou CP, Pietzsch HJ, Friebe M, Spies H, Johannsen D, Chiotellis E. *Inorg Chim Acta* 1999;295:1.
29. Pelecanou M, Pirmettis IC, Papadopoulos MS, Raptopoulou CP, Terzis A, Chiotellis E, Stassinopoulou CI. *Inorg Chim Acta* 1999;287:142.
30. Pelecanou M, Pirmettis IC, Nock BA, Papadopoulos M, Chiotellis E, Stassinopoulou CI. *Inorg Chim Acta* 1998;281:148.
31. Papadopoulos M, Tsoukalas C, Pirmettis I, Nock B, Maina T, Abedin Z, Raptopoulou CP, Terzis A, Chiotellis E. *Inorg Chim Acta* 1999;285:97.
32. Babich, JW.; Ferill, K.; Rose, DJ.; Graham, W.; Maresca, KP.; Zubieta, J.; Fischman, AJ. 43rd Meeting of the Society of Nuclear Medicine; Denver, CO. 3–6 June 1996; abstract
33. Chatt J, Rowe GA. *J Chem Soc* 1962:4019.
34. Spencer DJE, Blake AJ, Parsons S, Schröder M. *J Chem Soc, Dalton Trans* 1999:1041.
35. Sheldrick, GM. SADABS, Program for Empirical Absorption Corrections. University of Gottingen; Germany: 1996.
36. Sheldrick, GM. SHELXL96, Program for Refinement of Crystal Structures. University of Gottingen; Germany: 1996.
37. Cromer, DT.; Waber, JT. *International Tables for X-ray Crystallography*. Vol. IV. Kynoch; Birmingham, UK: 1974.
38. Creagh, DC.; McAuley, JWJ. *International Tables for X-ray Crystallography*, vol C, Table 4. Kluwer Academic; Boston: 1992.
39. The Sadtler standard spectra. Sadtler Research Laboratories; Philadelphia, PA: 1991.
40. Addison AW, Rao TN, Reedijk K, Verschoor GC. *J Chem Soc, Dalton Trans* 1984:1349.
41. Papadopoulos M, Pirmettis I, Pelecanou M. *Inorg Chem* 1996;35:7377. [PubMed: 11666932]
42. Lever SZ, Baidoo KE, Mahmood A. *Inorg Chim Acta* 1990;176:183.
43. Davison A, Orvig C, Trop HS, Sohn M, DePamphilis BV, Jones AG. *Inorg Chem* 1980;19:1988.
44. Bryson N, Dewan JC, Lister-James J, Davison A. *Inorg Chem* 1988;27:2154.
45. deVries N, Jones AG, Davison A. *Inorg Chem* 1991;30:2662.
46. Blower PJ, Dilworth JR, Hutchison JP, Nicholson T. J Zubieta, *J Chem Soc, Dalton Trans* 1986:1339.
47. Davison A, Dephamphilis BV, Faggiani R, Jones AG, Lock CJL, Orvig C. *Can J Chem* 1985;63:319.
48. Bandoli G, Gerber TIA. *Inorg Chim Acta* 1987;126:205.
49. Glegg W, Boyde S, Garner CD. *Acta Crystallogr, Sect C* 1988;44:172.
50. Jones AG, Davison A, LaTegola MR, Brodak JW, Orvig C, Sohn M, Toothaker AK, Lock CJL, Franklin KJ, Costello CE, Carr SA, Biemann K, Kaplan ML. *J Nucl Med* 1982;23:809.
51. Smith JE, Byrne EF, Cotton FA, Sekutowski JC. *J Am Chem Soc* 1978;112:5571.
52. Francesconi LC, Graczyk G, Wehl S, Shaikh SN, Mc-Clinton D, Liu S, Zubieta J, Kung HF. *Inorg Chem* 1993;32:3114.
53. Chrysou K, Pelecanou M, Papadopoulos MS, Raptopoulou CP, Pirmettis IC, Chiotellis E, Stassinopoulou CI. *Inorg Chim Acta* 1998;268:169.
54. Rose DJ, Chang YD, Chen Q, Kettler PB, Zubieta J. *Inorg Chem* 1995;34:3973.

Fig. 1. A view of the structure of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{N(CH}_2\text{C}_6\text{H}_5)\}(\eta^1\text{-C}_6\text{H}_4\text{Br-4-S})]$ (**1**), showing the atom-labeling scheme and 50% thermal ellipsoids.

Fig. 2.
A view of the structure of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{-N(CH}_2\text{C}_5\text{H}_4\text{N)}\}(\eta^1\text{-C}_6\text{H}_4\text{Br-4-S})]$ (**2**), showing the atom-labeling scheme and 50% thermal ellipsoids.

Fig. 3.

A view of the structure of $[\text{ReO}\{\eta^3\text{-(SCH}_2\text{CH}_2)_2\text{-N(CH}_2\text{CH}_2\text{C}_5\text{H}_4\text{N)}\}(\eta^1\text{-C}_6\text{H}_4\text{Br-4-S})]$ (**7**), showing the atom-labeling scheme and 50% thermal ellipsoids.

Table 1

Summary of the crystallographic data for the compounds (1–10)

	1	2	3	4	5	6	7	9	10
Chemical formula	$C_{17}H_{19}NO_3BrRe$	$C_{16}H_{18}N_2O_3BrRe$	$C_{16}H_{18}N_2O_3ClRe$	$C_{16}H_{18}N_2O_3S_3Re$	$C_{17}H_{21}N_2O_2S_3Re$	$C_{18}H_{23}N_2O_2S_3Re$	$C_{17}H_{20}N_2O_3BrRe$	$C_{18}H_{23}N_2O_2S_3Re$	$C_{21}H_{29}N_2OS_3ClRe$
<i>a</i> (Å)	11.2989(6)	9.3051(7)	9.3247(4)	8.9720(5)	9.4042(4)	8.3855(4)	9.4161(4)	11.6331(4)	11.2621(7)
<i>b</i> (Å)	13.1804(7)	9.5482(7)	9.4485(4)	9.4770(5)	9.4183(4)	24.8281(12)	9.5641(4)	10.4400(4)	12.0955(7)
<i>c</i> (Å)	13.5621(8)	11.4612(8)	11.3743(5)	11.6194(6)	11.5443(5)	9.7181(5)	12.0349(5)	16.5754(6)	16.5589(10)
<i>α</i> (°)	80.791(1)	69.709(1)	70.320(1)	72.876(1)	111.975(1)	90	109.8550(10)	90	85.1660(10)
<i>β</i> (°)	75.088(1)	74.940(1)	74.615(1)	70.865(1)	100.927(1)	104.9880(10)	104.4130(10)	105.5100(10)	74.3980(10)
<i>γ</i> (°)	78.042(1)	86.287(1)	84.808(1)	81.471(1)	90.201(1)	90	91.9330(10)	90	85.7420(10)
<i>V</i> (Å ³)	1897.15(18)	921.94(12)	909.78(7)	890.54(8)	927.90(7)	1954.44(17)	978.80(7)	1939.77(12)	2161.7(2)
<i>Z</i>	4	2	2	2	2	4	2	4	4
Formula weight	615.62	616.61	572.15	555.70	567.74	581.76	630.64	581.76	643.29
Space group	<i>P</i> <i>1</i>	<i>P</i> <i>1</i>	<i>P</i> <i>1</i>	<i>P</i> <i>1</i>	<i>P</i> <i>1</i>	<i>P</i> 2 ₁ / <i>n</i>	<i>P</i> <i>1</i>	<i>P</i> 2 ₁ / <i>n</i>	<i>P</i> <i>1</i>
<i>T</i> (K)	89(5)	87(5)	87(5)	88(5)	90(5)	87(5)	89(5)	90(5)	93(5)
<i>λ</i> (Å)	0.71073	0.71073	0.71073	0.71073	0.71073	0.71073	0.71073	0.71073	0.71073
<i>D</i> _{calc} (g cm ^{−3})	2.155	2.221	2.089	2.072	2.032	1.977	2.140	1.992	1.977
<i>μ</i> (mm ^{−1})	8.846	9.103	7.176	7.190	6.899	6.554	8.577	6.603	6.052
<i>R</i> ^a	0.0439	0.0665	0.0289	0.0426	0.0315	0.0641	0.0345	0.0342	0.0560
<i>wR</i> ₂ ^b	0.0832	0.1158	0.0710	0.1127	0.0710	0.0744	0.0753	0.0534	0.1238

$$\alpha_{\Sigma}|F_o| - |F_c|/\Sigma|F_o|.$$

$$\frac{1}{\Sigma|F_o|^2} \left[\sum [w(F_o^2 - F_c^2)] / \sum [w(F_o^2)] \right]^{1/2}.$$

Table 2
Comparison of selected bond angles ^a for the rhenium-mixed thiolate complexes of this study

Angle	1	2	3	4	5	6	7	9	10
O(1)–Re–N(1)	96.3(2)	93.9(2)	93.71(13)	93.10(18)	94.60(14)	94.47(17)	96.23(14)	97.41(10)	97.8(2)
O(1)–Re–S(2)	118.23(17)	121.6(2)	122.19(11)	118.37(14)	117.24(12)	119.65(13)	118.12(11)	117.78(9)	117.36(16)
N(1)–Re–S(2)	83.31(15)	84.01(18)	83.97(9)	84.09(12)	84.50(10)	83.11(11)	83.58(10)	83.57(8)	82.86(14)
O(1)–Re–S(1)	120.07(17)	118.6(2)	118.14(11)	122.05(14)	122.55(12)	117.31(13)	118.15(11)	117.55(9)	117.34(16)
N(1)–Re–S(1)	83.80(15)	83.86(18)	83.77(9)	83.63(12)	83.16(10)	83.71(11)	83.90(11)	83.46(7)	83.43(14)
S(2)–Re–S(1)	121.23(7)	119.05(8)	118.94(4)	118.74(5)	119.60(4)	122.21(5)	123.26(4)	124.27(3)	124.84(6)
O(1)–Re–S(3)	105.66(17)	102.8(2)	102.86(10)	102.89(14)	103.53(10)	105.62(13)	104.07(11)	104.65(8)	105.22(16)
N(1)–Re–S(3)	157.91(15)	163.35(17)	163.43(9)	164.00(12)	161.85(10)	159.56(12)	159.42(10)	157.78(7)	156.50(14)
S(2)–Re–S(3)	84.05(7)	87.34(8)	87.28(4)	88.03(5)	87.83(4)	90.02(5)	83.86(4)	88.44(3)	82.34(6)
S(1)–Re–S(3)	87.37(7)	87.97(8)	88.21(4)	88.03(5)	86.39(4)	83.75(5)	89.45(4)	84.03(3)	90.19(6)

^a Angles reported in degrees with e.s.d. values in parentheses.

Table 3

The τ index for the five-coordinate rhenium complexes of this study, as a measure of geometric distortion from idealized geometries

Compound	α^a	β^b	τ^c
[ReO{SN(R ₁)S}(η^1 -C ₆ H ₄ Br-4-S)] (1)	121.23(7)	157.91(15)	0.61
[ReO{SN(R ₂)S}(η^1 -C ₆ H ₄ Br-4-S)] (2)	119.05(8)	163.35(17)	0.74
[ReO{SN(R ₂)S}(η^1 -C ₆ H ₄ Cl-4-S)] (3)	118.94(4)	163.43(9)	0.74
[ReO{SN(R ₂)S}(η^1 -C ₆ H ₄ F-4-S)] (4)	118.74(5)	164.00(12)	0.75
[ReO{SN(R ₂)S}(η^1 -C ₆ H ₄ OCH ₃ -4-S)] (5)	119.60(4)	161.85(10)	0.70
[ReO{SN(R ₂)S}(η^1 -C ₆ H ₄ OCH ₃ -4-CH ₂ S)] (6)	122.21(5)	159.56(12)	0.62
[ReO{SN(R ₃)S}(η^1 -C ₆ H ₄ Br-4-S)] (7)	123.26(4)	159.42(10)	0.60
[ReO{SN(R ₃)S}(η^1 -C ₆ H ₄ OCH ₃ -4-S)] (9)	124.27(3)	157.78(7)	0.56
[ReO{SN(R ₃)S}(η^1 -C ₆ H ₄ Cl-4-CH ₂ S)] (10)	124.84(6)	156.50(14)	0.53

^a $\alpha \equiv$ the second largest basal angle.

^b $\beta \equiv$ the largest basal angle.

^c $\tau \equiv (\beta - \alpha)/60$. SN(R)S= η^3 -{(SCH₂CH₂)₂N(R)} (R₁ = CH₂C₆H₅; R₂ = CH₂C₅H₄N; R₃ = (CH₂)₂C₅H₄N).

Table 4

Comparison of selected bond lengths ^a for rhenium-mixed thiolate complexes

Complex	Re=O	η^1 -Thiolate		η^3 -Donor [S(1)-N(1)-S(2)]		ΔRe^b
		Re-S(3)	Re-S(1)	Re-S(1)	Re-N(1)	Re-S(2)
1	1.699(5)	2.3097(19)	2.2794(18)	2.208(6)	2.2824(18)	0.0817
2	1.695(6)	2.313(2)	2.270(2)	2.221(7)	2.281(2)	0.0977
3	1.702(3)	2.3087(10)	2.2821(10)	2.229(3)	2.2757(10)	0.1011
4	1.705(4)	2.3074(14)	2.2793(13)	2.231(5)	2.2772(14)	0.1087
5	1.697(3)	2.3119(11)	2.2876(11)	2.234(4)	2.2704(11)	0.0926
6	1.707(4)	2.2983(15)	2.2766(14)	2.215(5)	2.2929(14)	0.1084
7	1.694(3)	2.3144(11)	2.758(12)	2.217(4)	2.2918(11)	0.0811
9	1.703(2)	2.3189(8)	2.2889(9)	2.220(3)	2.3189(8)	0.0753
10	1.688(5)	2.2954(15)	2.2875(15)	2.226(5)	2.2962(16)	0.0815

^aDistances reported in Å with e.s.d. values in parentheses.

^b ΔRe , distance of the Re from the least square plane of the equatorial donors.