Published in final edited form as:

Tetrahedron Lett. 2010 June 9; 51(23): 3053–3056. doi:10.1016/j.tetlet.2010.04.006.

Syntheses of carbocyclic aminonucleosides and (–)-*epi*-4'-carbocyclic puromycin: Application of palladium(0)/indium iodide-allylations and tethered aminohydroxylations

Cara Cesario, Lawrence P. Tardibono Jr., and Marvin J. Miller*

The University of Notre Dame, Department of Chemistry and Biochemistry, 251 Nieuwland Science Hall, Notre Dame, Indiana 46556

Abstract

Carbocyclic aminonucleosides and *epi*-4'-carbocyclic puromycin were prepared from an acylnitroso-derived hetero Diels–Alder cycloadduct. Pd(0)/InI-mediated allylations of a formyl species were used to install the 4'-hydroxymethyl group. A tethered aminohydroxylation strategy was employed to install the *cis*-2',3'-aminoalcohol moiety with complete regio- and diastereocontrol.

Puromycin is an aminonucleoside natural product that demonstrates broad spectrum antibiotic and antitumor activities (Figure 1).¹ The 3'-aminoacyl moiety structurally resembles the aminoacyladenyl region of an aminoacyl tRNA² and plays a crucial role in terminating protein biosynthesis. Specifically, the α -amino group is transferred to a carboxyl-activated peptidyl-tRNA to form a new peptide bond.³ However, subsequent aminoacyl-group transfer reactions are precluded due to the presence of an unreactive 3'-amide in the ribosomal acceptor site rather than an activated 3'-ester linkage. As a result, the nascent polypeptide chain is prematurely released from the ribosome.⁴

Although puromycin effectively inhibits transpeptidation, the ribonucleoside is rapidly metabolized *in vivo* to inactive and toxic components.⁵ In order to circumvent enzymatic degradation and toxicity issues, carbocyclic puromycin and related analogs have been prepared and studied extensively.⁶ The carbocyclic aminonucleosides have the same mode of action as puromycin and retain antimicrobial activity.⁷ Also, carbocyclic nucleosides lack an enzymatically labile glycosidic bond and provide a more metabolically robust scaffold.⁸ Structure-activity relationships (SAR) revealed that the 4'-hydroxymethyl group was not required for activity⁹ and the 1',2',3',4',-diastereomer lacked biological function. The *cis*-2',3'-aminoalcohol stereochemistry was essential for activity. Additionally, replacement of the functionalized cyclopentane core with a cyclohexyl ring resulted in decreased inhibition of protein synthesis.¹⁰

Our research group's continued interest in the syntheses of diverse carbocyclic nucleosides¹¹ led us to investigate carbocyclic puromycin derivatives. By employing Pd(0)/InI-mediated allylations of a formyl species, 4'-hydroxymethyl groups may be installed in the carbocyclic scaffold with regio- and diastereocontrol.¹² Additionally, a *N*-

© 2010 Elsevier Science. All rights reserved.

*Corresponding author. Tel.: +1-574-631-7571; fax: +1-574-631-6652; mmiller1@nd.edu..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

pentafluorobenzoyl-oxy carbamate linker may be constructed from the 4'-hydroxymethyl substituent. This substrate may serve as a precursor for a tethered aminohydroxylation¹³ to install the *cis*-2',3'-aminoalcohol moiety. Application of these key synthetic transformations allows complete functionalization of the cyclopentane core and provides aminonucleosides (±)-**1a** and (–)-**1b** and (–)-*epi*-4'-carbocyclic puromycin **2** (Figure 1).

We have previously reported direct hydroxymethylations at the C-4' position by using Pd(0)/InI allylation chemistry with carbocyclic platforms (±)-**3a** and (+)-**3b**.¹² Functionalized cyclopentene systems (±)-**3a** and (+)-**3b** serve as key intermediates in the syntheses of diverse carbocyclic nucleosides¹⁴ as well as starting materials in our route to carbocyclic puromycin derivatives.

Treatment of *syn*-1,4 scaffold (±)-**3a** with HCl at 80 °C removed the Boc group (Scheme 1). The crude amine salt was treated with 5-amino-4,6-dichloropyrimidine in refluxing *n*-BuOH for 4 d to afford the S_NAr product (±)-**4a**. Cyclization of compound (±)-**4a** to 6-chloropurine (±)-**5a** was achieved with triethylorthoformate in the presence of catalytic camphor sulfonic acid. The *anti*-1,4 diastereomer (+)-**3b** was subjected to identical reaction conditions to provide chloropurine analog (+)-**5b** (Scheme 2).

We developed a method to key homoallylic alcohols (±)-**5a** and (+)-**5b** and turned our attention to installing the *cis*-2',3'-aminoalcohol by using a tethered aminohydroxylation reaction.^{15,16} *N*-Sulfonyloxy carbamates¹⁷ and *N*-pentafluorobenzoyloxy carbamates^{13b} have emerged as reoxidants in tethered aminohydroxylation reactions and avoid the use of chlorinating agents and basic reaction conditions. Although this powerful methodology has already been used in total synthesis,¹⁸ few examples of tethered hydroxyaminations with functionalized homoallylic alcohols have been reported.^{13b,16c-e,17}

In order to prepare the appropriate tethered aminohydroxylation precursor, we employed a two-step sequence¹⁹ to *N*-hydroxycarbamate (±)-**7a**. When chloropurine (±)-**5a** was treated with CDI followed by an aqueous solution of *O*-*p*-methoxybenzylhydroxylamine,²⁰ *N*-benzyloxycarbamate derivative (±)-**6a** was isolated in 66% yield. Removal of the *p*-methoxybenzyl group with neat TFA provided *N*-hydroxycarbamate (±)-**7a** (Scheme 3).

Initial attempts to synthesize *N*-pentafluorobenzoyloxy carbamate (±)-**8a** were thwarted by undesired by-products. When *N*-hydroxycarbamate (±)-**7a** was treated with pentafluorobenzoyl chloride in the presence of Et₃N, the *bis*-benzoylated product dominated the reaction mixture (ca. 42% by LCMS). Although *bis*-benzoylation has not been reported in the syntheses of *N*-pentafluorobenzoyloxy carbamates,^{13b} we suspected that deprotonation of the NH with Et₃N readily occurred after *mono*-benzoylation²¹ and the stabilized anion reacted with another equivalent of the acid chloride. In order to minimize the unwanted *bis*-benzoylation reaction, we exchanged Et₃N for a weaker base. When a CH₃CN solution of *N*-hydroxycarbamate (±)-**7a** was treated with pentafluorobenzoyl chloride in the presence of pyridine, *mono*-benzoylated derivative (±)-**8a** was isolated as the major product in 67% yield.²²

The tethered aminohydroxylation was the key step en route to carbocyclic aminonucleoside core structures. We expected complete regiocontrol and anticipated diastereofacial selectivity to favor the same side where the tether originated. Dihydroxylation of *syn*-1,4-disubstituted cyclopentenes has been studied²³ and the π -facial selection of these reactions was governed by sterics and the Cieplak effect.²⁴

A *t*-BuOH/H₂O (3:1) solution of homoallyl *N*-pentafluorobenzoyloxy carbamate (±)-**8a** was treated with catalytic K₂OsO₄ and hydroxyamination product (±)-**9a** was obtained. However, two competing side reactions contributed to the low conversion to (±)-**9a**.

Hydrolysis of *N*-pentafluorobenzoyloxy carbamate ($\pm$)-**8a** provided *N*-hydroxycarbamate ($\pm$)-**7a**. Additionally, the presence of a homoallyl carbamate²⁵ suggested that hydrolysis of the imidotrioxosmium (VII) species²⁶ was also competitive. When the tethered aminohydroxylation reaction was conducted in *n*-BuOH/H₂O (3:1), the homoallyl carbamate was the dominant product and hydroxyamination product ($\pm$)-**9a** was also observed. Although LCMS and ¹H NMR of the crude product mixtures indicated the formation of an exclusive diastereomer, unwanted by-products precluded the use of these solvent conditions.

Early work on the catalytic asymmetric aminohydroxylation reaction reported the use of CH₃CN/H₂O as an effective solvent system.²⁷ When a CH₃CN/H₂O (1.5:1) solution of homoallyl *N*-pentafluorobenzoyloxy carbamate ($\pm$)-**8a** was treated with catalytic K₂OsO₄, tethered aminohydroxylation product ($\pm$)-**9a** was the major component of the reaction mixture (Scheme 3). Cyclic carbamate ($\pm$)-**9a** was isolated in 58% yield as the exclusive regio- and diastereoisomer. The relative stereochemistry was determined by NOE correlations in the ROESY spectrum.²⁸

Treatment of compound ($\pm$)-**9a** with dimethylamine afforded the S_NAr product and the cyclic carbamate was cleaved with LiOH in refluxing MeOH to provide unprecedented *lyxo*-carbocyclic aminonucleoside ($\pm$)-**1a**.

We had successfully developed methodology to prepare the *N*-pentafluorobenzoyloxy carbamate linker and identified optimal conditions for the tethered aminohydroxylation reaction by starting with *syn*-1,4-disubstituted cyclopentene diastereomer ($\pm$)-**5a**. We applied this optimized route to synthesize carbocyclic aminonucleoside precursor (–)-**9b** (Scheme 4).

Chloropurine analog (+)-**5b** was treated with CDI for 2 h. Then, a solution of *O*-*p*-methoxybenzylhydroxylamine was introduced to provide *N*-(*p*-methoxybenzyl)oxycarbamate analog (+)-**6b**. *N*-Hydroxycarbamate (+)-**7b** was revealed when (+)-**6b** was treated with neat TFA to remove the *p*-methoxybenzyl group. The tethered aminohydroxylation precursor (+)-**8b** was obtained by treatment of substrate (+)-**7b** with an CH₃CN solution of pentafluorobenzoyl chloride in the presence of pyridine. On the first attempt to effect aminohydroxylation, homoallyl *N*-pentafluorobenzoyloxy carbamate (+)-**8b** was treated with catalytic K₂OsO₄ in CH₃CN/H₂O (1.5:1) to afford hydroxyamination product (–)-**9b** in 83% isolated yield. In this case, the 1'-nucleobase is orientated on the opposite side of the tether. The reduced steric bulk may contribute to the increased yield of (–)-**9b** compared to reaction of diastereomer ($\pm$)-**8a**.

Carbocyclic aminonucleoside (–)-**1b** was synthesized by installation of the dimethylamino moiety followed by cleavage of the cyclic carbamate with LiOH (Scheme 5). The carbocyclic derivative (–)-**1b** was coupled to *N*-benzyloxycarbonyl-*p*-methoxyphenyl-L-alanine to provide 3'-derivative (–)-**10**. Hydrogenation in the presence of Pd/C removed the CBz protecting group and revealed targeted carbonucleoside, *epi*-4'-carbocyclic puromycin (–)-**2**.

In summary, we have synthesized biologically relevant carbocyclic aminonucleosides ($\pm$)-**1a** and (–)-**1b** and *epi*-4'-carbocyclic puromycin (–)-**2** from an acylnitroso-derived hetero Diels–Alder cycloadduct. Our route highlighted two key synthetic transformations. Pd(0)/In-mediated allylations were used to install the requisite hydroxymethyl group. A tethered aminohydroxylation was employed to prepare a highly functionalized carbocyclic core with complete diastereo- and regiocontrol.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We would like to thank Dr. Jed Fisher (University of Notre Dame) for helpful discussions, Dr. Bill Boggess (University of Notre Dame) and Nonka Sevova (University of Notre Dame) for mass spectroscopic analyses, Dr. Jaroslav Zajicek (University of Notre Dame) for NMR assistance. We acknowledge The University of Notre Dame and NIH (GM068012) for support of this work.

References

1. Porter JN, Hewitt RI, Hesseltine CW, Krupka J, Lowery JA, Wallace WS, Bohonos N, Williams JH. *Antibiot. Chemother.* 1952; 2:409–410.
2. Yarmolinsky MB, de la Haba GL. *Proc. Natl. Acad. Sci. U.S.A.* 1959; 45:1721–1729. [PubMed: 16590564]
3. Allen DW, Zamecnik PC. *Biochim. Biophys. Acta.* 1962; 55:865–874. [PubMed: 13860529]
4. (a) Nathans D. *Proc. Natl. Acad. Sci. U.S.A.* 1964; 51:585–592. [PubMed: 14166766] (b) Smith JD, Traut RR, Blackburn GM, Monro RE. *J. Mol. Biol.* 1965; 13:617–628.
5. Nathans, D. *Antibiotics*. Gottlieb, D.; Shaw, PD., editors. Vol. 1. Springer-Verlag; New York: 1967. p. 259–277.
6. (a) Daluge S, Vince R. *J. Org. Chem.* 1978; 43:2311–2320. (b) Vince R, Daluge S, Brownell J. *J. Med. Chem.* 1986; 29:2400–2403. [PubMed: 3783599]
7. Vince R, Daluge S, Palm M. *Biochem. Biophys. Res. Comm.* 1972; 46:866–870. [PubMed: 4550700]
8. (a) Casu F, Chiacchio MA, Romeo R, Gumina G. *Curr. Org. Chem.* 2007; 11:999–1016. (b) Jeong LS, Lee JA. *Antivir. Chem. Chemoth.* 2004; 15:235–250. (c) Rodriguez JB, Comin MJ. *Mini-Rev. Med. Chem.* 2003; 3:95–114. [PubMed: 12570843] (d) Marquez VE, Lim MI. *Med. Res. Rev.* 1986; 6:1–40. [PubMed: 3512934]
9. (a) Vince R, Daluge S. *J. Med. Chem.* 1974; 17:578–583. [PubMed: 4598021] (b) Pestka S, Vince R, Daluge S, Harris R. *Antimicrob. Agents Chem.* 1973; 4:37–43. (c) Daluge S, Vince R. *J. Med. Chem.* 1972; 15:171–177. [PubMed: 4621466]
10. Vince R, Daluge S. *J. Med. Chem.* 1977; 20:930–934. [PubMed: 327067]
11. (a) Cesario C, Miller MJ. *J. Org. Chem.* 2009; 74:5730–5733. [PubMed: 19514728] (b) Jiang MX, Jin B, Gage JL, Priour A, Savelle G, Miller MJ. *J. Org. Chem.* 2006; 71:4164–4169. [PubMed: 16709056] (c) Li F, Brogan JB, Gage JL, Zhang D, Miller MJ. *J. Org. Chem.* 2004; 69:4538–4540. [PubMed: 15202916]
12. Cesario C, Miller MJ. *Tetrahedron Lett.* 2010 In Press.
13. (a) Morales-Serna JA, Diaz Y, Matheu MI, Castillon S. *Synthesis.* 2009; 5:710–712. (b) Donohoe TJ, Bataille CJR, Gattrell W, Kloesges J, Rossignol E. *Org. Lett.* 2007; 9:1725–1728. [PubMed: 17388605]
14. For the synthesis of **3a**, see: (a) McGuigan C, Hassan-Abdallah A, Srinivasan S, Wang Y, Siddiqui A, Daluge SM, Gudmundsson KS, Zhou H, McLean EW, Peckham JP, Burnette TC, Marr H, Hazen R, Condreay LD, Johnson L, Balzarini J. *J. Med. Chem.* 2006; 49:7215–7226. [PubMed: 17125274] (a) Vince R, Hua M. *J. Med. Chem.* 1990; 33:17–21. [PubMed: 2296018] (b) Daluge SM, Martin MT, Sickles BR, Livingston DA. *Nucleos. Nucleot. Nucl.* 2000; 19:297–327. For the synthesis of **3b**, see: (c) Grumann A, Marley H, Taylor RJK. *Tetrahedron Lett.* 1995; 36:7767–7768.
15. Compound ($\pm$)-**5a** was first reacted with trichloroacetyl isocyanate followed by basic work-up to afford a homoallyl carbamate. In an early attempt to affect hydroxyamination, the carbamate was treated catalytic K_2OsO_4 in the presence of *t*-BuOCl and NaOH. The substrate was not compatible with the harsh oxidizing conditions and a complex product mixture resulted within 10 min; compound ($\pm$)-**9a** was not observed by this method.

16. For tethered aminohydroxylations with allyl carbamates as substrates, see: (a) Donohoe TJ, Johnson PD, Cowley A, Keenan M. J. Am. Chem. Soc. 2002; 124:12934–12935. [PubMed: 12405805] (b) Donohoe TJ, Johnson PD, Pye RJ. Org. Biomol. Chem. 2003; 1:2025–2028. [PubMed: 12945888] For tethered aminohydroxylations with homoallyl carbamates as substrates, see: (c) Kenworthy MN, McAllister GD, Taylor RJK. Tetrahedron Lett. 2004; 45:6661–6664. (d) Curtis KL, Fawcett J, Handa S. Tetrahedron Lett. 2005; 46:5297–5300. (e) Curtis KL, Evinson EL, Handa S, Singh K. Org. Biomol. Chem. 2007; 5:3544–3553. [PubMed: 17943217]
17. Donohoe TJ, Chughtai MJ, Klauber DJ, Griffin D, Campbell AD. J. Am. Chem. Soc. 2006; 128:2514–2515. [PubMed: 16492017]
18. In the synthesis of dysiherbaine, see: de la Pradilla RF, Lwoff N, Viso A. Tetrahedron Lett. 2007; 48:8141–8144.
19. Attempts to directly prepare *N*-hydroxycarbamate ($\pm$)-**7a** from ($\pm$)-**5a** occurred with limited success. When homoallylic alcohol ($\pm$)-**5a** was stirred with CDI for 2 h and then treated with a solution of hydroxylamine hydrochloride and triethylamine, starting material ($\pm$)-**5a** was recovered and low conversion to *N*-hydroxycarbamate ($\pm$)-**7a** was observed.
20. For preparation of *O*-*p*-methoxybenzylhydroxylamine hydrochloride, see: Zhu G, Yang F, Balachandran R, Hook P, Vallee RB, Curran DP, Day BW. J. Med. Chem. 2006; 49:2063–2076. [PubMed: 16539395]
21. The $pK_a < 7$ for the NH of *O*-acylhydroxamic acids, see: Bauer L, Exner O. Angew. Chem., Int. Ed. Engl. 1974; 13:376–385.
22. LCMS of the crude reaction mixture indicated that ($\pm$)-**8a** was the major component of the reaction mixture (74%). Unreacted ($\pm$)-**7a** (16%), *N*-mono-benzolyated product (3%) and *bis*-benzolyated product (7%) were also present.
23. Katagiri N, Ito Y, Kitano K, Toyota A, Kaneko C. Chem. Pharm. Bull. 1994; 42:2653–2655.
24. Diastereofacial addition will be directed *anti* to the electron rich bonds that stabilize the developing antibonding orbitals in the transition state, see: (a) Cieplak AS. Chem. Rev. 1999; 99:1265–1336. [PubMed: 11749447] (b) Cieplak AS. J. Am. Chem. Soc. 1981; 103:4540–4552.
25. Carbamates have been observed as by-products in tethered aminohydroxylations, see: ref. 13b, footnote, 7.
26. An imidotrioxosmium (VII) species is proposed in the tethered aminohydroxylation reaction mechanism, see: ref. 16b.
27. Li G, Chang H-T, Sharpless KB. Angew. Chem., Int. Ed. Engl. 1996; 35:451–454.
28. Refer to Supplemental Material for complete carbon and proton assignments for ($\pm$)-**9a** and (–)-**9b**.

Scheme 1.
Preparation of chloropurine (±)-5a.

Scheme 2.
Preparation of chloropurine (+)-**5b**.

Scheme 3.

Tethered aminohydroxylation to (±)-9a and elaboration to *lyxo*-carbocyclic aminonucleoside (±)-1a.

Scheme 4.
Tethered aminohydroxylation to (-)-9b.

Scheme 5.

Elaboration to carbocyclic aminonucleoside (-)-1b and *epi*-4'-carbocyclic puromycin (-)-2.

Figure 1.
Puromycin and related carbocyclic analogs.